

Nursing Health Assessment

A CRITICAL THINKING,
CASE STUDIES APPROACH

EDITION

2



Patricia M. Dillon

Nursing Health Assessment

A CRITICAL THINKING,
CASE STUDIES APPROACH

Patricia M. Dillon, DNSc, RN

Assistant Professor
Temple University
College of Allied Health Professions
Department of Nursing
Philadelphia, Pennsylvania

Illustrated by Dimitri Karenitkov

Photography by B. Proud

EDITION

2



F. A. DAVIS COMPANY • Philadelphia

F. A. Davis Company
1915 Arch Street
Philadelphia, PA 19103
www.fadavis.com

Copyright © 2007 by F. A. Davis Company

Copyright © 2007, 2003 by F. A. Davis Company. All rights reserved. This product is protected by copyright. No part of it may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without written permission from the publisher.

Printed in the United States of America

Last digit indicates print number: 10 9 8 7 6 5 4 3 2 1

Publisher: Lisa B. Deitch
Developmental Editor: William F. Welsh
Project Editor: Ilysa H. Richman
Design and Illustration Manager: Carolyn O'Brien

As new scientific information becomes available through basic and clinical research, recommended treatments and drug therapies undergo changes. The author(s) and publisher have done everything possible to make this book accurate, up to date, and in accord with accepted standards at the time of publication. The author(s), editors, and publisher are not responsible for errors or omissions or for consequences from application of the book, and make no warranty, expressed or implied, in regard to the contents of the book. Any practice described in this book should be applied by the reader in accordance with professional standards of care used in regard to the unique circumstances that may apply in each situation. The reader is advised always to check product information (package inserts) for changes and new information regarding dose and contraindications before administering any drug. Caution is especially urged when using new or infrequently ordered drugs.

Library of Congress Cataloging-in-Publication Data

Dillon, Patricia M.

Nursing health assessment : a critical thinking, case studies approach
/ Patricia M. Dillon ; illustrated by Dimitri Karenitkov, photography by
B. Proud. – 2nd ed.

p. ; cm.

Includes bibliographical references and index.

ISBN-13: 978-0-8036-1579-3

ISBN-10: 0-8036-1579-5

1. Nursing assessment. 2. Nursing assessment—Case studies.

I. Title.


[DNLM: 1. Nursing Assessment—Case Reports. 2. Medical History
Taking—methods—Case Reports. 3. Physical Examination—methods
—Case Reports. WY 100.4 D579n 2007]

RT48.D54 2007

616.07'5—dc22


2006033105

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by F. A. Davis Company for users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the fee of \$.10 per copy is paid directly to CCC, 222 Rosewood Drive, Danvers, MA 01923. For those organizations that have been granted a photocopy license by CCC, a separate system of payment has been arranged. The fee code for users of the Transactional Reporting Service is: 8036-1579/07 + \$.10.



Dedication

To my patients and students, who have taught me so much.



Preface to Students

Dear Students,

Welcome to F.A. Davis's *Nursing Health Assessment: A Critical Thinking, Case Study Approach*, second edition. Assessment is the **first step** of the nursing process and probably the **most important** because your assessment **directs the rest of the process**. It does not matter what your outcomes and goals are if you “missed it” on the assessment!

Assessment is a **thinking, doing, and feeling process**. You need to think as you act and interact with your patients. **Think critically** as you proceed through the assessment process and make clinical decisions.

Assessment is a skill, and as with any skill, the more you **practice**, the better you become. This book will walk you through the assessment process, but it is up to you to develop the skill. As a beginning practitioner, focus on learning the normal. Once you are able to identify the normal, if something abnormal occurs, you will know! Initially, you may not be able to identify the abnormal, but you will know that it does not belong. Don't worry; with practice you will be able not only to identify the normal, but also to differentiate it from the abnormal.


You will use assessment in every area of nursing. As you perfect your assessment skills, you will find that you can apply them to all levels of health care, in the community, in acute care, in the home, and in long-term care.

F.A. Davis has a variety of resources available to meet your learning needs and develop your assessment skills. Here is a brief guide on how to use them.

- **Interact** with the case study in the text. This is your patient.
- **Review** your anatomy and physiology, apply the assessment process, and identify your learning needs through the self-evaluation exercises in the applications text.
- **Listen** to the heart, breath, and bowel auscultatory sounds again and again until you can clearly hear each sound.
- **Visit** DavisPlus for additional resources.
- **Validate** your assessment findings and perfect your skills by using your pocket guide in the clinical area.

Consider this text your guide to assessment. Take ownership of it as you learn to assess. Consider the case studies as your patients. As you interact with them, you will learn assessment. Every patient encounter provides you with an opportunity to assess, so seize the opportunity. You can learn so much from your patients. Make each encounter with your patient a learning experience as you develop the skill—and art—of assessment. So practice, practice, practice! And have fun as you develop your assessment skills.

Patricia M. Dillon, DNSc, RN




Acknowledgments

I would like to begin by thanking you, my colleagues and students, for sharing your comments and suggestions with me. The changes to the second edition reflect your feedback, and I think the second edition is a better assessment text for it. I hope you like what you see in this revision.

I would like to tell my contributors and reviewers how much I appreciate their participation in this project. They helped me identify and correct the errors that inevitably creep into a new book, but perhaps more important they allowed me to see through other eyes, showing me how to make the book even more useful to a greater range of readers. I am particularly indebted to Dr. Daniel Mason who again so generously made available his unequaled collection of heart, lung, and abdominal sounds for inclusion on the student CD. Being able to repeatedly listen to these sounds will enable nursing students to develop a skill that will make a significant difference in their care of patients. And to all my colleagues who provided the photographs that will greatly enhance students' understanding of the assessment findings, thank you.

For believing in my vision and for their continued support throughout this project I would like to acknowledge the staff at F.A. Davis, especially Lisa Deitch, Nursing Publisher. I would also like to thank William Welsh, Nursing Developmental Editor; Ilysa Richman, Nursing Project Editor; and Darlene Pedersen, Nursing Content Development Manager for their editorial expertise, their logistical support, and their guidance through the revision process. Thanks, too, to everyone in the Production Department who made the second edition work well and look great: Carolyn O'Brien, Art Director; Sam Rondinelli, Production Manager; and David Orzechowski, Managing Editor. And I would especially like to acknowledge the F.A. Davis sales force for not only promoting but also believing in this educational approach to assessment.

Finally, and most importantly, a special thank you to my family—Joe, Joey, Katie, and Patty—for their understanding and unending support.



Contributors

**Noreen E. Chikotas, DEd, CRNP, FNP
(Motor-Musculoskeletal)**

Assistant Professor
Department of Nursing
Bloomsburg University
Bloomsburg, Pennsylvania

**Linda C. Curry, RN, PhD
(Wellness)**

Professor of Nursing
Harris School of Nursing
Texas Christian University
Fort Worth, Texas

**Denise Demers, RN, MS
(Abdomen)**

Assistant Professor of Nursing
St. Joseph's College
Standish, Maine

Patricia M. Dillon, DNSc, RN

Assistant Professor
Temple University
College of Health Professions
Department of Nursing
Philadelphia, Pennsylvania

**Pamela Jean Frable, RN, ND
(Teaching)**

Assistant Professor
Harris School of Nursing
Texas Christian University
Fort Worth, Texas

**Mary Jo Goolsby, MSN, EdD, ANP-C, FAANP
(Integumentary and Head, Face, and Neck)**

Director of Research and Education
American Academy of Nurse Practitioners
Austin, Texas
Patient Care Research Specialist
University Health Care System
Augusta, Georgia

**Diane Greslick, RNC, MSN
(Abdomen)**

Assistant Professor of Nursing
St. Joseph's College
Standish, Maine

**Annette Gunderman, RN, DED
(Motor-Musculoskeletal)**

Associate Professor
Department of Nursing
Bloomsburg University
Bloomsburg, Pennsylvania

**Shelton M. Hisley, RNC, PhD, WHNP, ACCE
(Mother-to-be)**

Assistant Professor
Graduate Clinical
Coordinator
School of Nursing
The University of North Carolina at Wilmington
Wilmington, North Carolina

**Mildred O. Hogstel, RNC, PhD
(Wellness)**

Professor Emeritus
Harris School of Nursing
Texas Christian University
Fort Worth, Texas

**Barbara Jones, RN, MSN, DNSc
(Elder Abuse)**

Associate Professor
Gwynedd-Mercy College
Gwynedd Valley, Pennsylvania

**Kathleen Kellinger, MSN, PhD, CRNP
(Adolescent)**

Chairperson of Department of Nursing
Slippery Rock University
Slippery Rock, Pennsylvania

**Judith Ann Kilpatrick, RN, MSN, DNSc
(Spirituality)**

Assistant Professor of Nursing
Widener University
Chester, Pennsylvania

**Maryanne Lachat, RNC, PhD
(Newborn and Infant)**

Associate Professor
Georgetown University
Washington, DC

**Daniel Mason, MD, FACC
(Cardiovascular)**

Professor of Medicine (Cardiology)
Hahnemann University School of Medicine
Philadelphia, Pennsylvania

**Carol Meadows, RNP, MNsc, APN
(Male and Female Genitourinary)**

Instructor
University of Arkansas
Eleanor Mann School of Nursing
Fayetteville, Arkansas

**Donna Molyneaux, RN, MSN, DNSc
(Breast)**

Associate Professor
Gwynedd-Mercy College
Gwynedd Valley, Pennsylvania

**Louise Niemer, RN, BSN, MSN, PhD, CPNP,
ARNP (Nutrition)**

Associate Professor, Nursing
Northern Kentucky University
Highland Heights, Kentucky

**Darlene D. Pedersen, MSN, APRN, BC
(Mental Health)**

Director and Psychotherapist
PsychOptions
Philadelphia, Pennsylvania

**Sandra G. Raymer Raff, RN, MS, FNP
(Putting It All Together)**

Perinatal Educator
Natividad Medical Center
Salinas, California

**Barbara Resnick, PhD, CRNP, FAAN, FAANP
(Older Adult)**

Associate Professor, University of Maryland
School of Nursing
Baltimore, Maryland

**Diane C. Seibert, PhD, CRNP, RN, WHNP, ANP
(Genetics)**

Assistant Professor
Uniformed Services University of the Health Sciences
Bethesda, Maryland

**S. Anne Stewart, ARNP, PhD, CS
(Homeless)**

Associate Professor
California State University
Sacramento, California


**Joanne L. Thanavaro, RN, MSN, ARNP, BC
(Peripheral-Vascular and Lymphatic)**

Assistant Professor of Nursing
Jewish Hospital College of Nursing and
Allied Health at Washington University
St. Louis, Missouri
Private Practice: Clayton Medical Consultants, Inc.
St. Louis, Missouri

**Rose Utley, RN, PhD
(Eye and Ear)**

Assistant Professor
Department of Nursing
Southwest Missouri State University
Springfield, Missouri

Case studies used in the companion *Nursing Health Assessment: Student Applications* were provided by N. Chitokas and A. Gunderman (Motor-Musculoskeletal), P. Frable (Teaching), M.J. Goolsby (Integumentary and Head, Face, and Neck), D. Greslick and D. Demers (Abdomen), J. Kilpatrick (Spirituality), C. Meadows (Male Genitourinary and Adolescent), L. Niemer (Nutrition and Toddler), S. Raff (Putting It All Together) and J. Thanavaro (Peripheral-Vascular and Lymphatic).



Reviewers

Katrina D. Allen, RN, MSN, CCRN

ADN Faculty
Faulkner State Community College
Bay Minette, Alabama

Heather Anderson, BSN

Instructor
Portage College
Lac La Biche, Alberta Canada

Ramona C. Anest, RN, MSN

Course Coordinator
Bob Jones University
Greenville, South Carolina

Rita C. Bergevin, RN, MA, BC

Clinical Assistant Professor
Binghamton University
Binghamton, New York

Judith Laver Bierschbach, MSN, CNP

Associate Professor
University of Cincinnati
Cincinnati, Ohio

Patricia Birchfield, ARNP, DSN

Professor
Eastern Kentucky University
Richmond, Kentucky

Anne L. Bishop, RN, BS, MSN

Coordinator, Health Occupations
Cayuga-Onondaga BOCES
Auburn, New York

Renee Byfield, MS, RN, FNP

Clinical Assistant Professor
Drexel University
Philadelphia, Pennsylvania

Catherine Caston, PHD, APRN, CNS

Nursing Instructor
VA Long Beach Healthcare System
Long Beach, California

Shelba Durston, RN, MSN, CCRN

Nursing Instructor
San Joaquin Delta College
Stockton, California

Brenda R. French, RN, MSN

Assistant Professor
J. Sargeant Reynolds Community College
Richmond, Virginia

Karen Hessler, RN, MSN, FNP-C- PH. D.,

Lecturer
University of Northern Colorado
Greeley, Colorado

Dottie Irvin, DNS, APRN, BC, CNE

Associate Professor
St. John's College
Springfield, Illinois

Janet L. Joost, RN, MSN

Program Director, Associate Degree Nursing
Front Range Community College, Boulder County
Campus
Longmont, Colorado

Katherine Kniest, RN, MSN

Professor, Nursing
William Rainey Harper College
Palatine, Illinois

Cathy Murray, MSN, RN, CNS, ONC

Assistant Professor
Ivy Tech Community College of Indiana
Muncie, Indiana

Darlene Pedersen, MSN, APRN, BC

Director and Psychotherapist
PsychOptions
Philadelphia, Pennsylvania

Connie Rockstad, RN, MSN

Director, Nursing
Whatcom Community college
Bellingham, Washington

**Elizabeth A. Reese, RN, MSN, MS ED,
APRN-BC**

Associate Clinical Professor
Binghamton University
Binghamton, New York

Cynthia Sanoski, PharmD

Associate Professor
University of the Sciences in Philadelphia
Philadelphia, Pennsylvania

Gale Sewell, RN, MSN

Assistant Professor
Kent State University
Ashtabula, Ohio

Kim Tucker, RN, MN, ANP

Nursing Instructor
Columbia Basin College
Pasco, Washington

Roberta Waite, EdD, RN, MSN, CS

Assistant Professor
Drexel University
Philadelphia, Pennsylvania

Jill E. Winland-Brown, EdD, MSN, APRN


Professor and Director
Florida Atlantic University,
Treasure Coast Campus
Port St. Lucie, Florida

Machelle Zink, RN, BSN

Nursing Instructor
Iowa Western Community College
Council Bluffs, Iowa

Michael Zychowicz, RN, MS, RNFA, NP-C

Assistant Professor
Mount Saint Mary College
Newburgh, New York



Contents

UNIT 1

Understanding Assessment 1

CHAPTER 1

Health Assessment and the Nurse 3

- Introduction to the Health Assessment 4
- The Nursing Process 4
- Communication 5
- The Assessment Process 11
- Levels of Preventive Healthcare 17
- Types of Assessment 17
- Collecting Data 17
- Methods of Data Collection 18
- Validating, Organizing, and Prioritizing Data 20
- Documenting Your Findings 23
- Summary 24
- Review Questions 25

CHAPTER 2

The Health History 27

- Purpose of the Health History 28
- Types of Health Histories 28
- Medical History versus Nursing History 30
- Setting the Scene 30
- Components of the Health History 30
- Documenting Your Findings 47
- Summary 51
- Review Questions 51

CHAPTER 3

Approach to the Physical Assessment 53

- Purpose of the Physical Assessment 54
- Types of Physical Assessment 54
- Tools of Physical Assessment 55
- Techniques of Physical Assessment 67
- Approach to the Physical Assessment 72
- Patients with Special Needs 73
- Positions for Physical Assessment 74
- Components of the Physical Assessment 74
- Summary 85
- Review Questions 85

CHAPTER 4

Assessing Pain 87

- Introduction to Assessing Pain 88
- Structure and Function of Pain 88
- Classification of Pain 90
- Nociceptive/Neuropathic Pain 92
- Developmental, Cultural, and Ethnic Variations 93
- Introducing the Case Study 95
- Performing the Pain Assessment 97
- Reassessment of Pain 107
- Case Study Analysis 107
- Research Tells Us 107
- Summary 108
- Review Questions 108

UNIT 2

Promoting Health 109

CHAPTER 5

Approach to the Mental Health Assessment 111

- Understanding Mental Disorders 112
- Biological Basis of Mental Illness 112
- Psychological Theories 114
- Behavioral Theories 114
- Cognitive Theories 114
- Developmental Theories 114
- Social Theories 114
- Developmental, Cultural, and Ethnic Variations 114
- People of Different Cultures or Ethnic Groups 118
- Introducing the Case Study 118
- Performing the Mental Health Assessment 120
- Case Study Evaluation 126
- Case Study Findings 137
- Case Study Analysis and Plan 137
- Research Tells Us 138
- Health Concerns 138
- Summary 142
- Review Questions 142

CHAPTER 6

- Teaching the Patient 145**
- Introducing the Case Study 146
 - Introduction to Teaching the Patient 146
 - Performing the Assessment 147
 - The Physical Assessment 150
 - Case Study Analysis and Plan 151
 - Summary 151
 - Review Questions 151

CHAPTER 7

- Assessing Wellness 153**
- Introducing the Case Study 154
 - Introduction to Assessing Wellness 154
 - Performing the Wellness Assessment 155
 - Case Study Analysis and Plan 162
 - Summary 162
 - Review Questions 162

CHAPTER 8

- Assessing Nutrition 163**
- Review of Nutrients 164
 - How a Nutritional Deficiency Develops 167
 - Introducing the Case Study 170
 - Performing the Nutritional Assessment 170
 - Case Study Evaluation 177
 - Case Study Analysis and Plan 182
 - Research Tells Us 182
 - Health Concerns 183
 - Summary 185
 - Review Questions 185

CHAPTER 9

- Spiritual Assessment 187**
- Brief Review of Spiritual Health 188
 - Developmental, Cultural, and Ethnic Variations 188
 - Introducing the Case Study 190
 - Performing the Assessment 190
 - Case Study Evaluation 194
 - Case Study Analysis and Plan 201
 - Research Tells Us 201
 - Summary 201
 - Review Questions 202

Systems Physical Assessment 203

CHAPTER 10

Assessing the Integumentary System 205

- Anatomy and Physiology Review 206
- Developmental, Cultural, and Ethnic Variations 209
- Introducing the Case Study 211
- Performing the Integumentary Assessment 212
- Case Study Evaluation 222
- Case Study Analysis and Plan 256
- Research Tells Us 256
- Health Concerns 257
- Summary 264
- Review Questions 264

CHAPTER 11

Assessing the Head, Face, and Neck 267

- Anatomy and Physiology Review 268
- Introducing the Case Study 275
- Performing the Head, Face, and Neck Assessment 275
- Case Study Evaluation 285
- Case Study Analysis and Plan 314
- Research Tells Us 314
- Health Concerns 315
- Summary 319
- Review Questions 319

CHAPTER 12

Assessing the Eye and Ear 321

- Anatomy and Physiology Review: The Eye 322
- Introducing the Case Study 330
- Performing the Eye Assessment 330
- Case Study Evaluation 337
- Case Study Analysis and Plan 360
- Research Tells Us 360
- Health Concerns 360
- Summary 364
- Review Questions 364
- Anatomy and Physiology Review: The Ear 365
- Introducing the Case Study 370

- Performing the Ear Assessment 370
- Case Study Evaluation 375
- Case Study Analysis and Plan 385
- Research Tells Us 387
- Health Concerns 388
- Summary 390
- Review Questions 390

CHAPTER 13

Assessing the Respiratory System 393

- Anatomy and Physiology Review 394
- Introducing the Case Study 399
- Performing the Respiratory Assessment 399
- Case Study Evaluation 408
- Case Study Analysis and Plan 430
- Research Tells Us 431
- Health Concerns 431
- Summary 435
- Review Questions 435

CHAPTER 14

Assessing the Cardiovascular System 437

- Anatomy and Physiology Review 438
- Introducing the Case Study 454
- Performing the Cardiovascular Assessment 454
- Case Study Evaluation 463
- Case Study Analysis and Plan 486
- Research Tells Us 487
- Health Concerns 487
- Summary 489
- Review Questions 489

CHAPTER 15

Assessing the Peripheral-Vascular and Lymphatic Systems 491

- Anatomy and Physiology Review 492
- Introducing the Case Study 497
- Performing the Peripheral-Vascular and Lymphatic Systems Assessment 497
- Case Study Evaluation 504
- Case Study Analysis and Plan 519
- Research Tells Us 519
- Health Concerns 520
- Summary 525
- Review Questions 525

CHAPTER 16

Assessing the Breasts 527

- Anatomy and Physiology Review 528
- Introducing the Case Study 533
- Performing the Breast Assessment 533
- Case Study Evaluation 539
- Case Study Analysis and Plan 548
- Research Tells Us 549
- Health Concerns 549
- Summary 553
- Review Questions 553

CHAPTER 17

Assessing the Abdomen 555

- Anatomy and Physiology Review 556
- Introducing the Case Study 562
- Performing the Abdominal Assessment 562
- Case Study Evaluation 576
- Case Study Analysis and Plan 603
- Research Tells Us 603
- Health Concerns 604
- Summary 610
- Review Questions 610

CHAPTER 18

Assessing the Female Genitourinary System 611

- Anatomy and Physiology Review 612
- Introducing the Case Study 617
- Performing the Female Genitourinary Assessment 617
- Case Study Evaluation 628
- Case Study Analysis and Plan 643
- Research Tells Us 643
- Health Concerns 643
- Summary 646
- Review Questions 647

CHAPTER 19

Assessing the Male Genitourinary System 649

- Anatomy and Physiology Review 650
- Introducing the Case Study 657
- Performing the Male Genitourinary Assessment 657
- Case Study Evaluation 665
- Case Study Analysis and Plan 678

- Research Tells Us 679
- Health Concerns 679
- Summary 684
- Review Questions 684

CHAPTER 20

Assessing the Motor-Musculoskeletal System 685

- Anatomy and Physiology Review 686
- Introducing the Case Study 693
- Performing the Musculoskeletal Assessment 693
- Case Study Evaluation 701
- Case Study Analysis and Plan 730
- Research Tells Us 731
- Health Concerns 732
- Summary 735
- Review Questions 735

CHAPTER 21

Assessing the Sensory-Neurologic System 737

- Anatomy and Physiology Review 738
- Introducing the Case Study 751
- Performing the Sensory-Neurologic and Cranial Nerve Assessment 751
- Case Study Evaluation 761
- Case Study Analysis and Plan 787
- Research Tells Us 787
- Health Concerns 789
- Summary 793
- Review Questions 794

CHAPTER 22

Putting It All Together 795

- Introducing the Case Study 796
- Developmental Considerations 796
- Performing the Assessment 797
- Case Study Evaluation 800
- Case Study Analysis and Plan 808
- Research Tells Us 808
- Summary 808
- Review Questions 809

UNIT **4****Adapting Assessment to Special Populations 811**

CHAPTER 23

Assessing the Mother-to-Be 813

- Anatomy and Physiology Review 814
- Performing the Assessment 823
- Introducing the Case Study 823
- Case Study Evaluation 832
- Case Study Analysis and Plan 842
- Case Study Follow-Up 843
- Research Tells Us 845
- Health Concerns 846
- Summary 846
- Review Questions 847

CHAPTER 24

Assessing the Newborn and Infant 849

- Anatomy and Physiology Review 850
- Introducing the Case Study 853
- Performing the Initial Newborn Assessment 853
- Case Study Follow-Up 875
- Performing the Infant Assessment 876
- Case Study Evaluation 878
- Case Study Analysis and Plan 883
- Research Tells Us 884
- Health Concerns 884
- Summary 885
- Review Questions 885

CHAPTER 25

Assessing the Toddler and Preschooler 887

- Developmental Changes 888
- Health Issues and Risk Factors 888
- Introducing the Case Study 889
- Performing the Toddler and Preschooler Assessment 889

- Back to the Case Study 897
- Research Tells Us 897
- Health Concerns 898
- Common Problems 898
- Summary 898
- Review Questions 898

CHAPTER 26

Assessing the School-Age Child and Adolescent 901

- Assessing the School-Age Child 902
- Introducing the Case Study 904
- Performing the School-Age Child Assessment 904
- Back to the Case Study 911
- Research Tells Us 912
- Health Concerns 912
- Common Problems 912
- Summary 913
- Assessing the Adolescent 913
- Introducing the Case Study 915
- Performing the Adolescent Assessment 915
- Back to the Case Study 924
- Research Tells Us 924
- Health Concerns 924
- Common Problems 925
- Summary 925
- Review Questions 925

CHAPTER 27

Assessing the Older Adult 927

- Anatomy and Physiology Review 928
- Introducing the Case Study 933
- Performing the Assessment 933

- Case Study Analysis and Plan 958
- Research Tells Us 958
- Health Concerns 958
- Summary 960
- Review Questions 960

APPENDICES

APPENDIX A

Chapter Feedback 961

APPENDIX B

Growth Charts 1007

APPENDIX C

Guidelines 1018

- Guidelines for Cultural Assessment 1019
- Guidelines for Environmental Assessment 1022
- Guidelines for Assessing the Homeless Person 1030
- Guidelines for Assessing Abuse 1033

APPENDIX D


Genetics 1037

- Family History, the First Genetic Test 1038
- Genetic Counseling: Who, When, How, and Why 1040
- Genetic Resources 1041

APPENDIX E

Bibliography 1042


INDEX 1057



UNIT

1

Understanding Assessment



Health Assessment and the Nurse

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Define assessment
- Discuss the role of assessment as part of the nursing process
- Identify skills needed for assessment
- Differentiate nursing assessment from medical assessment
- Identify the role of assessment in all levels of preventive healthcare
- Differentiate subjective and objective data
- Differentiate primary and secondary data sources
- Describe factors that affect communication
- Discuss cultural influences on communication
- Demonstrate communication techniques
- Identify phases of the interview
- Describe data collection methods
- Define the four techniques of physical assessment
- Demonstrate the four techniques of physical assessment
- Discuss methods used to validate assessment data
- Describe various documentation methods for charting assessment data

Introduction to the Health Assessment

You are doing blood pressure screening at a health fair. You take the blood pressure of a middle-aged man. Your reading is 170/100.

You are working in the emergency department (ED) when a father comes in with his 9-year-old daughter. He states that she fell off her bike and hit her head but did not lose consciousness. But she has a terrible headache and feels sick.

You are making a postpartum follow-up visit to the home of a young mother who had her first baby 2 days ago.

You are making an initial hospice visit to a 74-year-old woman with pancreatic cancer.

What do you do? Where do you begin? You begin with assessment. How well you perform your assessment will affect everything else that follows. You will ask questions, and you will use four of your senses to collect data. This chapter introduces you to the assessment process.

The Nursing Process

Nursing is the diagnosis and treatment of human responses to actual or potential health problems. Diagnosis and treatment are achieved through a process, called the *nursing process*, that guides nursing practice. The **nursing process** is a systematic problem-solving method that has five steps:

- Assessment
- Nursing diagnoses
- Planning
- Implementation
- Evaluation

The nursing process (Fig. 1.1) is used to identify, prevent, and treat actual or potential health problems and promote wellness. It provides a framework in which to practice nursing. Think of it as a continuous, circular process that revolves around your patient. You begin with assessment, collect data, cluster the data, and then formulate nursing diagnoses. Once you have identified the nursing diagnosis, you will develop a plan of care, determine the goals and expected outcomes, implement your plan, and then evaluate it. Then you will begin the nursing process again.

Characteristics of the Nursing Process

<ul style="list-style-type: none"> ■ Dynamic and cyclic ■ Patient centered ■ Goal directed ■ Flexible ■ Problem oriented 	<ul style="list-style-type: none"> ■ Cognitive ■ Action oriented ■ Interpersonal ■ Holistic ■ Systematic
---	---

Step 1: Assessment

Assessment is the process of collecting, validating, and clustering data. It is the first and most important step in the nursing process. The assessment phase sets the tone for the rest of the process, and the rest of the process flows from it. If your assessment is off the mark, then the rest of the process will be too. Assessment identifies your




FIGURE 1.1. The nursing process.

patient's strengths and limitations and is performed not just once, but continuously throughout the nursing process. After performing your initial assessment, you establish your baseline, identify nursing diagnoses, and develop a plan. Then, as you implement your plan, you also assess your patient's response. Finally, you assess the effectiveness of your plan of care for your patient.

Step 2: Nursing Diagnoses

Formulating a **nursing diagnosis** involves identifying and prioritizing actual or potential health problems or responses. Once you collect your data, you need to analyze them and then identify actual and potential health problems or responses to life processes and state them as nursing diagnoses. Nursing diagnoses can be actual, potential, possible, or collaborative problems as well as wellness issues. An **actual nursing diagnosis** identifies an occurring health problem for your patient. A **potential nursing diagnosis** identifies a high-risk health problem that most likely will occur unless preventive measures are taken. A **possible nursing diagnosis** is one that needs further data to support it. A **collaborative problem** is a potential medical complication that warrants both medical and nursing interventions. Because nursing's focus is not limited to illness, wellness diagnoses focus on promoting or enhancing a patient's level of wellness. Once you have identified the diagnoses, you need to prioritize them in order to develop a plan of care. (See *NANDA Nursing Diagnosis*.)

Step 3: Planning

The planning phase involves setting goals and outcomes. Once you have prioritized your diagnoses, you are ready to develop an individualized plan of care for your patient. First, you establish goals and determine measurable outcomes. Next, you identify nursing interventions needed to meet the goals and outcomes. Then, you communicate your plan to both the patient and all members involved in the plan of care to maintain continuity of care and ensure success.

Step 4: Implementation

Implementation (also called *intervention*) involves carrying out your plan to achieve goals and outcomes. This is the “doing” phase of the nursing process, in which you actually implement the nursing interventions in the plan. As you implement your plan, you continue to assess your patient's responses and modify the plan as needed. Be sure to document your care.

Step 5: Evaluation

Evaluation involves determining the effectiveness of your plan. Did you meet your goals and outcomes? Once again, assess your patient's response based on the criteria you set for the outcome. If the goals and outcomes have not been met, you'll need to rethink the plan and work through the process again to develop a more effective plan of care for your patient.

Communication

To assess, you must be able to communicate and communicate well. The relationship you establish with your patient directly affects your ability to collect data. **Communication** is a process of sharing information and meaning, of sending and receiving messages. The messages we communicate are both verbal and nonverbal. The sections below describe the many factors that influence communication. You need to consider all of them when communicating with your patient to be sure that the message you want to send is the one your patient actually receives.

Nonverbal Messages

Nonverbal behavior is an important source of data. Often, the nonverbal message being sent is more accurate than the verbal one. So examine both your nonverbal behavior and your patient's. Be conscious of your

Sample Nursing Diagnoses

A postoperative cholecystectomy patient might have the following nursing diagnoses:

- *Actual nursing diagnosis*: Ineffective breathing pattern r/t pain as evidenced by shallow respirations, tachypnea, and reluctance to cough.
- *Risk nursing diagnosis*: Risk for ineffective breathing pattern r/t pain.
- *Collaborative problems*: Potential for postoperative atelectasis: Ineffective breathing pattern.

A person seeking healthy nutritional instructions for weight control might have the wellness nursing diagnosis of Health-Seeking Behavior.

NANDA Nursing Diagnoses

Activity Intolerance [specify level]	Family Processes, readiness for enhanced
Activity Intolerance, risk for	Fatigue
Adjustment, impaired	Fear [specify focus]
Airway Clearance, ineffective	Fluid Balance, readiness for enhanced
Allergy Response, latex	Fluid Volume, deficient [hyper/hypotonic]
Allergy Response, risk for latex	Fluid Volume, deficient [isotonic]
Anxiety [specify level]	Fluid Volume, excess
Anxiety, death	Fluid Volume, risk for deficient
Aspiration, risk for	Fluid Volume, risk for imbalanced
Attachment, risk for impaired parent/infant/child	Gas Exchange, impaired
Autonomic Dysreflexia	Grieving, anticipatory
Autonomic Dysreflexia, risk for	+Grieving, dysfunctional
Body Image, disturbed	*Grieving, risk for dysfunctional
Body Temperature, risk for imbalanced	Growth, risk for disproportionate
Bowel Incontinence	Growth and Development, delayed
Breastfeeding, effective	Health Maintenance, ineffective
Breastfeeding, ineffective	Health-Seeking Behaviors (specify)
Breastfeeding, interrupted	Home Maintenance, impaired
Breathing Pattern, ineffective	Hopelessness
Cardiac Output, decreased	Hyperthermia
Caregiver Role Strain	Hypothermia
Caregiver Role Strain, risk for	Identity, disturbed personal
Communication, impaired verbal	Infant Behavior, disorganized
Communication, readiness for enhanced	Infant Behavior, readiness for enhanced organized
Conflict, decisional (specify)	Infant Behavior, risk for disorganized
Conflict, parental role	Infant Feeding Pattern, ineffective
Confusion, acute	Infection, risk for
Confusion, chronic	Injury, risk for
Constipation	Injury, risk for perioperative positioning
Constipation, perceived	Intracranial Adaptive Capacity, decreased
Constipation, risk for	Knowledge, deficient [Learning Need] (specify)
Coping, compromised family	Knowledge (specify), readiness for enhanced
Coping, defensive	*Lifestyle, sedentary
Coping, disabled family	Loneliness, risk for
Coping, ineffective	Memory, impaired
Coping, ineffective community	Mobility, impaired bed
Coping, readiness for enhanced	Mobility, impaired physical
Coping, readiness for enhanced community	Mobility, impaired wheelchair
Coping, readiness for enhanced family	Nausea
Death Syndrome, risk for sudden infant	Neglect, unilateral
Denial, ineffective	Noncompliance [ineffective adherence] [specify]
Dentition, impaired	Nutrition: less than body requirements, imbalanced
Development, risk for delayed	Nutrition: more than body requirements, imbalanced
Diarrhea	Nutrition: more than body requirements, risk for imbalanced
Disuse Syndrome, risk for	Nutrition, readiness for enhanced
Diversional Activity, deficient	Oral Mucous Membrane, impaired
+Energy Field, disturbed	Pain, acute
Environmental Interpretation Syndrome, impaired	Pain, chronic
Failure to Thrive, adult	Parenting, impaired
Falls, risk for	Parenting, readiness for enhanced
Family Processes: alcoholism, dysfunctional	Parenting, risk for impaired
Family Processes, interrupted	

Peripheral Neurovascular Dysfunction, risk for	Sorrow, chronic
Poisoning, risk for	Spiritual Distress
Post-Trauma Syndrome [specify stage]	+ Spiritual Distress, risk for
Post-Trauma Syndrome, risk for	Spiritual Well-Being, readiness for enhanced
Powerlessness [specify level]	Suffocation, risk for
Powerlessness, risk for	Suicide, risk for
Protection, ineffective	Surgical Recovery, delayed
Rape-Trauma Syndrome	Swallowing, impaired
Rape-Trauma Syndrome: compound reaction	Therapeutic Regimen Management, effective
Rape-Trauma Syndrome: silent reaction	Therapeutic Regimen Management, ineffective
*Religiosity, impaired	Therapeutic Regimen Management, ineffective community
*Religiosity, readiness for enhanced	Therapeutic Regimen Management, ineffective family
*Religiosity, risk for impaired	Therapeutic Regimen Management, readiness for enhanced
Relocation Stress Syndrome	Thermoregulation, ineffective
Relocation Stress Syndrome, risk for	Thought Processes, disturbed
Role Performance, ineffective	Tissue Integrity, impaired
Self-Care Deficit: bathing/hygiene	Tissue Perfusion, ineffective (specify type: renal, cerebral, cardiopulmonary, gastrointestinal, peripheral)
Self-Care Deficit: dressing/grooming	Transfer Ability, impaired
Self-Care Deficit: feeding	Trauma, risk for
Self-Care Deficit: toileting	Urinary Elimination, impaired
Self-Concept, readiness for enhanced	Urinary Elimination, readiness for enhanced
Self-Esteem, chronic low	Urinary Incontinence, functional
Self-Esteem, situational low	Urinary Incontinence, reflex
Self-Esteem, risk for situational low	Urinary Incontinence, risk for urge
Self-Mutilation	Urinary Incontinence, stress
Self-Mutilation, risk for	Urinary Incontinence, total
Sensory Perception, disturbed (specify: visual, auditory, kinesthetic, gustatory, tactile, olfactory)	Urinary Incontinence, urge
Sexual Dysfunction	Urinary Retention [acute/chronic]
Sexuality Pattern, ineffective	Ventilation, impaired spontaneous
Skin Integrity, impaired	Ventilatory Weaning Response, dysfunctional
Skin Integrity, risk for impaired	Violence, [actual/] risk for other-directed
Sleep, readiness for enhanced	Violence, [actual/] risk for self-directed
Sleep Deprivation	Walking, impaired
Sleep Pattern, disturbed	Wandering [specify sporadic or continuous]
Social Interaction, impaired	
Social Isolation	

Used with permission from NANDA International: Definitions and Classification, 2005–2006. NANDA, Philadelphia, 2005.

+ Revised NDs.

*New to the 3rd NANDA/NIC/NOC (NNN) Conference.

Information that appears in brackets has been added to clarify and enhance the use of NDs.

beliefs and values and do not let them influence your verbal or nonverbal communication. **Nonverbal behavior** includes vocal cues or paralinguistics, action cues or kinetics, object cues, personal space, and touch. For example, a patient who tells you he is having “crushing chest pain” should look like he is having “crushing chest pain.” His nonverbal behavior should be consistent with what he is telling you. So you would expect the tone in

his voice to convey “crushing pain” as he is clutching his chest.

Vocal Cues or Paralinguistics

Vocal cues describe the quality of your voice and its inflections, tone, intensity, and speed when speaking. These voice characteristics usually reflect underlying

feelings. If your patient is in pain, moaning or the inflections in her or his speech may alert you to the pain's severity. Someone who is afraid may speak softly, with trepidation. Anger, doubt, disbelief, or disapproval can all be conveyed by the tone of your voice.

Vocal characteristics may also reflect underlying physiological or psychological problems. For example, older patients may speak loudly because of hearing loss. Patients with hyperthyroidism may have rapid speech. Rapid speech is also associated with bipolar disorder, mania type.

Be aware of the characteristics of your own speech. What message are you sending to your patient? Your speech should be calm, reassuring, and without inflections that convey judgment or disapproval. Speak slowly enough and loudly enough so that your patient can hear you and respond appropriately.

Action Cues and Kinetics

Action cues or **kinetics** are body movements that convey messages. Posture, arm position, hand gestures, body movements, facial expressions, and eye contact all convey a message. The message may reflect a feeling, a mood, or an underlying physiological or psychological problem. A relaxed posture with arms at the sides conveys openness, whereas a tense posture with arms crossed may reflect anger, discomfort, or mistrust. A tense, guarded posture may also indicate pain. A tripod position (sitting position, leaning forward with elbows on table and hands supporting head) is assumed to ease breathing.

Hand gestures can convey approval, but they may signal anxiety when overused. Abnormal movements may indicate an underlying neurological problem. Facial expressions reflect a variety of emotions, including happiness, sadness, fear, anger, approval, and disapproval. Facial expressions may also signal an underlying problem, such as anguish from excruciating pain or fear and anxiety from shortness of breath. Eye contact usually conveys interest, caring, and trust. Lack of eye contact may reflect shyness, depression, disinterest, or mistrust.

Your body language is as important as your patient's. Remember, once a person assumes a "patient" role, she or he relinquishes a certain amount of role responsibilities and moves from an independent to a dependent role. During history taking, keep your body posture relaxed to convey a calm, reassuring attitude. Work at the same level as your patient. Standing over her or him conveys a sense of dominance and authority, enhancing her or his sense of dependence. Give reaffirming gestures such as nodding. Your facial expression should convey care and concern. Maintain eye contact to show the patient you are interested in what she or he has to say.

Object Cues

Your patient's dress and grooming reflect his or her identity and how he or she feels about himself or herself. Poor grooming or disheveled clothing may indicate a psychological problem such as depression. A depressed patient may not take much interest in how he or she looks. Poor grooming may also signal an underlying physical problem that has affected the patient's ability to care for himself or herself.

Your appearance is equally important. Your grooming and dress should be neat and appropriate for the situation and reflect your professionalism. Be sure to wear a tag with your name and title. Object cues can also be furnishings or possessions—for example, your jewelry or the seating arrangement for a meeting. These cues are usually selected on a conscious level to convey a specific message, so they are not as revealing as nonverbal communications.

Personal Space

Personal space is the territory surrounding a person that she or he perceives as private or the physical distance that needs to be maintained for the person to feel comfortable. When our personal space is invaded, we feel uncomfortable and anxious, often backing away to regain control of our territory. Personal space is often defined by culture and is also influenced by the situation. Public space, such as in a classroom or when giving a speech, is about 12 feet or more. Social-consultative space, such as in impersonal conversations or interviews, is about 4 to 12 feet. Personal distance, such as in personal conversation, is about 18 inches to 4 feet. Intimate distance, such as in intimate conversations and maximum sensory stimulation, is 0 to 18 inches. A person who is withdrawn or mistrusting may widen her or his personal space. An aggressive, angry person may invade your personal space.

Be aware of your position in relation to your patient. Assessment entails working in close proximity to your patient—about 3 to 4 feet when taking the health history, and much closer, within your patient's personal or intimate space, when performing the physical assessment. Be sure to explain what you are doing, and look for clues that your patient is becoming uncomfortable. If so, you may need to back off and try another approach.

Touch

Touch is a means of communication. An array of feelings, including anger, caring, and protectiveness, can be conveyed through touch. The interpretation of touch is often culturally prescribed. Touch can be easily misconstrued. It may be seen as an invasion of one's personal space or a threat. How a person responds to touch often depends on the trust established within a relationship.

Nursing involves touch. We often use touch to convey a sense of caring to our patients. Assessment involves touch. We must touch patients in order to collect data. Gaining your patient's trust can help you obtain an accurate, comprehensive health history. It can also make the physical assessment quicker and easier for both of you.

Cultural Considerations

Cultural communication patterns need to be considered when obtaining a health history. Culture can influence every aspect of communication, so consider not only the language but also the vocal, action, and object cues, personal space, and touch.

Obviously, if you and your patient do not speak the same language, you will need an interpreter to communicate. Even if your patient speaks English, he or she still might misinterpret some words or expressions. They may convey a totally different meaning to him or her.

Also consider the context in which the language is spoken. For example, German, English, and French are not very contextual because so many words are needed to express a thought. However, Chinese and Native American languages are very contextual because they use fewer words with more emphasis on what is unspoken. For example, Navajo Indians may say very little during a visit to a hospitalized family member. This may be interpreted as showing little support, but on the contrary, their very presence is the support.

Vocal characteristics also vary with different cultures. For example, European Americans and African Americans talk loudly compared with people of Chinese and Hindu backgrounds, who might misinterpret relative loudness as anger. Arabs often speak excitedly or passionately, another vocal characteristic that can easily be misinterpreted as anger.

Different cultures differ in the extent to which they feel comfortable expressing thoughts and feelings. European Americans tend to be open and willing to communicate their feelings and thoughts. Japanese often appear shy and withdrawn to strangers, whereas Jews and Italians are often seen as assertive. Appalachians and Mexican Americans may not be as willing to share information with a healthcare provider until a relationship is established.

Use of touch as a means of communication also varies with culture. For example, Egyptian Americans do not usually accept touch except between spouses and family. Even though Mexican Americans often use touch in communicating, they may be modest during a physical examination performed by someone of the opposite sex.

Personal space and eye contact also vary with culture. European Americans keep a much greater distance between one another (about 18 inches) than do people of Arabian or Turkish descent, who are comfortable with much shorter personal space. Direct eye contact is usually acceptable for European Americans, but not for Asian Americans when speaking with someone they consider a superior. Sustained eye contact is considered rude in Bolivia. Sustained eye contact between a child and an adult is considered an "evil eye" or a "bad eye" in many cultures.

Facial expression is also influenced by culture. Smiling conveys a warm, friendly message for most cultures, but no rule applies across the board. For instance, the Cofer Indians of Ecuador view smiling as a sign of aggression.

Acceptable greetings likewise vary from culture to culture. The handshake is accepted in most cultures as a friendly sign. However, since it is unacceptable for devout Muslim men and women to touch, the handshake is not permitted. Vietnamese Americans view the handshake as an acceptable practice between men, but women do not usually shake hands. And, the Navajo Indians consider the handshake as the traditional greeting, but the strength of the handshake is light rather than firm.

Tips for Using Interpreters

- Use an interpreter rather than a translator. A translator only restates the words; an interpreter translates the words with meaning.
- Use dialect-specific interpreters when possible.
- Allow time for the interpreter and the patient to become acquainted.
- Allow extra time for the health history.
- Be aware that some of the data may be lost through interpretation.
- Avoid using relatives—they may distort the interpretation.
- Avoid using children, especially for sensitive topics.
- Match interpreter with patient by age and gender when possible.
- Maintain eye contact with both the patient and the interpreter and read nonverbal cues.
- Be aware of your body language. The patient may understand more than she or he can express verbally.
- Speak clearly and slowly, but do not exaggerate or speak loudly.
- Use active rather than passive tense and wait for feedback.
- Use a reference book with common phrases such as *Taber's Cyclopedic Medical Dictionary*.
- Use as many words as possible in the patient's language to convey your message.
- Use nonverbal communication when you are unable to understand the language.

Source: Data from Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.

Body language also conveys different messages in different cultures. Finns consider it rude to speak with your hands in your pockets. A relaxed posture usually conveys confidence and competence in America, but a tense posture conveys the same meaning in Japan and Korea.

Patient Interview Communication Techniques

Because there is a purpose to your interaction with your patient, your interview will be focused. You will direct the conversation in order to obtain the data needed to complete the assessment. Even though you want your patient to be as open as possible, you may still need to redirect her or his conversation so that you can complete the task at hand. At the same time, follow cues that she or he gives you that may lead you to a better understanding of her or his health status.

The following communication techniques will help you direct the interview:

- **Affirmation/Facilitation.** Acknowledge your patient's responses through both verbal and nonverbal communication to reassure him that you are paying attention to what he is saying. Nonverbal gestures, such as nodding or sitting up and leaning forward, encourage your patient to continue. Verbal cues, such as saying "ah ha," "go on," or "tell me more," send the same message.
- **Silence.** Although silence is difficult to maintain at times, it can be very effective at facilitating communication. Periods of silence allow your patient to collect her or his thoughts before responding and help prevent hasty responses that may be inaccurate. Silence is usually more uncomfortable for you than for your patient. You may feel compelled to fill the silence by asking more questions or even by trying to answer the question for your patient. Resist the urge. You may end up losing valuable data and sending the wrong message to your patient. Silence also gives *you* more time to think and plan your responses. Use this "quiet time" to focus on your patient's nonverbal behavior and what she or he is feeling.
- **Clarifying.** If you are unsure or confused about what your patient is saying, rephrase what she said and then ask the patient to clarify. Use phrases like "Let me see if I have this right," or "I want to make sure I'm clear on this," or "I'm not sure what you mean."
- **Restating.** Restating the patient's main idea shows him that you are listening, allows you to acknowledge your patient's feelings, and encourages further discussion. It also helps clarify and validate what your patient has said and may help identify teaching needs. For example, if the patient states, "I take a water pill every day for my blood pressure," your response might be, "I see—you take Lasix every morning for your blood pressure." If the patient replies, "No, I take a water pill every morning," this identifies a teaching need.
- **Active Listening.** Pay attention, maintain eye contact, and really listen to what your patient tells you both verbally and nonverbally. As you listen, keep in mind what you are telling your patient nonverbally. Active listening conveys interest and acceptance.
- **Broad or General Openings.** This technique is effective when you want to hear what is important to your patient. Use open-ended questions such as, "What would you like to talk about?"
- **Reflection.** Reflection allows you to acknowledge your patient's feelings, encouraging further discussion. When your patient expresses a thought or feeling, you echo it back, usually in the form of a question. For example, if the patient states, "I am so afraid of having surgery," your response would be, "You're afraid of having surgery?"
- **Humor.** Humor can be very therapeutic when used in the right context. It can reduce anxiety, help patients cope more effectively, put things into perspective, and decrease social distance.
- **Informing.** Giving information allows your patient to be involved in his or her healthcare decisions. An

How You Communicate

Always be aware of the messages you are sending your patient, both verbally and nonverbally. How you respond is critical in establishing the nurse-patient relationship. Qualities that help establish and maintain this relationship include genuineness, respect, and empathy.

- **Genuineness:** Be open, honest, and sincere with your patient. Your patient can detect a less-than-honest response or inconsistencies between your verbal and your nonverbal behavior.
- **Respect:** Everyone should be respected as a person of worth and value. You need to be nonjudgmental in your approach. You may not always agree with your patient's decisions or like or approve of his or her behavior, but everyone needs to feel accepted as a unique individual.
- **Empathy:** Empathy is knowing what your patient means and understanding how she or he feels. Showing empathy acknowledges your patient's feelings; shows acceptance, care, and concern; and fosters open communication. Phrases that recognize your patient's feelings help build a trusting relationship—for example, "That must have been very difficult for you."

example would be explaining the postoperative course and the importance of coughing and deep breathing to your patient preoperatively.

- **Redirecting.** Redirecting your patient helps keep the communication goal-directed. It is especially useful if your patient goes off on a tangent. To get your patient on track again, you might say, “Getting back to what brought you to the hospital . . .”
- **Focusing.** Focusing allows you to hone in on a specific area, encouraging further discussion. Examples include: “You said your mother and sister had breast cancer?” or “Do you do BSE, and have you ever had a mammogram?” In this case, you have identified a risk factor and a potential area for health education.
- **Sharing Perceptions.** With this technique, you give your interpretation of what has been said in order to clarify things and prevent misunderstandings. You may need to question your patient if there is a discrepancy in the message sent. For example, you might say, “You said you weren’t upset, but you’re crying.”
- **Identifying Themes.** Identifying recurrent themes may help your patient make a connection and focus on the major theme. For example, you might say, “From what you’ve told me, it sounds like every time you were discharged from the hospital to home you had a problem.”
- **Sequencing Events.** If your patient is having trouble sequencing events, you may need to help her or him place the events in proper order. Start at the beginning and work through the event until the conclusion. You might say, “What happened before the problem started?” “Then what happened?” “How did it end?”
- **Suggesting.** Presenting alternative ideas gives your patient options. This is particularly helpful if the patient is having difficulty verbalizing his or her feelings. Suggesting is also a good teaching tool. For example, if the patient says, “I’ve tried so hard to lose weight, but I can’t,” you might say, “Have you tried combining diet and exercise?”
- **Presenting Reality.** If your patient seems to be exaggerating or contradicting the facts, help her or him reexamine what has already been said and be more realistic. For example, if she or he says, “I waited all day for someone to answer my call light,” you might respond, “All day?”
- **Summarizing.** Summarizing is useful at the conclusion of a major section of the interview. It allows the patient to clarify any misconceptions you may have. For example, you might say, “Let me see if I have this correct: You came to the hospital with chest pain, which started an hour ago, after eating lunch.”

The Assessment Process

The American Nurses’ Association (ANA) has identified assessment as the first Standard of Nursing Practice (ANA, 1998). The Standard describes assessment as the

systematic, continuous collection of data about the health status of patients. Nurses are responsible not only for data collection but also for making sure that the data are accessible, communicated, and recorded. Assessment is an ongoing process. Every patient encounter provides you with an opportunity for assessment.

Purpose of Assessment

The purpose of assessment is to collect data pertinent to the patient’s health status, to identify deviations from normal, to discover the patient’s strengths and coping resources, to pinpoint actual problems, and to spot factors that place the patient at risk for health problems.

Skills of Assessment

Assessment requires cognitive, problem-solving, psychomotor, affective/interpersonal, and ethical skills.

Cognitive Skills

Assessment is a “thinking” process. **Cognitive skills** are needed for critical thinking, creative thinking, and clinical decision making. Your theoretical knowledge base enables you to assess your patient holistically. The knowledge base includes not only biophysical knowledge but also developmental, cultural, psychosocial, and spiritual knowledge. This knowledge base enables you to differentiate normal from abnormal findings and to identify and prioritize actual and potential problems.

Critical Thinking

Critical thinking is a complex thinking process that has been defined in many ways. Critical thinking is reflective, reasonable thinking (Ennis, 1985). It is not just doing, it is asking “why?” Critical thinking involves inquiry, interpretation, analysis, and synthesis. It is the art of thinking about thinking that enables you to think better (Paul, 1990). Paul (1990) has identified attitudes or “traits of the mind” that are needed for critical thinking. Think of them as a mindset that enables you to use your cognitive skills to critically think. (See *Critical Thinking Skills*.)

Essentials of Critical Thinking

Intellectual Humility

Know what you don’t know.
Know your limits.
Be aware of your biases and prejudices.

Intellectual Courage

Be open and listen to other points of view.

(continued)

Essentials of Critical Thinking (continued)	
Intellectual Empathy	
Put yourself in another's shoes, so that you can understand his or her perspective.	
Intellectual Integrity	
Maintain high standards. Hold yourself to the same standards as you do others.	
Intellectual Perseverance	
Persist to find the answer or reach understanding.	
Faith in Reason	
Look out for the best interest for all involved. Consider the consequences.	
Intellectual Sense of Justice	
Be open to all viewpoints, with a sense of impartiality and fairness.	

Source: Green, C. (2000). *Critical Thinking in Nursing*. Upper Saddle River, NJ: Prentice Hall.

Clinical Decision Making

Assessment also requires clinical decision making. As you collect data, you will make clinical decisions as to its relevance. You will look for cues and make inferences. With experience, you will be able to identify patterns and recognize what differs from the norm and then use the data to make decisions that will best meet your patient's need. Use your knowledge, experience, and what the patient says to validate the data.

Problem-Solving Skills

Various problem-solving methods can be used as you assess your patient and work through the nursing process. With experience, you will develop your problem-solving skills. Do not limit yourself to one method; instead, select the method that best suits your patient's needs.

Reflexive thinking is automatic, without conscious deliberation, and comes with experience. For example, if you are working in the ED and a 45-year-old man is admitted with "crushing chest pain," you will automatically take his vital signs and do an electrocardiogram to assess his cardiac status as you ask him questions.

The **trial-and-error approach** is hit-or-miss thinking—random, not systematic and inefficient. This would never be your primary problem-solving approach, but it may prove helpful at times because it allows you to "think outside of the box." Trial-and-error thinking fosters creativity and allows you to formulate new ideas. Use it for situations that don't quite fit the picture, look beyond the obvious, and keep looking until you find an answer.

The **scientific method** is a systematic, critical-thinking approach to problem solving. It involves identifying a problem and collecting supporting data, formulating a hypothesis, planning a solution, implementing the plan, and then evaluating its effectiveness. Assessment corresponds with the "collection of supporting data phase" of the scientific method. The nursing process is similar to the scientific method, but the nursing process offers nurses the flexibility to practice both the art and the science of nursing.

Intuition is a problem-solving method that develops through experience, in which theory and practice are intertwined. This is how the "expert" nurse solves problems. Through experience, she has a well-established experiential base of pattern recognition, an intuitive grasp of the situation, and the ability to zero in on problem areas. For example, suppose a patient says, "I just don't feel right," even though her or his vital signs are stable and everything seems fine on the chart. The objective data may not support it, but the expert nurse senses that something is not quite right and returns to assess her patient further, only to find her or him in distress. She intervenes swiftly and saves the patient's life.

Psychomotor Skills

Assessment is "doing." Psychomotor skills are needed to perform the four techniques of physical assessment: inspection, palpation, percussion, and auscultation. As a beginning practitioner, you may feel unsure of your technique and your findings, but practice will hone your skills. Input from your colleagues will help you perfect your skills and interpret your findings. Through experience, you will become competent at performing the physical assessment and confident in interpreting your findings.

Affective/Interpersonal Skills

Assessment is also a "feeling" process. Affective skills are needed to practice the "art" of nursing. Affective skills are essential in developing caring, therapeutic nurse-patient relationships. **Interpersonal skills** include both verbal and nonverbal communication skills. The quality of your assessment depends on your communication skills and the relationship that you develop with your patient. Establishing trust and mutual respect is essential before you begin the assessment. Seeing your patient as an individual and being sensitive to his or her feelings conveys a message of caring and promotes human dignity. Illness often makes a patient very vulnerable, but the power of a caring relationship can have a major impact on the patient's sense of worth and well-being. Such a relationship can be mutually rewarding, affecting both you and your patient personally and professionally.

Your interpersonal skills are also needed to communicate with family members and members of the healthcare team. Communicating effectively with the healthcare team is essential in order to successfully meet your patient's goals.

Critical Thinking Skills

Divergent Thinking

Divergent thinking involves analyzing various viewpoints, differentiating relevant/irrelevant data, and making inferences (conclusions).

Reasoning

Reasoning is the ability to differentiate fact (truth) from assumptions (beliefs). It can be deductive or inductive. You will use both deductive and inductive reasoning to assess your patient and interpret the data. For example, suppose your patient is admitted to the hospital with the diagnosis of congestive heart failure (CHF). Knowing this enables you to look for specific signs and symptoms as you perform the assessment and determine your patient's response to this illness. In this case, you will use deductive reasoning, looking for specific clues that support the diagnosis of decreased cardiac output.

Inductive reasoning would be appropriate when assessing a postoperative patient who states, "It hurts too much to take deep breaths," and who has decreased breath sounds and crackles at the bases, pulse oximetry of 90 percent on room air, temperature of 101°F, pulse of 106 BPM, and respirations 26 and shallow. When you piece together all of the pertinent assessment data, you formulate diagnoses of Impaired Gas Exchange and Ineffective Airway Clearance.

Reflection

Reflection allows you to step back and think, considering "if . . . then" possibilities. Experience and new evidence may result in changing opinions. Integrating what you know from past experience with new evidence allows for change.

Creativity

Creativity allows you to "think outside of the box." Creativity involves identifying multiple solutions/approaches, realizing there is more than one way to approach a problem. Creativity is also listening to your "gut"—intuitive thinking. Intuitive thinking develops over time with experience.

Clarification

Clarification involves addressing assumptions, comparing and contrasting. Ask yourself, "What is similar?" "What is different?"

Basic Support

Basic support is based on sound, credible, known facts. Evidence-based practice that is research based leads to best practice.

Source: Green, C. (2000). *Critical Thinking in Nursing*. Upper Saddle River, NJ: Prentice Hall.

Ethical Skills

Part of assessment is being responsible and accountable. You are responsible and accountable for your practice. You are also an advocate for your patient. You must respect your patient's rights and ensure patient confidentiality. The ANA describes the ethical standards that guide nursing

practice in its Code for Nurses. The data collected through assessment are used to plan the patient's care, but it is important to remember that the data are the patient's information. The Health Insurance Portability and Accountability Act (HIPAA) has established rules to protect patient privacy. (See *Understanding HIPAA*.)

The Code for Nurses

1. The nurse, in all professional relationships, practices with compassion and respect for the inherent dignity, worth, and uniqueness of every individual, unrestricted by considerations of social or economic status, personal attributes, or the nature of health problems.
2. The nurse's primary commitment is to the patient, whether an individual, family, group, or community.
3. The nurse promotes, advocates for, and strives to protect the health, safety, and rights of the patient.
4. The nurse is responsible and accountable for individual nursing practice and determines the appropriate delegation of tasks consistent with the nurse's obligation to provide optimum patient care.
5. The nurse owes the same duties to self as to others, including the responsibility to preserve integrity and safety, to maintain competence, and to continue personal and professional growth.

(continued)

The Code for Nurses (continued)

6. The nurse participates in establishing, maintaining, and improving healthcare environments and conditions of employment conducive to the provision of quality healthcare and consistent with the values of the profession through individual and collective action.
7. The nurse participates in the advancement of the profession through contributions to practice, education, administration, and knowledge development.
8. The nurse collaborates with other health professionals and the public in promoting community, national, and international efforts to meet health needs.
9. The profession of nursing, as represented by associations and their members, is responsible for articulating nursing values, for maintaining the integrity of the profession and its practice, and for shaping social policy.

Source: American Nurses Association, Code for nurses with interpretive statements (Kansas City, MO: American Nurses Association, 2004), Reprinted with permission.

Understanding HIPAA

The American Health Insurance Portability and Accountability Act (HIPAA) has established rules that protect a patient's privacy of health information. This federal law gives patients rights over their health information and sets limits on who can review and receive patient information. The HIPAA rules apply to all health plans and healthcare providers.

A patient's rights with respect to his health information include a right to:

- Ask to see and get a copy of health records.
- Have corrections added to health information.
- Receive a notice informing him how health information is used and shared.
- Decide whether to give permission before information can be used or shared for certain purposes.
- Get a report on when and why health information is shared.
- Ask to be contacted somewhere other than home.
- Ask that information not be shared.
- File complaints.

Role of the Nurse and Assessment

The role of the nurse has changed drastically over the years. So have the nurse's responsibilities. The importance of assessment can be traced to the beginning of modern nursing. Florence Nightingale (1859) stressed the importance of observation and experience as essential in maintaining or restoring one's state of health. The scope of assessment has also expanded from simple observation to a holistic view of the patient that includes biophysical, psychosocial, developmental, and cultural assessments. The skills of assessments have also expanded—from simple observations to detailed use of physical assessment skills—as the scope of practice has expanded.

Nursing Assessment versus Medical Assessment

Assessment is not unique to nursing. It is also an integral part of medical practice. Although the assessment process may be the same for nursing and medical practice, the outcomes differ. The goal of medical practice is to diagnose and treat disease. The goal of nursing practice is to diagnose and treat human responses to actual or potential

health problems. Nursing assessment focuses not only on physiological and psychological responses but also on the psychosocial, cultural, developmental, and spiritual dimensions. It identifies patients' responses to health problems as well as their strengths. Nursing's aim is to help the patient reach her or his optimal level of wellness.

Medical and nursing assessments should complement, not contradict, each other in promoting the patient's health and wellness. Often, data obtained through the nursing assessment contribute to the identification of medical problems. By working together in a collaborative relationship, nursing and medicine ensure the best possible care for patients.

The nurse-physician relationship has changed through the years. In the late 1800s, nurses were subservient to physicians; their primary duty was to serve physicians. By 1936, the nurse-physician relationship was that of coworkers with the primary goal of restoring the patient to health. This relationship was further refined with the formation of the National Joint Practice Commission (NJPC) in 1972. Established by the American Medical Association (AMA) and the ANA, this

The Nurse's Changing Role

In the past 70 plus years, the nurse's role has changed dramatically, and so has the scope of nursing practice. A mirror of the times, the nurse's role has grown from that of a skilled observer in the 1930s to that of a healthcare professional who performs holistic assessment. The following excerpts from actual nursing notes reflect this change in the nurse's role.

1930s

"Patient admitted to ward in a wheelchair. Stated that he is unable to walk due to pinched nerves of right foot. Patient is crying, says he is homesick. Condition of skin good. Made as comfortable as possible."

This note includes observations only. The nurse's role is limited to that of a skilled observer.

1950s

"25-year-old female admitted ambulatory. Past history of ulcerative colitis. Now in because of abdominal cramps and vomiting \times 4 days. Is 8 months pregnant. TPR 99.4, 80, 20. Ht 5'4". Wt 116 $\frac{1}{4}$. Urine to lab."

This note traces the past and present history of illness as well as observations. The nurse's role has expanded to include interviewing skills that assess the patient's past and current health status.

1970s

"12-year-old white female admitted to Room 203 via stretcher from E.R. with leukemia. Parents don't seem to know of diagnosis. TPR 102.8, 120, 24. Ht 62 $\frac{1}{4}$. Wt 100 lb. No known allergies. Has not been eating much for last few days. Appears extremely pale. BP 150/70. No urine obtained. I.V. started. Blood started. Vital signs relatively stable. T 103° when blood started."

This note records intravenous and blood therapy and includes observations as well as information about past illnesses and diet. The nurse's role now includes observation, interviewing, performing procedures (venipuncture), and monitoring.

1990s to Present

"Young, obese Caucasian female states she came here to 'get the sugar out of her blood.' States she found out about her sugar 3 months ago by glucose tolerance test results (in Jan. 1990 miscarried 2-month pregnancy and GTT was part of workup); states she has seen her husband test his blood (fingerstick method) and give himself insulin but has done neither herself; has tried to prepare both 1,800 and 2,200 calorie American Diabetic Association diets as ordered for husband but 'he doesn't stick to it'; has noted increased hunger, increased thirst, increased urination for several months and occasional blurred vision.

Lives with husband and daughter next door to sister-in-law; husband makes playground equipment; states they are able to 'get by' on his salary, sometimes borrow money and have difficulty paying it back; have Blue Cross/Blue Shield which should pay for this hospitalization; husband has diabetes mellitus (takes insulin), is losing weight but refuses to see doctor—she is concerned about him; states husband will not be able to visit her because of work."

This note records observation and assessment of the patient as well as the biophysical, psychosocial, and cultural factors that influence the patient's health problem. The nurse's role has grown to include holistic health assessment.

interprofessional organization recognized the importance of a collaborative nurse-physician relationship in improving patient care. The NJPC identified five components that promote collaboration: primary nursing,

integrated patient records, individual decision making by nurses, a joint practice committee, and joint care reviews. Today, nurses practice both independently and collaboratively.

Steps in Decision Making

Identifying the need to make a decision (the why and what)

- Setting goals or outcomes, considering what needs to be done and what should be avoided
- Identifying alternatives, considering benefits and consequences
- Selecting and implementing alternatives
- Evaluating the outcome

Source: Kelly-Heidenthal, P., and Marthaler, M. (2005). *Delegation of Nursing Care*. United States: Clifton Park, N.Y., Thomson Delmar Learning.

The NJPC's Elements of Collaboration

According to the National Joint Practice Commission (NJPC) study, five elements foster nurse-physician collaboration and provide the following benefits:

Element and Description	Benefits
<p>Primary Nursing Nursing with little or no delegation of nursing tasks to others. One nurse is responsible for a patient's comprehensive care.</p>	<ul style="list-style-type: none"> ■ Direct communication between the primary nurse and the physician ■ Patient care coordinated by one person ■ Nursing autonomy ■ Shared accountability with the physician ■ Fewer errors in patient care ■ Increased patient satisfaction
<p>Integrated Patient Records Interdisciplinary patient progress notes that combine nurse and physician observations, judgments, and actions and provide a formal means of nurse-physician communication about patient care</p>	<ul style="list-style-type: none"> ■ Substantive and concise nursing notes ■ Opportunity for healthcare professionals to read each others' notes ■ Less duplication of charted information ■ Validation of observations and judgments by cosigning of nurse's and physician's notes
<p>Individual Decision Making by Nurses Ability to exercise independent judgment and initiate care based on judgment. Decisions must lie within the scope of nursing practice.</p>	<ul style="list-style-type: none"> ■ Greater nursing control of the environment ■ Greater job satisfaction for nurses
<p>Joint Practice Committee A committee of nurses and physicians that monitors nurse-physician relationships and recommends actions that support collaboration</p>	<ul style="list-style-type: none"> ■ Opportunity for nurses and physicians to discuss collaborative practice and work out problems ■ Improved understanding between nurses and physicians
<p>Joint Care Reviews Monthly review of patient charts by a review committee of nurses and physicians to evaluate collaborative care</p>	<ul style="list-style-type: none"> ■ Identification of ways to improve patient care and collaborative practice

Source: Morton, P (1993). *Health Assessment in Nursing*, ed. 2 Philadelphia: F.A. Davis.

Levels of Preventive Healthcare

Preventive healthcare can be classified as primary, secondary, and tertiary. Nurses have a vital role in all three levels of health care, and assessment skills are needed at all levels.

Primary preventive care focuses on health promotion and guards against health problems. The focus is not only wellness but also to minimize the risk of health problems. Examples include health fairs, immunizations, and nutritional instructions. Nurses use assessment skills to screen and identify patient education needs about wellness and the risk factors for health problems.

Secondary preventive care focuses on early detection, prompt intervention, and health maintenance for patients with health problems. It addresses acute health problems seen with inpatient hospitalization. Accurate assessment is crucial to establish the patient's baseline and to continually monitor his or her condition and response to treatment.

Tertiary preventive care deals with rehabilitative or extended care. Although the patient's health status is generally more stable than in secondary care, patients usually have a chronic illness. Skilled nursing care facilities, rehabilitative hospitals, long-term care facilities, home care, and hospice are examples of tertiary-level care. Assessment skills are needed to continually monitor health status.



HELPFUL HINT

Remember: At every level of preventive care, you have opportunities to practice assessment. You are often the one who decides when the patient's condition warrants medical intervention.

Types of Assessment

Assessments can be comprehensive or focused. A comprehensive assessment is usually the initial assessment. It is very thorough and includes a detailed health history and physical examination. A **comprehensive assessment** examines the patient's overall health status. A **focused assessment** is problem oriented and may be the initial assessment or an ongoing assessment. If a patient's condition does not warrant a comprehensive assessment, a focused assessment of the patient's present health problem is done. Once the patient's condition improves and stabilizes, the comprehensive assessment can be completed. A focused assessment is frequently performed on an ongoing basis to monitor and evaluate the patient's progress, interventions, and response to treatments.

Even when a focused assessment is performed, it is important to look at the entire picture. A problem in one system will affect or be affected by every other system.

So scan your patient from head to toe and note any changes in other systems. Look for clues or pertinent data that will help you formulate your diagnosis.

Collecting Data

Data can be classified as subjective and objective. Subjective data are covert and not measurable. They reflect what the patient is experiencing and include thoughts, beliefs, feelings, sensations, and perceptions. **Subjective findings** are referred to as *symptoms*. The health history is an example of subjective data. Objective data are overt and measurable. **Objective data** are referred to as *signs*. The physical examination and diagnostic studies are examples of objective data. Data sources are either primary or secondary. The patient is a primary data source. Secondary data sources are anyone or anything aside from the patient, including family members, friends, other health-care providers, and old medical records. Both primary and secondary data can also be subjective or objective.



CRITICAL THINKING ACTIVITY # 1

Identify the following data as subjective or objective:

- Headache
- Blood pressure (BP) 170/110
- Nausea
- Diaphoresis
- Equal pupillary reaction
- Tingling sensation
- Dizziness
- Decreased muscle strength
- Slurred speech
- Numbness, left arm



CRITICAL THINKING ACTIVITY # 2

Elaine Ploransky is a pregnant 29-year-old married woman, gravida 1, para 0. She is having contractions 5 minutes apart, which she describes as "severe cramps." Her husband states, "I think her water broke on the way to the hospital."

Physical examination reveals that Mrs. Ploransky is 6 cm dilated. Fetal monitor reveals a fetal heart rate of 120 beats per minute (BPM). The patient's vital signs include BP 140/80, pulse 90 BPM, respirations 22/min. Prenatal records reveal hemoglobin 12.0, hematocrit 45, and blood type AB+.

Identify:

- Primary data source
- Secondary data source
- Subjective data
- Objective data

Methods of Data Collection

Assessment requires sharp observation skills and the ability to distinguish relevant from irrelevant and important from unimportant data. Data collection can occur through interviews, observation, and physical assessment. Once the data are collected, they need to be organized, categorized, and validated.

Interviews

The interview is usually structured communication intended to obtain subjective data. An interview is most useful when taking the health history. For an interview to be successful, you need good interpersonal communication skills. Many factors, including cultural or developmental differences or life experiences, can impair effective communication with your patient. You need to examine your values and beliefs to minimize any bias that may interfere with the interview.

When you use your interpersonal skills in a healing way to help your patient, this is known as *therapeutic use of self*. Showing empathy, demonstrating acceptance, and giving recognition are three techniques that enhance therapeutic use of self. Being empathetic is the ability to understand another's feelings. Maintaining a neutral, nonjudgmental position and demonstrating acceptance of your patient's verbal and nonverbal communication are essential to developing a trusting relationship. Acknowledging the patient's verbal and nonverbal communication conveys true interest and encourages further communication.

Types of Interviews

Interviews can be either directive or nondirective. **Directive interviews** are structured with specific questions and are controlled by the nurse. These interviews require less time and are very effective for obtaining factual data. **Nondirective interviews** are controlled by the patient, although the nurse often needs to summarize and clarify the data. These interviews require more time than directive interviews but are very effective at eliciting the patient's perceptions and feelings. Nondirective interviews help you to identify what is important to the patient.

Types of Questions

Questions are classified as closed or open. **Closed questions** are often those that elicit a "yes" or "no" response, for example, "Do you have pain?" This type of question takes little time and is very effective for factual data. Directive interviews frequently use many closed questions. **Open questions** elicit the patient's perceptions, for example, "What brought you to the hospital?" More

time is needed for this type of question. They are frequently used in nondirective interviews.

Interviewing Techniques

Begin the interview by establishing trust and conveying a caring attitude. Make sure that the environment is comfortable and that privacy is ensured with minimal distractions. The following tips will help you get the most from your interviews:

- Introduce yourself.
- Don't rush. Allow enough time for the interview.
- Avoid interruptions.
- Explain that information from the interview is confidential.
- Actively listen to what your patient is saying.
- Maintain eye contact.
- Work at the same level as your patient. Pull up a chair and sit next to her or him.
- Don't invade your patient's personal space. Two to 4 feet away is a comfortable distance for most patients.
- Explain what you are doing and why.
- If the patient presents with a problem, begin by asking questions about that.
- Begin with nonsensitive issues. Leave more sensitive topics until the end.
- Consider your patient's cultural background. How does it affect the interview and your interpretation of the data?
- Consider your patient's developmental level. How does it affect the interview and your interpretation of the data?
- Don't become preoccupied with writing. You may convey to the patient that the forms you are completing are more important than he or she is.
- Be nonjudgmental.
- Avoid "why?" questions; they tend to put patients on the defensive.
- Nonverbal behavior is more accurate than verbal. Take a look at yours—What is it telling your patient?
- Take a good look at your patient's nonverbal behavior. Is it consistent with what she or he is telling you?
- Now look at your patient's nonverbal behavior another way. Does it indicate health problems?
- Never pass up an opportunity to teach.
- Present reality.
- Be honest.
- Provide reassurance and encouragement.
- Be respectful.

Interviewing Pitfalls

Avoid the following traps when interviewing a patient:

- *Leading the patient.* People will tell you what you want to hear. So don't lead the patient. Having him or her describe what is happening in his or her own words is much more helpful.

- **Biasing yourself.** You can bias yourself because of the patient, a particular disease, or a particular physician. But if you fail to check your biases, you limit your objectivity during assessment.
- **Letting family members answer for patient.** You will learn a lot more by having the patient describe things in her or his own words.
- **Asking more than one question at a time.** If you do, the patient may not know which one to answer. Or, you may not be sure which question is being answered.
- **Not allowing enough response time.** Give the patient time to think through his or her answer. This is especially important with older patients.
- **Using medical jargon.** Express your questions in lay terms to make sure your patient understands you.
- **Assuming rather than clarifying and validating.** Assuming can lead to inaccurate interpretations and incorrect conclusions.
- **Taking the patient's responses personally.** An angry or frustrated patient may verbally attack you or the healthcare facility. Realize that the patient is displacing her or his feelings on you and using you as a sounding board.
- **Feeling personally uncomfortable.** This often happens with beginning students. Stay focused. It will get easier with experience.
- **Using clichés.** Interviewers often resort to clichés when they feel uncomfortable or unsure of an answer. No one expects you to have all the answers. If you do not have an answer, tell the patient that you will try to find one and get back to him or her. Clichés like, "You'll feel better in the morning," show a disregard for your patient's feelings.
- **Offering false reassurance.** Telling the patient that everything will be fine is condescending. It may not be.
- **Asking persistent or probing questions.** Persistent or probing questions make your patient uncomfortable. Remember, the patient has a right to not answer a question.
- **Changing the subject.** Some nurses change the subject when the interview is making them uncomfortable. This is not very helpful for the patient. Remember: You are attending to the patient's needs, not your own.
- **Taking things literally.** Literal responses may lead to misunderstanding. Patients who have difficulty expressing feelings directly may use figurative language. So always consider the context in which the statement is made, and clarify if you are still unsure. For example, an extremely frustrated patient might say, "If I don't get out of here soon, I'll go crazy!" Don't rush to have a psychiatric consultation; recognize the patient's feelings of frustration and focus on them.
- **Giving advice.** The patient needs to be informed, but can make her or his own decisions.
- **Jumping to conclusions.** Make sure you have all the facts before drawing conclusions.

Phases of the Interview

The interview is divided into three phases: the introductory phase, the working phase, and the termination phase. Each phase has a specific purpose and different communication patterns.

Introductory Phase

The **introductory phase** is the time to introduce yourself to your patient, put him or her at ease, and explain the purpose of the interview and the time frame needed to complete it. Questions should be nonprobing and patient centered. Explain that you will be taking notes, but keep your writing to a minimum. Reassure your patient that the information collected is confidential.

The Working Phase

The **working phase** is often where data collection occurs. It is usually very structured; it is also the longest phase. Make sure you allot enough time for the working phase. Although you will need to take notes, stay focused on your patient. Listen to what she or he is saying both verbally and nonverbally. With experience, you will become skilled at taking minimal notes and then documenting your data *after* the interview rather than during it.

The Termination Phase

The end of the interview is the **termination phase**. During this phase, you need to summarize and restate your findings. This provides an opportunity to clarify the data and share your findings with the patient. Based on this information, both you and the patient can discuss follow-up plans.

Observation

The second method of data collection is observation. **Observation** entails deliberate use of your senses of sight, smell, and hearing to collect data. Look at both your patient and his or her environment to detect anything out of the ordinary. Your initial observations may provide clues to underlying problems. These clues can be further investigated during the physical examination.

Ask yourself:

- Does the patient show signs of physical or psychological stress?
- Does the patient seem comfortable?
- What is the patient doing?
- What position is she or he assuming?
- Are there any abnormal movements?
- What is the patient's body language telling you?
- Is the patient's verbal language consistent with his or her nonverbal language?
- Do you notice any unusual odors?
- Do you hear any unusual sounds?

- Is there anything unusual, unsafe, or risky in the patient's environment?

Look for:

- Facial expression, color changes, breathing problems
- Grimacing, guarding, diaphoresis
- Eye contact
- Tone of voice and flow of speech
- Position, orthopnea
- Grooming and dress
- Nervousness, restlessness, voluntary/involuntary movements
- Unusual odors, such as fruity smell associated with diabetic ketoacidosis, or foul odors
- Drainage associated with infection
- Unusual sounds, such as grunting, wheezing, rhonchi, or stridor associated with respiratory problems, or swishing sounds associated with murmurs

Physical Assessment

Physical assessment provides the objective database. It helps you assess your patient's health status and identify actual or potential problems. During physical assessment, you use your senses to collect data through the techniques of inspection, palpation, percussion, and auscultation.

During **inspection**, look at your patient and compare her or his appearance with what you know as nor-

mal. Use your sense of smell to detect any unusual odor that may warrant further investigation into a possible health problem. During **palpation**, use light touch to assess surface characteristics, to put your patient at ease, and to convey concern and caring. Use deep palpation to assess organs and masses. During **percussion**, use direct, indirect, and fist percussion to assess organ size and areas of tenderness. During **auscultation**, listen to your patient directly and indirectly to hear sounds produced by the body.

The extent of physical assessment depends on the patient's condition and your expertise. Physical assessment skills can be used with patients at all levels of healthcare. Your skills will develop over time with practice and provide invaluable, objective data essential in planning your patient's care. Chapter 3, Approach to the Physical Assessment, details the physical assessment.

Validating, Organizing, and Prioritizing Data

Once you have collected the data, you need to validate them to ensure their accuracy, organize them, and prioritize them so that they are useful.

Validation

Validation can occur simultaneously with the assessment process. Validating every piece of data is unrealistic, but you do need to validate any time you notice an inconsistency or are unsure of your findings. There are numerous ways to validate data. You can use the same sources that you used to collect data to validate them. For example, while you perform the health history, validate the subjective findings as you perform the physical examination by comparing them with the objective findings. The subjective and objective findings should support each other.

You can also call on your patient to validate the assessment data. Often, the patient can be very helpful. Other sources for validation include family members, other healthcare providers, past health records, and diagnostic tests.

As a beginning practitioner, you may initially feel unsure of your findings. Take advantage of the expertise of your colleagues to validate them. For example, maybe you think you hear an abnormal heart sound, but you are not sure. Ask a colleague to double-check. With experience, you will develop confidence in the accuracy of your assessment. When assessment findings are way out of line from the normal, even experienced nurses will rely on colleagues to validate findings.

Assessment Tips

- Involve your patient in planning care.
- Common problems occur commonly, rare problems, rarely. So concentrate on common problems.
- On the other hand, be open and expect the unexpected.
- Determine if the onset is acute or chronic.
- Determine if the problems are life threatening and need immediate attention.
- Know what is normal before you try to identify abnormal. Experience will help you refine your ability to differentiate the two.
- Look at your patient holistically and identify relevant health risk factors. For example, a middle-aged, African American male is at risk for hypertension because of his biographical data.
- Validate any inconsistencies.
- Realize that multiple problems will lead to varied human responses.
- Identify problems, but also look for strengths.
- Remember that every system is related. So if you identify a problem in one area, realize that this problem can affect every other system.




FIGURE 1.2. Maslow's Hierarchy of Human Needs.

Organizing Data

After you have validated the assessment data, you need to organize them. Begin by identifying pertinent data—any findings that are out of the norm and any findings that identify patient strengths. The pertinent data provide cues that will help you identify any actual or potential problems. Once you have identified these cues, “cluster” the data, grouping all cues that support a problem. Clustering relevant data gives meaning to the data. These clustered data will be the basis for formulating nursing diagnoses for your patient.

Several frameworks can be used for organizing data. A **framework** provides a way of looking at your patient and the data. All frameworks have a common thread—they address the physical, psychosocial, and spiritual needs of a patient in a holistic approach.

Common frameworks include the following:

- **Maslow's Hierarchy of Human Needs:** Organizes data according to the patient's basic human needs: physiological, safety and security, love and belonging, self-esteem, and self-actualization (Fig. 1.2).
- **Roy's Adaptation Theory:** Organizes data according to the patient's adaptation to physiological, self-concept, social role, and interdependence demands.
- **Gordon's Functional Health Patterns:** Organizes data into 11 functional groups that contribute to a person's overall health and well-being, quality of life, and attainment of human potential.

Correlating Gordon's Functional Patterns to Dillon's Assessment

GORDON'S FUNCTIONAL PATTERNS

DILLON'S ASSESSMENT

Health perception/health management	Health history Wellness Developmental
Nutritional-metabolic	Nutrition Integumentary Neck (thyroid) Mouth Abdominal Lymphatic
Elimination	Abdominal Genitourinary Rectal
Activity-exercise	Cardiovascular Peripheral vascular Respiratory Musculoskeletal
Sleep-rest	Sinuses, nose, and throat Health history Wellness
Cognitive-perceptual	Sensory neurological Eyes/ears
Role relationship	Health history
Value belief	Wellness
Self-perception/self-concept	Mental health
Coping/stress	
Sexuality/reproductive	Male/female genitourinary Breast Mother-to-be

- **NANDA–Unitary Person Framework:** Organizes data based on the concept that a person is an open system in constant interaction with the environment. Nine human responses result from this interaction: exchanging, communicating, relating, valuing, choosing, moving, perceiving, knowing, and feeling.

No matter what framework you choose to organize your data, be consistent. The framework allows you to view your patient in a particular way and makes your data meaningful.

Prioritizing Data

Organizing data will assist you in prioritizing the patient's problems. When prioritizing, consider the acuity of the problem, the patient's perception of the problem, and the

Common Data Collection Pitfalls

Because assessment affects every step of the nursing process, the accuracy of your information is critical. To help ensure accurate data collection, avoid the following common pitfalls:

- *Omitting data.* You are not thorough enough, are in a hurry, fail to follow clues that your patient gives you, or simply lack experience. If you do not know what to look for, you will never see it.
- *Misinterpreting data.* You fail to validate the data and then misinterpret them, or you lose your objectivity and allow your biases to enter into your interpretation of the data.
- *Hasty interpretation.* You make hasty interpretations before you have completed a thorough assessment.
- *Irrelevant data.* You have difficulty deciding which data are important. Experience and knowledge will help you distinguish what is relevant and what is not.
- *Failure to follow up.* You fail to assess continually for changes in your patient's responses. Assessment is not a one-shot deal; it is an ongoing process.
- *Poor communication.* You fail to communicate effectively with your patient verbally, nonverbally, or both. Your communication skills directly affect the quality of your assessment.

Avoid these problems by listening carefully to what your patient is telling you, both verbally and nonverbally. Be thorough and systematic in your approach, be objective, use your knowledge, and develop a strong experiential base for assessment.

situation at hand. Put life-threatening problems that require immediate attention at the top of the list, then problems affecting basic needs that require prompt attention, and finally, problems affecting psychosocial needs.

Top-priority or primary problems—such as airway problems—are life-threatening. Secondary problems—such as pain—require prompt attention to prevent further progression or deterioration in your patient's condition. Although important, third-level problems—such as teaching needs—do not require immediate attention and can be addressed once your patient's condition has stabilized.

Although actual problems usually take precedence over potential ones, potentially serious problems may also have high priority. Also consider the nature and context of the problem. For example, if a new mother is being discharged with her baby, providing instruction regarding care for her and the newborn would be a priority.



HELPFUL HINT

A person's basic needs must be met before addressing higher-level needs that involve self-esteem or self-actualization.

Identifying what is important to your patient must be considered when prioritizing problems. If your patient does not believe that he or she has a problem that you have identified, it will be difficult to develop and implement an effective plan to resolve the problem. Working together with your patient to identify

actual or potential problems is the best way to develop an effective plan, ensure compliance, and resolve the problem.



CRITICAL THINKING ACTIVITY #3

Give an example of a nursing diagnosis that would correspond to Maslow's Hierarchy of Human Needs.

- Physiological needs
- Safety needs
- Love and belonging
- Self-esteem needs
- Self-actualization



CRITICAL THINKING ACTIVITY #4

Prioritize the problems below, assigning primary, secondary, and tertiary levels of priority.

- Bleeding
- Fatigue, pallor, hemoglobin 7.0
- Nausea and vomiting
- Grieving
- Sleep disturbance, fever, night sweats
- Knowledge deficit on self-administration of insulin
- Impaired mobility, fractured hip
- Irregular heart rate >140 BPM
- BP 70/40
- New parent role

Documenting Your Findings

Once you have completed your assessment, document your findings. Your patient's initial assessment may require documenting a detailed health admission record that contains both the history and the physical assessment data. Every assessment that follows also needs to be documented. To ensure continuity of care, your patient's assessment needs to be communicated to all members of the healthcare team involved in her or his care.

Documentation Methods

The approach to documentation is usually source-oriented or problem-oriented. **Source-oriented documentation** is done by department, so each healthcare group has a section to document findings. This method easily identifies each discipline, but it tends to fragment the data, making it difficult to follow the sequencing of events. With **problem-oriented medical records (POMR)**, everyone involved in the care of the patient charts on the same form. This allows for better communication of data to resolve the patient's problems collaboratively.

No matter which approach you use, a variety of methods and forms are available for documentation. Many healthcare facilities use standardized nursing assessment forms in a checklist format, which is efficient and time-saving. Computerized documentation is also available in a standardized checklist format. The narrative format may also be used, but this is more time-consuming.

Two popular methods of problem-oriented documentation are SOAPIE and PIE. A third popular method, DAR, is used with FOCUS® charting, which focuses more holistically on the patient, identifying both strengths and problems that can be incorporated into his or her care. These methods are explained below. Another method—charting by exception (CBE)—is a shorthand documentation method frequently used to save time. It includes only significant data, that is, findings that deviate from well-defined standards. A CBE system usually includes flow sheets and standard of care checklists that are kept at the bedside. Any exception is then documented on the chart on either the nurse's note or the progress note.

Documentation Methods

Consider the following assessment of Mary Rutherford:

Mary Rutherford, age 43, is 1 day postoperative after a cholecystectomy. Her assessment data include the following:

- “It hurts to take a deep breath.”
- Pain rated 8/10
- Guarding abdomen

- Vital signs: BP 144/90; pulse 108; respirations 24 and shallow; temperature 100.8°F
- Pulse oximeter 92 percent on room air
- Decreased breath sounds at bases owing to poor ventilatory effort, also few crackles noted at bases
- Receiving patient-controlled analgesia (PCA) morphine
- Dressings dry and intact

Here is how you would document Mrs. Rutherford's findings using each method:

SOAPIE Method

■ Subjective data

“It hurts to take a deep breath”: Pain increases with activity and breathing, PCA helps; pain is sharp; pain is located in right upper quadrant (RUQ) and epigastric region; pain is rated 8/10; pain only when moving.

■ Objective data

First-day postop cholecystectomy; vital signs: BP, 144/90; pulse, 108; respirations, 24 and shallow; temperature, 100.8°F; pulse oximeter, 92 percent on room air; patient guarding abdomen; decreased breath sounds at bases because of poor ventilatory effort, also few crackles noted at bases; receiving PCA morphine.

■ Assessment/clinical judgment

Ineffective breathing pattern related to incisional pain.

■ Plan

Patient will establish effective breathing pattern; patient will experience no signs of respiratory complications.

■ Interventions

Encourage coughing and deep breathing; teach patient to splint incision; control pain with PCA; encourage ambulation; provide instruction on use of incentive spirometer; maintain adequate hydration.

■ Evaluation

Patient coughing and deep breathing, splinting incision; using incentive spirometer; ambulating; pain 5/10, using PCA morphine as needed; lungs clear; vital signs: BP 130/86, temperature 99°F, pulse 80, respirations 20, pulse oximeter 96% on room air.

DAR Method

■ Data

“It hurts to take a deep breath”; pain increases with activity and breathing, PCA helps; pain is sharp; pain is located in RUQ and epigastric region; pain is rated 8/10; pain only when moving; first-day postop cholecystectomy;

vital signs: BP 144/90; pulse 108; respirations 24 and shallow; temperature 100.8°F; pulse oximeter 92 percent on room air; patient guarding abdomen; decreased breath sounds at bases because of poor ventilatory effort, also few crackles noted at bases; receiving PCA morphine.

■ Action

Encourage coughing and deep breathing; teach patient to splint incision; control pain with PCA; encourage ambulation; provide instruction on use of incentive spirometer; maintain adequate hydration.

■ Response

Patient coughing and deep breathing, splinting incision; using incentive spirometer; ambulating; pain 5/10, using PCA morphine as needed; lungs clear; vital signs: BP 130/86, temperature 99°F, pulse 80, respirations 20, pulse oximeter 96 percent on room air.

PIE Method

■ Problem

Ineffective breathing pattern related to incisional pain.

Documentation Tips

- Be brief and to the point.
- Use acceptable abbreviations.
- If documentation is handwritten, make sure writing is legible.
- No need to write in complete sentences.
- State the facts. Avoid interpretations.
- Avoid terms such as “normal,” “good,” “usual,” and “average.”
- Avoid generalizations.
- Document sequentially, in chronological order.
- Do not leave blanks or skip lines.
- Use correct spelling and grammar.
- No erasures or whiting out.
- Record date and time and sign your full signature.

■ Interventions

Encourage coughing and deep breathing; teach patient to splint incision; control pain with PCA; encourage ambulation; provide instruction on use of incentive spirometer; maintain adequate hydration.

■ Evaluation

Patient coughing and deep breathing, splinting incision; using incentive spirometer; ambulating; pain 5/10, using PCA morphine as needed; lungs clear; vital signs: BP 130/86, temperature 99°F, pulse 80, respirations 20, pulse oximeter 96 percent on room air.

Narrative Method

- 10/3/05 8 A.M. Patient stated, “It hurts to take a deep breath.” Rates pain 8/10, + guarding, vital signs BP 144/90, temperature 100.8, pulse 108, respirations 24 and shallow, pulse oximetry 92 percent on room air. Decreased breath sounds at bases owing to poor inspiratory effort. Dressings dry and intact. Reviewed use of PCA morphine and incentive spirometer with patient. Patient instructed to cough and deep breathe with splinting. Patricia Dillon, RN
- 9 A.M. Patient coughing and deep breathing, using incentive spirometer. ambulating with assistance, using PCA morphine prn pain, 5/10, lungs clear. Vital signs: BP 130/86, temperature 99°F, pulse 80, respirations 20, pulse oximetry 96 percent on room air. Patricia Dillon, RN

Whatever format or method is used, you need to document accurately and concisely. Documentation is a part of your patient’s permanent record, and the information is confidential. Because the data should be available to all healthcare members involved in your patient’s care, they should be readily accessible and easy to read. Other members of the healthcare team should be able to peruse the data quickly and identify pertinent findings. Remember, if your plan of care is to be successful, everyone involved in your patient’s care needs to have access to the data.

S U M M A R Y


- The nursing process is a systematic, problem-solving method that guides nursing practice. Assessment is the first step of the nursing process. From your assessment of your patient, the rest of the process flows.
- Assessment is a process of continuous data collection—a thinking, doing, feeling process with ethical and legal ramifications.
- As a thinking process, it requires cognitive skills needed for critical thinking, creative thinking, and clinical decision making.
- As a doing process, assessment requires psychomotor skills to perform a physical assessment.
- As a feeling process, assessment requires affective and interpersonal skills to develop caring, therapeutic nurse-patient relationships.
- Assessment entails responsibility and accountability for your practice and to your patient.
- Assessment has always been a part of nursing practice, but the nurse’s role, particularly in physical assessment, has expanded through the years.

Assessment is now an expected part of every nurse's practice.

- Even though the assessment process may be the same for nursing and medical practice, the outcome differs.
- The goal of medical practice is to diagnose and treat disease.
- The goal of nursing practice is to diagnose and treat human responses to actual or potential health problems. Nursing assessment focuses not only on the patient's physiological and psychological responses but also on her or his psychosocial, cultural, developmental, and spiritual dimensions. It identifies the patient's responses to health problems as well as her or his strengths. Nursing's aim is to assist the patient to reach her or his optimal level of wellness.
- You will use interviewing, observation, and physical assessment to collect data. Once the data have been collected, you will need to validate them. The data are then organized into clusters and prioritized. From this, you will formulate your nursing diagnosis and develop a plan of care.
- Document your findings to communicate the plan of care to all members involved in your patient's care.
- Initially, as you begin to develop your assessment and clinical decision-making skills, your focus is limited. You have difficulty looking at your patient as a whole; instead, you tend to focus on the parts while you strive to develop skills. With experience, as you develop your assessment and clinical decision-making skills, you will begin to recognize patterns and to see the entire picture while zooming in on the acute problems.
- With practice, you will develop competence in your skills and confidence in your findings. So practice and learn from your patients, for they can teach you so much!

R E V I E W Q U E S T I O N S

1. You are performing a preschool examination on your patient. You have just taken his vital signs, height, and weight. Which phase of the nursing process do these activities describe?
 - a. Assessment
 - b. Planning
 - c. Implementation
 - d. Evaluation
2. What activities are performed during the assessment phase of the nursing process?
 - a. Identifying actual/potential health problems
 - b. Determining short- and long-term goals
 - c. Establishing patient outcomes
 - d. Collecting data
3. You are working as a community health nurse, teaching a drug awareness program to fifth-grade students at a local school. Which level of preventive health care does this situation exemplify?
 - a. Primary
 - b. Secondary
 - c. Tertiary
4. You are working in home care, caring for a hospice patient. Which level of preventive healthcare does this situation exemplify?
 - a. Primary
 - b. Secondary
 - c. Tertiary
5. In which situation would a focused assessment be most appropriate?
 - a. New admission scheduled for a total knee replacement
 - b. Patient admitted with chest pain
 - c. College student health exam
 - d. New employee physical exam
6. When obtaining a health history, what would be the appropriate distance between you and the patient?
 - a. 0–18 inches
 - b. 1–2 feet
 - c. 3–4 feet
 - d. 5–12 feet
7. Which of the following is the best example of an open-ended question?
 - a. Are you having chest pain?
 - b. What does the pain feel like?
 - c. Do you have pain in your arm?
 - d. Did you take any medications?
8. You are obtaining an admission history, and you say, "As you were saying, you came to the hospital for ..." This statement is an example of:
 - a. Redirecting
 - b. Restating
 - c. Focusing
 - d. Reflection
9. Which of the following tasks are performed during the working phase of an interview?
 - a. Explaining purpose of the interview
 - b. Collecting data
 - c. Setting the time frame
 - d. Summarizing data
10. You are performing a cardiac assessment on your patient and you think you hear a murmur, but you are not sure. What would be the best way to validate your finding?
 - a. Listen again
 - b. Check the patient chart
 - c. Have colleague or physician listen
 - d. Ask family member if patient has murmur



The Health History

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Discuss purpose of the health history
- Define the components of the health history
- Differentiate between a complete vs. focused health history
- Differentiate a nursing health history from a medical health history
- Identify the key questions for a symptom analysis
- Obtain a health history
- Document your health history

The **health history** provides the subjective database for your assessment. Often, the health history is your first major interaction with your patient—so make it count. You have only one chance to make a good first impression, and it is often the first impression that your patient will remember. This first encounter provides the foundation and sets the tone for your nurse-patient relationship.

The health history is subjective, allowing you to see your patient through his or her “eyes.” It consists of what the patient tells you, what the patient perceives, and what the patient thinks is important. It provides a holistic, qualitative picture of your patient. Clues that you obtain from the health history will direct your physical assessment and are essential in developing a successful plan of care for your patient.

- Identify supports.
- Identify teaching needs.
- Identify discharge needs.
- Identify referral needs.

Types of Health Histories

A health history may be either complete or focused. A **complete health history** includes biographical data, reason for seeking care, current health status, past health status, family history, a detailed review of systems, and a psychosocial profile. A **focused health history** focuses on an acute problem, so all of your questions will relate to that problem.

Purpose of the Health History

The purpose of the health history is to identify not only actual or potential health problems but also your patient’s strengths. It should also identify discharge needs. In fact, a successful discharge plan begins on admission with the health history. To create a successful plan of care, you must take a holistic view of your patient and all that affects her or him. Remember: The plan you develop will be successful only if your patient is able to follow through with it after discharge.

The purpose of the health history is to:

- Provide the subjective database.
- Identify patient strengths.
- Identify patient health problems, both actual and potential.

Complete Health History

The complete health history begins with biographical data, including the patient’s name, age, gender, birth date, birthplace, marital status, race, religion, address, education, occupation, contact person, and health insurance/social security number. It should also include the source of the health history and his or her reliability, who referred the patient, and whether or not the patient has an advance directive. Once you have obtained this information, you should then identify the reason for seeking healthcare, followed by a description of current health status.

The **past health history** includes childhood illnesses, surgeries, injuries, hospitalizations, adult medical problems, medications, allergies, immunizations, travel, and military service. The family history will identify familial or genetically linked disorders. The review of systems provides a comprehensive assessment to determine your

Components of Health Histories

COMPLETE HEALTH HISTORY

Biographical data
 Reason for seeking care
 Current health status and symptom analysis if indicated
 Past health history

 Family history
 Review of systems
 Psychosocial profile
 Developmental considerations
 Ethnic considerations

FOCUSED HEALTH HISTORY

Biographical data
 Reason for seeking care
 Current health status and symptom analysis

 Past health history only as it relates to specific reason for seeking care
 Check for history of the most prevalent diseases in the United States: heart disease, hypertension, cancer, diabetes, and alcoholism.
 Family history only as it relates to specific reason for seeking care
 Review of systems only as it relates to specific reason for seeking care
 Psychosocial profile only as it relates to specific reason for seeking care
 Developmental considerations only as they would affect the acute problem
 Ethnic considerations only as they would affect the acute problem

patient's physiological status. Past or current problems may be identified and warrant further investigation.

The **psychosocial profile** gives you a picture of your patient's health promotion and preventive patterns. It includes a description of health practices and health beliefs, a typical day, nutritional patterns, activity/exercise patterns, recreational patterns, sleep/rest patterns, personal habits, occupational and environmental risk factors, socioeconomic status, developmental level, roles and relationships, self-concept, religious and cultural influences, supports, sexuality patterns, and finally, the emotional health status of your patient. Once you have completed the health history, summarize any pertinent data.

A complete health history provides a comprehensive, holistic picture of your patient. It screens for actual or potential problems and identifies your patient's strengths and health promotion patterns. A complete health history may be obtained in a primary setting as a screening tool, in a secondary setting once your patient's condition stabilizes, or in a tertiary setting to establish a baseline from which to develop your plan of care.

Focused Health History

A focused health history contains necessary biographical data, including the patient's name, age, birth date, birthplace, gender, marital status, dependents, race, religion, address, education, occupation, contact person, and health insurance/social security number. It also includes the source of the health history and her or his reliability, who referred the patient, and whether or not the patient has an advance directive. You should then identify the reason for seeking care, followed by a complete symptom analysis.

In your past health history, address any areas that relate to the reason for seeking care, including diseases of high incidence in the United States, such as heart disease, hypertension, cancer, diabetes, and alcoholism. In your review of systems, ask questions about every system and how it relates to the presenting health problem. The questions in the psychosocial profile identify the impact of the presenting health problem on your patient's life.

A focused health history may be indicated when your patient's condition is unstable or when time constraints are an issue. Focused health histories may also be used for episodic follow-up visits for your patient. In this case, you have already obtained a detailed health history at an earlier point and have established the subjective database. During the follow-up visits, you need to obtain further subjective data to monitor and evaluate your patient's progress. Once you have completed the focused health history, remember to document any pertinent findings.

Focused versus Comprehensive History

Deciding which type of health history to do depends on two factors: your patient's condition and the amount of time you have.

Patient's Condition

First, determine the condition of your patient. This condition may prohibit a detailed health history upon admission. For example, if you are working in the emergency department and John Harrison, a 49-year-old man, presents with acute chest pain, a comprehensive health history is not indicated. Instead, you should obtain a focused history while you perform a physical assessment, draw laboratory studies, obtain an electrocardiogram, and connect your patient to a cardiac monitor. When a patient is in acute distress, trying to obtain a complete health history not only is detrimental but also provides little valuable or accurate information. So ask key questions that focus on the acute problem; once your patient's condition stabilizes, obtain a more detailed health history.



CRITICAL THINKING ACTIVITY # 1

Suppose you were caring for Mr. Harrison. What questions would you ask him to assess his chest pain?



CRITICAL THINKING ACTIVITY # 2

What question(s) would you ask Mr. Harrison related to his past health history, family history, review of systems, and psychosocial profile?

Amount of Time

Allot at least 30 minutes to an hour to obtain a complete health history. Be sure to let your patient know why you are asking these questions and that it will take time. If you do not have enough time to complete the history, do not rush. Instead, perform a focused history first, and then complete the history at later sessions.



HELPFUL HINT

To save time: Ask the patient to fill out a standard history form; then review it with him or her. If necessary, use secondary sources, such as family members, to obtain data. Ask questions while you're bathing the patient or performing other care.

Some information may already have been provided; for example, biographical data from the admission sheet. Do not repeat questions unless you need your patient's perspective. If the patient is asked the same questions several times by different members of the healthcare team, she or he may become frustrated. So, if you need to repeat questions, be sure to tell her or him the reason why.

Medical History versus Nursing History

The areas addressed and the questions asked during a medical health history are very similar to those in a nursing health history. However, some important differences exist. These differences are defined by the focus and scope of medical versus nursing practice. Although the history questions are similar, the underlying rationale differs. Remember: Physicians diagnose and treat illness. Nurses diagnose and treat the patient's response to a health problem.

For example, Mary Johnson, an 81-year-old woman, is admitted to the hospital with a fractured right hip. The focus of the medical history would be to identify what caused the fracture in order to determine the extent of injury. The history would also try to identify any preexisting medical problems that might make her a poor surgical risk. The physician will use the data that he or she obtains to develop a treatment plan for the fracture.

Although the nursing health history also focuses on the cause of the injury, the purpose is to determine Mary Johnson's *response* to the injury, or what effect it has on her. You will look at much more than the fractured hip. You will consider how the injury affects every aspect of her health and life. Your history will provide clues about the impact of the injury on her ability to perform her everyday activities and help you identify strengths she has that can be incorporated into her plan of care. You will also identify supports and begin your discharge plan. Then you will use the data to develop a care plan with Mary Johnson that includes not only her perioperative phase but also her discharge rehabilitative planning.

Setting the Scene

Before you begin your assessment, look at your surroundings. Do you have a quiet environment that is free of interruptions? A private room is preferred, but if one is not available, provide privacy by using curtains or screens. Prevent interruptions and distractions so that both you and your patient can stay focused on the history. Make sure that the patient is comfortable and that the room is warm and well lit. If the patient uses assistive devices, such as glasses or a hearing aid, be sure that she or he uses them during the assessment to avoid any misperceptions.

Before you begin asking questions, tell your patient what you will be doing and why. Inform him or her if you will be taking notes, and reassure the patient that what he or she says will be confidential. However, avoid excessive note taking—it sends the message to your patient

Key Points to Remember When Obtaining a Health History

- Listen to what your patient is telling you both verbally and nonverbally.
- Don't rush. Allow enough time to obtain the data.
- Ensure confidentiality.
- Provide a private, quiet, comfortable environment.
- Avoid interruptions.
- Tell your patient how long the interview will take and why you need to ask these questions.
- Do not be so concerned about completing forms that you neglect the patient.
- Start with what the patient perceives as the problem.
- Use open-ended questions to elicit the patient's perspective.
- Attend to any acute problems, such as pain, before obtaining a detailed history.
- Remember that quality is more important than the quantity of information obtained.

that the health history form is more important than he or she is. Also, if you are too preoccupied with writing and continually break eye contact, you may miss valuable nonverbal messages. Excessive note taking may also inhibit your patient's responses, especially when discussing personal and sensitive issues such as sexuality or drug or alcohol use.

Be sure to work at the same level as your patient. Sit across or next to her or him. Avoid anything that may break the flow of the interview. If the interview is being recorded or videotaped, be sure to get your patient's permission before starting. Position the equipment as unobtrusively as possible so that it does not distract you or your patient.

Your approach to your patient depends on his or her cultural background, age, and developmental level. Ask yourself, "Are there any cultural considerations that might influence our interaction?" "What approach is best, considering my patient's age?" (See Chapter 1, Health Assessment and the Nurse.)

Components of the Health History

A complete health history addresses health and illness patterns, health promotion and protective patterns, and roles and relationships. The parts of the health history that focus on health and illness patterns include the biographical data, reason for seeking healthcare, current health status, past health history, family history, and review of systems. You identify not only current health problems but

also past health problems and any familial factors that place your patient at risk for health problems. Your patient's health promotion activities, protective patterns, and role and relationship patterns are assessed through the psychosocial component of the health history. Here, you assess for risk factors that pose a threat to your patient's health in every aspect of her or his life. Also, you need to consider your patient's cultural and developmental status as it affects her or his health status.

Biographical Data

The **biographical data** provide you with direct information related to a current health problem, alert you to risk factors for health problems, and point out the need for referrals. Your patient's ability to provide biographical data accurately reflects his or her mental status.

Biographical data include the patient's name, address, phone number, contact person, age/birth date, place of birth, gender, race, religion, marital status, educational level, occupation, and social security number/health insurance. They also include the person who provided the history and her or his reliability as well as the person who referred the patient. Information on advance directives may also be obtained for hospital admissions. Also note any special considerations, such as the use of an interpreter. (See *Biographical Data*.)

Reason for Seeking Healthcare

Ask your patient why he or she is seeking healthcare; then document his or her direct quote. The patient's reason for seeking care is usually related to the level of preventive healthcare—primary, secondary, or tertiary. If the setting is a primary level of healthcare, there is usually no acute problem. The reason generally relates to health maintenance or promotion. For example, the patient states, "I am here for my annual physical examination."

If there is an acute problem, ask the patient to state what the problem is and how long it has been going on. For example, "I have had chest pain for the last hour." If your patient identifies more than one problem, she or he may be confusing associated symptoms with the primary problem. Help her or him clarify and prioritize the problems by asking questions such as, "Which problems are giving you the most difficulty?"

Usually, patients identify problems that affect their ability to do what they usually do. In an acute-care setting, the reason for seeking care is called the *chief complaint*. The chief complaint gives you the patient's perspective on the problem, a view of the problem through his or her eyes.

At the tertiary level, the problem may be well defined, a chronic problem, or an acute problem that is resolving. In this case, the problem does not have the acute or life-threatening urgency of an acute problem.

Current Health Status

Once you have identified the patient's reason for seeking healthcare, assess her or his current health status. At a primary level of healthcare (no acute problem), the current health status should include the following:

- Usual state of health.
- Any major health problems.
- Usual patterns of healthcare.
- Any health concerns.

For example: Patient is Maryanne Weller, age 42, married, mother of three, full-time teacher. Usual state of health good. Has yearly physical with pelvic examination and dental examination. Last eye examination 1 year ago. Expresses concern regarding family history of hypertension and ovarian cancer.

Patients in secondary or tertiary healthcare settings have an existing problem. So you will need to perform a symptom analysis to assess your patient's presenting symptoms thoroughly. Although many questions come to mind, your patient's condition and time constraints may preclude you from going into too much detail. If so, you'll need to zero in on several key areas to evaluate your patient's symptoms. As you perform the symptom analysis, try to determine how disabling this problem is for your patient. Also ask if he or she has any medical problems related to the current problems and if he or she is taking any medications for this current problem.



HELPFUL HINT

Perform a symptom analysis for any positive symptom that your patient reports.

The helpful mnemonic PQRST provides key questions that will give you a good overview of any symptom. Although you can ask additional questions, the following ones provide a thorough analysis of any presenting symptom:

■ Precipitating/Palliative Factors

Ask: What were you doing when the problem started? Does anything make it better, such as medications or certain positions? Does anything make it worse, such as movement or breathing?

■ Quality/Quantity

Ask: Can you describe the symptom? What does it feel like, look like, or sound like? How often are you experiencing it? To what degree does this problem affect your ability to perform your usual daily activities?

Biographical Data

DATA	SIGNIFICANCE/CONSIDERATIONS
Name (Full Name and Maiden Name)	Prevents confusion when patients have similar names. Formats of names may differ culturally (e.g., Cambodians put last name first; married Hispanic women take husband's surname while retaining both parents' last names).
Address and Phone Number	May provide socioeconomic information and identify environmental health risks (e.g., urban vs. suburban or rural neighborhoods).
Contact Person	May be a source for further data and support for patient. Essential in case of medical emergency.
Age/Birth Date/Place of Birth	Allows you to compare stated age with apparent age (e.g., chronic illness can make a person appear older). Identifies age-related risk factors for health problems (e.g., children more at risk for accidents; incidence of cardiovascular disease and cancer increases with age). Age helps identify patient's developmental stage. Place of birth may correlate with environmental risks.
Gender	Identifies risk for gender-related health problems (e.g., women have a much higher incidence of breast cancer than men).
Race/Ethnicity/Nationality	Identifies risk for health problems associated with a particular race or ethnic group (e.g., African American males have high incidence of hypertension, Native Americans have high incidence of diabetes). Identifies possible ethnic influences on healthcare practices.
Religion	Alerts you to religious beliefs that may influence health practices (e.g., Judaism and diet, Jehovah's Witnesses and blood products). May identify sources of support for your patient.
Marital Status (Never Married, Married, Widowed, Separated, Divorced)	Identifies possible support people. Present health status may affect family relationships, patient's economic status, and ability to support family. Referral may be necessary.
Number of Dependents	Do not assume that unmarried patients live alone. Ask if they live with someone.
Educational Level	Helps determine teaching approaches. Do not assume that educational level correlates with knowledge and understanding (e.g., newly diagnosed diabetic may have a Ph.D. but be totally unfamiliar with disease and its management). Do not talk down to patient who has had little formal education.
 HELPFUL HINT Ask your patient what teaching method she or he prefers (e.g., videos, pamphlets, group, one on one).	
Occupation	Identifies possible occupational risk factors. Provides clues about socioeconomic status. Current health status also may affect patient's occupational status. Referral may be necessary.
Social Security Number/Health Insurance	Social Security Number may help in retrieving records. Insurance type may influence length of stay. Referrals may be needed depending on type of health insurance.
Source of History/Reliability	Ideally, patient is source of history. Secondary source, usually a family member or friend, is necessary with children or when patient is unable to provide history. Establish reliability of person providing history by noting consistency in responses and willingness to communicate.
Referral	Identifies primary care physician/practitioner. If no referral source identified, may need to make referral for follow-up.
Advance Directives	Allows you to comply with patient's healthcare wishes if Advance Directives exist.

■ Region/Radiation/Related Symptoms

Ask: Can you point to where the problem is? Does it occur or spread anywhere else? (Take care not to lead your patient.) Do you have any other symptoms? (Depending on the chief complaint, ask about related symptoms. For example, if the patient has chest pain, ask if she or he has breathing problems or nausea.)

■ Severity

Ask: Is the symptom mild, moderate, or severe? Grade it on a scale of 0 to 10, with 0 being no symptom and 10 being the most severe. (Grading on a scale helps objectify the symptom.)

Pain Scales

Several scales are used for assessing pain, but they can be adapted to assess any symptom. Besides the numeric (0 to 10) scale, you may use scales that rate pain/symptoms with words like “little pain” or “worst pain possible.” For children, scales are available with drawings of happy to sad faces or photographs of children’s faces representing “no hurt” to “worst hurt ever.” Crayons may also be used, with different colors representing different degrees of symptoms or pain. For more information, see Chapter 4, Assessing Pain.

■ Timing

Ask: When did the symptom start? How often does it occur? How long does it last?

Past Health History

The **past health history** assesses childhood illnesses, hospitalizations, surgeries, serious injuries, adult medical problems (including serious or chronic illnesses), immunizations, allergies, medications, recent travel, and military service. The purpose is to identify any health factors from the past that may have a direct relationship to your patient’s current health status. For example, a history of rheumatic fever as a child may explain mitral valve disease as an adult.

The past health history also identifies any chronic preexisting health problems, such as diabetes or hypertension, which may directly affect the current health problem. For example, patients with diabetes often have poor wound healing. Also, even though the chronic disease may be well controlled, the current health problem may cause an exacerbation. For instance, a patient with well-controlled diabetes may need to adjust his or her medication when scheduled for surgery, because the stress of surgery can elevate blood glucose. In addition, the past health history can identify additional health risks caused by preexisting conditions.

The past health history may also explain your patient’s response to illness, healthcare, and healthcare workers. If she or he has a history of multiple medical problems requiring frequent hospitalizations, these experiences may affect her or his perception of healthcare either positively or negatively.

When obtaining the past health history, be sure to ask for dates, physicians’ names, names of hospitals, and reasons for hospitalizations or surgeries. This information is important if past records are needed. Also avoid using terms such as “usual,” “general,” or “routine.” For example, “usual” childhood illnesses vary depending on the age of your patient and available immunizations.

Past Health History

DATA

SIGNIFICANCE/CONSIDERATIONS

Childhood Illnesses	Positive history of mumps, chickenpox, rubella, frequent ear infections, frequent streptococcal infections or sore throats, rheumatic fever, scarlet fever, pertussis, or asthma may have a direct link to current health problem (e.g., history of chickenpox explains current shingles).
Hospitalizations	Previous hospitalizations may have a direct link to current problem or provide clues to preexisting problems. Knowing name of hospital and dates facilitates record retrieval. Ask about hospitalizations for both physical and psychological problems.
Surgeries	Knowing past surgical procedures may rule out certain problems or explain others. For example, a patient with right lower quadrant pain cannot have appendicitis if his or her appendix has been removed, but pain may be caused by adhesions.
Serious Injuries	History of serious injuries (fractures, head injuries with loss of consciousness, motor vehicle accidents, burns, or lacerations) may relate to current problem or

(continued)

Past Health History (continued)

DATA	SIGNIFICANCE/CONSIDERATIONS
Serious/Chronic Illness	<p>explain findings during physical examination. For example, past motor vehicle accident (MVA) may cause lingering musculoskeletal problems or scars.</p> <p>Encourages patients with well-controlled chronic illnesses (e.g., heart disease, hypertension, diabetes, cancer, seizures) to identify these illnesses. Otherwise, they may fail to mention them if they are not currently causing problems.</p> <p>Consider patient's age and ask about most prevalent diseases for her or his age group.</p>
Immunizations	<p>Lack of immunization may explain current problem. Consider patient's age: In the United States today, children receive many more immunizations than they did 20 years ago. Also, the need for some immunizations (e.g., smallpox) no longer exists. Ask if children have had the following immunizations: measles, mumps, rubella, chickenpox, hepatitis B, diphtheria, tetanus, polio, and <i>Haemophilus influenzae</i> B (HIB). Ask older adults if they have had a pneumococcal vaccine (Pneumovax) and influenza (flu shot). If not vaccinated against tuberculosis (TB), ask about last purified protein derivative (PPD) test. Consider where patient lives: In the United States, people are not routinely immunized against TB, but in other countries where incidence of TB is high, bacille Calmette-Guérin (BCG) vaccine may be used.</p>
<div style="border: 1px solid #0070C0; padding: 5px;">  <p>HELPFUL HINT</p> <hr style="border: 0; border-top: 1px solid #0070C0; margin: 5px 0;"/> <p>A patient vaccinated against TB with BCG will have a positive PPD.</p> </div>	
Allergies	<p>Allergy may explain current problem.</p> <p>If patient has an allergy, note type of reaction. Allergic reactions include hives, pruritus, and respiratory problems. Side effects include gastrointestinal (GI) upset, nausea, and diarrhea. Remember that immune systems change, so patient may become sensitive to something he or she was not allergic to before, or vice versa. Ask if he or she has ever received penicillin, and if there was a reaction.</p>
<div style="border: 1px solid #0070C0; padding: 5px;">  <p>HELPFUL HINT</p> <hr style="border: 0; border-top: 1px solid #0070C0; margin: 5px 0;"/> <p>Patients often confuse drug side effects with allergic reactions.</p> </div>	
Medications	<p>Medications may be causing current problem. For example, over-the-counter (OTC) medication may be interacting with a prescribed medication, causing adverse effects or negating desired effects. Allows you to assess patient's understanding of her or his medications, which may identify teaching needs. Ask about prescribed and OTC medications, including vitamins, supplements, and herbs. Obtain name of medication, dose, frequency, and last time taken.</p>
Recent Travel	<p>May identify exposure to health hazards and explain presenting symptoms (e.g., traveler's diarrhea).</p>
Military Service	<p>Recent or past military service may identify exposure to health hazards. For example, exposure to Agent Orange during Vietnam War is risk factor for cancer, and exposure to chemical toxins during Operation Desert Storm is risk factor for later health problems.</p>




FIGURE 2.1. Family history by genogram

Family History

The **family history** provides clues to genetically linked or familial diseases that may be risk factors for your patient. Ask about the health status and ages of your patient's family members. Family members include the patient, spouse, children, parents, siblings, aunts and uncles, and grandparents. Ask about genetically linked or common diseases, such as heart disease, high blood pressure, stroke, diabetes, cancer, obesity, bleeding disorders, tuberculosis, renal disease, seizures, or mental disease. If the patient's family members are deceased, record the age and cause of death.

The family history may be recorded in one of two ways. You can list family members along with their age and health status (see *Family History by Listing Family Members*), or you can use a genogram (family tree). A genogram allows you to identify familial risk factors at a glance. When developing a genogram, use symbols to represent family members, and include a key to explain the symbols and abbreviations (Fig. 2.1). Another tool that can be effective in taking a patient's family history is an ecomap (see *Understanding Ecomapping*).

Family History by Listing Family Members

The following history is for a 37-year-old female patient.

Patient: Age 37, alive and well
 Spouse: Age 40, divorced, alcoholism
 Daughter: Age 12, alive and well
 Son: Age 8, alive and well
 Brother: Age 32, alive and well
 Sister: Age 30, alive and well
 Father: Age 66, hypertension (HTN)
 Mother: Age 60, mitral valve prolapse (MVP)
 Paternal Aunt: Age 65, breast cancer
 Maternal Uncle: Age 62, HTN
 Maternal Uncle: Deceased age 28, tuberculosis (TB)
 Maternal Aunt: Age 64, MVP
 Maternal Aunt: Age 58, HTN
 Maternal Aunt: Deceased age 9, ruptured appendix
 Paternal Grandfather: Deceased age 68, cancer
 Paternal Grandmother: Age 80, HTN
 Maternal Grandmother: Age 77, HTN, breast cancer
 Maternal Grandfather: Deceased age 70, cardiovascular disease (CVD)

Understanding Ecomapping

An ecomap is a family assessment tool that identifies the needs, patterns, and relationships among family members and the environment, such as school, work, church, healthcare system. The ecomap presents a visual representation as a genogram does for a family history. Characteristics of an ecomap include:

- Identifies major systems that are a part of the family's life.
- Depicts the nature of the relationships with the external systems.
- Identifies both positive and negative relationships with the external systems.
- Identifies the flow of energy and resources between family members and the environment.
- Allows you to see each member within the whole family unit and the relationship with the environment, identifying areas of strengths and conflict.

Source: <http://www.uic.edu/nursing/genetics/Lecture/Family/ecomap.htm>

Review of Systems

The **review of systems (ROS)** is a litany of questions specific to each body system. The questions are usually about the most frequently occurring symptoms related to a specific system. The ROS is used to obtain the current and past health status of each system and to identify health problems that your patient may have failed to mention previously. Remember, if your patient has an acute problem in one area, every other body system will be affected, so look for correlations as you proceed with the ROS. Then perform a symptom analysis for every positive finding and determine the effect of, and the patient's response to, this symptom. The ROS also provides clues to health promotion activities for each particular system. Identify health promotion activities and provide instruction as needed.



CRITICAL THINKING ACTIVITY #3

After studying the boxed family history on page 35, what familial health risks would you identify for this patient?



HELPFUL HINT

As you proceed with the ROS, consider any prescribed or over-the-counter medications your patient is taking and how they affect every system. This may help explain some of your findings.

Review of Systems

AREA/SYSTEM

ASK ABOUT

General Health Survey

Unusual problems or symptoms, fatigue, exercise intolerance, unexplained fevers, night sweats, weakness, difficulty doing activities of daily living (ADLs), number of colds or illnesses per year.

Integumentary

Skin diseases, such as psoriasis, itching, rashes, scars, sores, ulcers, warts, and moles; changes in skin lesions; skin reaction to hot and cold. Changes in hair texture, baldness, usual patterns of hair care (e.g., shampooing, coloring, permanents). Changes in nails (e.g., color, texture, splitting, cracking, breaking); usual patterns of nail care (e.g., use of polish, acrylic nails).



HELPFUL HINT

Recommend an annual skin examination for all adults.

Head and Neck

Headaches; lumps; scars; recent head trauma, injury, or surgery; history of concussion or loss of consciousness; dizzy spells; fainting; stiff neck; pain with movement of head and neck; swollen glands, nodes, or masses.

AREA/SYSTEM	ASK ABOUT
Eyes	<p>Wearing glasses or contact lenses, visual deficit, last eye examination, last glaucoma check, eye injury, itching, tearing, drainage, pain, floaters, halos, loss of vision or parts of fields, blurred vision, double vision, colored lights, flashing lights, light sensitivity, twitching, cataracts or glaucoma, eye surgery, retinal detachment, strabismus, or amblyopia.</p> <div data-bbox="574 525 1212 672" style="border: 1px solid #0070C0; padding: 5px;">  HELPFUL HINT Recommend a routine eye examination and glaucoma check for adults every 2 years. </div>
Ears	<p>Last hearing test, difficulty hearing, sensitivity to sounds, ear pain, drainage, vertigo, ear infections, ringing, fullness in ears, ear wax problems, use of hearing aids, ear-care habits, such as use of cotton-tipped swabs.</p> <div data-bbox="574 871 1212 1039" style="border: 1px solid #0070C0; padding: 5px;">  HELPFUL HINT Test your patient's hearing during the physical examination and screen for hearing losses routinely in elderly patients. </div>
Nose and Sinuses	<p>Nosebleeds, broken nose, deviated septum, snoring, postnasal drip, runny nose, sneezing, allergies, use of recreational drugs, difficulty breathing through nose, problem with ability to smell, pain over sinuses, sinus infections.</p>
Mouth and Throat	<p>Sore throats, streptococcal infections, mouth sores, oral herpes, bleeding gums, hoarseness, changes in voice quality, difficulty chewing or swallowing, changes in sense of taste, dentures and bridges, description of dental health, dental surgery, last dental examination, dental hygiene patterns.</p> <div data-bbox="574 1365 1212 1480" style="border: 1px solid #0070C0; padding: 5px;">  HELPFUL HINT Recommend an annual dental examination. </div>
Respiratory	<p>Breathing problems; cough; sputum (color and amount); bloody sputum; shortness of breath (SOB) with activity; noisy respirations such as wheezing (as is seen with asthma); pneumonia; bronchitis; tuberculosis (TB); last chest x-ray and results; purified protein derivative (PPD) and results; history of smoking.</p>
Cardiovascular	<p>Chest pain; palpitations; murmurs; skipped beats; hypertension (HTN); awakening at night with SOB; dizzy spells; cold or numb hands and feet; color changes in hands and feet; pain in legs while walking; swelling of extremities; hair loss on legs; sores that do not heal; results of electrocardiogram, if ever done.</p> <div data-bbox="574 1837 1212 1974" style="border: 1px solid #0070C0; padding: 5px;">  HELPFUL HINT Recommend a blood pressure reading every 1 to 2 years. </div>

(continued)

Review of Systems (continued)

AREA/SYSTEM	ASK ABOUT
Breasts	<p>Breast masses, lumps; pain, discharge, swelling; changes in breast or nipples; cystic breast disease; breast cancer; breast surgery, reduction, enhancements; breast self-examination (BSE) (when and how); date of last clinical breast examination; date of last mammogram, if ever done.</p>
	<p> HELPFUL HINT</p> <p>Recommend BSE monthly, clinical breast examination yearly, and mammogram yearly for patients over age 40, unless indicated earlier.</p>
Gastrointestinal	<p>Loss of appetite; indigestion; heartburn; gastroesophageal reflux disease (GERD); nausea; vomiting; vomiting blood; liver or gallbladder disease; jaundice; abdominal swelling; regular bowel patterns; changes in bowel patterns; color of stool; diarrhea; constipation; hemorrhoids; weight changes; use of laxatives and antacids; date and results of last fecal occult blood test, if ever done.</p>
	<p> HELPFUL HINT</p> <p>Recommend annual fecal occult blood test and sigmoidoscopy/colonoscopy every 5 to 10 years for patients over age 50.</p>
Genitourinary	<p>Pain on urination; burning; frequency; urgency; dribbling; incontinence; hesitancy; changes in urine stream, color of urine; history of urinary tract infections, kidney infections, kidney disease, kidney stones, frequent nighttime urination.</p>
Female Reproductive	<p>Menarche; description of cycle; last menstrual period; painful menstruation; excessive bleeding; irregular menses; bleeding between periods; last Pap test and results; satisfaction with sexual performance; painful intercourse; use of contraceptives; history of sexually transmitted disease (STD); knowledge of prevention of STDs including human immunodeficiency virus (HIV); infertility problems; obstetrical history including pregnancies, live births, miscarriages, abortions.</p>
	<p> HELPFUL HINT</p> <p>Recommend a yearly pelvic examination with Pap test for all sexually active women.</p>
Male Reproductive	<p>Lesions; discharge; pain on urination; painful intercourse; prostate or scrotal problems; history of STDs, infertility problems, impotence, or sterility; satisfaction with sexual performance; knowledge of prevention of STDs including HIV; use of contraceptives; frequency and technique for testicular self-examination, if ever done; date and results of last prostate examination, if ever done.</p>
	<p> HELPFUL HINT</p> <p>Recommend monthly testicular self-examination and annual digital rectal examination of prostate for men age 50 and over.</p>

AREA/SYSTEM	ASK ABOUT
Musculoskeletal	Fractures, sprains, muscle cramps, pain, weakness, joint swelling, redness, limited range of motion, joint deformity, noise with movement, spinal deformities, low back pain, loss of height, osteoporosis, degenerative joint disease, or rheumatoid arthritis; impact on ability to do ADLs; use of calcium supplements.
	 HELPFUL HINT Recommend a dexascan for bone density for high-risk postmenopausal women.
Neurological	Loss of consciousness; fainting; seizures; head injury; changes in cognition or memory; hallucinations; disorientation; speech problems; sensory disturbances such as numbness, tingling, or loss of sensations; motor problems; problems with gait, balance, or coordination; and impact on ability to do ADLs.
Endocrine	Endocrine disorders such as thyroid disease or diabetes; unexplained changes in weight or height; increased thirst, hunger, or urination; heat and cold intolerance; goiter; weakness; hormone therapy; changes in hair or skin.
Immune/Hematologic	Anemia, bleeding disorders; recurrent infections; cancers; HIV; fatigue; blood transfusion; bruising; allergies; unexplained swollen glands.

Developmental Considerations

The last part of your health history is taking a psychosocial profile, but before you do this, consider the developmental stage of your patient. A person's development crosses the life span. Developmental assessments are often performed on children because the developmental changes that occur at this age are very observable and measurable. Yet adults also go through developmental changes that you need to consider during the assessment. Illness and hospitalization can have a major impact on a child's growth and development, by either halting its progression or regressing it to an earlier stage. For example, when Johnny, age 4, is admitted to the hospital for a hernia repair, he begins wetting the bed during the night, even though his

mother assures you that he has been "potty trained" since age 3.

Several developmental theories exist and will provide a framework for your psychosocial profile. These theories focus on specific areas of development, such as physical, psychosocial, cognitive, and behavioral. Identifying your patient's developmental stage will help you determine the relationship between the patient's health status and his or her growth and development. Because many of these theories do not cross the life span, do not limit yourself to one developmental model. Each theorist views development from a different perspective. So be open and choose the theory or theories that will best help you assess your patient's development. (See *Developmental Stages and Theories* and *Developmental Tasks*.)

Summary of Developmental Theories

Knowledge of the various developmental theories provides the nurse with a framework for cultural and psychosocial assessment. This chart lists selected developmental theorists and summarizes the focus of their theories.

Theorist	Theory Focus
Sigmund Freud	<i>Psychosexual</i> : Biological drives influence a person's psychological and personality development.
Erik Erikson	<i>Psychosocial</i> : The human life cycle is composed of eight developmental stages, each containing a developmental crisis to be resolved. Psychosocial strengths emerge with resolution of the crisis.
Abraham Maslow	<i>Self-actualization</i> : People are innately motivated toward psychological growth, self-awareness, and personal freedom. Basic needs must be met before a person can advance to higher needs.

(continued)

Summary of Developmental Theories (continued)	
Theorist	Theory Focus
Jean Piaget	<i>Cognitive development:</i> An individual's knowledge comes from the interaction between genetic potential and culturally influenced environmental experiences.
Lawrence Kohlberg	<i>Moral development:</i> Cognitive development and emotional growth affect the individual's ability to make autonomous decisions.
Carol Gilligan	<i>Moral development from a female perspective:</i> Women have moral concern for others based on their innate nurturing instincts. They maintain social rules and the expectations of families, social groups, or culture.
Robert Butler	<i>Life review:</i> Elderly patients spend time reviewing past events and concentrating on past conflicts. This life review correlates with the elderly patient's good long-term memory.
Robert Peck	<i>Psychosocial:</i> This theory emphasizes aspects of development from middle age through old age by dividing Erikson's last stage into two parts. There is a developmental crisis to be resolved in each part.
Robert Havighurst	<i>Activity during aging:</i> Elderly patients who stay active and maintain or find substitutes for activities of middle age are more satisfied with life than elderly patients who do not. Diminished activity is equated with increased social isolation and accelerated physical decline.
Elaine Cumming and William Henry	<i>Disengagement during aging:</i> A person naturally gives up roles (a career, for example) and undergoes losses (through death) as aging occurs. As losses occur, a person withdraws from the high activity level of middle age. At the same time, society withdraws from the elderly patient to avoid assigning crucial roles to a group with a high death rate. This mutual disengagement helps society and the elderly patient prepare for death.
Evelyn Duvall	<i>Family development:</i> Family goes through identifiable stages with tasks to be mastered.

Developmental Stages and Theories				
STAGE	FREUD	ERIKSON	PIAGET	KOHLBERG
Infancy: Birth to 1 Yr	Oral-sensory	Trust vs. mistrust	Sensorimotor	
Toddler: 1–3 Yr	Anal	Autonomy vs. shame	Preoperational thought—preconceptual phase	Preconventional level—punishment and obedience
Preschool: 3–6 Yr	Phallic	Initiative vs. guilt	Preoperational—intuitive phase	Preconventional level
School Age: 6–12 Yr	Latency	Industry vs. inferiority	Concrete operations—inductive reasoning and beginning to think logically	Conventional level—conformity and loyalty; responds to authority
Adolescence: 12–18 Yr	Genital	Identity vs. role diffusion	Formal operation—deductive and abstract reasoning	Postconventional—behavior corresponds with society
Young Adult: 18–40 Yr		Intimacy vs. isolation		
Middle Adult: 40–64 Yr		Generativity vs. stagnation		
Older Adult: 64 Yr to Death		Integrity vs. despair		

Developmental Tasks

STAGE	FREUD	ERIKSON	PIAGET	KOHLBERG	HAVIGHURST
Infancy	Receives pleasure (oral gratification) through sucking.	Begins to trust caregiver. Develops drive and hope.	Understanding of the world is through overt physical actions. Moves from simple reflexes to organized set of schemes of permanent objects.		<i>Between infancy to school age:</i> Learns to walk, talk, eat, control elimination. Learns sex differences and modesty. Attains psychological stability. Learns to relate emotionally. Learns right from wrong. Develops a conscience.
Toddler	Delays gratification through bowel and bladder control.	Becomes independent and learns to cooperate with others. Achieves self-control and willpower.	Develops symbolic play, graphic imagery, mental imagery, and language. Egocentric thinking, inflexible, semilogical reasoning, and limited social cognition.	<i>Stage 1:</i> Punishment/obedience. Decisions based on avoiding punishment.	
Preschooler	Curious about gender differences and genitals.	Achieving confidence in own abilities. Has direction and purpose.		Naïve instrumental orientation—actions are directed to meet his or her own needs. Very concrete sense of justice/fairness.	
School age	Transition period, develops peer relationships, identifies with parent or caregiver of same sex.	Accomplishments are source of pleasure. Develops methods and competence.	Develops the understanding of conservation of quantity.	<i>Stage 2:</i> Instrumental/relativist. More self-reliant. Behavior directed to satisfy needs. Negotiate and trade off to meet needs. <i>Stage 3:</i> Approval seeking. Behavior is directed to gain approval of significant others.	Develops physical skills; positive self-image; personal independence, relationships with peers, appropriate roles, cognitive skills of reading, writing, and calculations; concepts for everyday living (conscience, values, and morality) and opinions of social and institutional groups.

Developmental Tasks (continued)

STAGE	FREUD	ERIKSON	PIAGET	KOHLBERG	HAVIGHURST
Adolescent	Attains sexual maturity and becomes interested in sex.	Realizes a stable sense of identity. Develops devotion and fidelity.	Able to combine systems. Develops hypothetical-deductive thinking.	<i>Stage 4: Law and order.</i> Behavior guided by law and order, the rules of society.	Acceptance and development of gender role. Develops more mature relationships. Attains emotional, financial independence from parents. Begins to think of choosing a career and a partner for marriage. Intellectual skills and socially acceptable behavior develop further.
Young adult		Develops personal and professional relationships. Attains affiliation and love.		<i>Stage 5: Social contract.</i> Respects individual differences. Realizes that social laws may contradict moral principles. Tries to better the social condition.	Selects mate. Learns to live together with partner. Starts career and family. Rears family. Manages career, family, and home. Assumes civic responsibility. Has social group of friends.
Middle adult		Productive and creative in both work and relationships. Develops productivity and caring for family, friends, and community.		<i>Stage 6: Universal-ethical.</i> Universal-ethical principles guide moral judgment with underlying good of humanity as driving force even at expense of the individual. (This stage is rarely met by most individuals over the course of a lifetime.)	Assumes more civic and social responsibility. Maintains career and family. Develops leisure activities. Strengthens spousal relationship. Accepts midlife changes. Cares for aging parents.
Older adult		Has had purpose in life and fulfillment. Achieves sense of fulfillment and wisdom.	Adjusts to physical changes, retirement, reduced income, loss of spouse, family members, and friends.		Develops explicit identity with own age group. Modifies living arrangements. Meets social and civic obligations.

Source: Data from Craven, R., and Hirnle, C. (2006). *Fundamentals of Nursing: Human Health and Function*, ed. 3. Philadelphia: Lippincott Williams & Wilkins.

Assessing the Family

Assessment of how and to what extent the patient's family fulfills its functions is an important part of the patient history. The nurse should assess the family into which the patient was born (family of origin) and, if different, the current family.

Use this guide to assess how the patient perceives family functions. Because the questions target a nuclear family—that is, mother, father, and children—they may need modification for single-parent families, families that include grandparents, patients who live alone, or unrelated individuals who live as a family.

Affective Function

Assessing how family members feel about, and get along with, each other provides important information. In some families, one person performs the “sick role,” and the other family members support the illness and keep the member sick. For example, a child whose parents have marital problems may be sick to get attention. The parents may allow the child to be sick so they can focus their attention on the child and avoid dealing with problems. To assess affective function, ask the following questions:

- How do the members of your family treat each other?
- How do family members regard each other?
- How do the members of your family regard each other's needs and wants?
- How are feelings expressed in your family?
- Can family members safely express both positive and negative feelings?
- What happens in the family when members disagree?
- How do family members deal with conflict?

Socialization and Social Placement

These questions provide information about the flexibility of family responsibilities, which is helpful for planning a patient's discharge. For example, a mother of small children who has just had minor surgery may need household help when she goes home if the husband is not expected to help or does not want to. To assess socialization and social placement, ask the following questions:

- How satisfied are you with your role and your partner's role as a couple?
- How did you decide to have (or not to have) children?
- Do you and your partner agree about how to bring up children? If no, how do you work out the differences?
- Who is responsible for taking care of the children? Is this mutually satisfactory?
- How well do you feel your children are growing up?
- Are family roles negotiable within the limits of age and ability?
- Do you share cultural values and beliefs with the children?

Healthcare Function

This assessment will uncover many cultural beliefs. Identify the family caregiver and then use that information when planning care. For example, if the patient is the caregiver, then the patient may need household help when discharged.

To assess healthcare function, ask the following questions:

- Who takes care of family members when they are sick? Who makes doctor appointments?
- Are your children learning skills, such as personal hygiene, healthful eating habits, and the importance of sleep and rest?
- How does your family adjust when a member is ill and unable to fulfill expected roles?

Family and Social Structures

The patient's view of the family and of other social structures affects healthcare. For example, if the patient needing home healthcare belongs to an ethnic group with a strong sense of family responsibility, then the family probably will care for the patient. To assess the importance of family and social structures, ask the following questions:

- How important is your family to you?
- Do you have any friends that you consider family?
- Does anyone other than your immediate family (for example, grandparents) live with you?
- Are you involved in community affairs? Do you enjoy these activities?

Economic Function

Financial problems frequently cause family conflict. Ask these questions to explore money issues and how they relate to power roles within the family:

- Does your family income meet the family's basic needs?
- Does money allocation consider family needs in relation to individual needs?
- Who makes decisions about family money allocation?

Psychosocial Profile

The **psychosocial profile** is the last section of the health history. This section focuses on health promotion, protective patterns, and roles and relationships. It includes questions about healthcare practices and beliefs, a description of a typical day, a nutritional assessment, activity and exercise patterns, recreational activities, sleep/rest patterns, personal habits, occupational risks, environmental risks, family roles and relationships (see *Assessing the Family*), and stress and coping mechanisms.

In a primary healthcare setting, the psychosocial assessment enables you to identify how your patient

incorporates health practices into every aspects of her or his life. You can then teach and reinforce health promotion activities that your patient can incorporate into her or his everyday life. If she or he has an acute problem, the psychosocial assessment helps you determine the impact of this illness on every facet of the patient's life and assists you in determining discharge planning needs. For your plan of care to be successful, the patient must be able to follow through with it after discharge. Help ensure success by identifying clues as you perform the assessment and then making appropriate referrals.

Psychosocial Profile

DATA	SIGNIFICANCE/CONSIDERATIONS
Health Practices and Beliefs	Your patient's overall health is affected by her or his values, beliefs, financial status, expectations of healthcare, and other factors. How does the patient perceive her or his role in maintaining health? Does she or he get a yearly physical examination or seek healthcare only when ill? Does she or he perform self-examinations and other self-care measures? Health is a component of every aspect of life. Determine whether your patient's life positively or negatively affects her or his health.
Typical Day	Describing a typical day may identify health risk factors. Or, if the patient has a health problem, it helps determine what effect this problem has on his or her everyday life.
Nutritional Patterns	Eating habits can positively or negatively affect your patient's health. Religious beliefs or culture may influence eating habits. Financial problems, such as being on a fixed income, may limit food purchases. Ask about special diets, food preferences, food allergies, weight changes, happiness with weight, and history of eating disorders. Ask who does the cooking, who does the shopping, and who dines with the patient. All of these factors have an effect on nutrition. Screen for nutritional deficits by doing a 24-hour recall. Ask the patient, "What did you eat yesterday?" Do the recall on a weekday rather than a weekend because it is more reflective of the usual diet. After you complete the 24-hour recall, compare the diet with the food groups (meat, poultry, and fish; grains; dairy; vegetables and fruits) and note any deficits. Take note of number and size of servings. Also note if your patient is taking vitamins, mineral or herbal supplements. Make nutritional referrals as indicated. (See Chapter 8, Assessing Nutrition.)
Activity and Exercise Patterns	Activity and routine exercise can help maintain both physical and mental health. Ask about type and amount of activity or exercise. If your patient participates in contact sports, assess use of protective equipment and provide instruction as needed. If she or he has a health problem, determine its effect on her or his ability to maintain usual activity and exercise patterns.
Recreation, Hobbies, Pets	Recreational activities, hobbies, and pets usually enhance health by reducing stress. But they can also pose health risks. For example, stained glass work exposes one to lead, and pets may trigger allergies or carry diseases such as toxoplasmosis.
Sleep/Rest Patterns	Many physical and psychological problems can affect sleep or are affected by lack of sleep. Ask your patient how many hours of sleep she or he gets, if sleep is interrupted, how many hours she or he needs to feel rested, any medication taken to aid sleep, or if patient has any sleeping disorders such as narcolepsy or sleep apnea.

DATA	SIGNIFICANCE/CONSIDERATIONS
Personal Habits	<p>Unhealthy personal habits, such as use of tobacco, alcohol, caffeine, and drugs, can adversely affect health. Ask for specific information about the amount and length of use. (See Chapter 5, Approach to the Mental Health Assessment.)</p> <p><i>Tobacco:</i> Ask about cigarettes (filtered/unfiltered), pipe, cigars, or chewing tobacco.</p> <p><i>Alcohol:</i> Ask what kinds (wine, beer, mixed drinks), when patient drinks, if he or she drinks to help deal with stress, and whether patient notices a change in his or her drinking patterns (more or less). The CAGE questionnaire can be used to screen for alcohol abuse.</p> <p><i>Caffeine:</i> Ask what kinds (coffee, tea, chocolate) and if patient has trouble sleeping, nervousness, or palpitations.</p> <p><i>Drugs:</i> Ask what kinds (prescription, over-the-counter, recreational, or street drugs), when patient last took drug, and method of use (e.g., inhalation, intravenous, oral).</p> <div data-bbox="638 766 1276 882" style="border: 1px solid red; padding: 5px; margin-top: 10px;">  <p>ALERT IV drug users are at risk for hepatitis B and HIV.</p> </div>
Occupational Health Patterns	<p>Certain occupations pose health risks. Ask your patient if her or his job requires exposure to toxins such as asbestos, pesticides, plastics, anesthetics, radiation, or solvents; protective gear (e.g., construction workers, welders, landscapers, and tree trimmers), or heavy physical activity, such as nursing. Job satisfaction can also affect health. Ask your patient if she or he is satisfied with the job, what she or he likes best and least, how many hours a week she or he works, how far she or he travels to and from work, if there are occupational health programs, if she or he gets work breaks, how much vacation she or he gets, if she or he considers the job stressful, and if she or he is satisfied with the salary received.</p>
Socioeconomic Status	<p>Socioeconomic status can have a major impact on your patient's health and healthcare. Ask your patient if he or she has health insurance, dental insurance, or a prescription plan. Limited financial resources may limit available healthcare services or prevent your patient from following through with a treatment plan. You may need to make referrals to social service agencies for assistance.</p>
Environmental Health Patterns	<p>The patient's home and type of community can have an impact on her or his health. Ask if the patient's community is urban, suburban, or rural. Urban dwellers are exposed to more noise and air pollution, whereas rural dwellers are exposed to more polluted water sources, septic systems, and greater contact with allergens and toxins associated with farming. Determine type of neighborhood. Ask the patient if she or he feels safe; are there adequate sidewalks and lighting; with police and fire departments and ambulance nearby; public transportation; accessibility to food and drug stores; easy access to churches, schools, healthcare facilities, and community supports, such as senior citizen centers? Determine type of home patient lives in. Ask if it is a single home or apartment; if she or he rents or owns it; if living space, heating, water, and plumbing are adequate; if it is one story, two stories, or more; if there are smoke detectors, fire extinguishers, carbon monoxide detectors, and a burglar alarm system; and what type of heating is used. Also ask if medications and cleaning supplies are kept out of the reach of children and if the family uses seat belts, car seats, and bike helmets. Identify teaching needs or areas that warrant referrals or further follow-up.</p>

(continued)

Psychosocial Profile (continued)

DATA	SIGNIFICANCE/CONSIDERATIONS
Roles, Relationships, Self-Concept	<p>Your patient's feelings, attitudes, past experiences, and relationships contribute to his or her sense of value and worth and affect his or her overall health. A person with a positive self-concept and an internal locus of control takes charge and assumes responsibility for his or her life and for achieving health goals. Find out if your patient has a positive or negative self-image by asking what he or she feels are his or her strengths and weaknesses; what he or she likes best and least about himself or herself; and if he or she considers himself or herself outgoing or shy. Find out if his or her locus of control is internal or external by asking if he or she feels in control of what happens to him or her or believes that "whatever will be will be." Does he or she assume responsibility or place the responsibility on others? Illness or changes in body image can pose a threat to self-concept. Be alert for risk factors such as serious illness, loss of function, or surgery that threaten your patient's self-image. Use the supports and resources necessary to help him or her maintain the integrity of his or her self-concept.</p>
Cultural Influences	<p>Culture can influence communication patterns, health beliefs and practices, dietary habits, family roles, and life-and-death issues. Ask yourself: Is there a language barrier? Cultural practices that conflict with the prescribed treatment plan? Dietary preferences that need to be considered? Cultural practices that can be included in the patient's plan of care? As long as cultural beliefs will not harm the patient's health, accepting them and incorporating them into the plan of care can only help ensure compliance. Individualizing the plan of care gives the patient a feeling of ownership and makes her or him more likely to follow through.</p>
Religious/Spiritual Influences	<p>Religion and spirituality influence health beliefs and practices, dietary preferences, family roles, and life-and-death issues. Ask about religious beliefs that conflict with prescribed treatment plan. For example, a Jehovah's Witness may refuse a blood transfusion even in a life-threatening situation. Ask about dietary preferences, such as kosher food for Jewish patients. Ask about religious rites that should be incorporated into the plan of care, such as receiving the sacrament of the sick for Catholic patients. As long as religious beliefs will not harm the patient's health, accepting them and incorporating them into the plan of care will only help ensure its success.</p>
Family Roles and Relationships	<p>The family is an important support system for most ill people. Also, when people become ill, their role in the family changes and the family unit may need to reorganize to sustain itself. Determine what effect this change has on your patient and his or her family; then plan how to meet their needs. You may need to rely on outside supports. Ask your patient about his or her role within the family and other social groups. Is he or she head of the household, responsible for parenting and housekeeping, or are the roles shared? Today's family structure varies greatly, from traditional nuclear family, to single-parent family, to restructured family through divorce or remarriage, to alternative families, to the extended family of the past. But all families share common functions. (See <i>Assessing the Family</i>.)</p>
Sexuality Patterns	<p>Illness can have both physical and psychological effects on your patient's sexuality. Changes in body image or self-concept, changes in ability to perform sexually, prescribed medications, and depression can all have adverse affects on sexuality. Developmental, cultural, and religious factors influence your patient's perspective and expression of sexuality and her or his willingness to discuss sexuality with you. Sexuality concerns also vary according to your patient's age and developmental stage. Sexuality is a sensitive topic that both you and your patient may feel uncomfortable</p>

DATA	SIGNIFICANCE/CONSIDERATIONS
Social Supports	<p>discussing. Pick up on clues from your patient and address sexuality issues when most appropriate, either during the review of systems (ROS) or the psychosocial history. Be open and nonjudgmental. Ask if your patient is sexually active and, if so, whether she or he is satisfied with her or his sexual role, performance, and relationship. Find out the source of her or his knowledge of sexuality, reproduction, birth control, and safe sex practices.</p> <p>Support systems outside the family are important during illness. Ask your patient if there is anyone aside from family that he or she can call on for help—for example, friends, coworkers, community agencies, or clergy. Ask if he or she belongs to a church or to community organizations or clubs and if he or she attends on a regular basis. If your patient's social supports are limited, tell him or her about resources available within the community and church, and help him or her access those that meet his or her needs.</p>
Stress and Coping Patterns	<p>The amount of stress in your patient's life and how she or he copes with it can affect her or his health. Illness only adds stress and anxiety. Ask your patient how she or he deals with everyday stress, what she or he does when feeling upset, if she or he has ever felt sad or depressed, and whom she or he talks to when upset. Keep in mind that normal developmental changes and changes associated with illness can threaten a person's ability to cope. If you determine that your patient's coping skills are ineffective, develop a plan of care, including sources of support, to help her or him deal with stress more effectively.</p>

Documenting Your Findings

Once you have completed the health history, summarize pertinent findings and share them with your patient to confirm their accuracy. Then document your findings and begin to formulate a plan of care.

Documentation of history findings varies from one healthcare facility to another. Many acute-care facilities use computerized programs that enable you to enter the history directly into the computer. Standardized nursing admission assessment forms that combine both history

and physical assessments are also commonly used. Regardless of the system, here are some helpful hints for documenting a health history:

- Be accurate and objective. Avoid stating opinions that might bias the reader.
- Do not write in complete sentences. Be brief and to the point.
- Use standard medical abbreviations.
- Don't use the word "normal." It leaves too much room for interpretation.
- Record pertinent negatives.
- Be sure to date and sign your documentation.

Sample Documentation of Health History

You are now familiar with each step in the history-taking process. If you were interviewing Janet Perry, here is how you would document your findings:

Biographical Data

Name: Janet Perry
 Address: 1234 Happy Valley Road, Abington, Pennsylvania 19001
 Home Phone: 215-555-1234
 Work Phone: 215-555-2468
 Sex: Female

(continued)

Sample Documentation of Health History (continued)

Age: 37
 Birth Date: 4/5/69
 Social Security Number: 123-45-6789
 Place of Birth: Philadelphia, Pennsylvania
 Race: Caucasian
 Nationality: American
 Culture: German American
 Marital Status: Divorced
 Dependents: 2—Kate, age 15; Joe, age 12
 Contact Person: Sister, Marge Jones: 215-885-2345
 Religion: Episcopalian
 Education: B.S. Chemistry
 Occupation: Lab Assistant, Acme Chemical Corp.
 Health Insurance: BC/BS
 Source and Reliability: Patient, seems reliable
 Referral: Self, currently has no primary care provider
 Advance Directives: None

Reason for Seeking Healthcare

"I've been feeling tired lately, for about a month now, so I wanted a health exam to make sure nothing is wrong."

Current Health Status

Says she's been feeling tired for about a month. Denies any other symptoms. Denies loss of appetite or history of anemia or blood loss. States that she has been busier at work than usual and hasn't had enough time to spend with her children because she's going to night school for her master's degree. Thinks fatigue may be related to a busy schedule rather than a physical problem. Last had a complete physical including a Pap smear, 4 years ago.

Past Health History

Childhood Illnesses: Positive for measles, rubella, mumps, chickenpox, frequent sore throats.
Hospitalizations and Surgeries: Hospitalized at age 6 for tonsillectomy and adenoidectomy, St. Mary's Hospital, Philadelphia, Dr. Smith. Hospitalized 1991 and 1994 for childbirth, St. Mary's Hospital, Dr. Ballay.
Serious Injuries: Fractured left arm and concussion from bicycle accident age 10; resolved with no complications.
Serious/Chronic Illness: No history of cardiovascular disease (CVD), hypertension (HTN), diabetes, cancer, seizures.
Immunizations: Up to date, had diphtheria-pertussis-tetanus (DPT), oral polio, smallpox vaccine, last tetanus shot 4 years ago.
Allergies: No known drug, food, or environmental allergies; has had penicillin without reaction.
Medications: Takes Tylenol occasionally for headaches and multivitamin daily.
Travel: No recent travel outside United States.
Military Service: Never served in military.

Family History

Documented by listing family members and health status:

Patient: Age 37, alive and well
Spouse: Age 40, divorced, alcoholism
Daughter: Age 15, alive and well
Son: Age 12, alive and well
Brother: Age 32, alive and well
Sister: Age 30, alive and well
Father: Age 66, HTN
Mother: Age 60, mitral valve prolapse (MVP)
Paternal Aunt: Age 65, breast cancer

Maternal Uncle: Age 62, HTN
 Maternal Uncle: Deceased age 28, tuberculosis (TB)
 Maternal Aunt: Age 64, MVP
 Maternal Aunt: Age 58, HTN
 Maternal Aunt: Deceased age 9, ruptured appendix
 Paternal Grandfather: Deceased age 68, colon cancer
 Paternal Grandmother: Age 80, HTN
 Maternal Grandmother: Age 77, HTN, breast cancer
 Maternal Grandfather: Deceased age 70, CVD

Figure 2.2 is the same information documented as a genogram.

Pertinent Findings: Positive family history for CVD, HTN, MVP, TB, and breast and colon cancer. No family history of stroke, diabetes, obesity, bleeding disorders, renal disease, seizures, or mental disease.

Review of Systems

General Health Survey: Complains of fatigue. Had two head colds in past year. No other illnesses. Good exercise tolerance.

Skin, Hair, and Nails: No suspicious lesions, no changes in hair or nails.

Head and Neck: Positive headaches, once a month caused by tension, 6/10 pain scale, relieved with Tylenol. Had concussion as a child, no seizures. No neck pain, masses, has full range of motion (ROM) of neck.

Nose and Sinuses: No history of nosebleeds, rhinorrhea, or sinus infections.

Mouth and Throat: Last dental exam 1 year ago, few fillings, brushes twice a day and flosses. Had tonsillectomy and adenoidectomy (T and A) as child, no further sore throats.

Eyes: Last eye exam 2 years ago. Wears contact lenses for myopia, wears protective goggles in the lab.

Ears: Reports no hearing problems or ear infections. Last hearing exam 5 years ago.

Respiratory: No breathing difficulties, no history of pneumonia, bronchitis, or asthma; exposed to secondhand smoke while married.

Family History By Genogram




FIGURE 2.2. Family history by genogram.

(continued)

Sample Documentation of Health History (continued)

Cardiovascular: No chest pain, palpitations, murmurs, HTN, coldness/numbness/color changes in hands or feet, varicose veins, or swelling. Never had an electrocardiogram (ECG).

Breasts: Reports left breast slightly larger than right, no pain, masses, or discharge. Breast self-exam (BSE) about three to four times a year. Never had a mammogram.

Gastrointestinal: No history of weight changes, appetite change, indigestion, ulcers, liver, or gallbladder disease. Had hemorrhoids during last pregnancy, but resolved after delivery. Regular daily bowel movement, formed and brown in color.

Genitourinary: No history of kidney stones. Had bladder infection 4 years ago, treated with Macrochantin. No problems with urination or incontinence; voids clear yellow urine several times a day.

Reproductive: Menarche at age 12, menses every 30 days, lasts 3–4 days, moderate flow, no discomfort. Obstetrical history: Para 3, gravida 2, 1 miscarriage before oldest child, no complications. No problems with subsequent pregnancies, vaginal deliveries, full term. No history of sexually transmitted diseases (STDs). Used contraceptive pills in past with no adverse reaction. Sexual relations during marriage described as “fine.” Currently not sexually active.

Neurological: History of concussion, occasional headache, no history of loss of cognition or memory, no balance or coordination problems, no loss of sensation.

Musculoskeletal: No muscle or joint pain/stiffness. History of fractured arm.

Immune/Hematologic: Anemic during last pregnancy even when taking iron. No history of blood loss, anemia, or blood transfusion. Blood type A+. No reported swollen lymph nodes or glands.

Endocrine: No history of thyroid disease or diabetes. No hot/cold intolerance, no increased thirst, urination, or appetite.

Developmental Considerations

In “young adult” stage of development. Divorced for 2 years but has a good relationship with ex-husband, a recovering alcoholic. Feels she is emotionally stable, but feels down and lonely now and then because she hasn’t had much time to socialize since divorce.

Psychosocial Profile

Health Practices and Beliefs: Believes each person is responsible for leading a lifestyle conducive to health. Realizes she’s overdue for physical exam, in search of new primary care provider. Satisfied with past experiences with healthcare providers.

Typical Day: Wakes at 6:30 A.M., gets children up and ready for school, has breakfast, takes children to school, arrives at work at 8:30 A.M. and works till 5:00 P.M., picks children up at after-school program, makes dinner, helps children with homework, gets children ready for bed by 9:00–10:00 P.M., studies after children are in bed, retires between 11:30 P.M. and 12 midnight. Goes to night school two nights a week from 6:00 to 9:00 P.M.

Nutritional Patterns: 24-hour recall indicates deficits in iron, protein, and calcium. Takes vitamin and mineral supplement. Frequently skips lunch. Happy with weight, no recent changes.

Activity and Exercise Patterns: Jogs 2 miles three times a week during lunch, would like to do more but can’t find time.

Recreation, Pets, Hobbies: Enjoys swimming, hiking, and camping; goes camping with children in summer. Would like to do more, but has no time. No pets.

Sleep/Rest Patterns: Says she is a morning person. Awakens at 6:30 A.M., goes to bed around 12 midnight. Has no trouble falling asleep, rarely awakens during the night. Would like 8 hours of sleep but rarely gets it.

Personal Habits: Never smoked or used recreational drugs, has two cups of caffeinated coffee a day, drinks one to two glasses of wine a week. Takes vitamin and mineral supplement daily and occasional Tylenol for headache.

Occupational Health Patterns: Works 7½ hours a day with 1-hour lunch. Works well with coworkers. Has had job for 3 years; before divorce was homemaker. Feels under a lot of pressure at work. Is not very satisfied with job and is trying to improve credentials to get better job. Is exposed to toxic chemicals at work but uses proper safety precautions (mask, gloves, goggles).

Socioeconomic Status: States she makes enough to make ends meet—around \$25,000 plus child support of \$800 a month. Has health insurance and retirement plan through work, children’s insurance covered by husband.

Environmental Health Patterns: Lives in three-bedroom single home in suburbs. Considers neighborhood safe. Police and fire departments within 5 miles; schools, stores, and church within 10 minutes’ driving distance from home. Has smoke and carbon monoxide detectors in home, has a burglar alarm system. Drives 20 minutes to work every day. Family uses seat belts and bicycle helmets.

Roles, Relationships, Self-Concept: Describes self as fairly attractive with good sense of humor. Divorce was very difficult, received counseling for 6 months, but is doing OK now.

Cultural Influences: No specific cultural influences that would affect healthcare practices.

Religious/Spiritual Influences: Practicing Episcopalian, but doesn't consider self very religious. Feels it is important for children to attend Sunday services and Sunday school weekly. No religious influences that would affect healthcare practices.

Family Roles and Relationships: Single parent, lab assistant, student with never enough time to do all she wants. Communicates with ex-husband on childcare issues. He is cooperative and provides some financial support. Children get along well for the most part with occasional arguments over her time with them. They help with light chores around the house. Her sister and brother-in-law are major source of support and help with child care. Relationship with parents is distant both geographically and emotionally.

Sexuality Patterns: Currently not sexually active. Says dating again is difficult and she doesn't really have the time.

Social Supports: Aside from sister and brother-in-law, has three close friends in the neighborhood. Doesn't belong to any organized groups, but is aware of such community resources as Parents Without Partners.

Stress and Coping Patterns: Says she usually deals with stress by avoiding problem until it is staring her in the face and she has to deal with it. Job is stressful, but she manages on a day-to-day basis, uses coworkers as support. Feels she is easygoing, but when she hasn't had a break in a while, occasionally "blows up" at the kids.

Summary of Pertinent Findings

Patient, 37-year-old divorced woman experiencing much stress from work, single parenting, and school. No significant physical problems noted. Positive family history of CVD, HTN, MVP, and breast cancer. Needs physical exam and lab studies to rule out iron deficiency anemia or other physical problems. Needs health promotion instruction re: diet, time and stress management, expanded use of community resources. Positive findings include self-motivated to seek healthcare; exercises routinely; has family, friends, and coworkers who are source of support; self-motivated to further education to expand career options.


S U M M A R Y

- The health history provides the subjective database for your patient's assessment. It is usually your first major interaction with your patient and often provides the foundation of your nurse-patient relationship.
- As you proceed with the history, listen to both your patient's verbal and nonverbal messages.
- Phrase your questions with your patient's developmental stage and cultural background in mind.
- After you complete the history, document your findings and summarize pertinent data.
- Sharing your findings with your patient allows you to validate your assessment. You will begin to identify not only actual and potential health problems but also your patient's strengths. Use these strengths in planning his or her care.
- The health history provides clues that will direct your physical assessment as you complete your database and develop a plan of care for your patient.

R E V I E W Q U E S T I O N S

1. You are obtaining a health history on a newly admitted patient with the diagnosis of congestive heart failure. He tells you, "I'm having trouble breathing since yesterday." Once the chief complaint is identified, which part of the history should be completed next?
 - a. Symptom analysis
 - b. Past health history
 - c. Review of systems
 - d. Family history
2. Which section of the health history can be documented as a genogram?
 - a. Chief complaint
 - b. Past health history
 - c. Family history
 - d. Current health status

3. Which section of the health history identifies environmental health risk factors?
 - a. Review of systems
 - b. Psychosocial profile
 - c. Past health history
 - d. Family history
4. When obtaining a family history, who does not need to be included in the genogram?
 - a. Spouse
 - b. Siblings
 - c. In-laws
 - d. Aunts/uncles
5. What is the purpose of the review of systems in the health history?
 - a. Identify actual problems
 - b. Determine cause of current problem
 - c. Identify current/past health of systems
 - d. Validate current symptoms
6. Mr. Brunner, 35 years old, is complaining of abdominal pain. You perform a symptom analysis of the abdominal pain. Which question would best assess the quality of the pain?
 - a. What does it feel like?
 - b. Is the pain sharp?
 - c. When did the pain start?
 - d. Does the pain go anywhere else?
7. Mr. Brunner, 35 years old, is complaining of abdominal pain. You perform a symptom analysis of the abdominal pain. Which question would best assess the severity of the pain?
 - a. Is the pain really bad?
 - b. On a scale of 0 to 10, rate the pain
 - c. Does the pain feel knifelike?
 - d. Does the pain go anywhere else?
8. Mr. Brunner, 35 years old, is complaining of abdominal pain. You perform a symptom analysis of the abdominal pain. What is the best way to assess the location of the pain?
 - a. Where does it hurt?
 - b. Does it hurt in your belly?
 - c. Does it go anywhere else?
 - d. Can you point to where it hurts?
9. What part of the health history identifies the source and reliability?
 - a. Biographical data
 - b. Current health status
 - c. Chief complaint
 - d. Past medical history
10. Circle all that describe a chief complaint.
 - a. Statement of problem as a direct quote from patient
 - b. Symptom analysis of presenting problem
 - c. Identifies duration of problem
 - d. Treatment of problem



Approach to the Physical Assessment

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Discuss the purpose of physical assessment
- Differentiate complete from focused physical assessment
- Differentiate nursing physical assessment from medical physical assessment
- Identify tools used during the physical examination
- Define physical assessment techniques
- Demonstrate physical assessment techniques
- Discuss variations in approaches for different age groups
- Define components of physical assessment

The physical examination is a process during which you use your senses to collect objective data. You will need all of the skills of assessment—cognitive, psychomotor, interpersonal, affective, and ethical/legal—to perform an accurate, thorough physical assessment. You also need to know normal findings before you can begin to distinguish abnormal ones. The best way to perfect your physical assessment skills is through practice. Effective communication skills are essential to establishing the trust needed to proceed with the examination. Finally, you need to remember your ethical and professional responsibility to your patient in respecting his or her right to privacy and confidentiality.

Physical assessment provides another perspective. Whereas the health history allows you to see your patient *subjectively* through *hers* or *his* eyes, the physical examination now allows you to see your patient *objectively* through *your* senses. The objective data complete the patient's health picture.

Purpose of the Physical Assessment

Like the health history, the goal of physical assessment is not only to identify actual or potential health problems but also to discover your patient's strengths. Data from the physical assessment can be used to validate the health history. For example, you can use the physical examination to assess clues you obtained from the history. Combined with the history data, your physical assessment findings are essential in formulating nursing diagnoses and developing a plan of care for your patient.

Types of Physical Assessment

Also like the health history, physical assessment may be either complete or focused. A **complete physical assessment** includes a general survey; vital sign measurements; assessment of height and weight; and physical examination of *all* structures, organs, and body systems. Perform it when you are examining a patient for the first time and need to establish a baseline.

On the other hand, a **focused physical assessment** zeros in on the acute problem. You assess only the parts of the body that relate to that problem. It is usually performed when your patient's condition is unstable, as a follow-up to a complete assessment, or when you are pressed for time.

Complete Physical Assessment

The complete physical examination begins with a general survey. A general survey includes your initial observations of the patient's general appearance and behavior, vital signs, and **anthropometric measurements**. Vital signs include temperature, pulse, respirations, blood pressure, and pulse oximetry, if available. Anthropometric measurements include height and weight.

Next, perform a head-to-toe systematic physical assessment. As you proceed from one area to another, remember that all systems are related, so a problem in one area eventually will affect or be affected by every other system. Therefore, look for the relationships between the systems as you proceed. For example, a skin lesion or a sore that is not healing may be the first sign of an underlying vascular problem or endocrine problem such as diabetes.

This assessment can be performed at any level of healthcare prevention—primary, secondary, or tertiary. In a primary setting, a complete physical examination is often

Components of Physical Assessment

COMPLETE PHYSICAL ASSESSMENT

General Survey
General observations of appearance and behavior

Measurements
Vital signs: Temperature, pulse, respirations, BP, and pulse oximetry, if available
Height and weight

Complete head-to-toe physical assessment of every system or area

FOCUSED PHYSICAL ASSESSMENT

General Survey
General observations of appearance and behavior as they relate to area of concern

Measurements
Vital signs: Temperature, pulse, respirations, BP, and pulse oximetry, if available
Height and weight if patient's condition permits; if patient's condition does not permit, may need to make an estimate

Head-to-toe scan as it relates to area of concern
Focused physical assessment of area or system of concern

performed to establish or monitor health status. For example, it may be required for school, camp, or a job. In an acute-care setting, a complete physical examination is often performed shortly after admission to establish a baseline and detect any other actual or potential problems. In a long-term-care setting, a complete physical examination is also helpful in establishing a baseline from which the patient's condition can then be monitored and evaluated.

Because a complete physical assessment takes from 30 minutes to an hour, save time by asking some of the history questions (especially the review of systems) as you perform parts of the physical examination. Perform the assessment as soon as possible because the findings help to establish a baseline.

Focused Physical Assessment

The focused physical assessment consists of a general survey, vital sign measurements, and assessment of the specific area or system of concern. It also includes a quick head-to-toe scan of the patient, checking for changes in every system as they relate to the problem at hand. This scan may reveal associated problems and help you determine the severity of the problem. For example, if your patient is having breathing problems, do not limit your focused assessment to the respiratory system alone, because detecting confusion or cyanotic skin color changes may reflect severe hypoxia. The extent of the head-to-toe and focused examinations will depend on your patient's condition and your findings.

A focused physical examination is indicated when your patient's condition is unstable, when time constraints exist, or for episodic follow-up visits. In the last case, you have already performed a complete physical examination, and so now you perform focused physical assessments to monitor or evaluate your patient's health status.

Focused physical assessments also can be performed at any level of healthcare prevention. In a primary setting, they may be used to monitor your patient's health status, for example, performing a breast examination and teaching breast self-examination. In a secondary setting, after you have performed the initial physical assessment, focused assessments are often used to monitor and evaluate your patient's health problem. For example, you might take vital signs and then auscultate the lungs of a patient who was admitted with pneumonia. In a long-term-care setting, a focused assessment is often used to monitor and evaluate your patient's progress. For instance, if your patient is recov-

ering from a total hip replacement, you will probably assess his or her musculoskeletal system, including gait, muscle strength, and prescribed range of motion (ROM) of joints.

Medical Physical Assessment versus Nursing Physical Assessment

The physical assessment techniques used by physicians and nurses are essentially the same, but some critical differences do exist. These differences are defined by the focus and scope of nursing and medical practice. While the techniques are similar, the underlying rationale differs. Remember: Physicians diagnose and treat illness, but nurses diagnose and treat the patient's response to a health problem in an effort to promote her or his health and well-being.

For example, recall the patient in Chapter 2, 81-year-old Mary Johnson, who was admitted to the hospital with a fractured right hip. The physician would perform a physical assessment to confirm the extent of Mrs. Johnson's injury and ensure that she was medically stable for surgery. The nurse would also perform a physical examination, but the nurse's focus would be to identify the extent to which this injury affects Mrs. Johnson's physical, psychological, and social well-being. The nurse would also use physical assessment to work collaboratively with the physician to identify any potential complications that might pose a threat to Mrs. Johnson.

The sharper your assessment skills, the better you will be able to detect subtle changes in your patient. Alerting the physician to these changes may prevent a more serious complication. You will also use your physical assessment skills to monitor Mrs. Johnson's progress through the perioperative phase to rehabilitation and discharge. Along with the history findings, the data you obtain through physical assessment will be used to develop and evaluate a nursing plan of care.

Tools of Physical Assessment

The most important tools that you have for physical assessment are your senses. You will use your *eyes* to inspect, looking for both physical changes and nonverbal clues from your patient. You will use your *ears* to listen, hearing both sounds produced by various body structures and also what your patient is saying. You will use your *nose* to detect any unusual odors that may indicate an underlying problem. You will use your *hands* to feel for physical changes and also to convey a sense of caring to your patient.

You will also use a variety of equipment to perform the physical assessment and enhance your assessment abilities. As with any equipment, assessment equipment, especially equipment that is used for measurement, needs to be periodically checked and calibrated for accuracy.

(Text continued on page 67)

Which Type to Do?

Which type of physical assessment you do depends on:

- Reason for performing the examination.
- Condition of your patient.
- Amount of time you have.

Physical Assessment Equipment

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p data-bbox="129 296 280 323">Thermometer</p> 	<p data-bbox="456 296 858 548">The thermometer measures body temperature. Measurements may be oral, rectal, temporal artery, tympanic, axillary, or skin. A rectal measurement is most reflective of core temperature, whereas skin or surface measurements are the least reflective.</p> <p data-bbox="456 554 863 646">Thermometers measure temperature in either degrees Fahrenheit (°F) or centigrade/Celsius (°C).</p> <p data-bbox="456 653 783 680">Types of thermometers include:</p> <ul style="list-style-type: none"> <li data-bbox="480 686 831 779">■ <i>Glass mercury thermometer:</i> Used for oral, rectal, or axillary temperature measurements. <li data-bbox="480 785 831 877">■ <i>Electronic digital thermometer:</i> Used for oral, rectal, or axillary temperature measurements. <li data-bbox="480 884 863 1005">■ <i>Tympanic thermometer:</i> Uses infrared sensors to sense temperature measurements of the tympanic membrane. <li data-bbox="480 1012 847 1131">■ <i>Temporal artery thermometer:</i> Measures arterial temperature through infrared scanning of the temporal artery. <li data-bbox="480 1138 799 1260">■ <i>Disposable paper strips with temperature-sensitive dots:</i> Used for oral or skin/surface temperature measurements. 	<p data-bbox="903 296 1398 388">Because mercury is very toxic, many healthcare agencies have stopped using glass mercury thermometers.</p> <p data-bbox="903 394 1382 455">Electronic thermometers are much faster than mercury thermometers.</p> <p data-bbox="903 462 1390 554">Disposable probe covers for electronic and tympanic thermometers minimize the risk of cross-contamination.</p> <p data-bbox="903 560 1382 621">When using a tympanic thermometer, pull the helix up and back for the adult patient.</p> <p data-bbox="903 627 1414 749">Although tympanic thermometers are frequently used in children, studies show conflicting results, especially in infants and children under age 6.</p> <p data-bbox="903 756 1406 938">When taking a temporal artery thermometer measurement, gently place the probe flat on the center of the forehead, midway between the eyebrow and the hairline. Lightly slide the thermometer across the forehead to the hairline as you scan the temperature.</p>  <p data-bbox="959 1243 1150 1270">Types of thermometers</p>
<p data-bbox="129 1339 268 1367">Stethoscope</p> 	<p data-bbox="456 1339 858 1751">The stethoscope is used to auscultate sounds produced by various body structures (e.g., breath sounds, bowel sounds, heart sounds, BP, and vascular sounds). It should have the ability to detect both high- and low-pitched sounds and may have a single tube, a double tube, or a double tube sealed as one. The average length of a stethoscope is 22 to 27 inches. The diameter of the chestpiece is approximately 1¾ inches.</p>	<p data-bbox="903 1339 1398 1400">Your ability to auscultate is only as good as the instrument you use.</p> <p data-bbox="903 1407 1366 1434">Always have the earpieces pointing forward.</p> <p data-bbox="903 1440 1390 1467">Double tubing is better for transmitting sound.</p> <p data-bbox="903 1474 1406 1535">The longer or lighter the scope, the less effective it is at transmitting sound.</p> <p data-bbox="903 1541 1398 1602">When using the bell portion of the stethoscope, apply light pressure.</p> <p data-bbox="903 1608 1334 1669">When using the diaphragm portion of the stethoscope, apply firm pressure.</p>

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
	<p>Types of stethoscopes include:</p> <ul style="list-style-type: none"> ■ <i>Combination stethoscope with bell and diaphragm:</i> The bell is the cupped part of the stethoscope. It is best for detecting low-pitched sounds, such as the third and fourth heart sounds, some murmurs, bruits, and Korotkoff sounds. The deeper the cup, the better its ability to detect low-pitched sounds. The diaphragm is the flat part of the stethoscope. It is best for detecting high-pitched sounds, such as breath sounds and some heart and bowel sounds. ■ <i>Pressure-sensitive single-head stethoscope:</i> Also has the ability to detect high- and low-pitched sounds by the amount of pressure applied. For high-pitched sounds, apply firm pressure. For low-pitched sounds, use light pressure. ■ <i>Electronic stethoscope:</i> Also has the ability to detect high- and low-pitched sounds. It also allows you to adjust the volume of the sound. This scope is ideal for anyone with a hearing impairment. ■ <i>Fetoscope:</i> Used to auscultate fetal heart sounds. It is designed to transmit sound through your ears (air conduction) and through your skull (bone conduction), thus increasing your ability to hear the sounds. 	<p>Use a smaller chest piece on infants and small children.</p>  <p>Types of stethoscopes</p>
<p>Doppler</p>	<p>The Doppler is used to detect fetal heart sounds. It is also used to locate pulses, such as the pedal pulses, that may be difficult to palpate.</p>	 <p>Doppler</p>

(continued)

Physical Assessment Equipment (continued)



EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p>Sphygmomanometer</p> 	<p>The sphygmomanometer or BP manometer is used to measure BP. Choose a cuff size according to the circumference of the patient's limb. Cuff width should be approximately 40 percent of the arm circumference, and the cuff's bladder should encircle about 80 percent of the arm.</p> <p>Types of BP manometers include:</p> <ul style="list-style-type: none"> ■ Mercury manometer. ■ Aneroid manometer. ■ Electronic-digital manometer. <p>This equipment may be wall-mounted or portable.</p>	<p>Incorrect cuff size can lead to inaccurate readings. Mercury manometers are more accurate than aneroid types and require less maintenance. But many healthcare facilities are replacing the mercury type with aneroid types because of the toxicity of mercury.</p>  <p>Types of BP manometers</p>
<p>Visual Acuity Charts</p>  <p>Snellen eye chart</p>	<p>Visual acuity charts are used to assess far and near vision. Far vision testing for adults is 20 feet from chart. Far vision for children is 10 feet from chart. Near vision testing is at a distance of about 14 inches. The red and green color bars on the Snellen eye chart can be used to screen for colorblindness.</p> <p>Types of visual acuity charts include:</p> <ul style="list-style-type: none"> ■ <i>Snellen eye chart</i>: The "E" chart is used to test children under age 6 or illiterate or non-English-speaking patients. The letter chart can be used for school-age children and literate adults. ■ <i>Stycar chart</i>: Uses commonly recognizable letters such as X and O to test vision. Used for children over age 2½ and illiterate adults. ■ <i>Allen cards</i>: Pictures of familiar objects, such as a car, house, or horse are used to test vision of children as young as 24 months. ■ <i>Pocket vision screener or printed material</i>: Can be used to test near vision. 	<p>Test each eye separately, then both eyes together. If your patient wears glasses or contact lenses, test him or her both with and without them. No more than two mistakes are allowed when using the Snellen chart.</p> <p>Snellen chart:</p>  <p>Types of equipment to assess the eyes and ears</p>

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p>Penlight or Pocket Light</p> 	<p>The penlight is used to assess eyes and hard-to-see places, such as the mouth, throat, and nose.</p>	<p>Check your batteries often.</p>
<p>Ophthalmoscope</p> 	<p>The ophthalmoscope is used to assess the internal structures of the eye. It contains two wheels—the light wheel and the lens wheel.</p> <p>The light wheel includes:</p> <ul style="list-style-type: none"> ■ Small white light, used for undilated pupil. ■ Large white light, used for dilated pupil. ■ Green light, to filter out the color red. ■ Blue light, used to detect lesions when fluorocine dye is used. ■ Grid, used to locate structures and lesions. ■ Slit of light, used to determine shape of lesions. <p>The lens wheel contains:</p> <ul style="list-style-type: none"> ■ Red or negative numbers that are concave lenses, which focus far. ■ Black or positive numbers that are convex lenses, which focus near. <p>Types of ophthalmoscopes include:</p> <ul style="list-style-type: none"> ■ Battery-operated ophthalmoscope (some with rechargeable batteries). ■ Penlight type or a combination oto/ophthalmoscope with interchangeable heads. ■ Portable or wall-mounted ophthalmoscope. <p>(For more information, see Chapter 12, Assessing the Eye and Ear.)</p>	<p>Always use the ophthalmoscope in a dark room. When assessing the patient's right eye, use your right eye. When assessing the patient's left eye, use your left eye.</p>

(continued)

Physical Assessment Equipment (continued)

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p>Otoscope</p> 	<p>The otoscope is used to illuminate and magnify the external ear canal and tympanic membrane. Disposable specula of 3, 4, or 5 mm and about ½ inch in length are used. When choosing a speculum, use the largest one the patient's ear can accommodate in order to seal the canal. Also use the shortest speculum possible to prevent trauma or discomfort because the inner two-thirds of the ear canal is over the temporal bone and is very sensitive. For an adult, insert the scope about ½ inch; for a child, about ¼ inch. Many otoscopes also have a pneumatic tube attachment to assess the mobility of the tympanic membrane. This is especially valuable when assessing children's ears.</p> <p>Types of otoscopes include:</p> <ul style="list-style-type: none"> ■ Battery-operated otoscope (some with rechargeable batteries). ■ Penlight type or a combination oto/ophthalmoscope with interchangeable heads. ■ Portable or wall-mounted otoscope. <p>(For more information, see Chapter 12, Assessing the Eye and Ear.)</p>	<p>Always palpate the tragus, helix, and mastoid process for tenderness before inserting an otoscope. If they are tender, proceed carefully. For an adult, pull the helix up and back to straighten the canal. For a preschool child, pull the ear lobe down and back to straighten the canal.</p>
<p>Tuning fork</p> 	<p>The tuning fork is used to assess hearing and vibratory sensations.</p> <p>Types of tuning forks include:</p> <ul style="list-style-type: none"> ■ <i>Low-frequency (256-Hz) tuning fork</i>: Best for testing vibratory sensation during the neurological examination. ■ <i>High-frequency (512-Hz) tuning fork</i>: Best for assessing hearing. 	<p>Strike the tuning fork firmly against a hard surface, being careful not to touch the tines—this dampens the vibrations.</p>
<p>Nasoscope</p>	<p>A nasoscope is used to illuminate the nostrils. Types of nasoscopes include:</p> <ul style="list-style-type: none"> ■ A metal nasal speculum attached to a penlight that illuminates and opens the nostrils to allow visualization. ■ An oto/ophthalmoscope with a special nasal tip to illuminate and open the nostrils, allowing better visualization. 	

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p>Transilluminator</p> 	<p>A transilluminator is used to assess for fluid in the sinuses, the fontanelles of the newborn, and the male scrotum. When pressed against a body surface, the light produces a red glow. You can then detect whether the underlying surface contains air, fluid, or tissue.</p> <p>The transilluminator is an attachment that fits on the combination oto/ophthalmoscope with interchangeable heads.</p>	<p>Transillumination should be performed in a dark room.</p> <p>If you do not have a transilluminator, use a penlight instead.</p>
<p>Tape Measure</p> 	<p>A tape measure is used to measure lengths and circumferences of extremities and abdominal girth. In pregnant patients, it is used to measure fundal heights. In newborns, it is used to measure head, chest, and abdominal circumference and length. A stationary wall-mounted tape measure is more accurate for measuring height in children.</p>	<p>Tape measures and rulers are usually calibrated in both centimeters and inches. Centimeters are used more often than inches.</p>
<p>Pocket Ruler</p> 	<p>A pocket ruler is used to measure liver size, respiratory excursion, jugular venous pressure, and any lesion found during the examination.</p>	 <p>Types of measuring equipment</p>
<p>Goniometer</p> 	<p>A goniometer is used during a musculoskeletal examination to assess the range of motion (ROM) of a joint. It is a protractor with a movable arm and a fixed arm (axis). The center (zero point) is placed at the joint with the fixed arm perpendicular to the joint. As the joint moves, the movable arm is moved to measure the joint's ROM in degrees.</p>	

(continued)

Physical Assessment Equipment (continued)

EQUIPMENT

PURPOSE

TYPES, HINTS, AND CAUTIONS

Triceps Skinfold Calipers



Triceps skinfold calipers are used to measure body fat. They measure the thickness of the subcutaneous tissue in millimeters.


Marking Pen



Marking the skin with a marking pen helps ensure accuracy. A marker is commonly used for liver sizes and diaphragmatic excursion measurements. Pulses that are difficult to find may also be marked for subsequent assessments.

Markers should be washable and nontoxic.

Scale



A scale is used to measure weight in pounds and kilograms. Be sure the scale is calibrated and balanced.



Types of scales include:

- *Platform scale with height attachment:* Appropriate for weighing a child or an adult. Height attachment appropriate for adults; measure children with a wall-mounted measure.
- *Portable floor scale:* Appropriate for a child or an adult.
- *Bed scale:* Appropriate for bedridden patients.
- *Smaller platform "baby" scale:* Appropriate for infants. Has curved sides to prevent infant from rolling off.

If obtaining daily weights, weigh patient at the same time with the same scale. Weights should be taken with the patient wearing as little clothing as possible.



Platform scale with height attachment

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p>Wooden Tongue Depressor</p> 	<p>A wooden tongue depressor allows better visualization of the pharynx. It is also useful in assessing the gag reflex, CN V (trigeminal cranial nerve), and the strength of the muscles of mastication.</p>	<p>Break tongue depressors after use.</p>
<p>Cotton Balls</p> 	<p>Cotton balls are used during the neurological examination to assess light touch. A wisp of cotton may also be used to assess the corneal reflex.</p>	 <p>Equipment used for neurological exam</p>
<p>Test Tubes</p>	<p>Test tubes are used during the neurological examination to assess hot and cold sensation.</p>	<p>If test tubes are not readily available, use the cold metal portion of your stethoscope to assess cold sensation.</p>
<p>Coffee</p> 	<p>A substance with a strong scent, such as coffee or vanilla, is needed to assess the sense of smell.</p>	
<p>Sugar and Lemon</p>	<p>Sugar and lemon are needed to assess the sense of taste.</p>	

(continued)

Physical Assessment Equipment (continued)

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p data-bbox="134 296 284 327">Cup of Water</p> 	<p data-bbox="459 296 823 390">A cup of water is needed to assess your patient's swallowing ability and thyroid.</p>	
<p data-bbox="134 726 245 758">Paper Clip</p> 	<p data-bbox="459 726 831 947">A familiar object, such as a paper clip, is needed to assess for stereognosis during the neurological examination. Stereognosis is the ability to identify a familiar object through the sense of touch.</p>	
<p data-bbox="134 1157 245 1188">Safety Pin</p>	<p data-bbox="459 1157 836 1283">A safety pin or other sharp object is used during the neurological examination to assess light touch and pain.</p>	<p data-bbox="903 1157 1422 1283">Discard safety pins or straight pins after use. Be careful not to break the skin. A toothpick can be used in place of a safety pin, minimizing the risk of breaking the skin.</p>
<p data-bbox="134 1335 209 1367">Gloves</p> 	<p data-bbox="459 1335 847 1556">Wear gloves at any point during the assessment when there is a risk of exposure to blood or body fluids, such as examination of the mouth, pelvic structures of a female patient, male reproductive structures, or rectum.</p>	<p data-bbox="903 1335 1406 1367">Wear nonlatex gloves. Change gloves as needed.</p>

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p data-bbox="159 294 335 325">Cytology Brush</p> 	<p data-bbox="494 294 893 483">A cytology brush and scraper and thin Pap test slides are used for specimens obtained during a pelvic examination. If a thin Pap test is not available, use a slide and fixative.</p>	<p data-bbox="925 294 1452 357">Make sure specimens are labeled and sent to the lab promptly.</p> 
<p data-bbox="159 682 255 714">Scraper</p> 		
<p data-bbox="159 1071 383 1102">Thin Pap Test Slides</p> 		
<p data-bbox="159 1459 255 1491">Fixative</p> 		

Equipment used for pelvic exam

(continued)

Physical Assessment Equipment (continued)

EQUIPMENT	PURPOSE	TYPES, HINTS, AND CAUTIONS
<p>Potassium Hydroxide (KOH)</p> 	<p>Slides and KOH may also be needed for specimen examination.</p>	
<p>Lubricant</p> 	<p>A lubricant is needed when performing pelvic and rectal examinations.</p>	<p>Use a water-soluble lubricant.</p>
<p>Speculum</p> 	<p>A vaginal speculum is used to assess the female reproductive system. Specula come in various sizes. After the speculum is inserted into the vaginal canal, it is opened to allow for better visualization of the canal and cervix. Some specula are equipped with a light to illuminate the canal and cervix.</p>	
<p>Hemoccult Test</p>	<p>A hemoccult test is used during a rectal examination to test the stool for blood.</p>	

Techniques of Physical Assessment

The four techniques of physical assessment are **inspection**, **palpation**, **percussion**, and **auscultation**. They are performed in this order, with the exception of the abdominal assessment. In this case, auscultation precedes palpation and percussion so as not to alter the bowel sounds.

Inspection

Always begin with inspection (Fig. 3.1). Inspection is the most frequently used assessment technique, but its value is often overlooked. With inspection, you use not only your sense of sight but also your senses of hearing and smell to inspect your patient critically. Do not rush the process; take your time and really look at your patient. Perform inspection at every encounter with your patient.

Be sure you have adequate lighting, and sufficiently expose the area being assessed. Be systematic in your approach, working from head to toe and noting key landmarks and normal findings. Use your patient as a comparative when possible. Ask yourself, “Does it look the same on the left side as the right?” Look for surface characteristics such as color, size, and shape. Ask yourself, “Are there color changes? Is the patient symmetrical?” Look for gross abnormalities or signs of distress. Do you notice any unusual odors or hear any unusual sounds that warrant further investigation? Always view your findings in light of the patient’s growth and developmental stage



FIGURE 3.1. Inspection.

and cultural background, which may influence your interpretation. For example, an older patient with acute appendicitis may not present with acute abdominal pain as is seen in a younger patient, or a Native American patient with the same problem may not complain of severe pain since pain is something to be endured in his or her culture.

Inspection may be direct or indirect. **Direct inspection** involves directly looking at your patient. **Indirect inspection** involves using equipment to enhance visualization. For example, the oto/ophthalmoscope allows better visualization of the ears and eyes; and specula, such as the nasal speculum and vaginal speculum, open and illuminate, allowing for better visualization.

Palpation

During palpation, you are using your sense of touch to collect data. Palpation is used to assess every system. It usually follows inspection, but both techniques are often performed simultaneously. Palpation allows you to assess surface characteristics, such as texture, consistency, and temperature, and allows you to assess for masses, organs, pulsations, muscle rigidity, and chest excursion. It also lets you differentiate areas of tenderness from areas of pain.



HELPFUL HINT

Tenderness is elicited on palpation; pain is present even without palpation.

Different parts of the hand are best suited for specific purposes (Fig. 3.2). For example, the dorsal aspect of the hand is best for assessing temperature changes (Fig. 3.3); the ball of the hand on the palm and ulnar surface is best for detecting vibration (Fig. 3.4), and the finger pads and tips are the most discriminating for detecting fine sensations, such as pulsations (Fig. 3.5).

Types of Palpation

Two types of palpation may be performed—light and deep. Always begin with light palpation (Fig. 3.6), if for no other reason than to put your patient at ease and convey a message of caring. **Light palpation** is applying very gentle pressure with the tips and pads of your fingers to a body area and then gently moving them over the area, pressing about $\frac{1}{2}$ inch. Light palpation is best for assessing surface characteristics, such as temperature, texture, mobility, shape, and size. It is also useful in assessing pulses, areas of edema, and areas of tenderness. Closing your eyes while palpating may help you concentrate better on what you are feeling.




FIGURE 3.2. Parts of hands used in palpation.

Deep palpation is applying harder pressure with your fingertips or pads over an area to a depth greater than ½ inch. Deep palpation can be single-handed (Fig. 3.7) or bimanual (Fig. 3.8). When using the bimanual technique, feel with your dominant hand. You can place your other hand on top to help control your movements or to stabilize an organ with one hand while you palpate it with the other.

Deep palpation is used to assess organ size, detect masses, and further assess areas of tenderness. To assess for rebound tenderness, press down firmly with your



FIGURE 3.3. Palpating temperature changes with dorsal part of hand.



FIGURE 3.4. Palpating vibratory sensations and tactile fremitus with balls of hands.



FIGURE 3.5. Palpating with fingertips to assess pulsations.

dominant hand and then lift it up quickly. An increase in the patient's pain when you release the pressure signals rebound tenderness. If your patient's muscles tense on palpation (voluntary guarding), have the patient take a few slow, deep breaths or place her or his hands over or under yours during palpation (Fig. 3.9).



HELPFUL HINT

Wash and warm your hands before palpating.

Ballottement

Ballottement (Fig. 3.10) is a palpation technique used to assess a partially free-floating object. Deep palpation is applied in one area, causing the partially attached object to become palpable in another area. It is frequently used to assess the fetus during pregnancy. It is also used to assess for fluid in the suprapatellar pouch, which, if



FIGURE 3.6. Light palpation.



FIGURE 3.7. Deep palpation, single hand.

present, results in a floating kneecap that bounces back on your finger when tapped.



HELPFUL HINT

If you identify an area of tenderness, palpate that area last.

Percussion

Percussion is used to assess density of underlying structures, areas of tenderness, and deep tendon reflexes (DTRs). It entails striking a body surface with quick, light blows and eliciting vibrations and sounds. The sound determines the density of the underlying tissue and whether it is solid tissue or filled with air or fluid. By determining the density, you can also “outline” the underlying structure, allowing you to determine the size and shape of the underlying structure. (See *Percussion Sounds*.)

Two factors influence the sound produced during percussion—the thickness of the surface being percussed and your technique. The more tissue you have to percuss through, the duller the sound. Percussion is a skill that usually requires practice to perfect. You also need to develop skill at identifying and differentiating the percussion sounds. (See *Percussion Sounds*.)

Types of Percussion

Direct or immediate percussion (Fig. 3.11) is directly tapping your hand or fingertip over a body surface to elicit



FIGURE 3.8. Deep palpation, bimanual.



FIGURE 3.9. Palpating hands using patient's hands.



FIGURE 3.11. Direct percussion.

a sound or to assess for an area of tenderness. Direct percussion may be used instead of indirect percussion on an infant's chest. It is also used to assess for sinus tenderness. To perform **indirect** or **mediate percussion** (Fig. 3.12), place your nondominant hand over a body surface, pressing firmly with your middle finger. Then place your dominant hand over it. Flexing the wrist of your dominant hand, tap the middle finger of your nondominant hand with the middle finger of your dominant hand. Do not rest your entire hand on the body surface because this dampens the sound. Keep only your middle finger on the body surface,



FIGURE 3.10. Ballottement.

and hyperextend it as you percuss. Tap lightly and quickly, removing your top finger after each tap. An alternative technique is to flex the fingers of the striking hand and use your proximal interphalangeal joint as the hammer.



HELPFUL HINT

Trim your nails. Percussing with long fingernails may injure your finger or the area being percussed.

Using a percussion hammer (Fig. 3.13) to test reflexes is also a type of indirect percussion. Instead of your finger, you use the hammer to tap a body surface. The purpose of this type of percussion is to elicit DTRs.

Fist or blunt percussion is used to assess for organ tenderness (Fig. 3.14). It may be direct or indirect. **Direct fist percussion** involves striking a body surface with the ulnar surface of your fist. **Indirect fist percussion** involves placing your nondominant hand over the body surface and then striking that hand with the ulnar surface of your other fist.

Percussion Sounds

Percussion produces sounds that vary according to the tissue being percussed. This chart shows important percussion sounds along with their characteristics and typical locations.

SOUND	INTENSITY	PITCH	DURATION	QUALITY	SOURCE
Resonance	Moderate to loud	Low	Long	Hollow	Normal lung
Tympany	Loud	High (musical)	Moderate	Drumlike	Gastric air bubble; intestinal air
Dullness	Soft to moderate	Medium	Moderate	Thudlike	Liver; full bladder; pregnant uterus
Hyperresonance	Very loud	Very low	Very long	Booming	Hyperinflated lung (as in emphysema)
Flatness	Soft	High	Short	Flat	Muscle



FIGURE 3.12. Indirect percussion.



FIGURE 3.14. Percussing for costo-vertebral angle tenderness.



HELPFUL HINT

When performing indirect percussion, be careful not to slap your stationary finger; striking your stationary finger like a hammer will get you the best sound.



HELPFUL HINT

Do not percuss over bone—this produces a dull sound and may be misleading. For example, when percussing the chest, percuss in the intercostal spaces rather than across the ribs.



FIGURE 3.13. Using a percussion hammer.

Auscultation

Auscultation involves using your sense of hearing to collect data. You will listen to sounds produced by the body, such as heart sounds, lung sounds, bowel sounds, and vascular sounds. Auscultation can be both direct and indirect. **Direct auscultation** is listening for sounds without a stethoscope, but only a few sounds can be heard this way. Two examples are respiratory congestion in a patient who requires suctioning and the loud audible murmur of mitral valve replacement. For most of the sounds produced by the body, you will need to perform **indirect auscultation** with a stethoscope (Figs. 3.15 and 3.16). Your ability to hear is affected by the quality of the stethoscope. The stethoscope should have the ability to detect both high- and low-pitched sounds.

Auscultation is also a skill that requires practice. You need to know what constitutes normal sounds before you can begin to identify abnormal sounds. Listen for the characteristics of sound—the pitch, intensity, duration, and quality. Pitch may be high, medium, or low. Ask yourself, “Which part of the stethoscope was I using when I heard the sound best?” Intensity can range from soft to loud or



FIGURE 3.15. Listening with diaphragm of stethoscope.

can be graded on a scale. Ask yourself, “Could I hear the sound easily, or did I have to listen closely?” Duration may be short or long. Ask yourself, “How long was the sound?” Quality describes the sound. Ask yourself, “What did it sound like? Was it harsh, blowing, etc.?”

Practice listening to normal sounds. Be a selective listener and focus only on the sounds that you are auscultating, filtering out other sounds. For example, when auscultating heart sounds, close your eyes and block out background breath sounds so that you can concentrate on heart sounds. Once you become skilled at identifying normal sounds, you will then be able to identify anything that sounds different. You may have difficulty identifying abnormal sounds at first, but with practice, you will be able to differentiate them. If you are not sure of what you are hearing, ask a more experienced nurse to validate your findings. Also, do not rely on just one assessment technique to assess your patient. If you hear something abnormal, look for other data to support or negate that finding.

Besides comparing the sounds you hear with what you know is normal, be sure also to consider your patient’s size. It may affect the sounds you hear. The thicker the chest wall, the softer the sound and the more difficult it is to hear. Use your patient as his or her own comparative. For example, it may be easier to hear breath sounds in a patient who weighs 110 pounds than in one who weighs 220 pounds. But both patients’ breath sounds may be normal.



FIGURE 3.16. Listening with bell of stethoscope.

Auscultation Tips

- Always have earpieces pointing forward to seal the ear canal. Warm your stethoscope.
- Work on the patient’s right side. This stretches your stethoscope across the patient’s chest and minimizes interference.
- Never listen through clothes.
- Make sure that the environment is quiet.
- If hair is a problem, wet it to minimize artifact.
- Use light pressure to detect low-pitched sounds
- Use firm pressure to detect high-pitched sounds.
- Close your eyes to help you focus.
- Learn to become a selective listener.
- Most of all—practice!

Approach to the Physical Assessment

Whether you are doing a complete physical assessment or a focused physical assessment, you need to be systematic in your approach. You can approach the physical examination either by systems or by region. With a systems approach, you perform all assessments related to one system before proceeding to the next system. With a regional approach, you perform a thorough assessment of one region, thus encompassing many systems, before proceeding from head to toe. Each approach is systematic and comprehensive and elicits the same data.

Which approach should you use? That depends on the patient’s health status, the purpose of the assessment, and your own preference. The systems approach is best suited for a focused physical assessment and corresponds with the documentation of findings. The regional approach is best suited for a complete physical assessment because it minimizes patient position changes and requires less time. No matter which approach you use, be consistent. You may find it helpful to develop a written checklist so that you do not miss any areas. If you use the same approach consistently and adjust it to meet your patients’ needs, it will eventually become second nature to you.

As you proceed, remember that every system is related, and every system has an effect on every other. Be systematic. Work from head to toe, and whenever possible, from side to side so that you can use your patient as her or his own comparative. Always consider the developmental stage of your patient, and adjust your approach to meet her or his needs. Also be aware of cultural influences that may affect the assessment and your findings. For example, Hindu and Mexican women are

very modest and may request a nurse of the same gender to perform the physical examination.

Begin by introducing yourself and telling your patient what you are going to do and why. The examination itself may be stressful for your patient, especially if he or she is concerned over health problems that may be validated by the examination. Because patients often equate the length of the examination with the severity of a problem, let your patient know that the examination normally takes time. Also be conscious of your nonverbal behavior; maintain a professional demeanor and caring attitude, and be sensitive to your patient's needs. Explain what you are doing every step of the way, and encourage the patient to ask questions.

Make sure that the examination room is quiet and private and that you will not be interrupted. The room also needs to be warm and well lit. Ask the patient if she or he needs to void before the examination. Provide privacy to allow your patient to change into a gown if needed. While she or he is changing into a gown, assemble all your equipment and make sure that everything is in working order. Designate one area as clean for the unused equipment and one area as dirty for the used equipment. If your patient is uncomfortable removing all of her or his clothing, allow her or him to leave undergarments on and remove them only during the parts of the examination when it is necessary. Wash your hands before you begin, and wear gloves if the possibility of exposure to blood or body fluids exists. Drape your patient. Work from the right side when possible, and expose only the area being assessed.

During the examination, you will use all four techniques of physical assessment: inspection, palpation, percussion, and auscultation. If your patient has identified an area of concern, begin there; otherwise, proceed from head to toe. Usually, the more private areas of the examination, such as the pelvic and rectal examination, are performed last. Do not rush. Pay attention to your patient's responses, both verbal and nonverbal, and respond accordingly. For example, if you see that your patient is tiring, you may need to provide a short rest period. If you have identified any health teaching needs, the physical examination is an ideal opportunity for health teaching.

Patients With Special Needs

Children, pregnant, elderly, or disabled patients need special consideration during physical assessment. The following paragraphs contain some tips for assessing these patients. In each case, remember to incorporate teaching into your assessment.

Children

- *Infants:* Allow parents to be present and to help, if appropriate.
- *Children 1 to 2 years old:* May be fearful, so use games or toys during examination. Leave eye, ear, and mouth assessment until last.
- *Children 2 to 3 years old:* Most difficult to examine. May cling to parents, so let parents help. Allow child to see and touch equipment. Demonstrate use of equipment on doll or parent.
- *Children 4 to 5 years old:* More cooperative; respond well when play is incorporated into examination.
- *School-age children:* Usually cooperative; can converse and follow instructions.
- *Adolescents:* May be sensitive and modest. May not want parent present during examination. Let patient know that it is okay to ask questions.
- *All ages:* Look for normal growth and developmental changes. (For more information, see Chapter 24, Assessing the Newborn and Infant; Chapter 25, Assessing the Toddler and Preschooler; and Chapter 26, Assessing the School-Age Child and Adolescent.)

Pregnant Patients

- Assess both the mother-to-be and the fetus.
- Include fundal heights and fetal heart tones in the examination.
- Assess for normal changes that occur during pregnancy.
- Pay special attention to nutritional assessment.
- Remember that patient may have difficulty changing or assuming positions during last trimester.
- Be aware that hormonal swings may exaggerate patient's responses. (For more information, see Chapter 23, Assessing the Mother-to-Be.)

Elderly Patients

- Do not rush.
- Look for developmental changes.
- Do not assume. For example, your patient may be elderly, but that does not mean he or she is hard of hearing.
- Conserve your patient's energy by minimizing position changes and helping her or him change positions as needed.
- Allow enough time for patient to respond to questions or instructions. (For more information, see Chapter 27, Assessing the Older Adult.)

Disabled Patients

- Identify the disability.
- Focus on the patient's functional ability and mental capacity.

- Modify your assessment based on the patient's assets and needs. For example, if he or she is deaf, you may need to write instructions or have someone available who can sign.
- Be alert and sensitive to your patient's needs, especially if she or he is unable to communicate verbally.



CRITICAL THINKING ACTIVITY #1

In what ways might you adjust your assessment if your patient were blind?

Positions for Physical Assessment

In order to perform a thorough physical assessment, you must ask the patient to assume various positions. The table *Examination Positions* describes the positions needed to assess each area.

Components of the Physical Assessment

Before you focus on a particular system or area, perform a general survey and measure vital signs, height, and weight. During the health history, you performed a review of systems to detect possible problems in other systems. This provided you with a subjective look at the relationship of these systems. Now take an objective look at each system as it affects each other before you focus on a specific system or area.

General Survey

Always begin the physical assessment with the general survey. This “first impression” of your patient begins as soon as you meet him or her. Use your senses and your observational skills to look, listen, and take note of any unusual odors. First, look for the obvious: apparent age, gender, and race. Then look closer for clues that might signal a problem. Also consider your patient's developmental stage and cultural background; these may influence your findings and interpretation. Make a mental note of any clues detected on the general survey that you may need to follow up during the physical examination. When looking for clues, watch for signs of distress and check facial characteristics, body size and type, posture, movements, speech, grooming, dress, and hygiene.

Signs of Distress

Ask yourself, “Are there any obvious signs of distress (e.g., breathing problems)?” If signs of acute distress are appar-

ent, do a focused physical assessment and address the acute problem.



CRITICAL THINKING ACTIVITY #2

What other signs would indicate acute distress?

Facial Characteristics

Ask yourself, “What is the patient's face telling me?” Do you see pain, fear, or anxiety? Does the patient maintain eye contact? Is her or his facial expression happy or sad? Facial expression may signal an underlying problem, such as masklike expression in Parkinson's disease. Look at the patient's facial features. Are they symmetrical? Ptosis of the eyelid may indicate a neurological problem; drooping on one side of the face may indicate a transient ischemic attack (TIA) or stroke; and exophthalmos suggests hyperthyroidism. Also look at the condition of your patient's skin. It may reflect your patient's age, but excessive wrinkling from excessive sun exposure or illness may make the patient appear older than her or his stated age.

Body Type, Posture, and Gait

View your patient's body size and build with respect to his or her age and gender. Is he or she stocky, slender, or of average build? Obese or cachectic? Proportional? Does your patient have abnormal fat distribution, such as the truncal obesity and buffalo hump seen with Cushing's syndrome? Greet your patient with a handshake. The handshake will convey that you care but also allow you to assess muscle strength, hydration, skin temperature, and texture. Then take a close look at your patient's hands. Do you see clubbing, edema, or deformities? Clubbing and edema may reflect a cardiopulmonary problem; deformities may reflect arthritic changes.

Watch how your patient enters the room. Are her or his movements smooth and coordinated, or does she or he have an obvious gait problem? Does she or he walk with an assistive device such as a cane or walker? Do you see a wide base of support with short stride length? Gait problems may suggest a musculoskeletal or neurological problem. Spastic movements or unsteady gaits may be seen in patients with cerebral palsy or multiple sclerosis. A shuffling gait is seen with Parkinson's disease. Patients with a wide base of support and short stride length may have a balance or cerebellar problem.

If your patient is bedridden, observe his or her ability to move from side to side, sit up in bed, and change positions. Determine how much assistance he or she needs with moving. Any problem detected during the general survey should be further evaluated during the physical examination.

(Text continued on page 77)

Examination Positions

POSITION	AREAS ASSESSED	ADVANTAGES/ DISADVANTAGES
<p>Supine</p> 	<ul style="list-style-type: none"> ■ Anterior chest for respiratory, cardiac, and breast exams (should be supine for breast exam). ■ Pulses and extremities. 	<ul style="list-style-type: none"> ■ If patient has trouble breathing in supine position, use semi-Fowler's position (semisitting with knees flexed and supported by pillows).
<p>Sitting</p> 	<ul style="list-style-type: none"> ■ Head and neck. ■ Anterior and posterior chest for respiratory, cardiac, and breast exams. ■ Vital signs and upper extremities. 	<ul style="list-style-type: none"> ■ Provides good visualization. ■ Allows full lung expansion for respiratory assessment. ■ Patients with weakness or paralysis may have difficulty assuming position and need assistance.
<p>Dorsal recumbent</p> 	<ul style="list-style-type: none"> ■ <i>Abdomen</i>: Basically supine position with knees slightly flexed to relax abdominal muscles. ■ Female pelvic area if patient is unable to assume lithotomy or Sims' position. ■ Lithotomy position for female pelvic and rectal areas essentially same as dorsal recumbent but legs and feet in stirrups. 	<ul style="list-style-type: none"> ■ If patient has abdominal pain, flexing knees is usually more comfortable. ■ Older patients may have difficulty assuming lithotomy position.
<p>Sims'</p> 	<ul style="list-style-type: none"> ■ <i>Female pelvic and rectal areas</i>: Best alternative if patient is unable to assume lithotomy position. 	<ul style="list-style-type: none"> ■ May be difficult to assume if patient has arthritis. ■ Contraindicated if patient has had total hip replacement.

(continued)

Examination Positions (continued)

POSITION	AREAS ASSESSED	ADVANTAGES/ DISADVANTAGES
<p>Prone</p> 	<ul style="list-style-type: none"> ■ Musculoskeletal system. 	<ul style="list-style-type: none"> ■ Difficult position for many patients, especially those with respiratory disease.
<p>Left Lateral Recumbent</p> 	<ul style="list-style-type: none"> ■ <i>Chest</i>: Best for cardiac auscultation, particularly of S₃, S₄, and some murmurs. 	<ul style="list-style-type: none"> ■ Patients with respiratory problems may have trouble assuming this position.
<p>Knee-chest</p> 	<ul style="list-style-type: none"> ■ <i>Male rectal and prostate areas</i>: Best position for these exams. 	<ul style="list-style-type: none"> ■ This position and its alternative (bending over a table) are very difficult and embarrassing for most patients.

POSITION	AREAS ASSESSED	ADVANTAGES/ DISADVANTAGES
<p>Standing</p> 	<ul style="list-style-type: none"> ■ <i>Spine and joints (ROM)</i>: Best for these musculoskeletal areas; used for both neurological exam and to assess gait and cerebellar function. 	<ul style="list-style-type: none"> ■ Patients who are weak, disabled, or paralyzed may need assistance or may not be able to assume this position.

The patient's posture may also reveal clues about her or his overall health status. Is she or he sitting upright or slumped? Can she or he assume a supine position? Is there a position that she or he prefers? A slumped position may indicate fatigue or depression. Patients with cardiopulmonary disease often cannot assume a supine position. They prefer a sitting or tripod position because it eases breathing. Patients with abdominal pain also may find it difficult to assume a supine position. Sitting hunched over or in a side-lying position with legs flexed often eases the pain.

Speech

Listen to your patient's speech pattern and pace. Speech reflects the mental state, thought process, and affect. Are your patient's responses appropriate? Speech patterns and appropriateness of responses reflect his or her thought process. Pressured speech, inappropriate responses, and illogical or incoherent speech may be associated with psychiatric disorders. Pressured, hurried speech may also be seen in patients with hyperthyroidism. Note the tone and quality of his or her voice. Do you hear anger? Sadness? Changes in voice quality may indicate a neurological problem, specifically, a problem with cranial nerve (CN) IX

(glossopharyngeal) or CN X (vagus). Notice whether his or her speech is clear or garbled. Aphasia can be expressive (motor/Broca's), receptive (sensory/Wernicke's), or global, a combination of these. Aphasia of any type is often associated with a TIA or stroke.

The patient's vocabulary and sentence structure offer clues to her or his educational level, which you will need to consider when developing teaching plans. Also be alert for foreign accents. You need to determine if there is a language barrier and solicit an interpreter, if needed.

Dress, Grooming, and Hygiene

The way your patient is dressed and groomed tells a great deal about his or her physical and psychological well-being. Is he or she neatly dressed and well groomed or is he or she disheveled? Disinterest in appearance may reflect depression or low self-esteem. Poor hygiene and a disheveled appearance may also reflect the patient's inability to care for himself or herself.

Also take note of the appropriateness of dress for the season and situation. Inappropriate dress may be a sign of hyperthyroidism, which causes heat intolerance. Worn clothing may indicate financial problems. Be sure to take

note of any unusual odors, such as alcohol or urine, that may indicate a problem and warrant further investigation.

Mental State

Determine if your patient is awake, alert, and oriented to time, place, and person. Are her or his responses appropriate? Bizarre responses suggest a psychiatric problem. Is your patient lethargic? Many conditions can affect the level of consciousness. Determine if this is a change in your patient's mental status. If so, investigate further. Also take note of your patient's medications, because they may be contributing to the change in mental status.

Cultural Considerations

Note any cultural influences that may affect your patient's physical characteristics, response to pain, dress, grooming, and hygiene. Patterns of verbal and nonverbal communication may also be culturally influenced. Keeping these factors in mind will help you avoid making hasty, inaccurate interpretations.

Here are a few examples of cultural differences:

- Asian Americans are often shorter than Westerners.
- European Americans and African Americans tend to speak loudly, whereas Chinese Americans speak softly.
- Asians may avoid eye contact with anyone considered a superior.
- Japanese and Korean individuals maintain a tense posture to convey confidence and competence, whereas Americans assume a relaxed posture to convey the same message.

For more information on cultural influences, see Chapter 1, Health Assessment and the Nurse.

Developmental Considerations

Consider the developmental stage of your patient. Here are some points to keep in mind when performing the general survey:

Children

- Behavior should correspond with the child's developmental level.
- Children tend to regress when ill.
- Take note of the relationship between child and parent.

Pregnant Patients

- General appearance should reflect gestational age.
- Look for normal changes that occur with pregnancy, such as wide base of support and lordosis.
- Look for swelling.
- Note patient's affect and response to pregnancy.

Older Adults

- Look for normal changes that occur with aging.

- Look for clues of decreasing ability to function, especially dress and grooming problems.
- Pay attention to your patient's affect, especially signs of depression.
- Note changes in mental status such as confusion, and then consider medications, hydration, and nutritional status or an underlying infection as a possible cause.

Documenting the General Survey

Documentation of the general survey records your first overall impression of your patient. It should include age (actual and apparent), gender, race, level of consciousness, dress, posture, speech, affect, and any obvious abnormalities or signs of distress.

Here is an example of documentation:

Joe Dunleavy, 50-year-old white male, looks younger than stated age; alert and oriented to time, place, and person. Neatly dressed, well groomed. Well developed. Speech clear; responds appropriately, affect pleasant. No signs of acute distress.

Vital Signs

Vital signs include measurements of temperature, pulse, respirations, and blood pressure. They are the most frequent assessments performed, reflecting cardiopulmonary function and the overall functioning of the body.

The purpose of taking vital signs is to:

- Establish a baseline.
- Monitor the patient's condition.
- Evaluate the patient's response to treatments.
- Identify problems.

The initial set of vital signs serves as the baseline. How often you will need to take vital signs depends on the patient's condition. Serial measurements are more reflective of health than one-time measurements. For example, a blood pressure of 150/98 in a patient who was hurrying to be on time for his appointment and was upset because he was late may not be a reflection of his usual blood pressure. Stress can cause abnormal elevations. Also check your equipment periodically to make sure that it is reliable and accurate. Do not rely on one assessment finding—look for other data to support or negate your finding. For example, if your patient's temperature is 102°F, his or her skin should be warm and flushed.

Temperature

Body temperature is the difference between heat produced and heat lost. The hypothalamus acts as the body's thermostat to maintain a constant body temperature. The balance is maintained between the body's heat-producing functions (metabolism, shivering, muscle contrac-



FIGURE 3.17. Taking a tympanic temperature.

tion, exercise, and thyroid activity) and the heat-losing functions (radiation, convection, conduction, and evaporation). When one becomes greater than the other, temperature changes are seen; greater heat-producing functions result in temperature elevations (fever/**hyperthermia**), and greater heat-losing functions result in temperature decreases (**hypothermia**).

Keep in mind that a fever can be a part of the inflammatory response that helps the body fight diseases. Diseases that affect metabolism, such as thyroid disease, can also

affect body temperature. Because the hypothalamus is the body's thermostat, disorders that affect the hypothalamus, such as a stroke, will affect the temperature.

Temperature is measured in either degrees Fahrenheit (°F) or degrees Centigrade, or Celsius (°C). The average normal oral temperature is 98.6°F (37°C). Rectal, temporal artery, and tympanic temperatures (Fig. 3.17) are more reflective of the body's core temperature, with the average temperature being 1 degree higher than oral temperature. Axillary temperature measures skin temperature and is on average 1 degree lower than the oral temperature.

Various factors can influence the normal temperature reading, including:

- **Age:** Newborns are **homeothermic**, stabilizing temperature within a small range, which makes them easily susceptible to temperature variations, especially hypothermia. Children have slightly higher normal temperatures than adults—99°F until age 3. Elderly people have decreased thermoregulation, also making them vulnerable to temperature changes. They tend to have lower-than-normal temperatures—97°F

Advantages and Disadvantages of Temperature Routes

ROUTE	ADVANTAGES	DISADVANTAGES
Oral (Normal: 98.6°F; 37°C)	<ul style="list-style-type: none"> ■ Easy, fast, accurate 	<ul style="list-style-type: none"> ■ Cannot be used for clients who are unconscious, confused, prone to seizures, recovering from oral surgery, or under age 6. ■ Need to wait 15–20 minutes after eating.
Rectal (Normal: 99.5°F; 37.5°C)	<ul style="list-style-type: none"> ■ More reflective of core temperature ■ Fast 	<ul style="list-style-type: none"> ■ Cannot be used for clients who have rectal bleeding, hemorrhoids, or diarrhea or who are recovering from rectal surgery. ■ Contraindicated for cardiac clients because it may stimulate the vagus nerve and decrease heart rate. ■ Not recommended for newborns because of risk of perforating anus.
Tympanic (Normal: 99.5°F; 37.5°C)	<ul style="list-style-type: none"> ■ More reflective of core temperature ■ Safe, good for children 	<ul style="list-style-type: none"> ■ Reports of accuracy are conflicting.
Axillary (Normal: 97.6°F; 36.5°C)	<ul style="list-style-type: none"> ■ Safe, good for children and newborns 	<ul style="list-style-type: none"> ■ Measures skin surface, which can be variable.
Forehead (Normal: 94°F; 34.4°C)	<ul style="list-style-type: none"> ■ Safe and easy 	<ul style="list-style-type: none"> ■ Measures skin surface temperature. ■ Least accurate method.
Temporal arterial (Normal: Close to rectal temperature, 1°F or 0.5°C higher than an oral temperature, and 2°F or 1°C higher than an axillary temperature)	<ul style="list-style-type: none"> ■ Noninvasive, safe, good for children and infants 	

- **Gender:** Women have greater fluctuations in body temperature than men because of hormonal changes. For example, increases in progesterone during ovulation can cause a woman's body temperature to increase by 0.5°F to 1°F.
- **Circadian rhythms (diurnal, sleep/wake cycle):** Body temperature changes at different times of the day. It is usually lowest in the morning and highest in the evening. Temperature fluctuations can range from 1°F to 2°F.



HELPFUL HINT

The conversion from °F to °C is $(^{\circ}\text{F} - 32) \times 5/9 = ^{\circ}\text{C}$.
The conversion from °C to °F is $(^{\circ}\text{C} \times 9/5) + 32 = ^{\circ}\text{F}$.

- **Exercise/stress:** Stress and exercise cause an increase in epinephrine and norepinephrine, increasing the metabolic rate and in turn increasing temperature.
- **Environmental factors:** Exposure to temperature extremes can lead to hypothermia (low temperature from exposure to excessive cold) and hyperthermia (high temperature from exposure to excessive heat).

Pulse

The **pulse** reflects the force of the heart contracting. This force creates a wave of pressure as blood circulates throughout the systemic circulation. The pulse reflects **stroke volume**, the amount of blood ejected with each contraction.



FIGURE 3.18. Taking a radial pulse.

Assess the patient's pulse for rate and rhythm. The radial pulse (Fig. 3.18) is generally used when assessing vital signs. Normal pulse rate varies with age. (See *The Effect of Age on Vital Signs*.) The average rate for a newborn is 125 beats per minute (BPM). By age 4, the rate decreases to about 100 BPM, and by adolescence, the rate is in the normal adult range of 60 to 100 BPM.

Cardiac output is equivalent to stroke volume times heart rate ($\text{CO} = \text{SV} \times \text{HR}$). The fastest way to increase CO is to increase the HR. HRs above 100 BPM are called *tachycardia*; below 60 BPM, *bradycardia*. Athletes' hearts work more efficiently, so they often have HRs lower than 60 BPM. To meet the increased oxygen demands associated with exercise, SV increases over time.

If you have trouble finding your patient's radial pulse, use a larger, more central pulse, such as the carotid or femoral. Whenever you obtain an irregular rhythm, auscultate an apical heart rate (Fig. 3.19) for 1 full minute. Children often have a rhythm that varies with

The Effect of Age on Vital Signs

Normal vital signs vary with age, as this chart shows.

AGE	TEMPERATURE		PULSE RATE	RESPIRATORY RATE	BLOOD PRESSURE (mm Hg)
	°F	°C			
Newborn	98.6–99.8	37–37.7	120–160	30–80	Systolic: 50–52 Diastolic: 25–30 Mean: 35–40
3 yr	98.5–99.5	36.9–37.5	80–125	20–30	Systolic: 78–114 Diastolic: 46–78
10 yr	97.5–98.6	36.3–37	70–110	16–22	Systolic: 90–132 Diastolic: 5–86
16 yr	97.6–98.8	36.4–37.1	55–100	15–20	Systolic: 104–108 Diastolic: 60–92
Adult	96.8–99.5	36–37.5	60–100	12–20	Systolic: <120 Diastolic: <80
Older Adult	96.5–97.5	35.9–36.3	60–100	15–25	Systolic: <120 Diastolic: <80



FIGURE 3.19. Taking an apical pulse.

respirations, causing a sinus arrhythmia. During inspiration, there is an increased venous return to the right side of the heart, causing a slight increase in rate to handle the increased load. The apical pulse is also used in infants and small children. (See Chapter 15, Assessing the Peripheral Vascular and Lymphatic Systems, for a more detailed assessment of the pulses.)

Respirations

Assessing respirations includes checking rate, rhythm, and depth. It includes assessing **inspiration** (taking oxygen into the lungs) and **expiration** (removing carbon dioxide from the lungs). Many factors—including the patient's metabolism and whether the cardiovascular, musculoskeletal, and neurological systems are intact—influence respirations. The **respiratory rate** is a count of one full inspiration/expiration cycle for 1 full minute.

The normal respiratory rate varies with age. The newborn's respiratory rate is quite rapid, averaging about 40 breaths per minute. The respiratory rate gradually decreases with age until it reaches the adult rate of 12 to 20 breaths per minute. Respiratory rates that are within normal range are termed **eupnea**; those above normal range are termed **tachypnea**; and those below normal range are called **bradypnea**. Absent breathing is **apnea**, and difficult breathing is **dyspnea**.

The rhythm of respirations for adults and children is usually regular, with an occasional “sigh,” or deeper breath. Infants normally have irregular respiratory rhythms, but be alert for prolonged periods of apnea. Apnea lasting for more than 15 seconds is cause for alarm and warrants further assessment. Irregular breathing patterns may also indicate an underlying problem such as a neurological disorder.

The depth of respirations reflects the **tidal volume** (the amount of air inhaled with each breath). In adults, the normal tidal volume is about 300 to 500 mL. Ask yourself, “Are the patient's respirations deep or shallow?” Deep and rapid respirations may indicate that your patient is hyperventilating. Conditions such as metabolic acidosis as well as anxiety can cause hyperventilation. Shallow respirations may indicate hypoventilation, which may be caused by conditions such as sedation and guarding from pain.

As you are counting the respirations, take note of signs of distress, such as cyanotic color changes and retraction or use of accessory muscles to breathe. Also listen for audible sounds such as wheezing or congestion that may signal distress and warrant further assessment. (See Chapter 13, Assessing the Respiratory System, for further information.)

Blood Pressure

Blood pressure (BP) is a measurement of the pressure within the vascular system as the heart contracts (systole) and relaxes (diastole). BP indirectly reflects your patient's overall cardiovascular functioning. It is equal to CO times peripheral vascular resistance ($BP = CO \times PVR$). Normal BP varies with age. Other factors that can affect BP include stress, genetics, medications, heavy meals, diurnal variations, exercise, and weight. (See *Measuring Blood Pressure*.)

Normal BP for an adult ranges from 100 to less than 120 mm Hg (systolic) and from 60 to less than 80 mm Hg (diastolic). Normal BP for children and infants is much lower.



ALERT

Because BP reflects the pressure within the vascular system, a person cannot maintain a high BP without serious effects.

A systolic reading 120 to 139 mm Hg and a diastolic reading 80 to 89 mm Hg is considered **prehypertension**; a systolic 140 to 159 mm Hg with a diastolic 90 to 99 mm Hg, **stage 1 hypertension**; a systolic 160 mm Hg or higher with a diastolic 100 mm Hg or higher, **stage 2 hypertension**; and a systolic reading lower than 90 mm Hg and a diastolic reading lower than 60 mm Hg is considered **hypotensive**.

An elevated BP reading should be followed up with a serial reading to see if the BP is consistently elevated. Keep in mind that the systolic pressure changes more readily in response to external stimuli, whereas the diastolic is more constant. Consequently, an elevated diastolic pressure may be more worrisome than an isolated elevation of the systolic pressure.

The **pulse pressure**—the difference between the systolic and the diastolic readings—is normally about 40 mm Hg. A widened pulse pressure is often seen in elderly people because of a decreased elasticity of the arterial walls, but it may also be a sign of increased intracranial pressure. A narrow pulse pressure can be seen when the patient is hypovolemic or going into shock and heart failure.

BP measurements are often taken with the patient in several positions—supine, sitting, and standing.

Measuring Blood Pressure

To assess a patient's BP accurately, the nurse should follow this procedure.

Preparing the Patient

Before beginning, make sure the patient is relaxed and has not eaten or exercised in the past 30 minutes. The patient can sit, stand, or lie down during BP measurement.

Applying the Cuff and Stethoscope

- Wrap the sphygmomanometer cuff snugly around the arm 1 inch (2.5 cm) above the antecubital area (the inner aspect of the elbow), with the cuff bladder centered over the brachial artery (Fig. 3.20).



FIGURE 3.20. Measuring cuff size.



HELPFUL HINT

Most cuffs have arrows that should be placed over the brachial artery.

- To determine the peak inflation point, palpate the brachial pulse just below and slightly medial to the antecubital area to determine the approximate systolic pressure.
- Rapidly inflate the cuff to about 30 mm Hg above the level at which the pulsations disappear.
- Deflate the cuff slowly until you feel the pulse again. Then, deflate the cuff completely.
- Place the earpieces of the stethoscope in your ears and position the stethoscope head over the brachial artery (Fig. 3.21). Although the diaphragm is more frequently used, the bell is best for detecting the low-pitched Korotkoff sounds.



FIGURE 3.21. Taking BP with bell of stethoscope.

- Make sure to use the proper-sized cuff for the patient.
- Keep the manometer at eye level; if using an aneroid gauge, place it level with the patient's arm.
- Keep the patient's arm level with the heart. *Do not* use the patient's muscle strength to hold up the arm; tension from muscle contraction can elevate systolic pressure. Have a recumbent patient rest the arm at the side.

Obtaining the Blood Pressure Reading

- Watching the manometer, pump the bulb until the column or aneroid gauge reaches about 30 mm Hg above the point at which the pulse disappears to be sure to hear phase I and avoid the auscultatory gap. The **auscultatory gap** is a period of silence that may occur between systolic and diastolic pressures and result in a false reading.

- Slowly open the air valve and watch the column drop or the gauge needle descend. Release the pressure at a rate of about 3 mm Hg per second and listen for pulse sounds (Korotkoff sounds).
- As soon as you hear blood begin to pulse through the brachial artery, note the reading on the aneroid dial or column. Reflecting phase I (the first Korotkoff sound), this sound coincides with the patient's systolic pressure.
- Continue deflating the cuff, noting the point at which pulsations diminish or become muffled—phase IV (the fourth Korotkoff sound)—and then disappear—phase V (the fifth Korotkoff sound).
- For children and highly active adults, many authorities consider phase IV the most accurate reflection of BP.

The American Heart Association and the World Health Organization recommend documenting phases I, IV, and V. However, usually only phases I and V are recorded in practice. If the three phases are charted, record phase I, IV, and V:

The Korotkoff sounds, which determine the BP measurement, are classified as follows:

- Phase I: Onset of clear, faint tapping, with intensity that increases to a thud or louder tap.
- Phase II: Tapping that changes to a soft, swishing sound.
- Phase III: Return of clear, crisp tapping sound.
- Phase IV (first diastolic sound): Sound becomes muffled and takes on a blowing quality.
- Phase V (second diastolic sound): Sound disappears.

Measuring BP in several positions allows you to check for **orthostatic hypotension** or **postural hypotension** (a drop in BP with position change). If the patient has orthostatic hypotension, the systolic reading will drop 20 mm Hg, and the pulse rate will increase by about 20 BPM, causing dizziness or lightheadedness. Certain medications and hypovolemia can cause orthostatic hypotension.



HELPFUL HINT

If you are unsure about a BP reading, double-check or have another nurse retake it.

If you are having difficulty hearing the BP, try using an electronic BP device. Or, if electronic equipment is not available, do a palpable BP. Inflate the cuff and palpate the brachial artery. As you deflate the cuff, note at what point the brachial pulse becomes palpable. This number represents the systolic reading. A diastolic reading cannot be obtained using the palpation method. An example of documentation of a palpable BP is “90/palpated.”

BP may also be difficult to hear in infants. If so, use the flush method. As you inflate the cuff and occlude the artery, the extremity will become pale; as you release the pressure, blood flow returns and the extremity will flush. The point at which the extremity flushes coincides approximately with the mean arterial pressure (MAP = diastolic pressure + $\frac{1}{3}$ of the pulse pressure).

Documenting Vital Signs

Vital signs are often documented on graphic charts, which allow you to monitor or plot the patterns of the

vital signs. This is very helpful in evaluating the effectiveness of treatment. Vital signs may also be documented in narrative form. For example:

T 98.8°F; P 88, regular; R 16 unlabored, regular and deep; BP 120/80 lying, 118/80 sitting, and 118/80 standing.

Height and Weight

Anthropometric measurements provide valuable data about your patient's growth and development, nutritional

Blood Pressure Tips

- Too-narrow cuff will cause a false high reading.
- Too-wide cuff will cause a false low reading.
- Cuff wrapped too loosely will cause a false high reading.
- Arm above heart level during reading will cause a false low reading.
- Muscle contracted on arm during BP reading will cause a false high reading.
- Arm pressures should not differ more than 10 mm Hg.
- Leg pressures should be the same or slightly higher than arm pressures.

Which Arm to Use?

You can usually use either arm for BP measurement but you should avoid doing BP on an:

- Arm with injury or cast.
- Arm with IV site.
- Arm of patient who has had a mastectomy.
- Arm that has a vascular access for dialysis.



FIGURE 3.22. Measuring height.

status, and overall general health. Accurate assessment of height (Fig. 3.22) and weight (Fig. 3.23) is also a vital assessment used to determine appropriate medication dosages. This is particularly true with children because medication dosages are calculated based on weight. For adults, weight is used to determine chemotherapy dosages and anesthesia administration. Weight can also be used to evaluate the effectiveness of a treatment, such as intravenous (IV) fluids, drugs, or nutritional therapy. For example, the goal of a patient receiving nutritional therapy might be maintenance or weight gain.



FIGURE 3.23. Measuring weight.

Weight and height for adults who can stand can be obtained using a platform scale with a sliding ruler to measure height. Be sure your scale is balanced. Weigh the patient while she or he is wearing minimal clothing (gown) and no shoes. If the patient cannot stand, use a bed scale. Children may also be weighed on a platform scale, but they should be measured with a wall-mounted measure to ensure accuracy. Weigh infants with a small table platform scale. To measure an infant's length, use a stationary measure or mark disposable examination paper at the top of the head and the heel of the extended foot and then measure the distance between the two. If serial weights are being done, weigh the patient on the same scale at the same time of day. Be sure to document the measurements in centimeters or inches, kilograms or pounds. Because children have frequent changes in growth and development, their measurements are often documented on growth charts for easy monitoring. (See Appendix B.)

Head-to-Toe Scan

Whether you are doing a complete or a focused physical assessment, remember that all body systems are interrelated. A problem in one system will eventually affect or be affected by every other system. When performing a complete physical assessment, perform a thorough-assessment of each system, looking for relationships. When you perform a focused physical assessment—an in-depth assessment of one system—begin by scanning the other systems, looking for changes as they relate to the system being assessed.

For example, suppose you were performing a focused assessment of the cardiovascular system. Here are some of the effects you should look for:

- **Integumentary:** Look for color changes, pallor, edema, and changes in skin and hair texture and growth. Patients with cardiovascular problems may have edema and thin, shiny, hairless skin.
- **Head, eyes, ears, nose, and throat (HEENT):** Look for periorbital edema. Funduscopic examination may reveal vascular changes.
- **Respiratory:** Look for cyanosis and breathing difficulty. Auscultate for adventitious sounds such as crackles that are associated with congestive heart failure (CHF).
- **Breast:** Assess for male breast enlargement that may occur with cardiac medications such as digoxin.
- **Abdominal:** Assess for ascites and hepatomegaly, which may be associated with CHF.
- **Musculoskeletal:** Assess for muscle weakness and atrophy associated with disuse.
- **Neurological:** Assess level of consciousness. Confusion, fatigue, and lethargy may be associated with decreased CO and hypoxia.

After you look for the relationships between the systems, you are ready to perform the focused physical assessment. This text is structured to emphasize the relationship of one system to another. The relationship between the systems is presented for every system assessed.

Documenting Your Physical Assessment Findings

Once you have completed the physical assessment, document your findings. Because writing should be kept to a

minimum during the examination, document your findings as soon as possible, while they are foremost in your mind. Document accurately, objectively, and concisely. Avoid general terms such as “normal.” Record your findings system by system, being sure to chart pertinent negative findings. Once you have completed your objective database, combine it with your subjective database, cluster your data, identify pertinent nursing diagnoses, and develop a plan of care for your patient. For more information on documentation, see Chapter 1, Health Assessment and the Nurse.


S U M M A R Y

- The physical assessment provides objective data for the database. It involves using your senses to collect data. You look, listen, smell, and feel.
- Specific equipment may be used to enhance the data collection process.
- The four techniques of physical assessment are inspection, palpation, percussion, and auscultation.
- The physical examination may be complete (an entire, thorough assessment) or focused (an assessment of one system or area).
- The physical assessment also helps you validate subjective data found in the history, identify any health problems, and evaluate the patient’s response to treatment.
- The physical examination should always begin with a general survey and a measurement of vital signs, height, and weight.
- For a focused physical examination, a head-to-toe scan identifies the system’s effects on every other system.
- Physical assessment requires the use of cognitive, psychomotor, interpersonal, affective, and ethical skills.
- As with any skills—practice. The more you practice, the more competent and confident you will be in performing an accurate physical assessment and in interpreting your findings—findings that are essential to the plan of care for your patient.

R E V I E W Q U E S T I O N S

1. Physical assessments can be either complete or focused. Which situation would warrant a focused physical examination?
 - a. Student physical examination
 - b. Pre-employment physical examination
 - c. Preoperative patient scheduled for hip replacement
 - d. Patient admitted with chest pain
2. You are auscultating your patient’s heart sounds. Which is the correct technique for using the bell portion of the stethoscope?
 - a. Hold bell lightly on chest
 - b. Hold bell firmly on chest
 - c. Tilt bell slightly off chest
3. When assessing the skin, which part of the hand is best for detecting temperature changes?
 - a. Palmar aspect
 - b. Ulnar aspect
 - c. Dorsal aspect
 - d. Finger tips
4. Which part of your hand is best for detecting vibratory sensations?
 - a. Balls of your hand
 - b. Dorsal surface of your hand
 - c. Ulnar surface of your hand
 - d. Palmar surface of your hand
5. Which part of your hand is the most discriminatory, able to detect fine sensations?
 - a. Dorsal surface
 - b. Finger pads
 - c. Ulnar surface
 - d. Palmar surface
6. How would you distinguish pain from tenderness?
 - a. Tenderness occurs with palpation
 - b. Tenderness is constant
 - c. Tenderness occurs with or without palpation
 - d. Tenderness is what the patient tells you it is
7. You are working in the nursery, assessing a newborn’s vital signs. What would you expect the normal heart rate to be?
 - a. 60 BPM
 - b. 80 BPM
 - c. 100 BPM
 - d. 120 BPM
8. The difference between the systolic and the diastolic pressure is referred to as:
 - a. Pulse pressure
 - b. Mean arterial pressure
 - c. Phase IV of Kortokoff
 - d. Pulse deficit

9. When taking your patient's BP, which technique is best for cuff inflation?
 - a. Inflate cuff to 200 mm Hg
 - b. Inflate cuff 30 mm Hg above point palpable pulse disappears
 - c. Inflate cuff to level of the prior reading
 - d. Inflate the cuff until the first sounds are heard
10. You are working night shift; while making rounds you note that your patient's breathing has a period of absent respirations lasting for 15 seconds. This is referred to as:
 - a. Apnea
 - b. Bradypnea
 - c. Eupnea
 - d. Dyspnea



Assessing Pain

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Define the mechanisms of pain
- Describe different types of pain
- Dispel myths associated with pain
- Identify developmental influences on pain
- Identify cultural influences on pain
- Perform pain assessment
- Perform symptom analysis
- Identify components of a pain assessment

Introduction to Assessing Pain

Although **pain** is referred to as the fifth vital sign, pain is a symptom. Subjective in nature, pain is “whatever the person says it is, whenever she or he says it does” (McCaffery, 1999). Pain has also been defined as an “unpleasant sensory and emotional experience associated with actual or potential tissue damage” (International Association for the Study of Pain, 2006). It is the most common reason patients seek healthcare.

Far too often, not only the public but also health-care providers undertreat pain owing to misconceptions surrounding it and its effective management. This is a serious issue, as undertreated pain can have serious physical, psychological, and financial consequences that affect the patient’s quality of life.

Accurate assessment of your patient’s pain is the first step in developing an effective treatment plan to deal with pain. Because pain is subjective, no two patients experience pain the same way. Since pain has physiological, psychological, sociocultural, behavioral, and affective components, assessment must address each component as it relates to your patient.

Structure and Function of Pain

Understanding the process of pain will help you to accurately assess and develop a plan of care to manage your patient’s pain. The process by which a painful stimulus is transmitted to the central nervous system (CNS) and perceived as pain is referred to as **nociception** (Fig. 4.1). The process of pain includes transduction, transmission,




FIGURE 4.1. Pain transmission. After nociceptors are activated, pain is transmitted along A-delta fibers or C-fibers to the dorsal horn of the spinal cord. From there, the pain message is sent to the brain for perception.

perception, and modulation. However, the exact mechanism of pain is unknown, and this process does not explain all types of pain, such as phantom limb. Also, not all nociception results in the perception of pain.

Transduction

Transduction begins with a response to a noxious (painful) stimulus that results in tissue injury. The noxious

Common Misconceptions About Pain

Common misconceptions about pain include:

- Physical or behavioral signs of pain are more reliable indicators of pain than patient self-report.
- Elderly or cognitively impaired patients cannot use pain intensity rating scales.
- Pain does not exist in the absence of physical or behavioral signs or detectable tissue damage.
- Pain without an obvious physical cause, or that is more severe than expected based on findings, is usually psychogenic.
- Comparable stimuli produce the same level of pain in all individuals (i.e., a uniform pain threshold exists).
- Prior experience with pain teaches a person to be more tolerant of pain.
- Analgesics should be withheld until the cause of the pain is established.
- Noncancer pain is not as severe as cancer pain.
- Patients who are knowledgeable about pain medications, are frequent emergency department patrons, or have been taking opioids for a long time are necessarily addicts or “drug seekers.”
- Use of opioids in patients with pain will cause them to become addicted.
- Patients who respond to a placebo drug are malingering.
- Neonates, infants, and young children have decreased pain sensation.

Types of Noxious Stimuli

TYPE	CAUSES	RESPONSE
Mechanical	Tissue trauma, such as surgery, laceration Alteration of tissue, such as edema Obstruction, such as biliary or bowel Abnormal tissue growth, such as tumor Abnormal contraction of muscle, such as spasm	Tissue damage directly stimulates pain receptors and activates inflammatory response. Edema causes pressure on pain receptors. Obstruction causes distension and pressure on pain receptors. Tumors cause irritation and pressure on pain receptors. Spasms stimulate pain receptors.
Thermal	Extreme heat or cold	Extreme temperatures cause tissue damage and activate thermosensitive pain receptors.
Chemical	Tissue ischemia, such as coronary artery disease or muscle spasms	Decreased blood supply leads to tissue ischemia and damage, with lactic acid production and release of tissue byproducts, such as bradykinin, that stimulate pain receptors.

Adapted from Kozier, B., Erb, G., Berman, A., and Snyder, S. (2004). *Fundamentals of Nursing*, ed. 7. Upper Saddle River, NJ: Pearson/Prentice Hall.

stimuli can be mechanical, thermal, or chemical. The noxious stimuli are then converted into a nerve impulse by sensory receptors called nociceptors.

Nociceptors are receptors found in the skin, subcutaneous tissue, joints, walls of arteries, and most internal organs (with the highest concentration found in the skin and the least in internal organs) that respond to painful stimuli. In response to tissue injury, cells break down and release byproducts and inflammation mediators (bradykinin, prostaglandins, substance P, histamine, serotonin, and cytokines). These substances activate or sensitize the nociceptors. Bradykinin also acts as a potent vasodilator, triggering the inflammatory response of redness, swelling, and pain. The inflammatory response is the greatest cause of pain. However, all pain that originates from the periphery is not nociceptive pain. Damaged peripheral nerves can result in **peripheral neuropathic pain**.

Transmission

Once the nociceptors are activated, the nerve impulse is transmitted to the spinal cord and brain. The sensory nerve impulses travel via afferent neurons to the dorsal horn of the spinal cord. **Primary afferent nerve fibers** travel along C-fibers, unmyelinated, slow-conducting fibers that transmit dull, aching pain, or A-delta fibers, myelinated, fast-conducting fibers that transmit sharp, localized pain. **C-fibers** transmit slow, constant pain from mechanical, thermal, and chemical stimuli. **A-delta fibers** transmit fast, intermittent pain from mainly mechanical stimuli.

Excitatory amino acids (e.g., glutamine, aspartate) and **neuropeptides** (e.g., substance P) facilitate transmission of impulse to the dorsal horn at the synapses from the primary afferent neuron to a second-order neuron in the dorsal horn. **Inhibitory amino acids** (e.g., γ -aminobutyric acid [GABA]) and **neuropeptides** (e.g., endogenous opioids) inhibit transmission of impulse. Once in the dorsal horn, the impulse can be transmitted to the brain or inhibited. Neuroregulators, neurotransmitters, or neuromodulators either inhibit or facilitate nociception.

From the dorsal horn, the impulses are then transmitted from (1) the spinothalamic tract to the thalamus, (2) the spinoreticular tract to the reticular formation, (3) the spinomesencephalic tract to the mesencephalon, and (4) the spinothalamic tract to the hypothalamus. Most impulses are transmitted to the thalamus from which they are directed to three areas: the **somatosensory cortex**, which perceives and interprets physical sensation; the **limbic system**, which allows for emotional response to stimuli; and the **frontal cortex**, which involves thought and reason and perception of pain.

Perception

Perception is an awareness of pain and involves both the cortical and the limbic system structures. Realize that **pain threshold**, the point at which a painful stimulus is perceived as painful, is consistent from one person to the

next; however, **pain tolerance**, the amount of pain one is able to endure, varies greatly. Psychosocial and cultural factors and past experiences influence pain perception, thereby accounting for such individual differences in pain perception even with similar painful stimuli.

Modulation

Nerve transmission from the dorsal horn is modulated by descending inhibitory input. Inhibition can also occur at the peripheral, spinal, and supraspinal levels. Inhibition occurs by analgesia or the gate-control theory of pain modulation. Inhibitory substances (e.g., GABA and neuropeptides [endogenous opioids, serotonin, norepinephrine]) bind to primary afferent receptors and dorsal horn neurons to inhibit transmission of impulse. Endogenous opioids include enkephalins, dynorphins, and beta endorphins. Endogenous opioids can also be produced through exercise, meditation, visualization, and music therapy. The brain also sends descending inhibitory input that modulates the transmission of nociceptive transmission in the dorsal horn.

The **gate-control theory of pain modulation** (Fig. 4.2) proposes that impulses can be blocked by non-painful somatic stimuli. Impulses compete for transmission. Since two impulses cannot be transmitted at the same time, one goes through while the other is blocked.

For example, if you stub your toe, you automatically rub your toe. The rubbing sensation is transmitted by the fast, A-delta fibers, while the impulse from stubbing the toe travels along the slow C-fiber and is blocked.

Sensitization

Sensitization to pain can be peripheral or central. **Peripheral sensitization** occurs with prolonged exposure to noxious stimuli. The result is a lower threshold of pain, leading to hyperalgesia (increased response to painful stimuli) and allodynia (painful response to nonpainful stimuli). **Central sensitization** also occurs with prolonged exposure to noxious stimuli with spinal neuron hyperexcitability and results in hyperalgesia and allodynia as well as persistent pain and referred pain. Sensitization can act as a protective mechanism during healing, but when it persists, chronic pain can develop.

Classification of Pain

Pain can be categorized by duration, site of pain, cause, or other qualities.




FIGURE 4.2. The gate-control theory of pain modulation. *A*, Normally, C-fibers carrying slow-pain signals block inhibitory interneurons and transmit their signals across the synapse unimpeded. *B*, A-delta fibers carrying pleasurable signals (e.g., from touch) excite inhibitory interneurons, which then block the transmission of slow-pain signals (+ equals transmission, - equals no transmission).

Types of Pain

TYPE	CAUSE	DURATION	EFFECTS	EXAMPLES
Acute	Injury or pathology Nociception and/or sensitized central neurons	Self-limiting, resolves with healing Continuous or intermittent	Activates autonomic nervous system, such as increased heart rate, respirations, blood pressure, diaphoresis Dilated pupils Protective responses, such as rubbing or guarding area of pain	Trauma Medical or surgical procedures Labor Acute disease
Chronic	May or may not be associated with pathology Weak connection-between cause and extent of pain May be nociceptive, neuropathic, or both	Persists beyond expected healing time Remote from the original cause Continuous, intermittent, with or without acute exacerbation	Parasympathetic response Normal vital signs Dry, warm skin Depressed and withdrawn No protective behaviors	Injury Malignant conditions Chronic non-life-threatening diseases such as arthritis
Cancer/malignant	Associated with underlying malignancy, diagnostic procedure, or disease treatment	May be acute or chronic Pain level strongly correlates with the degree of pathology	Variable depending on location and duration	Malignancy and associated diagnostic tests and treatments
Chronic noncancer pain	Weak connection between cause and extent of pain Associated with chronic disease or no identifiable cause Pain becomes disease	Prolonged Possibly lifelong	Pain, ranging from mild to excruciating and possibly affecting any system or region	Osteoarthritis Low back pain Myofascial pain Fibromyalgia Headaches Central pain Chronic abdominal pain Sickle cell disease Chronic pain syndrome Phantom limb pain Peripheral neuropathy Neuralgias
Chronic pain syndrome	Chronic pain that consumes and incapacitates patient	Prolonged Possibly lifelong	Psychosocial dysfunction, such as anger, depression, anxiety, substance abuse May stress personal and work relationships Life centers on obtaining pain relief	

Acute Pain

Pain is defined as **acute** when it lasts the expected recovery time. Acute pain serves as a protective mechanism in response to an actual or potential threat to injury, such as when you withdraw your hand from a hot surface. Injury also activates the “stress hormone response,” which, if left untreated, can produce negative physiological and psychological effects resulting in chronic pain. Acute pain is seen with injury and surgery and may last up to 6 months.

Chronic Pain

Chronic pain is defined as pain enduring for 6 months or longer. More specifically, chronic pain endures beyond expected recovery time. Chronic pain may be in response to a progressive illness or result from no apparent injury, as with neuropathic pain. Chronic pain often serves no adaptive purpose, and frequently results in depression. Chronic pain can be further classified as cancer/malignant pain or chronic noncancer pain.

Cancer/Malignant Pain

Pain associated with cancer may be categorized separately. **Cancer** or **malignant pain** may be acute pain or chronic pain that is associated with an underlying malignancy, diagnostic procedure, or disease treatment. The level of pain strongly correlates with the degree of pathology. **Intractable pain**, pain resistant to treatment, is often seen with advanced metastatic disease.

Chronic Noncancer Pain

Persistent pain not associated with malignancy may be categorized as **chronic noncancer pain**, a subtype of chronic pain. Pain levels and pathology have a weak link and may have no discernable cause. The pain becomes the disease. Any body system or region can be affected, with intensity ranging from mild to severe. Chronic noncancer pain has a major effect on every aspect of the patient’s daily life and is referred to as *chronic pain syndrome*. In this syndrome, the patient can no longer function and her or his entire life is centered on finding pain relief.




FIGURE 4.3. Most common areas of referred pain.

Visceral pain results from overdistension, spasms, ischemia, inflammation, or traction of organs, such as colic, appendicitis, peptic ulcer disease, and bladder distension. The pain can be localized or diffuse or referred. **Referred pain** is felt at a site other than at the site of origin. Referred pain occurs when internal organs and structures share nerve pathways within the CNS (Fig. 4.3). Pain can also radiate. **Radiating pain** begins in one area and extends to others, such as chest pain associated with myocardial infarction may be in the chest and radiate to the jaw or arm.

The quality of visceral pain is deep aching or sharp stabbing. Visceral pain is often accompanied by nausea, vomiting, malaise, sweating, tenderness, and muscle spasm.

Nociceptive/Neuropathic Pain

Pain can also be classified by underlying pathology.

Nociceptive Pain

Nociceptive pain results from exposure to noxious (painful) stimuli. The painful stimuli can occur in the viscera, resulting in **visceral pain**, or tissue, resulting in **somatic pain**.

Examples of Referred Pain

Origin of Pain	Site of Referred Pain
Appendicitis	Umbilical region
Angina pectoris	Arm, jaw
Aortic aneurysm	Back
Pleuritis	Shoulder
Cholecystitis	Right shoulder/scapular area
Gastroesophageal reflux	Chest
Urinary tract infection	Back, abdomen

Somatic pain can be superficial or deep. **Superficial somatic pain** originates in the skin or mucous membranes from external stimuli such as sunburn; chemical and thermal burns; or injury to the skin, such as lacerations or contusions. The pain is very localized and described as sharp, pricking, or burning. Cutaneous tenderness, hyperalgesia (increased sensitivity to pain), hyperesthesia (increased sensitivity to sensory stimuli), and allodynia (painful response to nonpainful stimulus) may be associated with superficial somatic pain.

Deep somatic pain originates in muscles, bones, and joints from overuse, injury ischemia, cramping, or inflammation such as arthritis, tendinitis, and sprains. The pain can be localized or diffuse and radiating and described as dull, achy, or cramping. Tenderness and muscle spasm with an autonomic response (increase pulse, blood pressure, and respiratory rate; sweating; pallor; dilated pupils; nausea; vomiting; dry mouth; and increased muscle tension) are often associated with deep somatic pain.

Neuropathic Pain

Neuropathic pain results from injury to the peripheral or central nervous system. Neuropathic pain serves no adaptive purpose and therefore is “pathological” pain. Neuropathic pain can be classified as mono/polyneuropathies, deafferentation, sympathetically maintained pain, and central pain (Clark, 2004).

Mono/Polyneuropathies

Mono/polyneuropathies involve pain along one or more damaged peripheral nerves. Neuropathies may be caused by metabolic disorders (diabetic neuropathy), toxins (alcoholic neuropathy or chemotherapy), infections (human immunodeficiency virus [HIV], postherpetic neuralgia), trauma, compression (compartment syndrome, carpal tunnel syndrome), and autoimmune and hereditary diseases. The pain associated with neuropathies may be described as continuous, deep, burning, aching or bruised, paroxysmal shocklike, or abnormal skin sensitivity.

Deafferentation

Deafferentation pain occurs with loss of afferent input from damage to a peripheral nerve, ganglion, or plexus, or the CNS. Phantom limb and postmastectomy pain are examples of deafferentation pain. The pain is often described as burning, cramping, crushing, aching, stabbing, or shooting. The patient may experience hyperalgesia (excessive sensitivity to pain), hyperpathia (hypersensitivity to sensory stimuli), dysesthesia (abnormal sensations of the skin such as numbness, tingling, burning, or cutting), or other abnormal sensations. **Phantom limb pain**, pain that is experienced in a missing limb or

body part, is the result of sensory representations of the missing limb that are still present in the brain.

Sympathetically Maintained Pain

Sympathetically maintained pain is a pain mechanism that results from sympathetic nervous stimulation. Peripheral nerve damage, sympathetic efferent innervation, or circulating catecholamines can result in sympathetically maintained pain. Chronic pain syndrome, phantom limb pain, postherpetic neuralgia, and some metabolic neuropathies can trigger sympathetically maintained pain.

Central Pain

Central pain results from a primary lesion or dysfunction of the CNS. Ischemia (stroke), tumors, trauma, syrinx (a pathological cavity in the spinal cord or brain), or demyelination (multiple sclerosis) can cause central pain. The quality of the pain varies from a burning, numbing, tingling to a shooting sensation. Onset may be spontaneous, steady, or reactive with positive or negative sensory loss, allodynia, and hyperalgesia (Clark, 2004).

Developmental, Cultural, and Ethnic Variations

A common misconception is that infants, young children, and older adults or cognitively impaired adults do not experience pain. Just because the patient cannot verbally describe pain, that does not negate its presence. Although self-report is the most reliable indicator of pain, if this is not possible, physiological and behavioral indicators are used to assess pain. The challenge then is to accurately assess for pain in all patients at any age and to develop an effective treatment plan.

Infants

At one time, it was believed that infants did not experience pain, but research has refuted that idea. The process of pain is intact even though the emotional response is unclear. Since the infant cannot verbalize pain, physiological and behavioral indicators are used to assess for pain in the infant.



HELPFUL HINT

A good rule of thumb to remember when assessing for pain in an infant who is nonverbal is “whatever is painful for you, would most likely be painful for the infant/child” (Wong & Hockenberry-Eaton, 2001).

Infant physiological changes associated with pain include (Wong & Hockenberry-Eaton, 2001):

- **Integumentary system:** Pallor or flushing, diaphoresis, palmar sweating.
- **Cardiovascular:** Increased heart rate, increased blood pressure.
- **Respiratory:** Rapid, shallow respirations, decreased arterial oxygen saturation, and transcutaneous oxygen saturation.
- **Musculoskeletal:** Increased muscle tone.
- **Neurological:** Increased intracranial pressure, dilated pupils, decreased vagal nerve tone.
- **Endocrine (hormonal release):** Increased catecholamines, growth hormones, glucagon, cortisol, corticosteroids, and aldosterone.
- **Metabolism:** Increased plasma lactate, pyruvate, ketone bodies, and fatty acids.
- **Laboratory values:** Increased blood glucose (hyperglycemia) and corticosteroid levels and decreased pH.

Infant behavioral changes associated with pain include:

- **Vocalization:** Intense, sustained crying, whimpering, and groaning.
- **Facial expression:** Eye squeeze, brow bulge, open mouth, taut tongue, chin quivering, and grimaces.
- **Body movements:** Limb withdraw, thrashing, rigidity or flaccidity, and fist clenching.
- **Sleep/wake cycle:** Increased wakefulness and irritability.
- **Feeding:** Loss of appetite, vomiting, loss of interest and/or energy in sucking.
- **Activity level:** Decreased activity level; fussiness, irritability, and listlessness.

Children

Depending on the age of the child, assessing pain in children can be a challenge. The mnemonic QUESTT is helpful in assessing pain in children (Wong & Hockenberry-Eaton, 2001). QUESTT stands for:

- Question the child.
- Use a pain rating scale.
- Evaluate behavioral and physiological changes.
- Secure parents' involvement.
- Take the cause of pain into account.
- Take action and evaluate results.

Question the Child

Self-report is the most accurate means for assessing pain, so ask the child to describe the pain. Self-report is more accurate than the child's behavior.



HELPFUL HINT

When questioning a child, these techniques are often helpful:

- Use language the child will understand, such as “boo-boo.”
- Use toys to enable the child to identify painful area by asking the child to show you where it hurts on a doll or puppet, having the child draw a picture of pain, or identifying the painful area with crayon on a drawing of human figure.

Use a Pain Rating Scale

Various pain rating scales are available for assessing pain in children. Be sure to select one that is age appropriate for your patient. These scales are discussed later in this chapter.

Evaluate Behavioral and Physiological Changes

If the child is nonverbal and unable to describe pain, detecting behavioral and/or physiological changes is essential. Physiological changes that may be seen with acute pain include increase in blood pressure, increased or decreased heart rate, increased respirations, flushing, sweating, dilated pupils. Behavioral changes can vary depending on the child's developmental level. Behavioral changes for a young child include crying, screaming, vocalizing hurt, thrashing of arms and legs, pushing away, clinging to parents, and increasing restlessness and irritability. Behavioral changes for a school-age child include crying, muscle rigidity, clenched fists, white knuckles, clenched teeth, closed eyes, and stalling techniques when anticipating a painful procedure. Behavioral changes for an adolescent include verbalization of pain and muscle tension.

Secure Parents' Involvement

Parents know their child, so rely on their assessment. Parents are more attuned to subtle changes in their child's behavior. Parents also usually know what will best comfort their child. Ask the parent:

- What is the child's past experience with pain?
- What is the child's response to pain?
- How do you know your child is having pain?
- What relieves the child's pain?

Take the Cause of Pain Into Account

Consider the pathophysiology of the underlying problem when you evaluate the child's pain.

Take Action and Evaluate Results

After assessing the pain, develop a plan to treat the pain. After treating pain, you need to assess the child to evaluate

Common Misconceptions of Pain With the Elderly

Older Patients' Misconceptions of Pain

Pain is unavoidable.
 Pain is a punishment.
 Pain medication is addictive.
 Pain is a sign of weakness.
 The nurse is too busy to get pain medication.
 Pain medication may cause confusion.
 The negative side effects of pain medication outweigh their positive effects.
 Pain medications are expensive and unaffordable.
 Pain is a sign that an underlying illness is worsening.

Healthcare Providers' Misconceptions of Pain in the Elderly

Confused elderly patients don't feel pain.
 Elderly patients have decreased pain sensation.
 Pain medication is addictive.
 Pain may hasten death.
 The older the patient, the more she or he will complain about pain.
 Elderly patients who are sleeping don't have pain.
 Elderly patients have poor tolerance for pain medications.
 Pain causes respiratory depression in elderly patients.
 Elderly patients experience more side effects from pain medications than do younger patients.

Adapted from Victor, K. (2001). Properly assessing pain in the elderly. *RN* 64(5):45–49.

the effectiveness of the treatment and revise the plan as needed.

Older Adults

Assessing pain in the older patient can be challenging because of the misconceptions both patients and healthcare providers have surrounding pain and the elderly.

It is important to realize that the elderly do experience pain both acute and chronic. Untreated pain increases the risk for complications such as pneumonia, constipation, deep vein thrombosis, impaired immune function, sleep disturbances, weight loss, social isolation, and depression. Often, the older patient experiences chronic pain and does not exhibit the typical signs of pain. Answers to questions about the patient's functional level and quality of life are good indicators of the effects of pain.

If the patient is cognitively impaired, assessing for pain becomes a greater challenge. The degree of impairment will determine the use of pain scales. Keep questions simple, specific, and in the here and now. If the patient is unable to verbally communicate, rely on physiological signs associated with pain, such as increased blood pressure, heart rate, and respirations; diaphoresis; and behavioral changes, such as agitation, restlessness,

facial expression of pain, and vocal sounds (moaning and groaning) of pain.

Family members or caregivers should be included in the assessment process. They can provide invaluable data of the patient's pain-related behavior and effective methods for relieving pain.

People of Different Cultures and Ethnic Groups

Although the physiological response to pain is consistent across cultures, your patient's cultural or ethnic background influences his or her psychological and behavioral responses to pain. (See *Cultural and Ethnic Variations in Response to Pain*.)

Introducing the Case Study



Mary Kane, age 74, has degenerative joint disease of the right hip. She is scheduled for a total right hip replacement. You are performing her preoperative assessment. Pain control would be a priority goal postoperatively; therefore, a thorough pain assessment is crucial to ensure effective pain management postoperatively.

Cultural and Ethnic Variations in Response to Pain

CULTURAL GROUP	RESPONSE TO PAIN
African American	<ul style="list-style-type: none"> ■ Pain is a sign of illness or disease. ■ Absence of pain may affect compliance with treatment (i.e., may not take medication as prescribed unless pain present). ■ Pain is seen as inevitable and must be endured. ■ Spiritual and religious beliefs account for high tolerance to pain. ■ Persistent pain associated with little faith. ■ Prayer and laying on of hands used to treat pain.
Appalachian	<ul style="list-style-type: none"> ■ Pain is endured and stoically accepted. ■ Patient expects to be cared for when experiencing pain. ■ Believes use of objects can rid pain (e.g., knife or ax under the bed to “cut out” pain).
Arab American	<ul style="list-style-type: none"> ■ Pain seen as unpleasant and something to be controlled. ■ Expression of pain more reserved with strangers while more expressive with family, sends conflicting perceptions of pain. ■ Confidence in Western medicine, expects immediate relief of pain.
Chinese American	<ul style="list-style-type: none"> ■ Description of pain more diverse in terms of body symptoms such as pain more dull and diffuse. ■ Explain pain from the traditional influences of imbalances in yin and yang. ■ Cope with pain using oils, massage, warmth, sleeping on affected area, relaxation, and aspirin.
Cuban American	<ul style="list-style-type: none"> ■ Pain sign of physical problem that warrants traditional or biomedical healer. ■ Response to pain very expressive (i.e., crying, moaning, groaning) and seen as pain-relieving function.
Egyptian American	<ul style="list-style-type: none"> ■ Avoids pain and seeks prompt treatment. ■ Expression of pain more reserved with strangers while more expressive with family, sends conflicting perceptions of pain. ■ Description of pain more generalized. ■ Age and birth order correlates with individual responses and description of pain (e.g., younger children and first-born more expressive about pain). ■ Helpful to have close family member present during pain episodes, preferably woman, who is seen as more nurturing, caring, and capable of comforting patient in pain.
Filipino American	<ul style="list-style-type: none"> ■ Pain seen as part of living honorable life. ■ Pain seen as means to atone for past transgressions and achieve fuller spiritual life. ■ Response to pain stoic and tolerant of high degree of pain. ■ May need to encourage use of pain interventions. ■ May use prayer in pain management.
French Canadian	<ul style="list-style-type: none"> ■ Pain described as more intense and more affectively described.
Greek American	<ul style="list-style-type: none"> ■ Pain, “ponos,” cardinal symptom of ill health. ■ Pain considered evil and needs to be eradicated. ■ Family relied on to find resources to relieve pain. ■ Physical pain expressed publicly, but emotional pain kept within privacy of family.
Iranian	<ul style="list-style-type: none"> ■ Expressive about pain. ■ Men more stoic than women in expressing pain. ■ May justify pain in light of later rewards after death.
Irish American	<ul style="list-style-type: none"> ■ Stoic response to pain; may ignore or minimize pain. ■ Denial of pain may delay treatment.

CULTURAL GROUP	RESPONSE TO PAIN
Jewish American	<ul style="list-style-type: none"> ■ Preservation of life paramount; therefore, would seek immediate treatment for pain. ■ Verbalization of pain is acceptable and common. ■ Not only want relief from pain but also want to know cause.
Mexican American	<ul style="list-style-type: none"> ■ Good health associated with being pain free. ■ Pain seen and accepted as necessary part of life. ■ Obligated to endure pain in performance of duties. ■ Type and amount of pain divinely predetermined. ■ Pain and suffering seen as consequence for immoral behavior. ■ Methods used to relieve pain maintain balance within person and environment. ■ Perceptions of pain may delay seeking treatment.
Navajo Indian	<ul style="list-style-type: none"> ■ Pain viewed as something to be endured, leading to inadequate pain control. ■ May use herbal medications to treat pain.
Vietnamese American	<ul style="list-style-type: none"> ■ Pain is endured, seen as part of life. ■ Cultural restraints against showing weakness limits use of pain medication.

Source: Purnell, L. & Paulanka, B. (2003). *Transcultural Health Care: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.

Performing the Pain Assessment

Pain assessment includes a history and physical examination. As with any assessment, always begin with a history. Since pain is subjective in nature, the history component of the pain assessment is the most important.

Health History

Self-report is the most accurate indicator of pain. Because pain is subjective, the patient's health history provides the best assessment of pain. The history also allows you to assess past experiences with pain, effective pain treatments, and the effects pain has on every aspect of the patient's life.

Biographical Data

Review your patient's biographical data. As described previously, your patient's age, ethnicity, and religion may affect her or his perception and behavioral response to pain. Even the patient's gender may influence her or his perception of pain. Also, your patient's occupation may be a direct cause of the pain, such as back pain related to heavy lifting; the presence of pain may prohibit a return to work.

Current Health Status

When your patient presents with pain, perform a symptom analysis. The mnemonic PQRST provides a thorough description of pain. Ask:

- **Precipitating/Palliative/Provocative Factors**
 - What were you doing when the pain started?
 - Does anything make it better, such as medication or a certain position?
 - Does anything make it worse, such as movement or breathing?
- **Quality/Quantity**
 - What does it feel like?
 - *Superficial somatic pain* is sharp, pricking, or burning.
 - *Deep somatic pain* is dull or aching.
 - *Visceral pain* is dull, aching, or cramping.
 - *Neuropathic pain* is burning, shocklike, lancinating, jabbing, squeezing, or aching.
 - How often are you experiencing it?
 - To what degree is the pain affecting your ability to perform your usual daily activities?
- **Region/Radiation/Related Symptoms**
 - Can you point to where it hurts?
 - Does the pain occur or spread anywhere else?
 - *Localized pain* is confined to the site of origin, such as cutaneous pain.
 - *Referred pain* is referred to a distant structure, such as shoulder pain with acute cholecystitis or jaw pain associated with angina.
 - *Projected (transmitted) pain* is transmitted along a nerve, such as with herpes zoster or trigeminal neuralgia.
 - *Dermatomal pattern* as with peripheral neuropathic pain.

- *Nondermatomal pattern* as with central neuro-pathic pain, fibromyalgia.
- No recognizable pattern as with complex regional pain syndrome.
- Do you have any other symptoms? (e.g., nausea, dizziness, shortness of breath)
 - *Visceral pain-related symptoms* include sickening feeling, nausea, vomiting, and autonomic symptoms.
 - *Neuropathic pain-related symptoms* include hyperalgesia and allodynia.
 - *Complex regional pain syndrome-related symptoms* include hyperalgesia, hyperesthesia, allodynia, autonomic changes, and shin, hair, and nail changes.
- **Severity**
 - Use appropriate pain scale. (See Pain Scales, below.)
- **Timing**
 - When did the pain begin?
 - How long did it last?
 - *Brief flash*: Quick pain as with needle stick.
 - *Rhythmic pulsation*: Pulsating pain as with migraine or toothache.
 - *Long-duration rhythmic*: As with intestinal colic.
 - *Plateau pain*: Pain that rises then plateaus such as angina.
 - *Paroxysmal*: Such as neuropathic pain.
 - How often does it occur?
 - *Continuous fluctuating pain*: As with musculoskeletal pain.
 - Do you have times when you are pain free?

When taking a patient’s current health status, pay attention to the words that the patient uses to describe her or his current pain level. They can provide important clues to her or his status.

Common Pain Descriptors

TERM	SENSORY WORDS	AFFECTIVE WORDS
Pain	<ul style="list-style-type: none"> ■ Burning ■ Crushing ■ Drilling ■ Penetrating ■ Piercing ■ Scalding ■ Searing ■ Sharp ■ Shooting ■ Wrenching 	<ul style="list-style-type: none"> ■ Agonizing ■ Exhausting ■ Frightful ■ Intense ■ Killing ■ Miserable ■ Punishing ■ Suffocating ■ Terrifying ■ Torturing ■ Unbearable
Hurt	<ul style="list-style-type: none"> ■ Hurting ■ Pressing ■ Pricking ■ Tender 	<ul style="list-style-type: none"> ■ Heavy ■ Throbbing
Ache	<ul style="list-style-type: none"> ■ Aching ■ Cold ■ Cramping ■ Dull ■ Flickering ■ Numb ■ Radiating ■ Sore 	<ul style="list-style-type: none"> ■ Annoying ■ Gnawing ■ Nagging ■ Sickening ■ Tender ■ Tiring ■ Troublesome ■ Uncomfortable

Source: Kozier, B., Erb, G., Berman, A., and Snyder, A. (2004). *Fundamentals of Nursing*, ed. 7. Upper Saddle River, NJ: Pearson/Prentice Hall.

Pain Scales

Various instruments are available to assess pain. Consider the patient’s age and developmental status along with his or her cultural background when selecting a pain scale. Select the one that will best meet your patient’s needs. Instruments may also be specified by your institution.

OLDCART

- Another mnemonic used to assess pain is OLDCART.
- Onset: When did the pain begin?
 - Location: Where does it hurt? Can you point to where it hurts?
 - Duration: How long does it last?
 - Characteristics: What does it feel like?
 - Aggravating factors: Does anything make it worse?
 - Radiation: Does the pain go anywhere else?
 - Treatment: Did anything make it better? (Pain medication, ice, heat?)

Source: Kozier, B., Erb, G., Berman, A., and Snyder, A. (2004). *Fundamentals of Nursing*, ed. 7. Upper Saddle River, NJ: Pearson/Prentice Hall; and LaDuke, S. (2002), Undertreated pain. *Nursing* 32:18.

Pain Scales for Adults

Unidimensional and multidimensional pain scales are available. The unidimensional scale assesses one dimension, usually intensity of pain, and is often used to assess acute pain. Multidimensional scales provide additional information about pain, such as the pain's characteristics and the effects on the patient's daily life. A multidimensional scale is useful in assessing chronic pain.

Unidimensional Scales


These scales generally use numeric, verbal, or visual descriptors to quantify pain. Examples include the Numeric Rating Scale, Visual Analogue Scale, and categorical scales.

Numeric Rating Scale

The Numeric Rating Scale rates pain on a scale of 0 (no pain) to either 5 or 10 (worst pain) by asking the patient to rate her or his current pain level.

Visual Analogue Scale

The Visual Analogue Scale utilizes a vertical or horizontal 10-cm line with anchors. One end of the line is labeled "No pain" and the opposite end of the line is labeled "Worst pain." The patient marks his or her current pain level on the line.



HELPFUL HINTS

The line can be either vertical or horizontal.

Categorical Scales

Categorical scales use verbal or visual descriptors to identify pain intensity. The patient selects the descriptor that she or he feels best represents the current pain level. Verbal descriptors include:

- Mild, discomforting, distressing, horrible, excruciating.
- No pain, mild pain, moderate pain, severe pain, very severe pain, worst possible pain.

Visual descriptors include the Faces Pain Scale for Adults and Children (FPS), which utilizes illustrated faces with facial expressions ranging from happy (no pain) to sad and crying (worst pain). The FPS has eight faces to select current pain level. The patient is asked to select the face that best represents his or her current pain level.

Multidimensional Pain Scales

These scales assess pain characteristics and its effects on patient's activities of daily living and include such scales as the Initial Pain Assessment Inventory (IPAI), Brief Pain Inventory (BPI), McGill Pain Questionnaire (MPQ), and the Neuropathic Pain Scale.

Initial Pain Assessment Inventory

The IPAI is used for initial assessment of pain. It assesses characteristics of pain; effects of pain on the patient's life, such as daily activities, sleep, appetite, relationships, and emotions; and the patient's expression of pain. This assessment tool includes a diagram to note pain location, a scale to rate pain intensity, and space to document additional comments and the treatment plan.

Brief Pain Inventory

The BPI is used to quantify pain intensity and associated disability. It assesses pain intensity, location, effects on life, type, and effectiveness of treatment over the last 24 hours. Benefits of the BPI include that it is quick and easy to use and available in multiple languages.

McGill Pain Questionnaire

The MPQ uses descriptive words to assess pain on three levels: sensory, affective, and evaluative. It can be used with other tools and is available in short and long forms.

Neuropathic Pain Scale

The Neuropathic Pain Scale assesses the type and degree of sensations associated with neuropathic pain. The patient rates eight common qualities of neuropathic pain (sharp, dull, hot, cold, sensitive, itchy, deep, or surface pain) on a scale of 0 (no pain) to 10 (worst pain). This scale is still in the developmental stages, but early testing holds diagnostic and therapeutic promise.

Pain Scales for Children

Many pain scales are available to assess a child's pain. Many of the pain scales are age specific. Instruments may also be specified by your institution since pain assessment forms are common use.

FACES Pain Rating Scale

The FACES Pain Rating Scale assesses pain for children ages 3 years and up. The Wong-Baker has five faces from which the child can select her or his current pain level (Fig. 4.4).

Oucher

The Oucher scale assesses pain for children ages 3 to 13 years with photos or a numeric scale. The photographic scale uses six photographs of children ranging from a child with "no hurt" to a child with "a lot of hurt." The photographs are arranged vertically from 0 to 5, with 0

Wong-Baker FACES Pain Rating Scale

Explain to the person that each face is for a person who feels happy because he has no pain (hurt) or sad because he has some or a lot of pain. Face 0 is very happy because he doesn't hurt at all. Face 2 hurts a little more. Face 3 hurts even more. Face 4 hurts a whole lot. Face 5 hurts as much as you can imagine, although you do not have to be crying to feel this bad. Ask the person to choose the face that best describes how he is feeling. Rating scale is recommended for persons age 3 and older.

FIGURE 4.4. Wong-Baker FACES Pain Rating Scale.

(no hurt) on the bottom and 5 (lot of hurt) on the top. This scale also has photographs of black and Hispanic children available.

Numeric Scale

The numeric scale ranges vertically from 0 to 100, with 0 being “no hurt” and 100 being “biggest hurt” (Beyer, Denyes, & Villaruel, 1992):


- 0 = no hurt.
- 1–29 = little hurt.
- 30–69 = middle hurt.
- 70–99 = big hurt.
- 100 = biggest hurt.

Poker Chip Tool

The Poker Chip Tool assesses pain in children 4 years of age and up. The nurse places red poker chips horizontally in front of the child, with the poker chips denoting “pieces of hurt.” She then asks the child to select how many pieces of hurt he or she has (Hester et al., 1998).


Word-Graphic Rating Scale

The Word-Graphic Rating Scale assesses pain in children ages 4 to 17 years. It uses words on a horizontal linear scale to assess pain. The child is asked to identify her or his current pain level on the scale (Tesler et al., 1991).

**Numeric Scale**

The Numeric Scale assesses pain for children ages 5 years and older. It uses a horizontal linear scale with numbers from 0 to 5 or 10, with 0 being “no pain” and 5 or 10 being “worst pain.” The child is asked to identify his or her current pain level on the scale. Although similar to a

scale used for adults, this provides the child with a visual to help assess his or her pain.

**Visual Analogue Scale**

The Visual Analogue Scale, which assesses pain in children ages 4½ and older, is similar to that used for adults. The child is asked to identify her or his pain level by marking the line in the area that represents her or his level of pain (Cline et al., 1992).

Color Tool

The Color Tool assesses pain for children as young as 4 years by having the child create a body outline using colored markers or crayons. The child selects four colors. The first color represents “most hurt,” the second represents “little hurt,” the third represents “least hurt,” and the last represents “no hurt.” Using all four colors, the child identifies areas and degree of hurt on the body outline (Eland & Banner, 1999).

Past Health History

The past health history can identify factors that may affect the patient’s pain, response to pain, and treatment plan. When conducting the past health history, make sure to identify:

- Cause of pain.
- Past and present medical problems that may influence pain and its management.
- Past psychiatric illnesses and chemical dependence.
- Past and present pain management strategies.
- Past experiences with pain.
- Chronic vs. acute pain.

Family History

The family history may identify genetically linked causes of pain, such as sickle cell anemia and cancer. The family history also identifies familial history of chronic pain or illness. Ask the patient about hereditary or familial health problems.

Review of Systems

If there is a problem in one system, eventually other systems will be affected. The underlying cause of the pain determines the effect the pain will have on other systems.

Psychosocial Profile

Assessing the psychosocial history identifies the effects that pain has on every aspect of the patient’s life and evaluates quality of life.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Health Practices and Beliefs

What are your goals for treatment?

Influences response to pain and expectations for treatment.

Typical Day

What is your typical day like?

How does the pain affect your day?

May affect ability to get through typical day or perform activities of daily living.

Nutritional Pattern

How is your appetite?

Do you have any nausea associated with pain?

Any changes in your weight?

May affect appetite, causing anorexia.

May have associated symptoms of nausea and vomiting.

May have weight loss.

Activity and Exercise Pattern

What is your activity level?

Has the pain affected your activity level or ability to exercise?

Does activity or exercise make the pain worse/better?

May decrease activity level or create an inability to exercise.

Activity or exercise might increase pain.

Recreation, Pets, Hobbies

Do you have any hobbies?

Has the pain affected your ability to perform in recreational activities or hobbies?

Do these activities make the pain better/worse?

May affect ability to participate in recreational activities.

Recreational activities, pets, and hobbies may act as distraction and relaxation in dealing with chronic pain.

Sleep/Rest Patterns

How is your sleep?

Does the pain interrupt your sleep?

May disrupt sleep pattern, resulting in decreased or interrupted sleep.

Personal Habits

Do you use drugs, alcohol, or nicotine to control pain?

May resort to use of drugs, alcohol, nicotine to control pain.

Occupational Health Pattern

Has pain affected your ability to work?

Was the pain the result of a work-related injury?

Occupation can be cause of pain owing to work-related injury.

May prevent return to occupation.

Socioeconomic Status

Are you able to fulfill your treatment plan, prescriptions?

May place financial strains owing to cost of treatment and lost wages (financial strains may be great if pain is chronic).

Environmental Health Patterns

Do you have any difficulty getting around in your home?

Is there a need for assistive devices?

Environmental changes in the home may be needed to adapt.

Role, Relationships, Self-concept

How has the pain affected your mood?

Uncontrolled pain can lead to anxiety, fear, anger, depression.

Cultural Influences

Are there any cultural practices that you use to treat pain?

Affect response to pain.

Influence treatment modalities.

Religious/Spiritual Influences

Do you use any religious practices to help you cope?

Affect meaning of pain.

Affect response to pain.

May be source of coping (e.g., prayer).

Family Roles and Relationships

How has your pain affected your family and relationships?

Affect role in family.

Inability to fulfill role functions.

Changes in role.

Affect roles of other family members.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Sexuality Patterns Has the pain affected your sexual ability?	May affect libido. May affect ability to perform sexually.
Social Supports Who would you consider your supports, family, friends?	Sexual activity may increase pain. Associated depression may cause patient to withdraw and isolate from others.
Stress and Coping Are there any recent stresses in your life?	Increases stress. Identify coping strategies for pain (e.g., distraction).



CRITICAL THINKING ACTIVITY #1

What history question would you want to ask Mrs. Kane prior to surgery that will help manage her postoperative pain?



CRITICAL THINKING ACTIVITY #2

Because Mrs. Kane has degenerative joint disease (DJD) and is scheduled for a total hip replacement, would it be possible for her to experience both acute and chronic pain? Why?



CRITICAL THINKING ACTIVITY #3

Select a pain scale that would be appropriate for use postoperatively to assess Mrs. Kane's pain.

Physical Assessment

The initial purpose of the physical examination is to identify the underlying cause of pain. As you perform the examination, assess the effects of physical movements (e.g., deep breathing, position changes) on the patient's pain level. The physical examination may also reveal complications associated with untreated pain. The history findings should direct the physical exam. Be alert for physical, behavioral, and psychological responses to pain.

The autonomic response of the sympathetic nervous system to painful stimuli accounts for many of the physiological changes that are seen. This response is acute and cannot be sustained for prolonged periods of time, as the body adapts and physical indicators are no longer apparent in patients with prolonged pain or chronic pain. Again, physical signs are not as reliable of an indicator as self-report, the most reliable indicator of pain. However, when patients are unable to verbally

communicate or who are cognitively impaired, you must rely on physical and behavioral indicators obtained by a head-to-toe assessment to evaluate your patient's pain. (See *Performing a Head-to-Toe Physical Assessment*.)

Behavioral Pain Assessment Scales

Although the patient's self-report of pain is the most accurate means to assess pain, there are times when the patient's age or condition does not permit self-report, such as when the patient is confused or very young. Pain scales have been developed to objectively assess pain by scoring behavioral and physiological responses to pain. A variety of objective pain scales are available. The scales are mainly geared for the infant and young child. You need to consider the age of your patient in selecting a scale that will best assess your patient's pain. Also, your institution may prescribe a particular scale as best for assessing patient pain.

Behavioral Pain Assessment Scales for Infants

Since an infant cannot verbalize feelings of pain, physiological and behavioral changes need to be used to assess pain. Many scales are available to help you assess your patient's pain. Identify the scale that is most appropriate in assessing your patient.

CRIES

CRIES (crying, requiring increased oxygen, increased vital signs, expression, sleeplessness) assesses postoperative pain from 32 weeks' gestation to 20 weeks' post-term. Each of the five categories is scored from 0 to 2. A score of 10 equates to the worst pain, and any score greater than 4 identifies significant pain (Krechel & Blidner, 1995). (See *CRIES*.)

Postoperative Pain Score

The Postoperative Pain Scale (POPS) assesses postoperative pain in infants ages 1 to 7 months. It scores each of its 10 categories (sleep, facial expression, quality of

Responses to Pain

A person's response to pain includes physiological (involuntary), behavioral (voluntary), and psychological (affective) responses.

Physiological

Sympathetic Response (Acute)

- Increased blood pressure
- Increased heart rate and force of contraction
- Increased respiratory rate
- Dilated cerebral blood vessels with increased alertness
- Dilated pupils
- Rapid speech

Parasympathetic Response (Deep or Prolonged Pain)

- Decreased systolic blood pressure with possible syncope
- Decreased pulse
- Variable breathing pattern
- Withdrawal
- Constricted pupils
- Slow, monotonous speech

Behavioral Responses

- Withdrawing from painful stimuli
- Facial grimacing
- Crying
- Agitation
- Guarding painful area

Psychological Responses

- Anxiety
- Depression
- Anger
- Fear
- Exhaustion
- Hopelessness
- Irritability

cry, spontaneous motor activity, spontaneous excitability, flexion of fingers and toes, tone, consolability, and sociability) on a scale of 0 to 2. A total score of 0 indicates worst pain, a score of 20, no pain (Attia, 1987).

Neonatal Infant Pain Scale

The Neonatal Infant Pain Scale (NIPS) is used to assess infants at an average gestational age of 33.5 weeks. It grades six categories: facial expression (0 to 1), cry (0 to 2), breathing pattern (0 to 1), arms (0 to 1), legs (0 to 1), and state of arousal (0 to 1). A final score of 0 identifies no pain while a score of 7 identifies worst pain (Lawrence et al., 1993).

Pain Assessment Tool

The Pain Assessment Tool (PAT) assesses pain from a gestational age of 27 weeks to full term. It grades 10 different categories: posture/tone (1 to 2), sleep pattern (0 to 2), expression (1 to 2), color (0 to 2), cry (0 to 2), respirations (1 to 2), heart rate (1 to 2), saturations (0 to 2), blood pressure (0 to 2), and nurse's perception (0 to 2). A final score of 4 identifies no pain, while a score of 20 identifies worst pain (Hodgkinson et al., 1994).

Pain Rating Scale

The Pain Rating Scale (PRS) assesses pain for infants ages 1 to 36 months. The infant is given a grade of 0 (no pain)

Performing a Head-to-Toe Physical Assessment

SYSTEM/AREA	ABNORMAL FINDINGS/RATIONALE
General Appearance Facial Expression Gross Abnormalities	<ul style="list-style-type: none"> ■ Abnormal body posture, splinting, guarding ■ Grimacing, frowning, crying ■ Weight loss, muscle atrophy, deformities ■ Change in vital signs, fever, increased heart and respiratory rate, shock due to altered release of multiple hormones (e.g., adrenocorticotropic hormone [ACTH], cortisol, catecholamines, insulin)
Integumentary Head, eyes, ears, nose, and throat (HEENT)	<ul style="list-style-type: none"> ■ <i>Pallor, diaphoresis</i>: Possible signs of shock ■ Pained facial expression ■ Clenched teeth ■ Abnormal sounds, such as moans and groans ■ Dilated pupils
Respiratory	<ul style="list-style-type: none"> ■ Increased respiratory rate ■ <i>Atelectasis and pneumonia</i>: Decreased air flow due to involuntary (reflex muscle spasm) and voluntary (splinting) mechanisms that limit respiratory effort
Cardiovascular	<ul style="list-style-type: none"> ■ Increased heart rate ■ Increased blood pressure ■ <i>Unstable angina (chest pain) or myocardial infarction (heart attack)</i>: Increased vascular resistance and increased myocardial oxygen demand ■ <i>Deep vein thrombosis (blood clot)</i>: Hypercoagulation
Gastrointestinal	<ul style="list-style-type: none"> ■ <i>Hypoactive bowel sounds</i>: Delayed gastric emptying and peristalsis resulting in constipation, anorexia, ileus
Genitourinary	<ul style="list-style-type: none"> ■ <i>Decreased urine output</i>: Abnormal release of hormones that affect urine output, fluid volume, and electrolyte balance ■ Hypertension (fluid retention) ■ Electrolyte disturbances
Musculoskeletal	<ul style="list-style-type: none"> ■ Limited mobility ■ Involuntary, reflexive movements ■ Muscle tensing/spasm ■ Muscle weakness and fatigue
Neurological	<ul style="list-style-type: none"> ■ Altered affect, depression ■ Screening neurological exam revealing sensory deficits, motor abnormalities, coordination problems, signs of sympathetic responses, and orientation and memory deficits
Immune Other regions	<ul style="list-style-type: none"> ■ <i>Infection</i>: Impaired immune function ■ Findings dependent on site of pain

through 5 (worst pain) based on the following guidelines. For a score of 0, the infant exhibits smiling, sleeping, and no change when moved or touched. Behaviors associated with a score of 1 include taking small amounts orally, restlessness, moving, and crying. A score of 2 includes behaviors of not eating or drinking and short periods of crying but distracted with rocking or use of pacifier.

With a score of 3, a definite change in behavior is noted. The baby is irritable with facial grimacing and arms and/or legs shake or have jerking movements. With a score of 4, the baby is inconsolable, flailing with a high-pitched wailing cry. A score of 5 is characterized by prolonged sleep periods interrupted by jerking movements, continuous crying, and shallow respirations (Joyce et al., 1994).

CRIS

	0	1	2
Crying	No	High pitched	Inconsolable
Requires oxygen	No	<30%	>30%
Increase in vital signs	Heart rate and blood pressure less than or equal to preoperative state	Heart rate and blood pressure increase less than 20% of preoperative state	Heart rate and blood pressure increase greater than 20% of preoperative state
Expression	None	Grimace	Grimace/grunt
Sleepless	No	Wakes at frequent intervals	Constantly awake

Source: Krechel, S., MD, and Bildner, J., RNC, CNS, 1995, from University of Missouri-Columbia, with permission.

Premature Infant Pain Profile (PIPP)

The Premature Infant Pain Profile (PIPP) assesses pain for gestational ages 28 to 40 weeks. It grades seven categories on a scale of 0 (no pain) to 3 (worst pain). The categories are gestational age, behavioral stage, heart rate, oxygen saturation, brow bulge, eye squeeze, and nasolabial furrow. A final score of 0 identifies no pain, while a score of 21 identifies worst pain (Stevens, 1996).

Modified Behavioral Pain Scale

The Modified Behavioral Pain Scale (MBPS) assesses pain for children ages 4 to 6 months. It assesses three categories: facial expression (0 to 3), cry (0 to 4), and movements (0 to 3). A final score of 0 identifies no pain, while a score of 10 identifies worst pain (Taddio et al., 1995).

Behavioral Pain Assessment Scales for Children

Several behavioral pain assessment scales have been developed to assist in assessing pain not only for the infant but also for the older child. Brief descriptions are provided below. Again, you need to select the instrument most appropriate to meet your patient's needs.

Objective Pain Score

The Objective Pain Scale (OPS) assesses pain for infants and children from ages 4 months to 18 years. It assesses five categories, each on a scale of 0 (no pain) to 2 (worst pain). The categories are blood pressure, crying, moving, agitation, and verbal evaluation/body language. A final score of 0 identifies no pain, and a score of 10 identifies worst pain (Hannallah et al., 1987).

Riley Infant Pain Scale

SCORE	FACIAL EXPRESSION	SLEEP	MOVEMENTS	CRY	TOUCH
0	Neutral Smiling, calm	Sleeping quietly	Moves easily	None	
1	Frowning Grimaces	Restless	Restless body movements	Whimpering	Winces with touch
2	Clenched teeth	Intermittent	Moderate agitation	Crying	Cries with touch Difficult to console
3	Crying expression	Prolonged with periods of jerking or no sleep	Thrashing flailing	Screaming, highpitched	Screams when touched Inconsolable

Source: Schade, J.G., et al. (1996). Comparison of three preverbal scales for post-operative pain assessment in a diverse pediatric sample. *Journal of Pain Symptom Management* 12(6):348-359.

Children’s Hospital of Eastern Ontario Pain Scale

The Children’s Hospital of Eastern Ontario Pain Scale (CHEOPS) assesses pain for children ages 1 to 5 years. It assesses six categories: crying (1 to 3), facial (0 to 2), child verbal (0 to 2), torso (1 to 2), touch (1 to 2), and legs (1 to 2). A final score of 4 identifies no pain, while a score of 13 identifies worst pain (McGrath et al., 1985).

Nurses Assessment of Pain Inventory

The Nurses Assessment of Pain Inventory (NAPI) assesses pain for infants and children from birth to age 16 years. It assesses three categories: body movement (0 to 2), facial (0 to 3), and touching (0 to 2). A final score of 0 identifies no pain, while the higher score identifies the worst pain (Stevens, 1990).

Behavioral Pain Score

The Behavioral Pain Score (BPS) assesses pain for children ages 3 to 36 months. It assesses three categories: facial expression (0 to 2), cry (0 to 3), and movements (0 to 3). A final score of 0 identifies no pain, while a score of 8 identifies worst pain (Robieux et al., 1991).

Riley Infant Pain Scale

The Riley Infant Pain Scale (RIPS) assesses pain in children younger than 36 months and children with cerebral palsy. It assesses five categories on a scale of 0 (no pain) to 3 (worst pain). (See *Riley Infant Pain Scale*.)

FLACC Postoperative Pain Tool

The FLACC Postoperative Pain Tool assesses pain for children ages 2 months to 7 years. It assesses five categories

SCORE	FACE	LEGS	ACTIVITY	CRY	CONSOLABILITY
0	Neutral or smile	Relaxed, normal	Moves easily	None	Content
1	Grimace or frown, withdrawn	Restless Tense	Squirming	Whimpers or moans, occasional complaint	Reassured with touching, can be distracted
2	Constant frown, clenched jaw, quivering chin	Flexed Kicking	Arched, rigid, jerking	Crying, screams, verbal complaints	Difficult to console

Source: Adapted from Wong, D., and Hockenberry-Eaton, M. (2001). *Wong’s Essentials of Pediatric Nursing*, ed. 6. Philadelphia: Mosby.

Pain Assessment in Advanced Dementia Scale

ITEM	0	1	2
Breathing, independent of vocalization	Normal	Occasional labored breathing, short periods of hyperventilation	Noisy, labored breathing; long period of hyperventilation; Cheyne-Stokes respirations
Negative vocalization	None	Occasional moan and groan; low-level speech with a negative or disapproving quality	Repeated, troubled calling out; loud moaning and groaning; crying
Facial expression	Smiling or inexpressive	Sad, frightened, frowning	Facial grimacing
Body language	Relaxed	Tense, distressed, pacing, fidgeting	Rigid, fists clenched, knees pulled up, pulling or pushing away, striking out
Consolability	No need to console	Distracted or reassured by voice or touch	Unable to be consoled, distracted, or reassured

Source: Lane, P. (2004). Assessing pain in patients with advanced dementia. *Nursing*, 34(8):17.



CRITICAL THINKING ACTIVITY #4

Mrs. Kane is in the postanesthetic care unit (PACU) post-op total hip replacement. Patient-controlled analgesia (PCA) with morphine sulfate has been ordered for pain control. What physiological and behavioral indicators would indicate postoperative pain?

using a scale of 0 (no pain) to 2 (worst pain). A final score of 0 indicates no pain, while a score of 10 indicates worst pain.

Behavioral Pain Assessment Scales for Older Adults

Since a confused patient may not be able to verbalize feelings of pain, physiological and behavioral changes need to be used to assess pain.

Pain Assessment in Advanced Dementia Scale

The Pain Assessment in Advanced Dementia Scale (PAINAD) can be used to assess pain levels in patients with advanced dementia. It assesses five categories on a scale of 0 to 2. The higher the score, the greater the pain.

Reassessment of Pain

Reassessment of pain is imperative to determine the effectiveness of treatment. Self-report and physical find-



Case Study Analysis

Identify actual or potential nursing diagnosis with supporting data.



CRITICAL THINKING ACTIVITY #5

How often would you reassess Mrs. Kane for pain?

ings will help you evaluate the effectiveness of treatment. Compare the patient's responses to the expected outcomes. Have the patient grade his or her current pain level and compare the grade with the previous rating. If relief has not been obtained, revise your plan and implement alternative interventions. Current recommendations for pain reassessment include:

- Within 30 minutes after parenteral administration of pain medication.
- Within 1 hour after oral administration of pain medication.
- After each and every report of new or changes in pain.

The frequency of reassessment depends on the patient, type of pain, and setting. In acute care settings, reassessment of pain frequently occurs when vital signs are obtained. Outpatients or long-term healthcare residents should be instructed to report any changes in pain or ineffective pain control. Patients with chronic pain should have periodic pain reassessments to ensure effective pain control.

RESEARCH TELLS US

Owing to the subjective nature of pain, assessing an infant's pain can be a challenge. Reyes' study (2003) of infant pain compared 24 neonatal intensive care unit (NICU) nurses' beliefs about infant pain and actual assessment of pain as documented on 107 infant records. Fifty percent to 90% of the nurses surveyed acknowledged that an infant's pain is difficult to recognize, individualized, and that nurses need to use established assessment tools to assess pain and document findings.

However, a retrospective review of infants' records from the day shift revealed that only 62% reflected

pain assessment, while 56.2% of records from the evening shift had no record of pain assessment. The study also revealed that less than 10% of the infants undergoing invasive procedures, such as arterial puncture or peripherally inserted central catheter (PICC) line insertion, had pain assessments.

These findings support the need to better educate nurses on infant pain assessment and support the recommendation of the Joint Commission on Healthcare Organization Pain Standard and the American Academy of Pediatrics to standardize infant pain assessment.


Source: Beal, J., and Freda, M. (2004). Nursing assessment of infant pain. *The American Journal of Maternal/Child Nursing* 29(5):330; and Reyes, S. (2003). Nursing assessment of infant pain. *Journal of Perinatal and Neonatal Nursing*, 17(4):291-303.

S U M M A R Y

- Pain is referred to as the fifth vital sign. Subjective in nature, self-report is the most accurate assessment indicator of pain.
- Pain assessment includes a detailed history and symptom analysis and physical examination. The history is the most important piece of the assessment.
- Realize that pain threshold, the point at which a painful stimulus is perceived as painful, is consistent from one person to the next; however, pain tolerance, the amount of pain one is able to endure, varies greatly.
- Developmental, psychosocial, and cultural factors and past experiences influence pain perception, thereby accounting for such individual differences in pain perception even with similar painful stimuli.
- The initial purpose of the physical examination is to identify the underlying cause of pain. The physical examination also identifies physical, behavioral, and psychological nonverbal responses to pain.
- Although self-report is the most accurate assessment tool of pain, the age and mental status of the patient may prevent self-report. Assessment then becomes dependent on assessing physical, behavioral, and psychological nonverbal responses to pain.
- Various pain scales are available to assess pain. Select the one best suited to meet your patient's needs.
- Be sure to document your findings, evaluate the effects of treatment, and revise the plan of care as needed to ensure pain relief for your patient.

R E V I E W Q U E S T I O N S


1. Patrick Connors, age 57, has prostate cancer that has metastasized to the bone. He is complaining of severe back pain. What is the best means for assessing his pain?
 - a. Ask what he is feeling
 - b. Assess for changes in his vital signs
 - c. Assess for behavioral changes
2. Which question would best assess the quality of Mr. Connors pain?
 - a. Is the pain sharp?
 - b. What does the pain feel like?
 - c. Does anything make the pain worse?
 - d. When did the pain first start?
3. Which question would best assess the region of Mr. Connors pain?
 - a. Tell me where the pain is?
 - b. Is the pain in your back?
 - c. Can you point to where it hurts?
 - d. Does the pain go anywhere else?
4. Which question would best assess the severity of Mr. Connors pain?
 - a. Is the pain really bad?
 - b. How bad is the pain?
 - c. Does anything make your pain better?
 - d. On a scale of 0 to 10, with 10 being the worst, how would you grade your pain?
5. Which question would best assess the timing of Mr. Connors pain?
 - a. When did the pain begin?
 - b. Did anything make the pain go away?
 - c. Did you awaken with pain?
 - d. Does the pain interfere with your ability to work?
6. Which question would best assess precipitating factors of Mr. Connors pain?
 - a. Did the pain awaken you from sleep?
 - b. Did the pain interfere with your ability to work?
 - c. When did the pain begin?
 - d. What were you doing when the pain started?
7. How would you differentiate acute pain from chronic pain?
 - a. Chronic pain persists for up to 1 month
 - b. Chronic pain persists for 1 to 3 months
 - c. Chronic pain persists for 3 to 6 months
8. While working in the chemistry lab, Joe Matthews spilled a caustic agent on his arm. Which pain description would most likely be associated with chemical burns?
 - a. Burning
 - b. Stabbing
 - c. Deep
 - d. Cramping
9. Joan Bock, age 80, has DJD of the knees and hips. Which pain description would most likely be associated with DJD?
 - a. Sharp
 - b. Stabbing
 - c. Aching
 - d. Burning
10. David Fowler, age 9, is admitted to the hospital with the diagnosis of acute appendicitis. Which pain description would most likely be associated with appendicitis?
 - a. Stabbing
 - b. Cramping
 - c. Burning
 - d. Pricking



UNIT

2

Promoting Health



Approach to the Mental Health Assessment

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Describe the various theories in understanding mental health/illness
- Describe developmental influences on mental health/illness
- Discuss various cultural influences on mental health/illness
- Identify history data as they relate to mental health/illness
- Identify physical assessment data as they relate to mental health/illness
- Describe additional assessment components specific to mental health/illness assessment
- Perform a mental health assessment
- Document assessment findings
- Identify actual/potential health problems stated as nursing diagnoses with supporting data

Mental health is a “state of successful performance of mental function, resulting in productive activities, fulfilling relationships with other people, and the ability to adapt to change and to cope with adversity” (Healthy People 2010). Jahoda (1958) identified six characteristics of positive mental health that address positive self-perception, personality integration, autonomy and independence, perception and reality, and growth and development leading to self-actualization. Mental health or wellness affects every aspect of one’s life, and the impact of this can be seen when mental illness occurs. The effects are not only mental but also physical.

Understanding Mental Disorders

Mental disorders affect thinking, mood, or behavior or a combination of all three, leading to impaired function, disability, pain, and even death. The term **mental illness** is applied to diagnosable mental disorders. All age groups, gender, racial, and ethnic groups are affected regardless of education or socioeconomic level. A goal of Healthy People 2010 is “to improve mental health and ensure access to appropriate, quality mental health services.” As a nurse, you are well positioned to assess the mental health status of your patients at every level of health prevention and in every area of healthcare.

Prevention

Primary prevention entails decreasing the incidence of mental disorders by identifying those at risk, those who are vulnerable to developing mental disorders. Identifying risk factors include biological, psychosocial, and environmental factors that may contribute to devel-

oping mental illness. A biological risk factor could be a family history of mental illness with a genetic link such as schizophrenia. A person’s developmental level may be a psychosocial risk factor. External stressors are examples of environmental factors.

Assessment entails not only identifying your client’s weaknesses but also identifying his or her strengths. So, you need to assess not only for risk factors but also for protective factors, such as coping mechanisms and resources.

Secondary prevention involves reducing the prevalence of mental disorders through screening and initiating prompt treatment of mental disorders. **Tertiary prevention** aims at reducing the severity of the illness by providing follow-up and rehabilitative services.

Biological Basis of Mental Illness

Current research is providing more and more evidence that there are organic and neurochemical bases for some mental illnesses and behaviors. Be sure to review the anatomy and physiology section in Chapter 21, Assessing the Sensory-Neurological System.

Genetic links have been identified for certain brain disorders such as schizophrenia. Although schizophrenia, depression, bipolar disorder, anxiety disorders, and Alzheimer’s are classified as mental illnesses, biological bases have been identified. Biochemical factors, specifically **neurotransmitters**, have been identified in depression and mania (Fig. 5.1). Another part of the brain is the limbic system, which is called the *emotional brain* and is associated with fear, rage, love, anger, aggression, hope, sexuality, and social behavior (Fig. 5.2).

Neurotransmitters: Functions and Effects

NEUROTRANSMITTER	FUNCTION	EFFECT
Dopamine	Inhibitory	Fine movement, emotional behavior. Implicated in schizophrenia and Parkinson’s.
Serotonin	Inhibitory	Sleep, mood, eating behavior. Implicated in mood disorders, anxiety, and violence.
Norepinephrine	Excitatory	Arousal, wakefulness, learning. Implicated in anxiety and addiction
Gamma-aminobutyric acid (GABA)	Inhibitory	Anxiety states
Acetylcholine	Excitatory	Arousal, attention, movement. Increase = spasms Decrease = paralysis

Source: Pedersen, D. (2005). *PsychNotes*. Philadelphia: F.A. Davis.




FIGURE 5.1. Impulse transmission at a synapse. The *arrow* indicates the direction of the electrical impulse. The entry of Na⁺ ions stimulates an electrical charge in the postsynaptic neuron.




FIGURE 5.2. Structures of the limbic system.

Psychological Theories

Psychodynamic theories focus on intrapersonal concepts. These theories look at the development of the mind over the course of a lifetime. Freud, the “Father of Psychiatry,” introduced the psychoanalytic model of the conscious and unconscious mind. Various interpersonal and humanist models followed. Freud identified **defense mechanisms**, which served to protect the individual from unwanted anxiety.

Behavioral Theories

Behavioral theories focus on normal behaviors rather than the causes of mental illness. Pavlov, Watson, Thorndike, and Skinner are behavioral theorists. The goal is to change behavior through conditioning, behaviorism (which includes frequency and recency), and positive reinforcement (Boyd, 2005).

Cognitive Theories

Cognitive theories provide an understanding by looking not only at behavior but also at an individual’s cognitions and how a person processes thoughts (distorted cognitions and maladaptive behaviors). Bandura and Beck are cognitive theorists. Cognitive theory is of value when teaching patients new coping skills; individual collaboration with the therapist and active involvement of the patient are needed for change to take place (Townsend, 2006).

Developmental Theories

Developmental theories provide an understanding of the individual’s growth and development over one’s lifetime. Erikson, Piaget, Gilligan, and Miller are developmental theorists. Various developmental theories are described in Chapter 2, The Health History.

Social Theories

Social theories include family dynamics, role theories, and sociocultural perspectives; thus, the assessment of

sociocultural aspects is critical in developing a plan of care for the patient.

Developmental, Cultural, and Ethnic Variations

Developmental Level

Mental disorders can occur with any age group, with the incidence of certain disorders higher for specific age groups. For example, learning and behavioral problems are seen with children. Adolescents, known risk takers, have a high incidence of substance abuse. Bipolar illness or psychoses such as schizophrenia more commonly become apparent during young adulthood. Depression is often seen in the older adult.

Children

Because rapid growth and developmental changes occur during childhood, delays are readily detectable. Learning disorders and behavioral disorders often become apparent by the time the child reaches school age. (See Chapter 2, The Health History, for more on developmental stages and theories.)

Mental Health Assessment for Children

Assessing the mental health of children should include both a detailed history and a physical examination. The assessment should include an interview with the child. If abuse is suspected, be sure to interview the child alone without the parents or caregiver present.

Health History

The health history should include a family history with attention to a history of mental health problems. A **maternal history** is also needed and should identify:

- Age and health of mother during pregnancy.
- Exposure to medications, alcohol or drugs during pregnancy.
- Course of labor and type of delivery.
- Infant’s state of health at birth.

Assess the child’s normal growth and development milestones, such as motor development, bowel and bladder control, speech and language development, and social development. Also, assess the child’s medical health, noting any major illness or injury.

The assessment should also include a current developmental assessment, a mental status examination, and a physical examination. Various instruments are available to assess a variety of mental health disorders in children. Be sure to select the one best suited to meet the child’s needs.

Children Rating Scales for Mental Health Disorders

INSTRUMENT

Behavior Assessment System for Children (BASC)
Child Behavior Checklist (CBCL)
Children's Depression Inventory (CDI)

Pediatric Anxiety Rating Scale (PARS)
Yale-Brown Obsessive Compulsive Scale (YBOCS)
Swanson, Nolan, Pelham (SNAP)-IV

ASSESSES FOR

Behavior and emotions for children ages 2–18
Psychopathology and competencies for children ages 4–16
Physical symptoms, harm avoidance, social anxiety, and separation/panic disorders for children ages 7–17
Severity of anxiety for children ages 6–17
Obsessive compulsive disorder (OCD) for children ages 6–17
Inattention and overactivity (attention deficit hyperactivity disorder [ADHD]) and oppositional defiant disorders

Adapted from Boyd, M.A. (2005). *Psychiatric Nursing Contemporary Practice*. Philadelphia: Lippincott, Williams & Wilkins.



HELPFUL HINTS

- Learning disorders and behavioral disorders often become apparent by the time the child reaches school age.
- Depending on the age of the child, play or drawing can be very helpful in eliciting the child's perceptions.

Child Abuse

Child abuse affects all areas of a child's life and may manifest as a mental or physical disorder. Child abuse takes many forms from physical abuse, neglect, sexual abuse, emotional abuse, factitious disorders by proxy (Munchausen syndrome), and secondary abuse (children of battered women). The assessment should include an interview with the child. If abuse is suspected, be sure to interview the child alone without the parents or caregiver present. (See Guidelines for Assessing Abuse, in Appendix C.)

Risk Taking

Adolescents are known risk takers. Experimenting with drug and alcohol is common, and suicide is the second leading cause of death among adolescents. Mental and substance abuse disorders or a combination of disorders are more often than not associated with suicide. A goal of Healthy People 2010 is to decrease the attempted suicide rate among adolescents from a 12-month average of 2.6 percent of adolescents in grades 9 through 12, to a 12-month average of 1 percent.

Another popular form of risk taking during childhood is the **choking game**, which produces a high or a state of euphoria by cutting off the blood supply to the brain. If the blood supply is compromised for too long, the child will pass out and feel a "rush" as consciousness returns. Depriving the brain of oxygen may result in permanent and cumulative death to brain cells, stroke, seizures, retinal damage, or death.

Risk Factors for Teen Suicide

- Depression
- Social isolation
- History of abuse
- Drug and alcohol abuse
- Not fitting in with peers
- School pressure

Source: Gorman, L., Raines, M., and Sultan, D. (2005). *Psychosocial Nursing for General Patient Care*, ed. 2. Philadelphia: F.A. Davis.

This behavior, which is also known as *fainting, American dream, flatliner, space cowboy, knockout, gasp, rising sun, airplaning, pass out, blackout, and space monkey*, is most common among children ages 10 through 16. It can be performed either alone or with others. Use of ligatures and performing this alone increases the risk of death.



HELPFUL HINT

Children who "play" the choking game are usually not troubled children and generally have no history of drug or alcohol abuse.

Signs and Symptoms of the Choking Game

- Unexplained marks or bruises on neck
- Severe headaches
- Red eyes
- Belts, leashes, ropes, and shoelaces tied in strange knots or in unusual locations
- Disorientation after being alone
- Locked bedroom doors

Pregnant Women

Although the birth of child is seen as a happy event, mental health disorders can occur, affecting not only the mother but also the entire family. Many mothers experience mild depression, also known as *baby blues*, usually within 4 weeks following childbirth. If the baby blues persist for more than a few weeks, postpartum depression may have developed. The depression is more serious and results in postpartum depression with or without psychosis. The mother may experience severe mood swings and pervasive sadness. The mother often has feelings of guilt and inadequacy as a mother. In addition to the feelings associated with postpartum depression, with postpartum psychosis, the mother experiences delusions and thoughts of harming herself or her baby.

If you detect at-risk mothers or postpartum depression, be sure to make the appropriate referrals for follow-up care. Assessment for postpartum depression is important. Screening tools for postpartum depression include the Postpartum Check List (Beck, 1995) and the Edinburgh Postpartum Depression Scale (Cox, Holden, & Sagovsky, 1989).

Older Adults

Older adults are more at risk for the development of mental health problems for several reasons, ranging

from the normal physiological changes associated with aging to acute and chronic illness to side effects associated with polypharmacy. Risk factors may be developmental, situational, internal, or external in nature.

Mental Health Assessment of the Older Adult

Approximately 25 percent of all older adults experience some type of mental disorder (depression, anxiety, substance abuse, and dementia) that is not related to the normal changes associated with aging (Healthy People 2010). Alzheimer's affects 8 percent to 15 percent of older adults over the age of 65 and accounts for 69 percent to 70 percent of all dementia (Healthy People 2010). Assessing the mental health of the older adult should include both a detailed history and a physical examination. The assessment should focus on mental status, behavioral responses, functional ability, general health, and social supports of the client.

Mental status assessment should include a Mini-Mental State Examination (MMSE), a mental status examination, and assessment for depression, anxiety, and psychosis. A change in mental status has to be carefully evaluated. Changes are often attributed to normal changes associated with aging. Changes in mental status may be physiological or psychological in nature. Because dementia increases with age, care must be taken to differentiate dementia from delirium and depression.

Risk Factors for Postpartum Depression

- History of mood disorders
- Low self-esteem
- Unwanted pregnancy
- Unemployment of mother or head of household
- Poor marital relationship
- Father depressed
- Poor support systems
- External stressors

Risk Factors for Development of Mental Health Problems in Older Adults

- Physical changes associated with aging that may affect functional ability
- Cognitive changes and memory loss
- Chronic medical illness
- Psychosocial changes, such as retirement, that occur with aging
- Limited financial resources
- Polypharmacy
- Loss and bereavement
- Loss of social supports



ALERT

Delirium should be treated as an emergency because its acute onset may have an underlying life-threatening cause. The MMSE and the Clock Scoring Test are screening instruments used to assess cognitive impairment and screen for dementia. (See Chapter 27, Assessing the Older Adult, for more information.)

Assessment of behavioral responses includes description of behavior and triggers, with special attention to behavioral changes and problem behaviors. A functional assessment should focus on the client's ability to perform activities of daily living, physiological functioning, mobility, and risk for fall assessment. (See Chapter 27, Assessing the Older Adult, for more information.) A nutritional assessment and history of substance abuse is included in the assessment of the client's general health. Finally, the assessment should identify social supports, family-patient interactions, and caregiver concerns.

Characteristics of Dementia, Delirium, and Depression

FEATURE	DEMENTIA	DELIRIUM	DEPRESSION
Onset	Gradual (months to years)	Abrupt (hours to a few weeks)	Either
Prognosis	Irreversible	Reversible	Variable
Course	Progressive	Worse in P.M.	Possibly worse in A.M.
Attention	Normal	Impaired	Variable
Memory	Impaired recent and remote	Impaired recent and immediate	Impairment
Perception	Normal	Impaired	Normal
Psychomotor behavior	Normal/apraxia	Hypo-/hyperkinetic	Retardation/agitation
Cause	Caused by many diseases, including alcoholism, acquired immunodeficiency syndrome (AIDS), cerebral anoxia, and brain infarcts	Caused by acute illness, fever, infection, dehydration, electrolyte imbalance, medications, and alcoholism	May coincide with life event, such as death in the family, loss of a friend or a pet, or a move

Depression

Depression is a major mental health problem for older adults, but it can be easily missed. If left untreated, depression can lead to suicide or self-neglect. In the United States, the suicide rate is highest among older adults, with older men over age 80 at highest risk and guns the most frequent method (Gorman, Raines, & Sultan, 2005). But the suicide rate is probably higher, because older adults can easily commit passive suicide by not complying with medical regimens or failing to eat.

Elder Abuse

Elder abuse, if taking place, affects all areas of the patient's life and may manifest as a mental or physical disorder. Elder abuse can take many forms: physical abuse, neglect, sexual abuse, emotional abuse, and financial abuse. (See Guidelines for Assessing Abuse, in Appendix C.)

Risk Factors for Suicide Among Older Adults

- Lives alone
- Widower
- Lack of financial resources
- Poor health
- Social isolation
- Drug and alcohol abuse

Source: Gorman, L., Raines, M., and Sultan, D. (2005). *Psychosocial Nursing for General Patient Care*, ed. 2. Philadelphia: F.A. Davis.

Geriatric Depression Scale

- Are you basically satisfied with your life? YES/NO
- Have you dropped many of your activities and interests? YES/NO
- Do you feel that your life is empty? YES/NO
- Do you often get bored? YES/NO
- Are you in good spirits most of the time? YES/NO
- Are you afraid that something bad is going to happen to you? YES/NO
- Do you feel happy most of the time? YES/NO
- Do you often feel helpless? YES/NO
- Do you prefer to stay at home rather than going out and doing new things? YES/NO
- Do you feel you have more problems with memory than most? YES/NO
- Do you think it is wonderful to be alive now? YES/NO
- Do you feel pretty worthless the way you are now? YES/NO
- Do you feel full of energy? YES/NO
- Do you feel that your situation is hopeless? YES/NO
- Do you think that most people are better off than you are? YES/NO

Bold answers = depression

GDS Scoring:

- 12–15 Severe depression
- 8–11 Moderate depression
- 5–8 Mild depression
- 0–4 Normal

Source: Yesavage, et al., 1983, pp. 37–47.

GDS website: <http://www.stanford.edu/~yesavage/>

Signs and Symptoms of Depression in the Older Adult

Typical

- Changes in appetite
- Decreased self-esteem
- Changes in sleep patterns
- Social withdrawal
- Feelings of helplessness
- Loss of motivation
- Constipation
- Hostility
- Pessimism
- Agitation
- Guilt
- Aggression

Atypical

- Vague somatic complaints—such as constipation, joint pain, fatigue, and memory changes—that seem to be out of proportion to the actual problem.
- Client may become obsessed with the problems and feel that if the problems are relieved, she or he will be fine.

People of Different Cultures or Ethnic Groups

Cultural and ethnic variations often influence a patient's view of mental health. Cultural beliefs influence perceptions, understanding, and treatment of mental illness.

Introducing the Case Study



Hollie Baker, age 40, is complaining of extreme fatigue and not being able to sleep for the past month. She is wondering if there is something physically wrong with her or maybe she just needs some vitamins. After further discussion, she states she has “no energy to do anything” and that she has a lot on her mind. She states that she is feeling overwhelmed and depressed by the whole situation stating, “maybe things would be better if I could just disappear.” Recently divorced and the mother of two adolescent children, she works full-time as an administrative assistant. Ms. Baker has a history of breast cancer, hypertension, hypothyroidism, and bipolar disorder. Her current medications include tamoxifen, metoprolol, Synthroid, lithium, and Risperdal.

Cultural Perceptions of Mental Health/Illness

CULTURAL GROUP

PRACTICES/BELIEFS

Amish	<ul style="list-style-type: none"> ■ Children with mental or physical differences are seen as “hard learners.” ■ The mentally ill are generally cared for at home.
Appalachian	<ul style="list-style-type: none"> ■ Usually takes care of own, so mentally deficient and handicapped are readily accepted. ■ Mentally handicapped are not crazy, but rather have “bad nerves” or are “quiet turned” or “bad turned.” ■ Behaviors that would warrant psychiatric treatment are seen as lazy, mean, immoral, criminal, or psychic and treated by punishment or tolerance.
Arab American	<ul style="list-style-type: none"> ■ Mental illness is seen as a social stigma, therefore psychiatric symptoms may be denied, attributed to “bad nerves” or supernatural beings, or caused by a physical ailment or emotional trauma. ■ Somatic orientation leads to patients' tolerance of emotional suffering and relatives' tolerance of behavioral disorders. ■ Patients with mental distress will somatize symptoms. Somatic treatment for psychiatric disorders is preferred over psychotherapy.
Chinese American	<ul style="list-style-type: none"> ■ Balance between yang and yin explains mental health/illness. ■ Mental illness results from metabolic imbalance and organic problem. ■ Stigma is associated with mental illness, so patient will seek folk healer first. ■ Does not readily seek treatment for emotional or nervous disorders.

CULTURAL GROUP	PRACTICES/BELIEFS
Cuban American	<ul style="list-style-type: none"> ■ Emotional crises are treated by santero, a practitioner of santería (a 300-year-old Afro-Cuban religion that combines Roman Catholic elements with ancient Yoruba tribal beliefs and practices). ■ May act out emotional problems in a non-threatening way, allowing person to maintain self-esteem.
Egyptian American	<ul style="list-style-type: none"> ■ Mental illness is considered a stigma. ■ More tolerance for emotional problems. ■ View mental health problems with a supernatural framework, including “curse” and “devil.” ■ Emotional problems of grief and losses are due to wrongdoings of others or weakness and inability to control and snap out of distress. ■ Mental and emotional issues are expressed somatically so therefore treated with psychosomatic interventions.
Filipino American	<ul style="list-style-type: none"> ■ Seeks family and friends for advice. ■ Stigma is associated with mental illness. ■ Mental illness is caused by heredity. ■ Takes care of mentally ill family member rather than seek mental healthcare.
French Canadian Greek American	<ul style="list-style-type: none"> ■ Federal Canadian laws protect the mentally disabled from discrimination. ■ Stigma is associated with mental illness. ■ Mental illness is seen as hereditary linked, therefore “polluting” the blood line, bringing shame to family. ■ Families of mentally ill may experience loss of friends and social isolation. ■ Mental illness often somatized. Wide range of acceptable behavior leads to delay of seeking treatment. ■ Folk model for “nerva” (nerves) is socially acceptable and treated with medication rather than with psychotherapy.
Iran American	<ul style="list-style-type: none"> ■ Stigma is associated with mental illness. ■ Mental illness is caused by genetic problem; more likely to be labeled a “neurological” problem. ■ Avoids psychotherapeutic treatment. ■ Symptoms somatized and treated with psychopharmacology.
Irish American	<ul style="list-style-type: none"> ■ High rate of mental illness. ■ Difficulty expressing emotions and feelings. ■ Family may take care of mentally ill family member. ■ Some attribute mental illness to sin and guilt.
Jewish American	<ul style="list-style-type: none"> ■ Mental health is as important as physical health. ■ Mental incapacity relinquishes one from all responsibilities.
Mexican American	<ul style="list-style-type: none"> ■ Mental illness is seen as God’s will. ■ Family prefers to care for patient at home.
Native American	<ul style="list-style-type: none"> ■ Mental illness is caused by witches; treated by healers. ■ May wear turquoise to ward off evil. ■ Some tribes view mentally ill as having special problems.
Vietnamese American	<ul style="list-style-type: none"> ■ Mental illness results from offending a deity. ■ Brings disgrace to family and is therefore concealed, which delays treatment. ■ Emotional disturbances are caused by malicious spirits, bad luck, or family inheritance. ■ Buddhists see mental illness as bad karma from previous misdeeds. ■ Sometimes nervous system is seen as cause of mental problems.

Performing the Mental Health Assessment

A thorough assessment includes assessment of the psychological health of the patient. Mental health reflects a person's positive attitudes toward self, growth and development, self-actualization, integration, autonomy, reality perception, and environmental mastery (Stuart & Laraia, 2005). As you perform your assessment, look for clues that reflect the mental health status of your patient.

Realize that your patient may prefer to have you believe she or he has a physical problem rather than a psychological problem by somatizing feelings. Also, medical problems or drug use (prescribed, over-the-counter [OTC], illegal) may present as mental disorders, such as hyperthyroidism, which may produce signs and symptoms of anxiety. Comorbidities can also result with patients. Chronic illness can lead to depression, especially as the disease progresses.

Health History

When obtaining a health history, ask yourself, "What can the health history reveal about the mental health of the patient?" Identify factors that may affect mental health in either a positive or a negative way. Ensure patient confidentiality, but confidentiality should be breached if there is suspected abuse or a threat of child or elder abuse or if the patient poses a threat to herself or himself or others.



ALERT

Report any suspected abuse or any threat of child or elder abuse. Also report threats of suicide or homicide. Threats to self or others must always be taken seriously.

Health History as It Relates to Mental Health

RISK/FACTORS/ QUESTIONS TO ASK

Biographical

Age

- How old are you?

Gender

- Is the patient male or female?

Socioeconomic and Educational Levels

- What is your educational level?

RATIONALE/SIGNIFICANCE

- Response to stressors differs with age.
- Identifies possible maturational crises related to developmental changes.
- Women have greater incidence of depression and affective and anxiety disorders.
- Men have greater incidence of psychosocial and substance abuse disorders.
- Age of onset of schizophrenia occurs later in women than in men.
- Men are more likely to be aggressive and self-destructive than women.
- The higher the socioeconomic and educational level, the lower the incidence of depression.
- The higher the educational level, the more likely the individual is to use mental health services if needed.
- The lower the income, the higher the incidence of psychological symptoms.

**RISK/FACTORS/
QUESTIONS TO ASK****RATIONALE/SIGNIFICANCE****Ethnicity**

- What is the patient's cultural background?

- African Americans and Hispanics have twice the rate of being diagnosed with severe mental illness than other groups.
- African Americans are diagnosed with the most severe types of psychopathology and fewer affective disorders and are overdiagnosed with schizophrenia.
- Ethnic groups have three times more hospitalizations for mental illness than the general population.

Marital Status

- Are you single, married, or divorced?

- Married and partnered adults report less stress than single or divorced adults.

Religion

- What is your religious affiliation?

- Can affect mental health in either a positive or a negative way.
- Influences perceptions of mental illness.

Contact Person

- Who is your contact person?

- Identifies supports.

Current Health Status

- How is your health?

- Identifies any signs or symptoms that may reflect mental illness.

Past Health History**Childhood Illnesses**

- Did you have any major health problems (physical or mental) while growing up?

- Identify any past mental health problems and treatments.

Past Hospitalizations

- Have you ever been hospitalized for mental health problems?

- Identifies previous psychiatric hospitalization.

Family History

- Does anyone in your family have mental health problems?

- Identifies familial/genetically linked psychiatric disorders. Familial/genetically linked disorders include:
 - Schizophrenia.
 - Depression.
 - Bipolar disorders.
 - Anxiety and panic disorders.

Medications

- Are you taking any medications, either prescribed, OTC, or herbal? If yes, what are you taking? (See *Drugs That May Adversely Affect Mental Health*.)

- Identifies any psychiatric drugs.
- Identifies any possible drug interactions.
 - OTC medications such as pseudoephedrine can cause anxiety symptoms.
 - Herbal supplements such as St. John's wort, ephedra, ginseng, kava kava, and yohimbe can interact with psychotropics or other medications, or cause anxiety, drowsiness, or other adverse psychological effects (Pedersen, 2005). (See *Psychotropic Drugs and Some Related Side Effects*.)

Military Service

- Are you now or did you ever serve in the military? If yes, when and what was your tour of duty?

- Identifies possible source of stressor related to post-traumatic stress syndrome

Drugs That May Adversely Affect Mental Health

Drugs	Effect	
Antihypertensives	Depression	
■ Reserpine		
■ Beta blockers		
■ Methyldopa		
Oral contraceptives		
Corticosteroids		
Benzodiazepine		
Cancer chemotherapeutic agents		
■ Vincristine		
■ Vinblastine		
■ Interferon		
■ Procarbazine		
Psychoactive agents		
■ Alcohol		
■ Amphetamine or cocaine withdrawal		
■ Opioids	Manic states	
Corticosteroids		
Levodopa		
Amphetamines		
Tricyclic antidepressants (TCAs)		
MAO inhibitors		
Methylphenidate		
Cocaine		
Thyroid hormone		Psychotic reactions
Amphetamines		
Antidepressants (particularly tricyclics)		
Anticholinergics (atropine)		
Anticonvulsants (carbamazepine, valproic acid)		
Antiparkinsonians (levodopa)		
Antituberculars (isoniazid)		
Antivirals (acyclovir, amantadine)		
Antiarrhythmics (lidocaine)		
Alcohol		
Beta blockers (propranolol)		
Corticosteroids		
H ₂ -receptor blockers (cimetidine)		
Cyclosporine		
Disulfiram (Antabuse)		
Anesthetics (ketamine)		
Antibiotics (cephalosporins, ciprofloxacin, sulfonamides)		
Opioids (morphine, hydromorphone)		

Source: Gorman, L., Raines, M., and Sultan, D. (2005). *Psychosocial Nursing for General Patient Care*, ed. 2. Philadelphia: F.A. Davis.



ALERT

Neuroleptic malignant syndrome (NMS) is a potentially fatal side effect from antipsychotic drugs, characterized by fever, tachycardia, sweating, muscle rigidity, tremors, incontinence, stupor, leukocytosis, elevated creatine phosphokinase (CPK), and renal failure

Psychotropic Drugs and Some Related Side Effects

DRUG	SIDE EFFECTS
Lithium	Nausea, diarrhea, polyuria, acne, rashes, alopecia, tremors; weight gain, hypothyroidism; can also precipitate psoriasis and psoriatic arthritis.
Anticonvulsants	Sedation, hepatotoxicity, rash, and Stevens-Johnson syndrome (SJS), which is a life-threatening mucocutaneous reaction.
Anticholinergics TCAs	Blurred vision, constipation, dry mouth Sedation, blurred vision, dry mouth, constipation, life-threatening arrhythmias and electrocardiogram (ECG) changes
Selective serotonin reuptake inhibitors	Gastrointestinal (GI) symptoms, insomnia, and agitation
Antipsychotics	Movement disorders (akathisia) and tardive dyskinesia (especially conventional antipsychotics [chlorpromazine]), dystonia, parkinsonism, gynecomastia, and lactation, and possibly treatment-emergent diabetes

Source: Adapted from Pedersen, D. (2005). *PsychNotes*. Philadelphia: F. A. Davis.

Review of Systems

As you proceed with the review of systems, note

any problems that may affect your patient's mental health.

Review of Systems

AREA/GENERAL QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
General Health Survey ■ How have you been feeling?	■ Fatigue may be associated with depression. ■ Restlessness may be associated with anxiety.
Integumentary ■ Do you have any problems with your skin? If yes, describe.	■ Sweating, itching associated with anxiety
Head, Eyes, Ears, Nose, and Throat (HEENT) ■ Do you have headaches? If yes, describe. ■ Do you have thyroid disease?	■ Headaches associated with depression. ■ History of migraines and tension headaches affected by psychological factors. ■ History of hyperthyroidism and diabetes can be affected by psychological factors, manifest with psychological signs or symptoms, or mimic psychological disorders.
Respiratory ■ Do you have any breathing problems? If yes, describe.	■ History of breathing problems, such as hyperventilation, associated with anxiety disorders. Sighing associated with depression.
Cardiovascular ■ Do you have any CV problems? If yes, describe.	■ Hypertension, angina affected by psychological factors ■ Palpitations, racing heart associated with anxiety ■ Heterocyclic antidepressants—use with caution with cardiovascular disease

(continued)

Review of Systems (continued)

AREA/GENERAL QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Gastrointestinal</p> <ul style="list-style-type: none"> ■ Do you have any GI problems? If yes, describe. ■ Have you experienced changes in weight? If yes, describe. ■ Have you experienced changes in appetite? If yes, describe. 	<ul style="list-style-type: none"> ■ Changes in appetite associated with depression, anxiety, eating disorders, substance abuse ■ History of peptic ulcer, irritable bowel syndrome, colitis affected by psychological factors ■ GI complaints such as nausea, abdominal pain, diarrhea seen with anxiety ■ Indigestion and constipation seen with depression ■ Nausea is a common side effect of many psychotropic medications.
<p>Genitourinary (GU)</p> <ul style="list-style-type: none"> ■ Do you have any GU problems? If yes, describe. ■ If your patient is female, ask if she has experienced any irregularities in her menstrual cycle. ■ Do you have any concerns about your sexual performance? If yes, describe. 	<ul style="list-style-type: none"> ■ History of sexual dysfunction, such as impotence, frigidity, and premenstrual syndrome (PMS), affected by psychological factors ■ Menstrual irregularities associated with eating disorders ■ Pressure and frequency of urination associated with anxiety ■ Use TCAs (amitriptyline), cautiously with benign prostatic hyperplasia (BPH)
<p>Musculoskeletal (MS)</p> <ul style="list-style-type: none"> ■ Do you have any MS problems? If yes, describe. 	<ul style="list-style-type: none"> ■ History of rheumatoid arthritis and idiopathic low back pain affected by psychological factors ■ Weakness associated with anxiety ■ Osteoporosis associated with eating disorders ■ Use of antipsychotics contraindicated with myasthenia gravis.
<p>Neurological</p> <ul style="list-style-type: none"> ■ Do you have any neurological disorders? ■ Have you experienced seizures, concentration, or memory problems? ■ Have you experienced depression, anxiety? If yes, describe. 	<ul style="list-style-type: none"> ■ Seizures associated with eating disorders, use TCAs cautiously with seizures and bupropion is contraindicated with seizures. ■ Neuroleptic malignant syndrome and movement disorders are a side effect of antipsychotic medications ■ Cognitive problems, difficulty focusing, inability to concentrate associated with anxiety, depression, dementia, and schizophrenia

Psychosocial Profile

As you perform the psychosocial assessment, look for clues that would reflect your patient's mental health.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Health Practices and Beliefs

- How would you describe your mental health?
- Do you or would you use mental health services?

- Identify perceptions of mental health, mental health practices, and use of mental health services.

Typical Day

- Can you tell me what your typical day is like?

- Identifies ability to maintain activities of daily living (ADLs). Mental health problems such as depression and schizophrenia, compulsive disorders may affect ability to perform ADLs.

Nutritional Patterns

- Can you tell me what you ate yesterday (24-hour recall)?

- Nutritional disorders are associated with eating disorders, anxiety, and depression.

Activity and Exercise Patterns

- Do you exercise regularly? If yes, describe.

- Excessive exercise is associated with eating disorders.
- Inactivity is associated with depression.

Recreation, Pets, Hobbies

- What do you do for fun?
- Do you have pets?
- Do you have hobbies?

- Inactivity is associated with depression.

Sleep/Rest Patterns

- How many hours of sleep do you get a night?
- Do you have any problems falling asleep, staying asleep?
- Do you take or do anything to help you sleep?

- Problems with sleep and rest are associated with many psychological disorders, such as anxiety, depression, bipolar disorders, and substance abuse.

Personal Habits

- Do you use alcohol, drugs, caffeine, or nicotine? If, yes, how much?

- Identifies history of substance abuse (use of alcohol, drugs, caffeine, nicotine).

Occupational Health Occupation?

- How do you feel about your work? Your coworkers?
- Do you find your work stressful? If yes, how do you deal with it?
- Are there any health risks associated with your work? If yes, describe.

- Identify ability to maintain job. Work can be a source of stress.
- Identify risk for head injury, such as construction work

Environmental

- Where do you live?
- Are you exposed to any pollutants or toxins?

- Identifies risk for exposure to toxic substances, such as lead, mercury, herbicides, solvents, cleaning agents, and lawn chemicals, that can affect cognitive ability.

Roles, Relationships, Self-Concept

- How do you see yourself?
- Can you identify your various roles and relationships?

- Identifies sense of worth and value, which may be affected by psychological factors such as low self-image associated with depression and eating disorders.
- Roles and relationships may affect or be affected by psychological factors.

Cultural Influences

- What is your cultural background?
- What influences your perception of health?

- Identifies culture's perception of mental health and illness.

Religious/Spiritual Influences

- What is your religious background?
- What influence does your religious beliefs have on your perception of health?

- Identifies religious/spiritual influences on mental health and illness.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Family Roles and Relationships <ul style="list-style-type: none"> ■ What is your role in your family? ■ How is your relationship with your family? 	<ul style="list-style-type: none"> ■ Identifies sources of support or stress on patient's mental health. ■ Mental illness may affect patient's role and relationship in family.
Sexuality Patterns <ul style="list-style-type: none"> ■ Do you have any concerns about sexual patterns? If yes, describe. ■ Do you practice safe sex? 	<ul style="list-style-type: none"> ■ Sexual problems are often associated with mental illness such as depression. ■ Unprotected sexual activity is associated with substance abuse.
Stress and Coping Patterns <ul style="list-style-type: none"> ■ How do you deal with stress? ■ What do you do when you are upset? 	<ul style="list-style-type: none"> ■ Identifies current coping skills. ■ Determines effectiveness of coping skills.



Case Study Evaluation

Before you proceed with the physical examination, document the key history information you have learned from Ms. Baker's history and mental health history. The information from your patient history will be important to consider before and during your physical examination.



CRITICAL THINKING ACTIVITY #1

What may account for Ms. Baker feeling fatigued? What additional questions would you need to ask Ms. Baker?



CRITICAL THINKING ACTIVITY #2

Based upon what you know about Ms. Baker, what would be a priority concern that needs to be addressed immediately?



CRITICAL THINKING ACTIVITY #3

Based upon what you know about Ms. Baker, which findings suggest depression?

Physical Assessment

As you perform a head-to-toe physical examination, consider how your patient's physical findings reflect his or her mental health.

Mental Status Assessment

A mental status assessment, involving a systematic approach to various components, is done to assess and evaluate a patient's cognitive and mental functions. Each assessment provides information about that patient at that point in time. (See *Mental Status Assessment as it Relates to Mental Health/Illness*.) Additional assessments will demonstrate improvement, regression, or stabilization and will provide a "progress report" as well as "patterns" of functioning.

The components of the mental status assessment include:

- General appearance.
- Behavior/activity.
- Speech and language.
- Mood and affect.
- Thought process and content.
- Perceptual disturbances.
- Memory/cognitive.
- Judgment and insight.

Additional Assessments

Depending on your assessment findings, additional mental health screening may be indicated. Various mental health problems and assessment instruments are available to assess for a variety of mental health problems. Additional mental health assessments are presented below.

(Text continued on page 131)

Head-to-Toe Physical Examination as It Relates to Mental Health/Illness

SYSTEM	ABNORMAL FINDINGS/RATIONALE
General Health Survey	<ul style="list-style-type: none"> ■ <i>Poor grooming and personal hygiene</i>: Associated with depression ■ <i>Bright colors or unusual dress</i>: Associated with mania ■ <i>Poor eye contact</i>: May indicate depression ■ <i>Inability to maintain attention</i>: Associated with schizophrenia and depression ■ <i>Labile affect</i>: May reflect mania ■ <i>Flat, incongruent affect</i>: May reflect schizophrenia ■ <i>Stooped posture</i>: Associated with depression ■ <i>Restlessness, tension</i>: Associated with anxiety ■ <i>Malnourished appearance</i>: May indicate an eating disorder ■ <i>Slurred speech</i>: May indicate drug and alcohol abuse ■ <i>Pressured speech</i>: Seen with mania ■ <i>Disorganized speech</i>: Seen with schizophrenia ■ <i>Irritability</i>: Associated with anxiety ■ <i>Suspiciousness</i>: Associated with paranoia
Integumentary	<ul style="list-style-type: none"> ■ <i>Flushed or pallid skin color</i>: Seen with anxiety ■ <i>Excessive sweating</i>: Seen with anxiety ■ <i>Injury or scarring</i>: From self-injury or past suicide attempts or self-mutilation
HEENT	<ul style="list-style-type: none"> ■ <i>Dilated or constricted pupils</i>: Seen with drug abuse ■ <i>Poor eye contact</i>: Seen with depression ■ <i>Dental caries, parotid swelling</i>: Seen with eating disorders ■ <i>Erosion of nasal or oral mucosa</i>: May be secondary to drug use ■ <i>Rope marks on neck</i>: In children, associated with “choking game”
Respiratory	<ul style="list-style-type: none"> ■ <i>Increased respiratory rate and hyperventilation</i>: Seen with anxiety
Cardiovascular	<ul style="list-style-type: none"> ■ <i>Increased pulse rate and blood pressure</i>: Seen with anxiety
Abdominal	<ul style="list-style-type: none"> ■ <i>Increased bowel sounds</i>: Seen with excessive use of laxatives in eating disorders
Musculoskeletal	<ul style="list-style-type: none"> ■ <i>Abnormal liver size</i>: Associated with substance abuse ■ <i>Generalized weakness and tremors</i>: Seen with anxiety ■ <i>Abnormal muscle movement</i>: May be adverse effect of psychotropic drugs ■ <i>Excessive body movements</i>: Associated with anxiety, mania, or stimulant abuse ■ <i>Minimal or no body movement</i>: Associated with depression, catatonic states, or drug-induced stupor ■ <i>Repeated movements</i>: Associated with compulsive disorders ■ <i>Repeated picking at clothes</i>: May be associated with hallucinations, delirium, or toxic conditions
Neurological	<ul style="list-style-type: none"> ■ <i>Cognitive changes, thought process disorders</i>: Seen with schizophrenia ■ <i>Loose association</i>: Associated with schizophrenia ■ <i>Flight of ideas</i>: Associated with mania ■ <i>Preservation</i>: Associated with brain damage and psychotic disorders ■ <i>Auditory hallucinations</i>: Associated with schizophrenia ■ <i>Visual hallucinations</i>: Often organic in nature ■ <i>Tactile hallucinations</i>: Seen with organic problems, drug abuse, or delirium tremens (DTs) ■ <i>Changes in mental status</i>: Seen with mental illness such as schizophrenia, substance abuse, bipolar disorder ■ <i>Increased reflexes</i>: Seen with anxiety ■ <i>Movement disorders</i>: Adverse effect associated with antipsychotic medications

Mental Status Assessment as It Relates to Mental Health/Illness

AREA/QUESTIONS TO ASK	NORMAL	ABNORMAL FINDINGS/RATIONALE
Grooming/dress ■ Does the patient dress appropriate for age, gender, season, and situation?	■ Dress neat and appropriate	■ <i>Disheveled appearance</i> : Associated with depression ■ <i>Bright-colored clothing</i> : Associated with mania
Hygiene ■ Are the patient and his or her clothing clean? ■ Does the patient present with any unusual odors?	■ Clean, no unusual odors	■ <i>Poor hygiene</i> : Associated with depression or schizophrenia
Eye Contact ■ Does the patient maintain eye contact?	■ Maintains eye contact	■ <i>Poor eye contact</i> : Associated with depression
Posture ■ Does the patient assume a specific position? ■ Is posture erect?	■ Comfortably positioned, posture erect	■ <i>Slumped posture</i> : Associated with depression ■ <i>Defensive posture</i> : Associated with paranoia
Identifying Marks/Scars/Tattoos ■ Are there any obvious marks or scars?	■ No obvious marks or scars	■ <i>Scars</i> : May indicate self-mutilation or past suicide attempts
Appearance vs. Stated Age ■ Does the patient appear stated age, or younger or older?	■ Appears stated age	■ <i>Older appearance than stated age</i> : May be associated with depression
Behavior/Activity ■ Do you notice any unusual activity? ■ Is patient's behavior appropriate for the situation? ■ What is the patient's level of consciousness?	■ Calm, relaxed, no unusual behavior or movements ■ AAO × 3 (awake, alert, and oriented to time, place, person)	■ <i>Hyperactivity</i> : Associated with anxiety, mania, or stimulant abuse ■ <i>Hypoactivity</i> : Lethargic, thinking slowed; associated with depression, alcohol or drug abuse. ■ <i>Altered orientation</i> : May be seen with organic disorders or schizophrenia ■ <i>Agitation</i> : May be seen with dementia or delirium ■ <i>Psychomotor retardation, slow movements</i> : May be associated with depression ■ <i>Tremors</i> : May indicate drug/alcohol withdrawal ■ <i>Tics</i> : May be an adverse effect of psychotropic medications ■ <i>Unusual movements such as jaw/lip smacking</i> : May be associated with tardive dyskinesia, an adverse effect of antipsychotic medications ■ <i>Catatonia</i> : May be seen with schizophrenia ■ <i>Akathisia (restlessness)</i> : Seen with extrapyramidal adverse effect from antipsychotic medications ■ <i>Rigidity</i> : May be a sign of NMS

AREA/QUESTIONS TO ASK	NORMAL	ABNORMAL FINDINGS/RATIONALE
<p>Speech</p> <ul style="list-style-type: none"> What are the quality, tone, volume, fluency, and pace of speech? 	<ul style="list-style-type: none"> Quality and pace of speech normal with no exaggeration Fluent, pleasant tone 	<ul style="list-style-type: none"> <i>Slow speech</i>: Associated with depression <i>Rapid and pressured speech</i>: Associated with mania <i>Mumbling</i>: Seen with Huntington's chorea <i>Slurred speech</i>: Seen with alcohol intoxication <i>Suspicious tone</i>: Associated with paranoia <i>Volume</i>: Soft associated with depression <i>Poor fluency (mute/hesitation/latency of response)</i>: Less likely to talk with depression <i>Hesitancy</i>: Seen with mistrust or paranoia
<p>Attitude</p> <ul style="list-style-type: none"> What is the patient's attitude? Is it friendly? Hostile? 	<ul style="list-style-type: none"> Cooperative 	<ul style="list-style-type: none"> <i>Uncooperative</i>: Associated with paranoia <i>Warm/friendly/distant</i>: Seen with personality disorders <i>Suspicious/combative</i>: Seen with paranoia <i>Guarded/aggressive</i>: Seen with psychosis <i>Hostile/aloof</i>: Seen with psychosis <i>Apathetic</i>: Seen with depression
<p>Mood (Including Self-Report of Emotional State)</p> <ul style="list-style-type: none"> How does patient appear? How does the patient describe her or his mood? 	<ul style="list-style-type: none"> Appropriate for situation 	<ul style="list-style-type: none"> <i>Sad</i>: Associated with depression <i>Elated</i>: Associated with mania <i>Irritable/anxious</i>: Associated with anxiety <i>Fearful/guilty</i>: Seen with phobias <i>Worried/angry</i>: Seen with personality disorder <i>Hopeless</i>: Associated with depression <i>Labile</i>: Associated with mania <i>Mixed (anxious and depressed)</i>: Seen with depression
<p>Affect (Apparent Emotional State)</p> <ul style="list-style-type: none"> What is the patient's affect (facial expression)? 	<ul style="list-style-type: none"> Appropriate for situation 	<ul style="list-style-type: none"> <i>Flat affect</i>: Seen with schizophrenia <i>Blunted or diminished affect</i>: Seen with psychosis <i>Inappropriate/incongruent (sad and smiling or laughing)</i>: Associated with schizophrenia
<p>Thought Process</p> <ul style="list-style-type: none"> Can you follow the patient's thinking? Is it coherent and logical? 	<ul style="list-style-type: none"> Thought process intact Responds appropriately 	<ul style="list-style-type: none"> <i>Thought process disturbances</i>: Often seen with psychosis or organic brain disorders. Examples of thought process disturbances are: <ul style="list-style-type: none"> <i>Concrete thinking</i>: Unable to abstract; thinks in concrete terms <i>Circumstantiality</i>: Excessive, irrelevant detail, but eventually gets to the point

(continued)

Mental Status Assessment as It Relates to Mental Health/Illness (continued)

AREA/QUESTIONS TO ASK	NORMAL	ABNORMAL FINDINGS/RATIONALE
<p>Thought Content</p> <ul style="list-style-type: none"> ■ Does the patient's content of thought make sense and seem reality based? ■ Does the patient have thoughts about hurting self or someone else? 	<ul style="list-style-type: none"> ■ Thought content (what the patient is thinking) reality based 	<ul style="list-style-type: none"> ■ <i>Tangentiality</i>: Digresses from topic to topic, never getting to the point ■ <i>Loose association</i>: Loose connection between thoughts that are unrelated ■ <i>Echolalia</i>: Repetition of words spoken by another ■ <i>Flight of ideas</i>: Rapidly going from one topic to another ■ <i>Preservation</i>: Involuntary, excessive repetition of a single response to different questions ■ <i>Clang association</i>: Association of words by sound ■ <i>Blocking</i>: Draws a blank ■ <i>Word salad</i>: Combination of words that have no meaning ■ <i>Derailment</i>: Off track ■ <i>Delusions (grandiose/persecution/reference/somatic)</i>: Associated with psychosis ■ <i>Suicidal/homicidal thoughts</i>: Associated with depression, anxiety, or schizophrenia ■ <i>Obsessions</i>: Seen with OCD
<p>Perceptual Disturbances</p> <ul style="list-style-type: none"> ■ Does the patient have perceptual disturbances? Auditory disturbances? Visual disturbances? Olfactory disturbances? Tactile disturbances? 	<ul style="list-style-type: none"> ■ No hallucinations, illusions, or depersonalization. 	<ul style="list-style-type: none"> ■ <i>Paranoia</i>: Seen with schizophrenia ■ <i>Phobias</i>: Seen with anxiety disorders ■ <i>Magical thinking</i> (primitive form of thinking that thinking about something will make it happen) ■ Poverty of speech ■ <i>Visual hallucinations</i>: Often organic in nature ■ <i>Auditory hallucinations (commenting/discussing/commanding/loud/soft/other)</i>: Associated with schizophrenia ■ <i>Tactile hallucinations</i>: Seen with organic problems, drug abuse, or DTs ■ <i>Illusions (misperception of a real external stimulus)</i>: Common with dementia of Alzheimer's and schizophrenia



ALERT

If the patient is having homicidal thoughts, identify those toward whom those thoughts are directed.

AREA/QUESTIONS TO ASK	NORMAL	ABNORMAL FINDINGS/RATIONALE
<p>Memory</p> <p>Immediate</p> <ul style="list-style-type: none"> Can patient repeat objects named minutes before? <p>Remote</p> <ul style="list-style-type: none"> Can patient recall anniversaries, past important, historical events? <p>Insight (Awareness of the Nature of Illness) and Judgment</p> <ul style="list-style-type: none"> Does patient have insight into his or her problem? (Ask, "Can you tell me what the problem is?") Is judgment appropriate? (Ask the patient to respond to a hypothetical situation, such as, "If you were walking down the street and saw smoke coming from a window of a home, what would you do?") 	<ul style="list-style-type: none"> Immediate, recent, and remote memory intact No confabulation Insight and judgment intact and appropriate 	<ul style="list-style-type: none"> <i>Depersonalization</i> (altered perception or experience that causes temporarily loss of self or personal identity): Seen with panic disorder <i>Memory problems and confabulation</i>: Seen with organic, dissociative, and conversion disorders <i>Level of alertness</i>: Altered with substance abuse <i>Poor insight</i>: Seen with psychosis <i>Poor judgment</i>: Seen with psychosis <i>Poor impulse control</i>: Seen with OCD, psychosis, mania

Crisis

A **crisis** results from an acute event that stresses a person's resources and ability to cope; a crisis can also result from a perceived threat to self. Crises may be maturational or developmental and situational. An example of a **maturational crisis** would be a child becoming an adolescent. An example of a **situational crisis** might be the loss of a loved one. A situational crisis could also be a major disaster, such as a hurricane or a terrorist attack.

An event that may be a crisis for one person may not be for another. A stressful event causes disequilibrium for the person. It is the ability of the person to restore equilibrium that determines the outcome. Whether or not a person adapts to the crisis or not depends upon several factors: perception of event, situational supports, and coping mechanisms. Once you have performed a crisis assessment, you can then develop interventions that will help your patient deal with the crisis and regain equilibrium. (See *Crisis Assessment*.)

BATHE Technique

When you have only limited time to perform an assessment, it is important to keep the interview focused. The

BATHE technique helps the patient identify problems and coping strategies and is supportive of the patient. The acronym is representative of the interview's components (Stuart & Lieberman, 1993):

- **Background**—What is going on? What brought you here?
- **Affect**—How does this make you feel?
- **Trouble**—What troubles you most in this situation?
- **Handling**—How are you able to handle this situation/problem?
- **Empathy**—Empathize with client, shows an understanding of client's view of situation.



ALERT

The BATHE technique is not intended for use with patients with severe problems, such as patients who are suicidal or those who have suffered severe abuse.

Crisis Assessment

FACTOR	QUESTIONS TO ASK
Perception of event	<ul style="list-style-type: none"> ■ Can you tell me what has happened? Please be specific. ■ What caused the crisis? ■ How has this affected you? ■ How did this make you feel?
Supports	<ul style="list-style-type: none"> ■ Is there anyone I can call? ■ Do you live alone? ■ Do you have family or friends who support you? ■ Are you active in any religious or community groups?
Coping mechanisms	<ul style="list-style-type: none"> ■ How have you handled stressful events in the past? ■ Can you talk about the stressful event? ■ What do you do to relieve tension? Cry? Talk? Exercise? Use alcohol?

Source: Stuart, G., and Laraia, M. (2005). *Principles and Practices of Psychiatric Nursing*, ed. 8. Philadelphia: Mosby.

Assessing Depression

Depression poses a major health problem. Major depression has been identified by the World Health Organization as the leading cause of disability for adults in developed countries, such as the United States (Healthy People 2010). In any given year, approximately 6.5 percent of women and 3.3 percent of men will have a major depressive episode. The incidence of major depression is twice as great in women as in men. Identification of those at risk and accurate assessment of depression can lead to prompt intervention and promotion of mental health.

If you suspect that your patient is depressed, use the mnemonic SIGECAPS to easily recall and review the *DSM-IV* criteria for major depression:

- Sleep (increase/decrease)
- Interest (diminished)
- Guilty/low self-esteem
- Energy (poor/low)
- Concentration (poor)
- Appetite (increase/decrease)
- Psychomotor (agitation/retardation)
- Suicidal ideation

A depressed mood plus four or more SIGECAPS for 2 weeks or longer indicates a major depressive disorder. A depressed mood plus three SIGECAPS most days for 2 years or longer indicates **dysthymia**, which is a depressive neurosis with no loss of contact with reality.



HELPFUL HINT

Symptoms of dysthymia are similar to those of a major depressive disorder, but milder.

Suicide Assessment

In the United States, suicide is a major public health problem (Healthy People 2010). It is often the end result of mental illness. More women attempt suicide than do men, but men are four and a half times more likely to succeed at suicide than are women. If you suspect that your client is suicidal, perform a suicide assessment. It is important to identify those at risk for suicide and intervene promptly. (See *Performing a Suicide Assessment*.)



ALERT

Suicide attempts are more likely to occur as the patient's symptoms improve because the patient now has the energy to commit suicide.

Risk Factors for Suicide

- Past suicide of family member, close friend, or peer
- Impulsivity
- Mood disorders
- Substance abuse
- Recent loss of a spouse, partner, friend, or job
- Expressed hopelessness (patient sees no future)
- Social isolation (patient lives alone, has few friends or supports)
- Stressful life event
- Previous or current abuse (emotional, physical, sexual)
- Sexual identity crises/conflict
- Available lethal methods (guns)
- Legal issues/incarceration

Source: USPHS, HHS, 1999.

Performing a Suicide Assessment

ASSESSMENT AREA	QUESTIONS TO ASK
Feelings of hopelessness	<ul style="list-style-type: none"> How are you feeling? What are your plans for tomorrow? Next week? What do you have planned for the rest of the year?
	<div style="border: 1px solid black; padding: 5px;">  <p>HELPFUL HINT</p> <p>A key element in assessing for feelings of hopelessness is determining whether the patient is able to see a future with herself or himself in that future.</p> </div>
Suicidal ideations Plan for suicide Possessions	<ul style="list-style-type: none"> Have you ever had thoughts about hurting yourself? Do you have a plan to commit suicide? Have you been putting affairs in order? Have you been giving things away? Have you been contacting old friends?
Auditory hallucinations Lack of support network	<ul style="list-style-type: none"> Have you heard a voice telling you to hurt or kill yourself? How often do you communicate with friends and family? Have you found yourself withdrawing from friends and family? Who are your supports?
Alcohol or substance abuse Precipitating event	<ul style="list-style-type: none"> Do you drink or do drugs while alone? Have you recently experienced the death of a loved one or the loss of a job? Have you recently had a tragedy or disaster in your life?
	<div style="border: 1px solid red; padding: 5px;">  <p>ALERT</p> <p>For many people, holidays can be a precipitating event to a suicidal episode.</p> </div>
Media	<ul style="list-style-type: none"> Take note if the local media has reported on the suicide of a famous person or local teenager.

Source: Adapted from Pedersen, D. (2005) *PsychNotes*. Philadelphia: F.A. Davis.

Groups at Risk for Suicide

- Elderly persons who are isolated or widowed or who have experienced multiple losses
- Males who are widowed and without close supports
- Adolescents and young adults
- Persons with serious or terminal illness who become depressed or hopeless
- Persons with mood disorders, depression, and bipolar
- Persons with schizophrenia, either newly diagnosed or those experiencing auditory command hallucinations
- Persons who abuse drugs or alcohol, especially persons with a mental disorder
- Persons under stress (sometimes multiple stressors) with recent loss or losses

Assessing Substance Abuse

Substance use includes use of prescribed or OTC drugs, alcohol, caffeine, nicotine, steroids, and illegal drugs. **Abuse** and **addiction** exist when substance use has social, professional, or legal consequences. There are two types of substance abuse disorders, substance use disorders and substance-induced disorders. **Substance use disorders** include:

- Substance dependence:** Repeated use despite substance-related cognitive, behavioral, and psychological problems. Tolerance, withdrawal, and compulsive drug-taking can result.
- Substance abuse:** Recurrent, persistent substance use with significant adverse consequences during a 12-month period. Adverse consequences can include

repeated work absences, spousal arguments and fights, and driving while intoxicated offenses (DUIs) (APA, 2000).

Substance-induced disorders include:

- **Substance intoxication:** Overuse of a substance that results in a reversible, substance-specific syndrome; can be indicated by behavioral and psychological changes.

- **Substance withdrawal:** Symptoms differ upon the substances being used and develop upon their withdrawal (Pederson, 2005; APA, 2000).

If substance abuse is a concern with your client, it is important to ask the right questions and obtain an accurate history. If there is no time to obtain a detailed assessment, a focused assessment tool such as the CAGE questionnaire can also identify a possible substance abuse problem.

Substance Abuse History and Assessment Tool

1. When you were growing up, did anyone in your family use substances (alcohol or drugs)? If yes, how did the substance abuse affect the family?
 2. When (how old) did you use your first substance (e.g., alcohol, cannabis) and what was it?
 3. How long have you been using a substance(s) regularly? Weeks? Months? Years?
 4. What is your pattern of abuse?
 - a. When do you use substances?
 - b. How much and how often do you use?
 - c. Where are you when you use substances and with whom?
 5. When did you last use, what was it, and how much?
 6. Has substance use caused you any problems with family, friends, job, school, the legal system, other? If yes, describe.
 7. Have you ever had an injury or accident because of substance abuse? If yes, describe.
 8. Have you ever been arrested for a DUI because of your drinking or other substance use?
 9. Have you ever been arrested or placed in jail because of drugs or alcohol?
 10. Have you ever experienced memory loss the morning after substance use (can't remember what you did the night before)? Describe the event and feelings about the situation.
 11. Have you ever tried to stop your substance use? If yes, why were you not able to stop? Did you have any physical symptoms such as shakiness, sweating, nausea, headaches, insomnia, or seizures?
 12. Can you describe a typical day in your life?
 13. Are there any changes you would like to make in your life? If so, what are they?
 14. What plans or ideas do you have for making these changes?
 15. History of withdrawal:
- Other comments:

Source: Pedersen, D. (2005). *PsychNotes*. Philadelphia: F.A. Davis; and modified from Townsend, M. (2005). *Psychiatric Mental Health Nursing*, ed. 5. Philadelphia: F.A. Davis.

CAGE Questionnaire

- Have you ever felt you should Cut down on your drinking/drug use?
- Have people Annoyed you by criticizing your drinking/drug use?
- Have you ever felt bad or Guilty about your drinking/drug use?
- Have you ever had an Eye opener (use of alcohol or drugs first thing in the morning) to steady your nerves or get rid of a hangover?

A positive (yes) response to two or more questions suggests that there is an alcohol/substance problem.

Note: The need to cut down is related to tolerance (needing more substance for same effect), and the eye opener is related to withdrawal syndrome (reduction/cessation of substance).

Source: Ewing, J. A.: Detecting alcoholism: The CAGE questionnaire. *Journal of the American Medical Association* 252:1905–1907, 1984.

Abused Substances: Effects From Use and Withdrawal

SUBSTANCE	INTOXICATION	OVERDOSE	WITHDRAWAL
Depressants			
<ul style="list-style-type: none"> ■ Alcohol (booze, brew, juice, spirits) 	<ul style="list-style-type: none"> ■ Depressed cognitive functioning ■ Impaired psychomotor functioning ■ Decreased reaction time ■ Decreased balance and coordination ■ Decreased REM sleep 	<ul style="list-style-type: none"> ■ Unconsciousness ■ Respiratory depression ■ Coma ■ Death 	<ul style="list-style-type: none"> ■ Tremors ■ Diaphoresis ■ Anxiety ■ Hallucinations ■ Delusions ■ Increased pulse and blood pressure ■ DTs ■ Sleep disturbances ■ Insomnia ■ Hand tremor ■ Agitation ■ Nausea and vomiting ■ Anxiety ■ Tinnitus (with benzodiazepines) ■ Seizures ■ Cardiac arrest
<ul style="list-style-type: none"> ■ Sedatives, hypnotics, and anxiolytics, including barbiturates (barbs, beans, black beauties, blue angel, candy, downers, goof balls, BB, nebbies, reds, sleepers, yellow jackets, yellow) 	<ul style="list-style-type: none"> ■ Slurred speech ■ Labile mood ■ Inappropriate sexual behavior ■ Loss of inhibition ■ Drowsiness ■ Impaired memory 	<ul style="list-style-type: none"> ■ Hypotension ■ Nystagmus ■ Stupor ■ Cardiorespiratory depression ■ Coma ■ Death 	<ul style="list-style-type: none"> ■ Insomnia ■ Hand tremor ■ Agitation ■ Nausea and vomiting ■ Anxiety ■ Tinnitus (with benzodiazepines) ■ Seizures ■ Cardiac arrest
<ul style="list-style-type: none"> ■ Benzodiazepine (downers) 			
Stimulants			
<ul style="list-style-type: none"> ■ Amphetamines (A, AMT, bam, bennies, crystal, diet pills, dolls, eye-openers, pep pills, purple hearts, speed, uppers, wake-ups) 	<ul style="list-style-type: none"> ■ Euphoria ■ High energy ■ Impaired judgment ■ Anxiety ■ Aggressive behavior ■ Paranoia ■ Delusions 	<ul style="list-style-type: none"> ■ Ataxia ■ High temperature ■ Seizures ■ Respiratory distress ■ Cardiovascular collapse ■ Coma ■ Death 	<ul style="list-style-type: none"> ■ Depression ■ Agitation ■ Confusion ■ Vivid dreams followed by lethargy
<ul style="list-style-type: none"> ■ Cocaine (Bernice, bernies, big C, blow, C, Charlie, coke, dust, girl, heaven, jay, lady, nose candy, nose powder, snow, sugar, white lady) 	<ul style="list-style-type: none"> ■ Euphoria ■ Grandiosity ■ Sexual excitement ■ Impaired judgment ■ Insomnia ■ Anorexia ■ Nasal perforation (inhaled route) ■ Psychosis 	<ul style="list-style-type: none"> ■ High temperature ■ Seizures ■ Transient vasospasms (may cause myocardial infarction [MI], cerebrovascular accident [CVA], coma, death) 	<ul style="list-style-type: none"> ■ Fatigue ■ Depression ■ Anxiety ■ Suicidal behavior
<ul style="list-style-type: none"> ■ Crack (Conan, freebase, rock, toke, white cloud, white tornado) 			

(continued)

Abused Substances: Effects From Use and Withdrawal (continued)

SUBSTANCE	INTOXICATION	OVERDOSE	WITHDRAWAL
Marijuana			
<ul style="list-style-type: none"> ■ Cannabis (marijuana, hashish, Acapulco gold, Aunt Mary, broccoli, dope, grass, weed, grunt, hay, hemp, J, joint, joy stick, killer weed, pot, ragweed, reefer, smoke weed) 	<ul style="list-style-type: none"> ■ Euphoria ■ Intensified perceptions ■ Impaired judgment and motor ability ■ Increased appetite, weight gain ■ Sinusitis and bronchitis with chronic use ■ Anxiety, paranoia ■ Red conjunctiva 	<ul style="list-style-type: none"> ■ Extreme paranoia ■ Psychosis 	<ul style="list-style-type: none"> ■ None
Opiates			
<ul style="list-style-type: none"> ■ Heroin (H, horse, harry, boy, scag, shit, smack, stuff, white junk, white stuff) ■ Morphine ■ Hydromorphone ■ Meperidine ■ Codeine ■ Oxycodone ■ Opium ■ Methadone 	<ul style="list-style-type: none"> ■ Euphoria ■ Drowsiness ■ Impaired judgment ■ Constricted pupils 	<ul style="list-style-type: none"> ■ Dilated pupils ■ Respiratory depression ■ Seizures ■ Cardiopulmonary arrest ■ Coma ■ Death 	<ul style="list-style-type: none"> ■ Yawning ■ Insomnia ■ Anorexia ■ Irritability ■ Rhinorrhea ■ Muscle cramps ■ Chills ■ Nausea and vomiting ■ Feelings of panic and doom
Hallucinogens			
<ul style="list-style-type: none"> ■ Hallucinogens (LSD, DMT, Mescaline, acid, big D, blotter, blue heaven, cap, D, deeda, flash, L, mellow yellow, microdots, paper acid, sugar, ticket, yello) ■ Club drug, MDMA (ectasy) 	<ul style="list-style-type: none"> ■ Dilated pupils ■ Diaphoresis ■ Palpitations ■ Tremors ■ Enhanced perceptions of colors and sound ■ Depersonalization ■ Grandiosity ■ Euphoria ■ Muscle relaxation 	<ul style="list-style-type: none"> ■ Panic ■ Psychosis with hallucinations ■ Cerebral damage ■ Death ■ Confusion ■ Hallucinations ■ Severe anxiety ■ Hypertension ■ Seizures ■ High temperature 	<ul style="list-style-type: none"> ■ None ■ Psychological dependence can cause depression, flashbacks
Phencyclidine			
<ul style="list-style-type: none"> ■ Phencyclidine (PCP, angel dust, DOA, dust, elephant, hog, peace pill, supergrass, tictac) 	<ul style="list-style-type: none"> ■ Impulsive behavior ■ Impaired judgment ■ Belligerent, violent behavior ■ Ataxia ■ Muscle rigidity ■ Nystagmus ■ Hypertension 	<ul style="list-style-type: none"> ■ Hallucinations ■ Psychosis ■ Seizures ■ Respiratory arrest ■ Death 	<ul style="list-style-type: none"> ■ None

SUBSTANCE	INTOXICATION	OVERDOSE	WITHDRAWAL
Inhalants <ul style="list-style-type: none"> ■ Gasoline, glue, aerosol sprays, paint thinners (spray, rush, bolt, huffing, bagging, sniffing) 	<ul style="list-style-type: none"> ■ Numbness or diminished response to pain ■ Euphoria ■ Impaired judgment ■ Blurred vision ■ Unsteady gait 	<ul style="list-style-type: none"> ■ Psychosis with hallucinations ■ Cardiac arrhythmias ■ Central nervous system depression ■ Coma ■ Cerebral damage ■ Death ■ None 	<ul style="list-style-type: none"> ■ None
<ul style="list-style-type: none"> ■ Nicotine (Cigarettes, cigars, bidis, kreteks, pipe tobacco, chewing tobacco, snuff, nicotine gum or patches) 	<ul style="list-style-type: none"> ■ Sense of anxiety reduction ■ Relief from depression ■ Satisfaction 		<ul style="list-style-type: none"> ■ Insomnia ■ Depression ■ Irritability ■ Anxiety ■ Poor concentration ■ Increased appetite

Source: Gorman, L., Raines, M., & Sultan, D. (2005). *Psychosocial Nursing for General Patient Care*, ed. 2. Philadelphia: F.A. Davis.



Case Study Findings

Your general health survey, physical assessment, and mental status assessment of Ms. Baker have concluded. The following are your findings:

■ Pertinent physical findings

- Weight gain of 10 pounds over past month
- Poor eye contact
- Slow to respond
- Disheveled appearance
- Slouched position
- Crying
- Difficulty staying focused
- Constipation



CRITICAL THINKING ACTIVITY # 4

How might you explain Ms. Baker's physical findings?

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Ms. Baker. Use this information to identify a list of nursing diagnoses. Some possible nursing diagnoses are provided below. Cluster the supporting data.

1. Risk for Suicide
2. Ineffective Individual Coping
3. Fatigue

Identify any additional nursing diagnoses.



Case Study Analysis and Plan

Now that your examination of Ms. Baker is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.



CRITICAL THINKING ACTIVITY # 5

Now that you have identified appropriate nursing diagnoses for Ms. Baker, select a nursing diagnosis and develop a plan of care.

RESEARCH TELLS US

Worldwide, mental health problems, with depression and anxiety being the most frequent, occur in approximately 24 percent of patients in primary care (WHO, 2005). Mood disorders with increasing rates of depression and anxiety may be associated with physical illness. Deteriorating physical health and cognitive functioning that is often associated with aging is a major risk factor for developing depression later in life.

The purpose of the following study was to examine staff contact and input with mental health problems and to determine their experience, training, and attitudes to mental health problems. A cross-sectional design was used in the United Kingdom. A 40-item questionnaire was mailed to over 300 staff members in primary care. The return rate was 66 percent ($n = 217$, 95 percent confidence interval, with two-thirds of the respondents RNs). The questionnaire focused on depression in identifying mental problems, interventions, and mental health training. Demographics were obtained to describe the sample. The Depression Attitude Questionnaire (DAQ), a 20-item instrument, was used to assess staff attitude toward depression.

The findings reported that 16 percent of their patients had associated mental health problems, with dementia, depression, and anxiety being the most common. The staff reported a willingness to develop an understanding and skills needed to address mental health problems with patients, but reported little training in the past 5 years that addressed this issue. The staff, as revealed by the DAQ, was optimistic about treating depression.

The study supports the need for mental health training in primary care as evidenced by the limited detection and treatment of mental health problems. The staff was willing and eager to develop the knowledge and skills to assess and treat mental health problems. The areas identified included recognition of mental disorders, anxiety management, crisis intervention, and pharmacologic treatment for depression.

Although this study was limited to the United Kingdom, the problem is universal. A better understanding of assessment and treatment of mental health problems would provide a holistic approach to meeting the patient's needs.

Source: Haddad, M., Plummer, S., Taverner, A., Gray, R., Lee, S., Payne, F., and Knight, D. (2005). District nurses' involvement and attitudes to mental health problems: A three-area cross-sectional study. *Journal of Clinical Nursing*, 14(8):976–985.

HEALTH CONCERNS

Healthy People 2010 has identified specific objectives that address mental health/disorders.

Improving Mental Health

- Improve mental health and ensure access to appropriate, quality mental health services.
- Reduce the suicide rate. Target: 5 per 100,000 from 11.3 per 100,000 in 1998.
- Reduce the rate of suicide attempts by adolescents. Target: 1 percent over 12 months from 2.6 percent from grades 9 through 12 in 1999.
- Reduce the proportion of homeless adults who have serious mental illness. Target: 19 percent from 25 percent age 18 and older in 1996.
- Increase the proportion of persons with serious mental illness who are employed. Target: 51 percent from 43 percent age 18 and older in 1994.

Improving Treatment

- Reduce the relapse rates for persons with eating disorders.
- Increase the number of persons seen in primary healthcare who receive mental health screening and assessment.
- Increase the proportion of children with mental health problems who receive treatment.

- Increase the proportion of juvenile justice facilities that screen new admissions for mental health problems.
- Increase the proportion of adults with mental disorders who receive treatment.
- Increase the proportion of persons with co-occurring substance abuse and mental disorders who receive treatment for both disorders.

Improving State and Local Involvement

- Increase the proportion of local government with community-based jail diversion programs for adults with serious mental illness.
- Increase the number of states (including the District of Columbia) that track consumers' satisfaction with the mental health services they receive.
- Increase the number of states and territories (including the District of Columbia) with an operational mental health plan that addresses cultural competence.
- Increase the number of states and territories (including the District of Columbia) with an operational mental health plan that addresses mental health crisis interventions, ongoing screening, and treatment services for elderly persons.

Assessment of Common Mental Health Problems

MENTAL HEALTH DISORDER

Anxiety

- Diffuse feelings of apprehension with feelings of uncertainty, helplessness
- Ranges from mild to panic state that can be paralyzing



HELPFUL HINT

Anxiety differs from fear in that fear has an identifiable source, anxiety may not.

Types: (DSM-IV)

Panic Disorders

- Extreme, overwhelming anxiety in response to real or perceived life-threatening situation
- Can lead to phobias, avoidance, and agoraphobia

Phobias

- Unreasonable fear response to a specific object or situation
- Causes anxiety
- Person realizes fear is unreasonable

Social Anxiety Disorder

- Fear of social or performance situations that may cause embarrassment

OCD

- Recurrent thought or ideas (obsession)
- Action person cannot refrain from doing (compulsion)
- Obsession and compulsion interfere with social and occupational functioning
- Recognizes thoughts and behaviors are unreasonable

PTSD

- Traumatic event
- Threat of harm or death, actual death, and helplessness
- Re-experiences event (flashback)
- Hypervigilant
- Recurring nightmares
- Anniversary reactions related to trauma
- Persistent anxiety
- Acute < 3 mo, chronic > 3 mo, delayed > 6 mo

Acute Stress Disorder

Anxiety, dissociation, and other symptoms after exposure to recent stressors

Anxiety Disorder related to Medical Condition

Generalized anxiety disorder

- Excessive worry and anxiety for at least 6 mo
- Difficult to control worry
- Hypervigilant

Substance-induced anxiety

ASSESS FOR

- Restlessness
- On edge
- Easily fatigued
- Concentration problems
- Irritability
- Sleep disturbances
- Dizziness
- Palpitations
- Hot/cold flashes
- Tightness of chest
- Nausea
- Decrease appetite
- Abdominal pain

Affect

- Nervous
- Anxious
- Fearful

Physical Findings

- Sweating
- Tremors, rigidity, spasms
- Dilated pupils
- Increased respirations
- Wheezes due to bronchial spasms
- Tachycardia, increased blood pressure
- Diarrhea
- Vomiting
- Increased reflexes
- Muscle tension
- Pacing, clumsy movements
- Confusion
- Easily distracted
- Short attention span

(continued)

Assessment of Common Mental Health Problems (continued)

MENTAL HEALTH DISORDER

ASSESS FOR

Personality Disorders

- Pattern of relating and perceiving the world that is inflexible and maladaptive

Types:

Cluster A includes paranoid, schizoid, schizotypal personality disorders

- *Paranoid*: Distrustful and suspicious of others, preoccupied with doubts of loyalty, holds grudges, unwilling to forgive, quick to react and counter perceived insults
- *Schizoid*: Detached from social interaction, restricted expression of emotions, lacks desire for intimacy, emotionally cold
- *Schizotypal*: Social and interpersonal deficits, odd beliefs or magical thinking, perceptual alterations, odd or eccentric behavior

Cluster B includes antisocial, borderline, histrionic, and narcissistic personality disorders

- *Antisocial behavior*: Pattern of disregard and violation of rights of others
- *Borderline*: Unstable behavior with changes in relationships, self-image, and mood
- *Histrionic*: Excessive expression of emotion and attention-seeking behavior
- *Narcissistic*: Inflated sense of self, need for attention and admiration, no concern for others

Cluster C includes avoidant, dependent, and obsessive compulsive personality disorders

- *Avoidant*: Avoids social interaction for fear of criticism and feelings of inadequacy
- *Dependent*: Needs to be taken care of, submissive, fear of separation, low self-confidence, difficulty making decisions and voicing disagreement
- *Obsessive compulsive*: Rigid way of functioning, excessive detail or controlling ways



HELPFUL HINT

Obsessive compulsive personality disorder differs from OCD in that the person with obsessive compulsive personality disorder has no problem with behavior, whereas the person with OCD is anxious and wants to change

Psychotic Disorders

Schizophrenia

- Caused by neurobiological factors; influenced by social and environmental factors

Four "A"s of Schizophrenia

- Inappropriate Affect

Cluster A

- Distrustful, emotionally detached, eccentric personalities

Cluster B

- Disregard for others, unstable and intense interpersonal relationships, excessive attention seeking, entitlement issues with lack of empathy for others

Cluster C

- Avoider of social situations, clinging, submissive personality; and person preoccupied with details, rules, and order

Borderline Personality Disorder

- Pattern of unstable relationships
- Fear of abandonment
- *Splitting*: Idealize and devalue (love/hate)
- Impulsive (in two areas: sex, substance abuse, binge eating, reckless driving)
- Suicidal gestures/self-mutilation
- Intense mood changes lasting a few hours
- Chronic emptiness
- Intense anger
- Transient paranoid ideation

Positive Symptoms = Excessive function/distortion in behavior

- Delusions
- Hallucinations (auditory/visual)
- Hostility
- Disorganized thinking and behavior

MENTAL HEALTH DISORDER

- Loose Association
- Autistic thoughts
- Ambivalence

Mood Disorders

- Extremes of moods (mania or depression)

Types

Depressive

- *Major depressive disorder (unipolar depression)*: At least 2 wk of depression/loss of interest and four additional symptoms with one or more major depressive episodes.
- *Dysthymic disorder*: On-going, low-grade depression of at least 2 years' duration for more days than not and does not meet the criteria for major depression
- *Depressive disorder not otherwise specified (NOS)*: Does not meet criteria for depressions described above.

Bipolar Disorders

- *Bipolar I disorder*: One or more manic or mixed episodes with a major depressive episode
- *Bipolar II disorder*: One or two major depressive episodes and at least one hypomanic episode
- *Cyclothymic disorder*: At least 2 years of hypomanic episodes that do not meet the criteria for other disorders
- *Bipolar disorder NOS*: Does not meet any of the other bipolar criteria

Eating Disorders

Types

Anorexia Nervosa

- Terrified of gaining weight

ASSESS FOR

Negative Symptoms = Deficits in behavior, such as reduced function or self-care deficits

At least for 1 mo, two or more:

- Delusions
- Hallucinations
- Disorganized speech
- Disorganized behavior
- Negative symptoms
- Functional disturbances at work or school disturbance continues for 6 mo



HELPFUL HINT

- Auditory hallucinations usually begin distant and soft, then become louder, but become softer and distant as client's condition improves.
- In North America, most hallucinations auditory. Not likely to have both auditory and visual together.

Depressive

Depressed mood or loss of interest for at least 2 wk and five or more of:

- weight loss/gain
- insomnia or hypersomnia
- psychomotor agitation or retardation
- fatigue
- worthless feelings or inappropriate guilt
- problem concentrating
- recurrent thoughts of death



ALERT

When working with patients who have depression, it is important to identify any suicidal/homicidal ideations.

Mania

Persistent elevated, irritable mood for 1 wk or more, plus three or more (irritable, four or more):

- high self-esteem
- decreased sleep
- increased talking/pressured speech
- racing thoughts/flight of ideas
- distractibility
- extreme goal-directed activity
- excessive buying/sex/business investments (painful consequences)

Anorexia Nervosa

- Emaciated appearance
- Below normal weight
- Hair loss, dry skin

(continued)

Assessment of Common Mental Health Problems (continued)

MENTAL HEALTH DISORDER	ASSESS FOR
<ul style="list-style-type: none"> ■ Weight below minimally accepted (weight < 85% of what would be expected for age and height) ■ Disturbed self-perception of size and shape of body; even though underweight, still fears becoming overweight ■ Self-esteem and self-evaluation based on weight ■ Introverted, socially isolated, high achiever <p>Bulimia Nervosa</p> <ul style="list-style-type: none"> ■ Recurrent binge eating of large amounts of food over short period ■ Lack of control ■ Self-induced vomiting, use of laxatives, purging, fasting, excessive exercising ■ Weight normal, under-/overweight ■ Fluid and electrolyte imbalance ■ Impulsive, acting out, more histrionic (pervasive, excessive expression of emotions and attention-seeking behaviors) 	<ul style="list-style-type: none"> ■ Amenorrhea ■ Pedal edema <p>Bulimia</p> <ul style="list-style-type: none"> ■ Normal or overweight ■ Weight fluctuations ■ Dehydration ■ Hoarseness ■ Parotid gland enlargement (chipmunk facies) ■ Tooth enamel erosion ■ Finger or pharynx bruising

Sources: Pedersen, D. (2005). *PsychNotes*. Philadelphia: F.A. Davis; Gorman, L., Raines, M., and Sultan, D. (2005). *Psychosocial Nursing for General Patient Care*, ed. 2. Philadelphia: F.A. Davis; and *DSM-IV* criteria adapted from American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*, ed. 4, Text Revision. Washington, DC: American Psychiatric Association, 2000.


S U M M A R Y

- Holistic nursing care includes assessment of the psychological health of your patient. The developmental level and cultural background of your patient needs to be considered when assessing your patient's mental health.
- When obtaining a health history and physical assessment, look for clues that reflect the mental health of your patient.
- A thorough mental health assessment includes a detailed mental status assessment. Specific mental health assessment instruments are available to further assess the mental health needs of your patient.
- Identify supports and resources and make referrals as needed to meet the mental healthcare needs of your patient.

R E V I E W Q U E S T I O N S

1. A possible side effect of lithium that needs to be evaluated through lab tests is:
 - a. Hypothyroidism
 - b. Tardive dyskinesia
 - c. Abdominal rigidity
 - d. Parkinsonism
2. Which neurotransmitter is associated with anxiety and anxiety states?
 - a. Dopamine
 - b. Acetylcholine
 - c. Gamma-aminobutyric acid (GABA)
 - d. Serotonin
3. Mary Brown, age 75, is admitted to the hospital with change in mental status, sudden onset of confusion. This behavior is associated with:
 - a. Delirium
 - b. Dementia
 - c. Psychosis
 - d. Personality disorder
4. What is the most common mental health problem experienced by older adults?
 - a. Personality disorder
 - b. Psychosis
 - c. Bipolar disorder
 - d. Depression

5. John Bowes, age 36, has a history of schizophrenia with hallucinations. Which type of hallucination is most frequently associated with schizophrenia?
 - a. Auditory
 - b. Visual
 - c. Tactile
 - d. Olfactory
6. Which thought process is used by alcoholics who have memory loss?
 - a. Confabulation
 - b. Flight of ideas
 - c. Loose association
 - d. Word salad
7. Your patient complains of “feeling down and can’t shake it.” Your assessment includes a medication history. Which of the following medications is associated with depression?
 - a. Steroids
 - b. Beta blockers
 - c. Digitalis
 - d. Antibiotics
8. Your patient has glaucoma. Which psychotropic drug would be contraindicated for use in patients with glaucoma?
 - a. Lithium
 - b. Tricyclics
 - c. Monoamine oxidase (MAO) inhibitors
 - d. Antipsychotics
9. Your patient has bipolar disorder and is in a manic state. Her thought process most likely would be:
 - a. Concrete thinking
 - b. Circumstantiality
 - c. Flight of ideas
 - d. Loose association
10. Susan Stewart, age 69, panics when driving on congested highways, feels as if she is “outside her body,” as if someone else is driving her car. This is an example of:
 - a. Illusion
 - b. Visual hallucination
 - c. Preservation
 - d. Depersonalization



Teaching the Patient

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Discuss the purpose of teaching
- Identify subjective assessment data that would influence teaching your patient
- Identify objective assessment data that would influence teaching your patient
- Discuss developmental factors that would influence teaching your patient
- Discuss cultural factors that would influence teaching your patient
- Identify teachable moments in caring for your patient
- Incorporate assessment data into a teaching plan for your patient



Introducing the Case Study

Lucas Hernandez, a 21-year-old college junior, arrives for his annual checkup. He has no health complaints, but he is concerned about juggling school and work responsibilities. Lucas is studying to become a teacher and also moonlights as a bartender 4 nights a week, from 7 P.M. to 3 A.M. He usually sleeps from 6 A.M. to noon on workdays, but he is about to start student teaching and must be at school from 7 A.M. to 4 P.M. He says he plans to sleep in two shifts.

Lucas's past health history, family history, and review of systems are negative. His psychosocial profile reveals that he is gay and has been living with a partner for 2 years. He drinks only occasionally and is a vegetarian who includes eggs and milk in his diet. His physical assessment findings are also negative. His immunizations are current, but he has not been immunized against hepatitis B, and he declines human immunodeficiency virus (HIV) testing, stating that he is in a monogamous relationship.

What kind of teaching plan would you develop for Lucas? Before you can begin to develop this plan, you need to consider how his assessment findings will affect his learning needs.

Tips for Effective Teaching

- Polish your verbal and nonverbal communication skills.
- Establish a relationship that is holistic, interactive, and focused on the individual.
- Identify factors that will affect learning needs and compensate for them.
- Involve the individual in identifying his or her learning needs and planning your teaching strategy.
- Be sure that the individual values learning and that you both agree on her or his learning needs.
- Engage as many of the person's senses as possible—he or she will be more likely to learn.
- Build on what the person already knows.
- Consider the person's lifestyle and resources available to her or him to develop a practical plan.
- Be realistic in your expectations, or your patient's goals and outcomes will not be achieved.
- Remember: No matter how well you teach, you cannot make someone learn.

Introduction to Teaching the Patient

Teaching is a key nursing intervention at all levels of healthcare. As a nurse, you teach individuals, families, groups, and communities. At the primary level, you teach healthy people to reinforce desired behaviors, promote continued wellness, and anticipate and prevent potential health problems. At the secondary level, you teach ill people to help them cope with their immediate illness, restore and promote optimal wellness, and work most effectively with their healthcare team. At the tertiary level, you teach people or caregivers how to manage chronic illnesses and maintain an optimal level of health.

The goals of teaching are to:

- Promote health and wellness.
- Prevent disease and illness.
- Restore and maintain health and wellness.
- Promote coping.

Developmental, Cultural, and Ethnic Variations

Developmental factors influence learning need and teaching approaches. Cultural factors reflect the patient's beliefs and values and affect your patient's learning needs and teaching approaches.

Developmental Level

To develop an effective teaching plan, you must identify the individual's developmental level and understand normal growth and developmental tasks. Determine the person's developmental readiness to learn. Be aware of how he or she compares with developmental standards physically, cognitively, emotionally, and socially. Anticipate teaching content related to current and future growth and development issues.

Provide opportunities for learners of all ages to ask questions or voice concerns. For example, if the person is a young adolescent, you might say, "People's bodies and feelings go through many changes from ages 9 to 20. Are there any particular changes that you would like to discuss?" Or, if she or he is a parent introducing a new baby to an older sibling, you might say, "Older siblings have conflicting feelings about a new baby. How have you prepared your older child for the baby's homecoming?"

Also consider the cognitive and physical development of the individual. For example, teaching a newly diagnosed diabetic who is 5 years old about diabetes

would be very different from teaching a newly diagnosed 25-year-old. If you are teaching skills, remember to consider the person's motor development level. Fine-motor-skill development is needed to give self-injections, and young and older patients may have difficulty mastering this.

Culture and Ethnicity

Culture refers to characteristics that create common patterns of living or belief among people. Working at the same company or having similar incomes creates common patterns of living just as much as ethnic origin does. Identifying a person's cultural and religious background will help you approach him or her in a respectful manner, more accurately interpret body language and oral communication, be aware of taboos or practices that could affect learning or teaching, and be aware of the priority that wellness has in his or her life. For example, teaching a Cuban American patient about low-fat weight-reduction diets, knowing that their diets are high in calories, starches, and saturated fats, would be important in developing a successful teaching plan.

Performing the Assessment

Before you can develop a teaching plan for an individual, you need to assess her or him. Collect both subjective and objective data. Identify the person's learning needs, factors that could affect learning, and her or his stressors, strengths, and deficits. Then plan your teaching around them.

Health History

Be mindful to reflect carefully on everything you learn during the health history. It provides you with the background data needed to develop an effective teaching plan and tells you who the learner is. As you gather information in the following categories, keep asking yourself, "How will this affect my teaching?"

Biographical Data

Asking for biographical information is a good place to start when developing a teaching plan. Age and gender allow comparison with standards for demographic cohorts. Age also helps identify developmental stage and appropriate teaching level. Occupation and address identify environmental concerns that may affect learning and community learning as well as resources near the person's home or job. Knowledge of religion and culture identifies beliefs that may influence your teaching plan.



HELPFUL HINT

Do not assume that the person's educational level corresponds with knowledge or understanding.

Current Health Status

Knowing the person's current state of health and reason for seeking care will help you identify whether significant or unexpected changes are occurring in his or her health status, whether he or she has a preventive or crisis focus, and whether he or she understands the health problem and perceives it as serious. If he or she has an acute problem, you may need to address it before you begin teaching. If the person has major teaching needs but does not consider his or her problem serious, you need to help him or her understand the severity of the illness and the necessity of a teaching plan. Otherwise, learning will be difficult, if not impossible.

Having the person describe the problem in her or his own words gives you her or his perception of it and may reveal teaching needs. For example, an older woman with congestive heart failure (CHF) may say that her problem is "swollen ankles" and make no association with the heart.

Other factors that affect learning include fear, anxiety, and medications. For example, a person who is depressed or anxious will have trouble concentrating. Or the person may be taking medications that make him or her drowsy and limit the amount of time you can teach per session.

Ask the person: "Is there anything going on in your life now that makes it particularly easy or difficult for you to manage your health or practice good health habits?" This question alerts you to current stressors, strengths, and deficits that could affect learning. It allows you to:

- Plan teaching to accommodate for current issues.
- Be sensitive to learners' priorities in planning.
- More effectively assist learners to prioritize learning needs for short- and long-term time periods.

Past Health History

Questions about the person's past health history shed light on her or his learning abilities, new learning needs, or misconceptions that need clarification. Ask her or him: "Do you have any medical problems?" If the answer is yes, ask: "Did you receive instructions from healthcare providers? What were they?" Multiple health problems may affect the person's ability to learn. Negative experiences with healthcare providers may make her or him less receptive to learning.

Family History

The family history helps identify health risks that, in turn, identify teaching needs. Ask your patient: “Are there any major medical problems in your family?” This question identifies genetic or familial risk for health problems and points out the need for further education. Past experience with specific illnesses also may influence the person’s perception of the disease.

Review of Systems

The review of systems (ROS) helps you recognize changes in one body system that may influence other body systems. As you perform the ROS, identify any areas that may affect learning.

Review of Systems

AREA/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>General Health Survey</p> <ul style="list-style-type: none"> How have you been feeling? 	<ul style="list-style-type: none"> Identifies problems that may interfere with learning and alerts you to adjust your teaching plan. For example, depression or anxiety may interfere with learning. Fatigue may limit the person’s tolerance, so short teaching sessions may be more effective.
<p>Integumentary</p> <ul style="list-style-type: none"> Do you have any problems with your skin, hair, or nails? <p>Head, Eyes, Ears, Nose, and Throat (HEENT)</p> <ul style="list-style-type: none"> Do you have headaches? Do you have hearing problems? Do you wear a hearing aid? When was your last hearing test? Do you have vision problems? Do you wear glasses or contact lenses? When was your last eye examination? 	<ul style="list-style-type: none"> Helps identify problems that can affect learning. Helps identify teaching needs. Headaches can interfere with learning. Vision or hearing problems need to be addressed when developing an effective teaching approach. Vision problems may affect ability to perform tasks requiring psychomotor skills, such as self-administration of insulin.
<p>Respiratory</p> <ul style="list-style-type: none"> Do you have a history of lung disease or breathing problems such as shortness of breath (SOB)? 	<ul style="list-style-type: none"> SOB may limit tolerance, so short teaching sessions may be more effective.
<p>Cardiovascular</p> <ul style="list-style-type: none"> Do you have any history of cardiovascular disease? 	<ul style="list-style-type: none"> Fatigue associated with chronic cardiac problems such as CHF limits tolerance, so short teaching sessions may be more effective.
<p>Gastrointestinal/Genitourinary</p> <ul style="list-style-type: none"> Do you have any gastrointestinal (GI) or genitourinary (GU) problems? 	<ul style="list-style-type: none"> GI complaints, such as nausea or diarrhea, and GU complaints, such as urinary urgency and frequency, may affect learning.
<p>Neurological</p> <ul style="list-style-type: none"> Do you have any memory problems, numbness, or tingling? 	<ul style="list-style-type: none"> Forgetfulness or difficulty concentrating affect ability to learn. Numbness and tingling may affect ability to learn psychomotor skills, such as self-administration of insulin.
<p>Musculoskeletal</p> <ul style="list-style-type: none"> Do you have a history of arthritis, joint pain, or weakness? 	<ul style="list-style-type: none"> Joint pain, deformity, stiffness, or weakness may affect ability to perform tasks requiring psychomotor skills.
<p>Endocrine</p> <ul style="list-style-type: none"> Do you have thyroid disease or diabetes? 	<ul style="list-style-type: none"> You may need to adjust the climate of the teaching environment if the person has heat or cold intolerance.
<p>Immune/Hematologic</p> <ul style="list-style-type: none"> Do you have cancer, HIV, or a bleeding disorder? 	<ul style="list-style-type: none"> History of immunosuppression may prohibit group teaching programs. Anemias can cause fatigue.

Psychosocial Profile

The psychosocial profile can provide clues that are vital

to preparing a successful teaching plan, including areas in which health education is needed.

Psychosocial Profile

AREA/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Health Practices and Beliefs/Self-Care Activities

- What are you doing now to stay healthy?
- Why? How did you learn to do it?
- How do you usually learn about health matters (e.g., nurse, physician, pharmacist, newspaper, radio, TV, friends, family member, Internet)?

Typical Day

- How do you usually spend your day?
- How does a typical week go for you?

Nutritional Patterns

- Are you on a special diet?
- What times do you usually eat?

Activity and Exercise Patterns

- What activities do you do involving your body (e.g., hobbies, playing instruments, tasks at home or work)?

Sleep/Rest Patterns

- When do you have the most energy? When do you feel the most relaxed? When are you most tired or uncomfortable?

Personal Habits

- Do you use tobacco, drugs, or alcohol?

Occupational Health Patterns

- Does your place of employment offer health and wellness programs?

Socioeconomic Status

- Does your health insurance program offer or cover health programs? Are you on a fixed income?

Environmental Health Patterns

- What is your usual means of transportation?

Roles and Relationships

- How do you balance your role responsibilities?

Social Supports

- Will other people be coming to class with you?

- Identifies patient's focus (preventive or crisis). Identifies current level of self-care activities.
- Identifies why learner practices and adheres to existing health behaviors.
- Identifies usual sources of health information.

- Identifies times of day or week when learners are most rested and comfortable and when they prefer to learn.

- Consider special diets or special eating times when scheduling teaching sessions.
- Hunger can affect learning.

- If possible, incorporate patient's activities into your teaching plan.

- Helps identify best times for teaching/learning.

- Use of drugs and alcohol can be a barrier to learning and identify additional learning needs.

- Identifies resources within the workplace.

- Makes you aware of time, responsibility, and economic barriers or assets. Identifies resources so that you can make appropriate referrals. Health programs or coverage may be available through health insurance programs.

- Lack of transportation may be a barrier to learning. May need to identify alternative means, such as public transportation.

- Learning must fit into your patient's current role responsibilities.

- Prepares you for other family members or supporters who may attend teaching session.

(continued)

Psychosocial Profile (continued)

AREA/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Are there family members or friends who need to know what you are learning? ■ Are there relationships in your life that will make it easier or harder for you to make changes? <p>Stress and Coping</p> <ul style="list-style-type: none"> ■ Do you have any stress in your life currently? ■ How do you usually deal with stress? 	<ul style="list-style-type: none"> ■ Alerts you to people who may reinforce or impede the patient's learning/behavior change. ■ Anticipates and addresses logistical barriers to learning, such as having unreliable transportation or being responsible for dependent care. ■ Be aware of current stressors, strengths, and deficits that could affect learning.

The Physical Assessment

Once you have completed the health history, identify physical assessment findings that could affect learning.

Further investigate areas identified in the health history that might affect learning. As you perform the head-to-toe assessment, ask open-ended questions to determine the learner's baseline knowledge of your teaching content.

Physical Assessment

AREA/ASSESSMENT	ABNORMAL FINDINGS/RATIONALE
<p>General Health Survey</p> <ul style="list-style-type: none"> ■ Inspect level of consciousness, posture, signs of acute distress. ■ Measure vital signs. <p>Integumentary</p> <ul style="list-style-type: none"> ■ Inspect for color changes such as cyanosis, pallor, jaundice. <p>HEENT</p> <ul style="list-style-type: none"> ■ Test visual acuity and hearing. <p>Respiratory</p> <ul style="list-style-type: none"> ■ Measure respiratory rate. ■ Auscultate lungs. <p>Cardiovascular</p> <ul style="list-style-type: none"> ■ Measure blood pressure and pulse rate. <p>Abdominal</p> <ul style="list-style-type: none"> ■ Palpate for tenderness and organomegaly. <p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Inspect joints for deformity. ■ Test muscle strength, ROM. <p>Neurological</p> <ul style="list-style-type: none"> ■ Test balance and coordination, fine-motor skills, and senses. 	<ul style="list-style-type: none"> ■ If patient is in acute distress or has altered level of consciousness, postpone teaching until his or her condition improves. ■ Cyanosis may indicate hypoxia, and jaundice may indicate liver disease. Both conditions may alter mental status. ■ Pallor is associated with anemia, which causes fatigue. ■ Identify impaired vision or peripheral field and hearing deficits and adjust teaching plan to meet person's needs. ■ Type and severity of deficit influence teaching strategies and tools. ■ Hypoxia can cause confusion and decrease ability to concentrate. ■ Cardiovascular problems such as CHF are often associated with fatigue. ■ Liver disease can result in a confused mental state and weakness. ■ Muscle weakness, joint deformity, and pain may affect ability to learn motor skills. ■ Problems in these areas may make learning motor skills difficult.



Case Study Analysis and Plan

Before you identify learning needs and develop a plan, you need to be aware of your own learning preferences and values so that you will not impose them on your patient. For example, if you are an independent learner, a self-study approach may be an effective learning method for you, but if your patient is a dependent learner, this approach is not likely to work. Now that you know what factors to consider when devising a teaching plan, write a plan for Lucas based on his learning needs related to sleep.



HELPFUL HINT

As a nurse, you may feel that you never have enough time to teach, but remember that teaching can be informal as well as formal. Every time you interact with your patient presents a potential for teaching.



CRITICAL THINKING ACTIVITY

Can you identify any additional learning needs for Lucas?



HELPFUL HINT


To assess motor skills, observe the learner dressing, undressing, moving around the room, or performing a task such as writing.

S U M M A R Y

- Like the nursing process, the teaching process begins with assessment.
- During the health history, consider cultural, developmental, and psychosocial factors.
- During the physical examination, assess the person for factors that could affect learning and identify both deficits and strengths.
- After you complete your assessment, identify the person's learning needs, readiness to learn, learning preferences, and learning styles. Then identify methods and strategies needed to develop and implement a successful teaching plan.
- Implement your plan, evaluating and revising as needed. Be sure to document every step of the teaching process.

R E V I E W Q U E S T I O N S

1. You are teaching your patient about wound care. She has hyperthyroidism. How might you adapt your teaching plan to meet her needs?
 - a. Adjust the temperature of the room
 - b. Speak louder so that she can hear you
 - c. Provide more light in the room
 - d. Provide visual clues
2. Mr. Walker, age 71, is a newly diagnosed diabetic. You are teaching Mr. Walker insulin self-administration. What area should you assess to ensure that Mr. Walker can safely self-administer insulin?
 - a. Skin
 - b. Eyes
 - c. Heart
 - d. Lungs
3. Since Mr. Walker is older, what area needs to be assessed to ensure that he has learned the correct administration procedure?
 - a. Memory
 - b. Sensation
 - c. Balance
4. Tom Mallard, age 55, has CHF. How might you adapt your teaching plan to meet his needs?
 - a. Shorten class time
 - b. Hold classes in the evening
 - c. Present all material at one class
5. Joe McDonough, age 47, has immunosuppression owing to chemotherapy. How would you adapt your teaching plan?
 - a. Shorten class time
 - b. Hold class in A.M.
 - c. Schedule one-on-one instruction



Assessing Wellness

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Define health
- Identify the nurse's role in promoting wellness
- Identify factors that affect health
- Identify normal sleep patterns for various age groups
- Identify factors that can affect normal sleep patterns
- Identify normal exercise patterns for various age groups
- Identify factors that can affect normal exercise patterns
- Identify stressors for various age groups
- Identify risks for injury for various age groups



Introducing the Case Study

Mrs. Evelyn Chen (age 40); her daughter, Jan (age 15); and Mrs. Chen's mother, Mrs. Emily Wong (age 60), are attending a women's health fair sponsored by several large hospital systems and pharmaceutical companies. Health screenings and educational materials are available for women of all ages. If these three women asked you which screenings to have, what would you tell them? Before you can offer advice, you need to have an understanding of what health is and what factors influence a person's decision to live a healthy lifestyle.

Introduction to Assessing Wellness

Health is a broad concept that is difficult to define. Older definitions viewed health as the absence of disease. But today health is seen as a goal, a dynamic process that involves the self and self-care ability; optimal functioning of body, mind, and spirit; the ability to adapt; feelings of well-being and wholeness; and growing and becoming. People also have their own definitions of health, which are affected by gender, sociocultural factors, previous experience, age, and personal goals. These personal definitions influence their health promotion goals.

The three basic levels of prevention are primary, secondary, and tertiary. Primary prevention is essential to maintaining a state of wellness. At this level, the assessment process should include screening procedures, immunizations, and especially, prevention education.

How do you teach people to stay well? Follow these steps:

- Set an example with a healthy lifestyle.
- Motivate them to change unhealthy behaviors.
- Propose strategies for behavior change.
- Show them how to care for themselves more effectively.
- Build on their strengths.
- Help them find and use available resources.
- Provide support through telephone, individual, and group counseling and continuing education.

Healthy People 2010 has identified leading health indicators that are used to measure the health of our nation over the next 10 years. These indicators were selected based upon the ability to motivate change, data availability to evaluate progress, and importance as public health issues (Healthy People 2010). The indicators are not disease specific, but rather, many are behavior specific. It is the unhealthy behaviors that increase the risk for disease, such as smoking increases the risk for lung disease (chronic obstructive pulmonary disease

[COPD]) and lung cancer. As healthcare providers, you are in prime position to promote wellness.

The leading health indicators are:

- Physical activity.
- Overweight and obesity.
- Tobacco use.
- Substance abuse.
- Responsible sexual behaviors.
- Mental health.
- Injury and violence.
- Environmental quality.
- Immunizations.
- Access to healthcare.

These leading health indicators are addressed throughout this text. This chapter focuses on assessing four key areas fundamental to maintaining wellness—rest and sleep, exercise, stress management, and injury prevention—and suggests questions to ask at each developmental stage.

Factors Affecting Health Behaviors

Many factors influence a person's choice to maintain healthy behaviors. Some of these factors include support systems, psychological state of mind, access to healthcare, and motivational level. Some of these factors may be beyond your patient's control, and your influence on these factors could lead to a positive outcome. Referrals may be needed to facilitate healthy behaviors, and providing support for your patient can be crucial to his or her wellness.

Support Systems

The encouragement of friends and family can mean the difference between achieving personal health goals and falling short of them. Ask the person about her or his support systems, and then use these systems in the assessment process when needed. Other types of supports include organized religious groups, nurses and other caregiving professionals, and self-help groups such as Weight Watchers.

Psychological State

A person's psychological state affects his or her physical state. For example, if a person knows that he or she needs to stop smoking, start eating more healthfully, and start exercising more, then why doesn't he or she do it? Part of the assessment process is determining these reasons and working with people to change harmful behavior.

Access to Healthcare

People may be motivated to maintain or improve their state of wellness but be prevented from doing so because they lack access to healthcare. Causes include lack of finances or insurance coverage; rural location; lack of transportation; crowding at available healthcare resources; age,

gender, ethnicity, and/or religion (possibly because of prejudice); and healthcare rationing based on these factors.

Motivational Level

Because people tend to resist change, convincing them to trade unhealthy behaviors for healthy ones can be difficult. Teaching them the importance of primary prevention does not ensure that they will practice what you preach. For example, people know that smoking, obesity, and lack of exercise are dangerous to their health, but they still have trouble changing long-standing behaviors unless they are truly motivated.



HELPFUL HINT

Individualize your patient's health promotion plan—it increases the chance of success.



HELPFUL HINT

Do not assume that people know how to stay well. We may be bombarded with health information in the media and on the Internet, but what seems simple and obvious to the healthcare professional may still seem complex to the layperson.

Assessing People's Strengths

Knowing your patient's strengths and weaknesses will help both of you set realistic goals and help you plan appropriate interventions. Ask some or all of the following questions:

- What abilities do you possess to take care of your health?
- What activities help you to maintain or improve your health?
- What changes have you made in your lifestyle in the past 2 years to improve your health?
- What goals have you set to improve your health? Do you have a plan for reaching them?
- What changes in your life do you see in the future?
- What people currently give you the most support?
- What current activities make you feel happy?
- When you were younger, what activities gave you strength, comfort, and support?
- What helps you cope in a crisis?
- What gives you direction in your life?
- How do you spend a typical day?
- What do you feel motivated to do in terms of a lifestyle change?
- What barriers do you think will inhibit your lifestyle change?
- What health problems have you successfully dealt with in the past?

Health Promotion Plan of Action

- Identify the person's healthcare goals.
- Identify behavioral or health outcomes.
- Develop a behavior change plan.
- Reiterate benefits of change.
- Address environmental and interpersonal facilitators and barriers to change.
- Determine a time frame for implementation.
- Ask the person to make a commitment to healthcare goals.

Performing the Wellness Assessment

The wellness assessment entails asking the person and/or caregiver first general questions about health promotion behaviors and then specific questions about sleep and rest, exercise, stress management, and injury prevention. Health promotion behaviors and needs differ according to the person's developmental stage. (For a description of developmental stages see Chapter 2, The Health History.)

General Questions to Ask at Each Developmental Stage

Tailor your assessment questions to the person's developmental level—infant, toddler/preschool/school age, adolescent, young adult, middle age, or older adult. Include younger children in the interview as maturational level permits, and definitely include them by late school age. For young children, also assess health promotion behaviors for the family and the ability of the family to meet the child's needs.

Infants/Toddlers/Preschoolers

Ask the parent(s) or caregiver(s): Has your child's blood pressure been checked? Does the child brush her or his own teeth? How often? Are her or his immunizations (polio, diphtheria-tetanus-pertussis, measles-mumps-rubella, *Haemophilus influenzae* type B, hepatitis A and B, chickenpox, pneumococcal disease) up to date? (See *Recommended Preventative Screenings*.) Has she or he had a vision and hearing assessment? If so, when? If the assessment was abnormal, what follow-up has occurred? What are your child's nutritional habits? (For more information, see Chapter 8, Assessing Nutrition.) Has she or he been tested for anemia or tuberculosis? What are your child care arrangements? Have you ever harmed your child or wanted to? Do you know what community resources are available related to child abuse?

Recommended Preventative Screenings

SCREENING	MEN	WOMEN	PREGNANT WOMEN	CHILDREN	COMMENTS
Alcohol Misuse	X	X	X		
Asymptomatic Bacteriuria			X		
Breast Cancer		X			Mammography every 1–2 yr for women 40 and older.
Cervical Cancer		X			For women who have been sexually active and have a cervix.
Chlamydial Infection		X	X		For sexually active women age 25 and younger, other asymptomatic women at increased risk for infection, asymptomatic pregnant women age 25 and younger, and others at increased risk.
Colorectal Cancer	X	X			Men and women age 50 and older.
Depression	X	X			In clinical practices with systems to ensure accurate diagnoses, effective treatment, and follow-up.
Diabetes Mellitus Type 2	X	X			Adults with hypertension and hyperlipidemia.
Hepatitis B Virus Infection			X		Pregnant women at first prenatal visit.
High Blood Pressure	X	X			
Lipid Disorder	X	X			Men age 35 and older, women age 45 and older, younger adults with other risk factors for coronary artery disease (CAD). Screening for lipid disorders to include measurements of total cholesterol and high-density lipoproteins (HDLs).

SCREENING	MEN	WOMEN	PREGNANT WOMEN	CHILDREN	COMMENTS
Obesity	X	X			Intensive counseling and behavioral interventions to promote sustained weight loss for obese adults.
Osteoporosis		X			Postmenopausal women (women 65 and older and women 60 and older at increased risk for osteoporotic fractures).
Rh (D) Incompatibility			X		Blood typing and antibody testing at first pregnancy-related visit. Repeat antibody testing for unsensitized Rh (D)-negative women at 24–28 wk gestation unless biological father is known to be Rh (D) negative.
Syphilis	X	X	X		Persons at risk and all pregnant women.
Tobacco Use	X	X	X		Tobacco cessation interventions for those who use tobacco. Augmented pregnancy-tailored counseling to pregnant women who smoke.
Visual Impairment (in Children Younger than Age 5)				X	To detect amblyopia, strabismus, and defects in visual acuity.

Source: U.S. Department of Health and Human Services. (November 2000) *Healthy People 2010*, ed. 2. With *Understanding and Improving Health and Objectives for Improving Health*. 2 vols. Washington, DC: U.S. Government Printing Office.

School-Age Children

Adapt the questions in the previous paragraph and also ask the following: Has your child been part of home, school, or community education programs related to alcohol, drugs, sexuality, acquired immunodeficiency disease syndrome (AIDS), birth control, or sexually transmitted diseases (STDs)? Does the child brush his or her teeth and practice other health promotion behaviors

without being reminded? How is he or she doing in school?

Adolescents

Ask the adolescent the same questions as you asked about the school-age child, and also ask the following: Who are your community role models?

All Adults

Ask adults of any age: What is your or your family's definition of health? Do you believe you are healthy? Do other family members believe they are healthy? What health practices are included in your and your family's lifestyle? How is the family affected when someone is ill? Do you and your family have access to healthcare? Do you all have health insurance? Do you all receive regular physical and dental examinations? What religious or cultural beliefs do you all have that guide your purpose in life and your health practices?

Young Adults

Primary physical growth is completed during the 20s, and body systems usually reach peak functioning. Although most young adults are healthy, some major threats include accidental injury, cancer, heart disease, suicide, AIDS, and homicide. Men are more likely to die than women, especially as a result of homicide involving handguns. Ask the following questions: Are your immunizations up to date? Do you visit a family doctor? Do you perform self-screening (e.g., breast or testicular self-examination)? How often? What do you eat in a 24-hour period? Are there behaviors affecting your health that you would like to change?

Middle-Aged Adults

At this stage, people are likely to be knowledgeable and assertive about their health and healthcare. However, they may not know what specific screening tests they should have. Middle-aged men are less likely to seek routine preventive care than women, which may be one reason why the life expectancy for women is approximately 7 years longer than that of men. Ask the following questions: How do you care for your teeth? What screening tests have you had and when (e.g., mammograms and prostate-specific antigen blood tests)? What were the results? Do you keep a written record of your physician visits, immunizations, and screenings? Do you get a yearly flu shot? How about pneumonia vaccinations, tetanus-diphtheria, and hepatitis B injections?

Older Adults

Many older people begin to develop health problems, but they need to know that it is never too late to begin living more healthfully. Social support is the key in maintaining wellness in the later years. Research shows that older people who regularly attend religious services are less likely to require hospitalization, and when they do, have shorter hospital stays. Ask the same questions you asked the middle-aged person, but stress the importance of

getting immunized, limiting medication use, eating nutritious foods, drinking enough fluids, interacting with family and friends, and keeping active in the church or community.

Assessing Areas of Wellness

The following four areas are critical to maintaining wellness: rest and sleep, exercise, stress management, and injury prevention.

Rest and Sleep

Sleep restores, rejuvenates, and sometimes, heals the body. Lack of sleep causes fatigue, stress, depression, and a decrease in lymphatic system functioning, which increases the risk of infection and disease. The following factors can directly affect how well and how long we sleep:

- *Circadian rhythms:* Also called the *biological clock*, these rhythms help regulate the sleep/wake cycle, body temperature, and hormonal levels within a 24-hour period. Disruption affects muscle strength and coordination, attention, memory, and concentration.
- *Age:* Infants sleep 16 to 20 hours a day; preschoolers, 10 to 12 hours a day; school-age children, 9 to 10 hours a day; adolescents 7½ hours a day; and adults and older adults, 6 to 8 hours a day. Babies spend more time in **rapid eye movement (REM) sleep** than adults. Young children may experience sleep problems such as **nocturnal enuresis** (bedwetting), **nightmares** (bad dreams), **night terrors** (nightmare from which child awakens screaming), and **somnambulism** (sleepwalking). (For more information on sleep problems in children, see Chapter 26, Assessing the School-Age Child and Adolescent.) Older adults may take more time to fall asleep (**sleep latency**), may have a fragmented sleep pattern with less deep sleep, and may awaken early. (For more information on sleep problems in older adults, see Chapter 27, Assessing the Older Adult.)
- *Exercise:* Moderate exercise has little or no effect on sleep; however, vigorous exercise before retiring may inhibit sleep.
- *Nicotine, caffeine, alcohol:* Smoking increases the time needed to fall asleep and causes lighter sleeping and more frequent awakening. Caffeine near bedtime can increase sleep latency and reduce total sleep time, especially in older adults. Alcohol affects REM sleep, causing a fragmented sleep pattern, and also exacerbates sleep apnea.
- *Diet and weight:* Sleep apnea is more common in obese people. High-protein foods increase alertness; carbohydrates promote relaxation. People who are gaining weight usually sleep more; those who are losing weight sleep less.

- **Medical problems:** COPD, congestive heart failure (CHF), and pain can affect sleep patterns.
- **Stress:** Stress often increases arousal and inhibits sleep.
- **Medications:** Some prescription and over-the-counter (OTC) drugs can affect the number of hours a person sleeps as well as the sleep process.

Questions to Ask at Each Developmental Stage

Infants

Ask the parent(s) or caregiver(s): Where does the infant sleep? In what position do you place her or him? Does the infant sleep through the night? Does she or he go to sleep with a bottle? If so, do you remove it once she or he is asleep? (Having a nipple drip milk into the mouth once the child is asleep can cause tooth decay and ear infections, in addition to being a choking hazard.)

Toddlers/Preschoolers/School-Age Children

Ask the parent(s) or caregiver(s): Where does the child sleep? How many hours does he or she sleep? Does the child take naps? Does he or she sleep through the night? If not, how do you console the child when he or she awakens? What bedtime rituals do you practice? Does the child act tired during the day?

Adolescents

Ask the adolescent: How many hours do you sleep? Are you tired during the day? Do you have trouble paying attention at school or at work because of fatigue? Do you have a regular time you go to bed, and do you stick to it? Do you usually sleep through the night? If not, what causes you to wake up? How do you get back to sleep?

Young Adults

Ask the young adult: How many hours of sleep do you need to feel rested? How many hours of uninterrupted sleep do you get each night? Do you usually sleep through the night? If not, what causes you to wake up? How do you promote sleep or get back to sleep? Do you have a bedtime routine? Do you take medications that interrupt a normal sleep cycle? Do you take medications to help you sleep? Do you maintain a physical fitness program? Do you have problems concentrating because of fatigue? What are your usual work hours?

Middle-Aged Adults

Ask the same questions as you asked the young adult, as well as the following: How would you describe your quality of sleep? What time do you usually go to bed? How long does it take you to go to sleep? What time do you usually wake up and get up? What do you do for relaxation and how often do you do it?

Medications That Affect Sleep

Medication	Effect on Sleep
Amphetamines	Difficulty falling asleep
Antidepressants	Difficulty falling asleep
Barbiturates	Shallow, fragmented sleep; "hangover" effect
Phenylpropanolamine	Difficulty falling asleep (found in OTC decongestants and cold and diet remedies)
Diuretics before sleep	Frequent awakening
Beta blockers (e.g., Inderal)	Nightmares

Older Adults

Adapt the questions you asked the young and middle-aged adults, and add the following: What activities do you engage in during late afternoon or early evening? If you have trouble falling asleep, have you tried a light, warm snack at bedtime?

Exercise

Today, with so many people working at sedentary jobs, exercise needs to be planned for. Research shows that women are less active than men; people with lower incomes and less education are less active than those with higher incomes and education; people with disabilities are less active than people without disabilities; African Americans and Hispanics are less physically active than Caucasians; and many children and youth are overweight and exercising less, except for those active in organized sports. Major barriers to increasing physical activity are lack of time, access to convenient facilities, and safe environments in which to be active.

Questions to Ask at Each Developmental Stage

Infants

Keep in mind what gross and fine motor skills are normal at this age. Ask the parent(s) or caregiver(s): Does the infant play with her or his hands and feet and make noises? What kinds of toys do you give her or him? Which toys does the infant prefer? How often do you change her or his toys and play environment? Do you check toys regularly for loose parts and other safety hazards?

Toddlers/Preschoolers

Keep in mind what gross and fine motor skills are normal at this age. Ask the parent(s) or caregiver(s): Does the child have any physical limitations? Does he or she tire easily? What activities does he or she like? Does the child have a safe environment to explore and play? Whom does

he or she play with? How many hours does your child watch TV or participate in other sedentary activities? What do you do to encourage physical activity during bad weather?

School-Age Children

Ask the same questions as you asked about the toddler/preschool child, as well as the following:

- To the child: Do you play organized sports? What precautions do you take, and what protective equipment do you wear?
- To the parent(s) or caregiver(s): Do you encourage physical activity? Do you engage in physical activities with your children?

Adolescents

Ask the adolescent: What competitive sports or other physical activities do you like? Do you schedule exercise during your week? How often and for how long? Do you exercise with friends? Do you have any physical limitations? Have you ever been injured during exercise? Does your school encourage participation in physical activities? What precautions do you take and what protective equipment do you wear? Do you gain satisfaction from exercising?

Young Adults

Ask the young adult: What physical activities do you include during an average week? How often and for how long? Do you have any physical limitations? Do you have any health conditions that should be evaluated before beginning an exercise program? Where do you exercise? Do you participate in activities that raise your heart rate? Do you include a warm-up and cool-down period? Do you participate in organized sports? What precautions do you take and what protective equipment do you wear? Have you had any exercise-related injuries? With whom do you exercise? Do you engage in physical activities as a family? Do you enjoy your exercise program?

Middle-Aged Adults

Ask the middle-aged adult: What kinds of exercise do you do? How often and for how long? Where do you exercise? With whom do you exercise? Do you warm up before and cool down after each exercise period? Have you had any unusual or uncomfortable feelings before, during, or after exercising? (If so, refer back to the person's cardiovascular, respiratory, and musculoskeletal history data.)

Older Adults

Ask the same questions as you asked the middle-aged patient, but emphasize that the older adult should check with her or his doctor before starting a new exercise program.



HELPFUL HINT

Teach older people to take their pulses when exercising. The recommended formula for maximum heart rate at this age group is: 220 minus age, multiplied by 60 to 80 percent. For example, in an 80-year-old, the range would be 84 to 112 beats per minute (BPM), with 84 BPM being the minimum heart rate and 112 BPM the maximum heart rate for cardiovascular functioning.

Stress Management

Multiple stressors can occur at any age, even infancy, although different age groups are subject to different stressors. Over time, stress can cause hypertension, cardiac arrhythmias, cardiovascular disease, gastrointestinal problems, headaches, and decreased immunological functioning, which can contribute to cancer and other diseases.

Questions to Ask at Each Developmental Stage

Infants

Keep in mind what social developmental milestones are normal for this age. Ask the parent(s) or caregiver(s) the following: What emotions have you seen the infant express? How does the infant calm himself or herself when crying? How do you calm the infant when he or she is unable to become calm? How would you describe his or her temperament? Does the infant entertain himself or herself when alone? How do you set limits for him or her? Does your home environment provide cognitive, physical, and psychosocial stimulation for the infant? Who provides child care when needed? Does the caregiver stick to the infant's usual routines?

Toddlers/Preschoolers

Adapt the preceding questions, and also ask the following: How does your child calm herself or himself after an emotional outburst? Children tend to regress when ill, so what do you do to support the child's developmental level when she or he is ill? How do you show affection to your child, and how does she or he respond? How well does your child play with other children? What kinds of conflicts occur with other children? How do you set disciplinary guidelines for your child? How do you encourage her or his development of autonomy and initiative? How does she or he interact with siblings? How does she or he express positive (love, affection, happiness, joy) and negative emotions (hate, jealousy, anger, fear)? Do you actively model healthy expression of emotions in the home? Does your child enjoy child care experiences?

School-Age Children

- To the child: Do you usually feel happy and contented? Do you like yourself? What do you do when you feel

bored or sick? Do you like to compete with others in organized or informal activities? Do you enjoy learning new things? Do you feel confident as you begin new projects? Do you enjoy the challenge of solving new problems? Do you feel like your parents or caregivers support your activities and enjoy your successes? Do they support you even when you do not meet their expectations? Do you think of the consequences of your behavior before acting? Who are your friends? Do you feel included in most peer-group activities? If you have conflicts with friends, what is the source of the conflict, and how do you resolve it? Do you receive an allowance or have an opportunity to earn money? How do you manage your money? Do you feel good about your progress at school? What do you like most and least about school? How often do you miss school or other activities because you do not feel well? What types of physical activities do you do? How often and for how long?

- To the parent(s) or caregiver(s): Do you give your child an allowance and provide general guidelines for money management? How do you display your interest in your child's school work/progress? How do you support your child when he or she does not feel well? How would you describe your child's friends? What stress-management techniques do you model or actively teach your child? How does your child act when he or she is tired or stressed out? How does he or she cope with emotional stress? What types of changes have affected your family and the child during the last year?

Adolescents

Adapt the questions for the school-age child, and also ask the following: Do you feel comfortable with the physical changes accompanying puberty? What accomplishments are most important to you? What stressors do you experience weekly? How do you reduce stress or the effects of stress? Are you able to be assertive when you need to be? Can you give an example? What risks have you taken in the last year? Do you use tobacco, alcohol, or street drugs? What about OTC and prescription drugs? If so, what kinds, how often, and how much? How would you describe your peer group and your relationship with the group? How would you describe your relationships with the same sex, the opposite sex, and adults? How do you usually make decisions? What plans do you have once you leave or complete high school? Who are your role models for stress management? How do your parents support your efforts to be an independent person? What are the most common sources of conflict with your family and your peers? How do you resolve conflicts when they occur? Whom do you go to for support when you have a problem? What steps would you take if you were depressed or had thoughts of suicide or if you saw these characteristics in a friend? Have you ever been the victim of violence? Have you ever abused an animal or another person?

Young Adults

What stressors do you experience weekly? How do you reduce stress or the effects of stress? What risks have you taken in the last year? How do you make decisions? How would you describe your relationships with the same sex and the opposite sex? What are the most common sources of conflict with your family, friends, and coworkers? How do you resolve conflicts when they occur? Do you use tobacco, alcohol, or street drugs? What about OTC and prescription drugs? If so, what kinds, how often, and how much? Have you ever been the victim of violence? Have you ever wanted to hurt or abuse another person? Are you satisfied with your career choice? If not, what are your plans for a change? Do you have problems with time management? If you have children, what parenting rewards and challenges do you encounter? Do you feel confident and satisfied with your parenting skills?

Middle-Aged Adults

Ask the same questions that you asked the young adult.

Older Adults

Adapt the questions you asked the young adult and middle-aged adult, and also ask the following: When do you plan to retire from a full-time position? Are your financial resources adequate? Do you plan to work part-time after retirement? What activities are you interested in (e.g., travel, hobbies, volunteer work in the community)?

Injury Prevention

Injuries can occur at any age, and most people have a significant injury at some time in their lives. Although most accidents are predictable and preventable, accidental injuries are the leading cause of death in the 1- to 34-year-old age group. Additional millions are incapacitated by accidental injuries.

Questions to Ask at Each Developmental Stage

Infants

Ask the parent(s) or caregiver(s): What have you done to make your home safe for the infant? Do you use an infant car seat? Is the infant regularly exposed to tobacco smoke? Has she or he been injured as a result of an accident in the home, in another's home, or while riding in a motor vehicle? How often do you check equipment and toys for possible hazards? Are there guns in your home or in your caretaker's home? If so, are they securely stored?

Toddlers/Preschoolers/School-Age Children

Adapt the preceding questions, and also ask the parent(s) or caregiver(s) the following: Have you taught your child personal safety guidelines? Does he or she use protective equipment when participating in physical activities like skating or bicycling? Who supervises your child when he or she is playing? Can he or she swim and does he or she know water safety guidelines?

Adolescents

Ask the adolescent: Do you like to take risks? Have you completed a driver education course? What safe-driving behaviors do you practice? Do you talk on a cell phone while driving? Do you consider yourself well informed regarding the transmission, signs, symptoms, and treatment of STDs? Are you sexually active, and if so, do you practice safe sex? Do you know where to get confidential medical attention if you believe you have an STD or may be pregnant? Do you use alcohol, tobacco, or street drugs? What about OTC and prescription drugs? If so, what kinds, how often, and how much? Do you ever drive while under the influence of alcohol or drugs? Do you have guns in your home, and if so, are they securely stored? Have you ever been injured as a result of participation in physical activities? What protective measures do you take? Would you recognize signs and symptoms of depression in yourself or a peer? What would you do if you or a peer were depressed or had thoughts of suicide?

Young Adults

Adapt the preceding questions, and also ask: Have you evaluated your occupational health risks? What resources are available to you at work related to health maintenance or injury prevention? Do you have smoke and car-

bon monoxide detectors in your home, and do you check them frequently?

Middle-Aged Adults

Adapt the preceding questions, and also ask: Are you aware of environmental hazards in the home (e.g., loose rugs, electrical cords, stairways, steps)?

Older Adults

Adapt the questions for young and middle-aged adults, but focus on the prevention of falls. Ask: Can you describe any hazards in your home, especially in the bathroom, kitchen, or outside steps and sidewalks? Do you use an assistive device, such as a cane, walker, or wheelchair? Do you keep them in good repair? Have you made any modifications to your home, such as a wheelchair ramp or grab bars in the tub and by the toilet?

**Case Study Analysis and Plan**

Now that you know what factors to consider when assessing health and wellness, what screening would you recommend for Jan, Mrs. Chen, and Mrs. Wong?


S U M M A R Y

- There is no uniform definition of health as it applies to individuals, families, and communities. Yet, a definition forms the foundation for personal perceptions and is crucial in determining individual health promotion behaviors.
- Some factors that affect health behaviors are the person's motivation, support system, psychological state, and access to healthcare.

- The key to assessing wellness is to identify the person's perspective of health and any factors that affect health behaviors, and then work with him or her to develop a plan that promotes healthy living.

R E V I E W Q U E S T I O N S

1. Mr. Brown, age 73, complains of feeling tired. He is overweight and snores. Mr. Brown is at risk for developing:
 - a. Sleep apnea
 - b. Somnambulism
 - c. Sleep latency
2. You obtain a health history on Mr. Brown and note that he is taking a beta blocker for hypertension. What effect do beta blockers have on sleep?
 - a. Nocturia
 - b. Nightmares
 - c. Difficulty falling asleep
 - d. "Hangover" effect
3. Sleep patterns vary depending on the age of the patient. What change in sleep pattern is seen in older adults?
 - a. Sleep apnea
 - b. Somnambulism
 - c. Sleep latency
4. Carol Jenkins, age 54, is beginning an exercise program. Calculate her target heart rate for her.
 - a. 80–100 BPM
 - b. 100–133 BPM
 - c. 125–166 BPM
 - d. 156–209 BPM
5. What age group is at greatest risk for injuries?
 - a. Older adults
 - b. Middle-age adults
 - c. Adolescents



Assessing Nutrition

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Describe primary nutrients
- Define dietary guidelines based on the Food Guide Pyramid
- Differentiate among nutritional deficiencies
- Differentiate nutritional requirements for different age groups
- Discuss various cultural dietary preferences
- Perform a nutritional assessment

Nutrition is the relative state of balance between nutrient intake and physiological requirements for growth and physical activity. Optimal nutrition helps protect against disease, facilitates recovery, and decreases complications during illness. Good nutrition helps people stay healthy.

Malnutrition has traditionally been defined as a deficit of appropriate nutrients. However, it literally means *bad nutrition* and can encompass any situation that contributes to an imbalance in nutrient intake relative to actual needs. Therefore, malnutrition can mean a nutrient deficit or excess. Although nutritional deficits remain a significant health problem in Third World countries, the major problem in the United States is nutritional excess, as is shown by the rising rates of obesity in all age groups.

As a nurse, you are in a unique position to assess people's nutritional status and provide information on proper nutrition. You can reinforce positive nutritional patterns, identify people at risk for malnutrition, and encourage more healthful eating habits.

Assessing nutritional status achieves the following:

- Identifies actual nutritional deficiencies.
- Illuminates dietary patterns that may contribute to health problems.
- Provides a basis for planning for more optimal nutrition.
- Establishes baseline data for evaluation.

This chapter covers basic screening considerations and the procedures for conducting a complete nutritional assessment.

Review of Nutrients

The goal of eating is to supply body cells with necessary nutrients. Ingestion, digestion, absorption, and metabolism are the processes that normally accomplish this goal. Interference with any of these functions can contribute to nutritional problems. (For more information on these functions, see Chapter 17, *Assessing the Abdomen*.)

Primary Nutrients

Nutrients are substances contained in food that are essential for optimal body functioning. The **primary nutrients** are carbohydrates, proteins, fats, vitamins, minerals, and water. Carbohydrates, protein, and fat are the body's major energy sources. Carbohydrates and protein each supply 4 calories per gram, and fat provides 9 calories per gram. Vitamins are essential to specific functions in the body. Minerals are inorganic elements that are essential to cell structure and physiological functions in the body. Water makes up 50 to 60 percent of the adult weight. It is required for many functions, and humans cannot survive for more than a few days without it.

Carbohydrates

Carbohydrates are the body's major energy source. Foods that contain the most carbohydrates are grains, legumes, potatoes, corn, fruits, and vegetables. Adult carbohydrate intake should range from 50 to 100 g daily, depending on level of activity. In a normal diet, this means that 50 to 60 percent of the daily caloric intake should be derived from carbohydrates.

Protein

Protein is the primary building block of all tissues and organs and serves an important function in cell structure and tissue maintenance. Integrity of the skin, internal organs, and muscles depends on adequate protein intake and metabolism. The body can synthesize most of the necessary amino acids from nonprotein dietary sources. However, there are nine essential amino acids that the body cannot synthesize and that must be obtained through dietary sources. For this reason, adults require 0.8 g/kg per day of protein (about 10 to 20 percent of the daily caloric intake). Primary sources of protein include meat, milk and milk products (e.g., cheese and yogurt), nuts, and legumes.

More protein is needed during tissue building—for example, in pregnancy and lactation, childhood, adolescence, postoperative recovery, tissue damage, and long-term illness. Athletes also require additional protein to build and maintain muscle. Animal products are the most common source of protein in industrialized countries. However, well-planned vegetarian diets can also provide ample dietary protein.

Vegetarian Diets

Vegetarian diets consist primarily of plant foods. Vegans avoid all animal products in the diet, lactovegetarians eat dairy products, and lacto-ovovegetarians also eat eggs. Although these diets provide less protein than nonvegetarian diets, they meet or exceed the RDAs. Whole grains, legumes, vegetables, seeds, and nuts contain all the essential and nonessential amino acids. As long as food intake is varied and calories are sufficient, protein deficiencies will not occur. Furthermore, soy protein provides all essential and nonessential amino acids and is nutritionally equivalent to animal sources of protein. It can serve as the sole source of protein if desired. However, because considerable variation exists, evaluate the adequacy of vegetarian diets on an individual basis.

Fats

Lipids or fats are insoluble in water and soluble in alcohol, ether, and chloroform. They include true fats, lipids,

and sterols, such as cholesterol. Fat supplies twice as much energy as carbohydrates or proteins. It provides essential fatty acids (linoleic and linolenic acids) and promotes the absorption of the fat-soluble vitamins A, D, E, and K. The typical American diet usually contains adequate fat, and recommended daily allowances in grams do not exist. Triglycerides and cholesterol are major contributors to heart disease, diabetes mellitus (DM), and obesity. The U.S. Department of Agriculture (USDA) and the American Heart Association recommend that fat intake not exceed 20 to 30 percent of a person's total daily calories.

Highly **saturated fat** (solid at room temperature) significantly contributes to elevated serum triglyceride and cholesterol levels. So USDA/U.S. Department of Health and Human Services (USDHHS) guidelines recommend that no more than 10 percent of daily calories be derived from saturated fats. Unsaturated and polyunsaturated fats are recommended because of their inverse relationship with heart disease.

Cholesterol occurs naturally and exclusively in all animal food products. The body needs cholesterol for cell structure, as a precursor for certain hormones and vitamins, and to aid in the digestive process. But even if no cholesterol were consumed in the diet, the body would synthesize the needed supply.

Lipoproteins

Serum cholesterol and lipids attach to proteins and are transported throughout the body as **lipoproteins**. The relative ratio of lipid to protein determines the density of the molecule. A low lipid-to-protein ratio results in a high-density lipoprotein (HDL). HDLs are produced during cellular metabolism. They lower serum cholesterol by transporting it from the cell to the liver for metabolism and excretion. Conversely, a high lipid-to-protein ratio produces a larger and low-density lipoprotein (LDL). LDLs are produced in the gastrointestinal wall after eating and transport dietary cholesterol and triglycerides to the cells. Thus, HDLs guard against heart disease by lowering cholesterol, and LDLs contribute to heart disease by raising cholesterol. The risk for heart disease is particularly high when total serum cholesterol exceeds 200 mg/dL. The risk also increases as the HDL level decreases. An HDL less than 35 mg/dL is considered a major risk factor for heart disease, and an HDL above or equal to 60 mg/dL is a negative risk factor.

Vitamins and Minerals

Vitamins are organic compounds that play a major role in enzyme reactions associated with the metabolism of carbohydrate, protein, and fat. Although vitamins are required in small amounts, the body does not synthesize them, so they must be present in the diet. Vitamins are classified as water soluble or fat soluble. The body does not store water-soluble vitamins (B complex and C vitamins), so any

surplus in daily intake is excreted in the urine. Fat-soluble vitamins (A, D, E, and K) are obtained through dietary fat and are stored in adipose tissue, where they can accumulate and become toxic. Toxicity generally results from self-administered large doses of a vitamin or excessive intake of foods that contain the vitamin.

Minerals are inorganic compounds found in nature. They play a wide variety of roles in human nutrition. Minerals are required in varying amounts and are divided into major and trace elements. *Major minerals* are required in excess of 100 mg per day and include calcium, chloride, magnesium, phosphorus, potassium, sodium, and sulfur. The remaining minerals are known as *trace elements* (e.g. aluminum, bromine, chromium, cobalt, copper, fluorine, iron, iodine, magnesium, nickel, silicon, zinc, and other rare minerals).

Water

We usually do not think of water as being related to nutrition. But without it, humans cannot survive more than a few days. Water helps regulate body temperature; serves as a solvent for vitamins, minerals, and other nutrients; acts as a medium for chemical reactions; serves as a lubricant; and transports nutrients to and wastes from the cells. Adults consume about 6 cups of water per day through beverages; another 4 cups are obtained through food; and about 1 cup is produced as a byproduct of metabolism. **Sensible water loss** occurs through excessive perspiration, urine, and gastrointestinal secretions. Evaporative or **insensible water loss** occurs via the lungs and skin.

More than half of the body's weight is composed of water, and therefore rapid weight changes are usually a reflection of fluid balance. This is especially true for infants who have a greater proportion of water weight and proportionately more extracellular fluid. Thirst is not an accurate indicator of hydration status because it does not occur until about 10 percent of the intravascular volume is lost, or 1 to 2 percent of the intracellular volume is depleted. Assessment of hydration and signs and symptoms of dehydration are discussed later in this chapter.

Nutritional Guidelines and Standards

Early guidelines addressed the nutritional needs of the entire population, so they exceed the needs of most normal, healthy people. A diet that meets roughly two-thirds of the Recommended Daily Allowances (RDAs) for each nutrient is considered adequate.

Dietary Guidelines and the Food Guide Pyramid

The RDAs provide guidelines for specific nutrient quantities for clinical applications. The six-group Food Guide Pyramid (Fig. 8.1) was developed to make these recommendations easier for the public to understand and




FIGURE 8.1. Food Guide Pyramid (Courtesy of U.S. Department of Agriculture and Department of Health and Human Services).

follow. The pyramid can be used to evaluate individual nutritional status and to educate people about nutrition.

How a Nutritional Deficiency Develops

Nutritional deficiencies have characteristics that are relatively unique to the specific nutrient that is lacking in the diet. For example, diminished night vision is a classic symptom of vitamin A deficiency, and fetal deformities are associated with folic acid deficiencies in pregnancy. A thorough nutritional assessment can identify deficiencies long before actual clinical symptoms occur. The following section describes the four stages of nutritional deficiency.

Stage 1: Nutritional Deficiency Occurs

Malnutrition occurs when the nutrient in question is not available for digestion, absorption, and metabolism. **Primary malnutrition** results when a specific nutrient is lacking in the diet. Iron-deficient diets that result in anemia in infants and young children and calcium-deficient diets that cause osteoporosis in postmenopausal women are examples. **Secondary malnutrition** results from impaired bioavailability of nutrients to the body. Intake of nutrients may be adequate, but physiological processes prevent them from being digested, absorbed, or metabolized, such as with malabsorption syndrome. Primary and secondary malnutrition can occur together.



CRITICAL THINKING ACTIVITY #1

Identify some other causes of primary malnutrition.



CRITICAL THINKING ACTIVITY #2

Identify some other causes of secondary malnutrition.

Stage 2: Tissue Reserves Decrease

When a nutritional deficiency occurs, the body mobilizes tissue reserves to sustain metabolic processes. Nutrient levels in the blood generally will remain within normal limits as long as there are tissue reserves that the body can depend on. However, if the intake deficiency persists, tissue reserves become depleted and blood levels of nutrients drop, causing biochemical abnormalities.

Stage 3: Biochemical Lesions Occur

Biochemical lesions are changes in serum values that signal depletion of tissue reserves. Biochemical testing is a

valuable adjunct to nutritional assessment and may reveal nutrient deficiencies well before clinical signs and symptoms occur.

Stage 4: Clinical Lesions Occur

Clinical lesions are physical changes that result from an inadequate supply of one or more nutrients necessary for tissue growth and maintenance.

Developmental, Cultural, and Ethnic Variations

As people grow and develop, their nutritional needs change. Developmental groups especially at risk for nutritional problems include pregnant and lactating women, infants and children, adolescents, and older adults.

Infants, Children, and Adolescents

The growth and development that occur in infancy, childhood, and adolescence determine nutritional needs. For example, brain development is at its peak from birth through the second year of life, but after this, improved nutrition will not enhance brain growth. Growth charts are the standard against which infant and child growth is evaluated (see Appendix B). The following tables list indications of good nutrition in school-age children and summarize pertinent growth and developmental factors and corresponding nutritional needs of infants, children, and adolescents.

Indications of Good Nutrition in the School-Age Child

General Appearance	<ul style="list-style-type: none"> ■ Alert, energetic ■ Normal height and weight
Skin and Mucous Membranes	<ul style="list-style-type: none"> ■ Skin smooth, slightly moist ■ Mucous membranes pink, no bleeding
Hair	<ul style="list-style-type: none"> ■ Shiny, evenly distributed
Scalp	<ul style="list-style-type: none"> ■ No sores
Eyes	<ul style="list-style-type: none"> ■ Bright, clear, no fatigue circles
Teeth	<ul style="list-style-type: none"> ■ Straight, clean, no discoloration or caries
Tongue	<ul style="list-style-type: none"> ■ Pink, papillae present, no sores
Gastrointestinal System	<ul style="list-style-type: none"> ■ Good appetite ■ Regular elimination
Musculoskeletal System	<ul style="list-style-type: none"> ■ Well-developed, firm muscles ■ Erect posture ■ Bones straight without deformities
Neurological System	<ul style="list-style-type: none"> ■ Good attention span for age ■ Not restless, irritable, or weepy

Source: Lutz, C., and Przytulski, K. (2001). *Nutrition and Diet Therapy*, ed. 3. Philadelphia: F.A. Davis.

Nutritional Needs of Infants, Children, and Adolescents

AGE	DEVELOPMENTAL AND GROWTH FACTORS	CORRESPONDING NUTRITIONAL NEEDS
Infants: Birth–12 Mo	<ul style="list-style-type: none"> ■ Energy needs largely a result of high metabolic rate, increased skin-to-body-size ratio (facilitates heat loss and requires more thermal energy), and sucking and crying activities. ■ Increased protein needs result from rampant growth in first year. ■ Fat is critical to developing nervous system and is also an important energy source at this age. ■ Fetal iron stores depleted around 4–6 mo. 	<ul style="list-style-type: none"> ■ 100–200 kcal/day at birth, decreases to 80–100 kcal/day by 12 mo. ■ 2.5–3.5 g protein at birth; decreases to 2 g by 12 mo. ■ 30%–55% of daily caloric intake should be derived from fat. ■ Solids usually introduced around 4–6 mo. Cereal is important source of iron. Provide iron supplements if solids not introduced. Do not give eggs and cheese until immune system has matured, around 12 mo.
Toddlers: 1–3 Yr	<ul style="list-style-type: none"> ■ Growth rate slows drastically and appetite decreases. Wish for autonomy also contributes to selective appetite or refusal to eat. Parents usually become concerned about decreased/erratic appetite. ■ Fat remains an essential nutrient for energy and neurological development. ■ Risk of iron deficiency anemia related to decreased intake of iron-rich foods. ■ Children usually dislike vegetables, so fiber intake throughout childhood tends to be low. 	<ul style="list-style-type: none"> ■ Require all essential nutrients. Intake should adhere to Food Pyramid guidelines. Serving size roughly 1 tablespoon per year of child's age (e.g., 2½ tablespoons of rice for a 2½-yr-old). ■ Whole milk is necessary through the second year. ■ Milk contains no iron, so limit it to 1 quart daily because more will diminish food intake. ■ Beginning at age 2, daily fiber intake should be equal to child's age + 5 g of fiber.
Preschoolers: 3–6 Yr	<ul style="list-style-type: none"> ■ Continue to be very active. Growth continues at approximately 4–5 lb and 1–2 inches a yr. ■ At risk for iron deficiency anemia, especially those from low-income families, who are also at risk for vitamin A and C deficiency. ■ Children are trying to master behaviors that emulate the social norm and are internalizing nutritional patterns and habits. 	<ul style="list-style-type: none"> ■ Needs average of 1800 cal/d. Requires same nutrients as adults but in lesser amounts. 1 tablespoon per age in years remains appropriate serving size. ■ Monitor for iron deficiency anemia. ■ Children should be learning optimal dietary practices.
School Age: 6–12 Yr	<ul style="list-style-type: none"> ■ Doubles first year's weight by age 6. Energy needs and requirements correspond to age and activity. ■ Breakfast is important for optimal school performance. ■ Calcium and phosphorus are critical to prepubertal skeletal development. 	<ul style="list-style-type: none"> ■ RDA for caloric intake is 2000 cal/d. ■ Breakfast should include 25%–30% of daily nutrient intake. ■ RDA for calcium is 800 mg/d. Calcium and phosphorus retention is greatest just preceding puberty, so encourage milk intake.
Adolescence: 12–20 Yr	<ul style="list-style-type: none"> ■ Growth is erratic and occurs in spurts. ■ Sex hormones influence growth differences between boys and girls: Boys' needs related to increased size, muscle mass, and sexual maturation. 	<ul style="list-style-type: none"> ■ 60 to 80 cal/kg needed to support dramatic growth and intense physical and cognitive activity. ■ Boys have greater calorie, iron, and zinc needs for sexual maturation. Girls' needs related to menstrual function.

AGE	DEVELOPMENTAL AND GROWTH FACTORS	CORRESPONDING NUTRITIONAL NEEDS
	<ul style="list-style-type: none"> Continued skeletal growth with enhanced needs for calcium and phosphorus. Development characterized by identity-seeking behaviors and need to fit in with peers. Consumption of soft drinks and fast food increases. White female adolescents are at greatest risk for anorexia nervosa and bulimia. 	<ul style="list-style-type: none"> Girls require even more iron for menstrual function. Both sexes (especially athletes) need vitamins B and C for protein synthesis. Need four servings of milk or dairy products per day. At risk for developing obesity or calcium deficiency if soft drinks are substituted for milk.

Pregnant Women

Good nutrition is critical to the developing fetus. Women who fail to receive adequate nutrition before and during pregnancy are at risk for premature birth, low-birth-weight (LBW) infants, or infants who are small for gestational age (SGA). These infants are at higher risk for mental and physical disabilities and other congenital anomalies. In addition, pregnant women who are poorly nourished endanger their own health because maternal nutrients are sacrificed to compensate for the increasing demands of the growing fetus.

Pregnant women need an additional 300 calories a day, and lactating women an additional 500 calories a day. Development of the placenta, amniotic fluid, and fetal tis-

ues, and increases in maternal blood volume during pregnancy require 10 g of protein a day. This requirement can be met simply by adding one additional serving each of milk and meat to the daily diet. Prenatal vitamins and mineral supplements during pregnancy and lactation are also critically important. The following table summarizes the vitamin and mineral needs of pregnant and lactating women.

Older Adults

Older adulthood begins at age 65. The senses become less acute, and diminished taste of sweet and salty foods may cause people to compensate by increasing sugar and salt. Decreased gastric acidity can impede vitamin B₁₂ absorption. Antacid use impedes it further. Skin changes

Vitamin and Mineral Needs During Pregnancy and Lactation

VITAMIN/MINERAL	FUNCTION
Vitamin C	Activates folic acid (folate), facilitates iron absorption, and helps build connective tissue.
Folic acid (400 µg)	Needed for fetal brain development. Deficiencies have been correlated with mental retardation and neural tube defects.
Iron (30 mg)	Important for development of red blood cells. Is supplied to fetus regardless of mother's iron status, increasing risk of maternal anemia.
Calcium (1200 mg)	Necessary for fetal skeletal tissue. Derived from mother's circulating supply, and unless calcium reserves are replenished, pregnant women will experience compensatory bone demineralization.
Fluoride (1 mg)	Fetal tooth bud development begins near end of first trimester. No recommendations exist for fluoride supplementation during pregnancy, but children whose mothers took fluoride daily during pregnancy have fewer cavities.
Zinc	A structural element of cells and an important mineral for a growing, developing fetus. Not mobilized from maternal tissues and must be supplied to fetus through mother's diet. Deficiencies associated with small, malformed infants and difficult labor.

Source: Data from Glenn F.B., Glenn, W.D., and Duncan, R.C. (1982). Fluoride tablet supplementation during pregnancy for caries immunity: A study of the offspring reduced. *American Journal of Obstetrics and Gynecology* 143(5):560-564; and Lutz, C. and Przytulski, K. (2001). *Nutrition and Diet Therapy*, ed. 3. Philadelphia: F.A. Davis.

associated with aging impair vitamin D synthesis; so does spending less time outdoors. Diminished physical strength and decreased activity predispose older adults to bone demineralization and loss. This problem is especially common in women who do not consume adequate calcium and in postmenopausal women who have lost the protective advantage of estrogen. In addition, loss of urinary sphincter muscle tone in women and urination difficulty associated with prostate changes in men may discourage fluid intake. Confusion, a common consequence of dehydration, may impair mental status. Decreased income, loss of the social context for eating (e.g., death of spouse), and lack of accessibility to food markets are social and economic aspects of aging that may adversely affect nutrition.

The RDA amounts beginning in early adulthood change only moderately through the middle adult years. The most significant changes beginning in midlife and extending into older adulthood are related to the physical changes associated with aging. Energy requirements are thought to decrease by about 5 percent per decade after age 40. However, without dietary modifications, weight gain will occur, with the risk for obesity-related diseases. The RDAs reflect a 200-mg increase in calcium and a twofold increase for vitamin D to 10 µg after age 50. Unless fluid intake is restricted for medical reasons, older adults still need to drink 6 to 8 glasses of water daily.

People of Different Cultural/Ethnic Groups

Sociocultural patterns of food intake also influence nutritional status. Cultural patterns of eating are learned and reinforced early in life and are difficult to change without conscious efforts and external support. Many cultures have healthy eating habits. For example, the traditional Asian diet provides adequate nourishment and is associated with lower rates of the chronic diseases that plague Western populations. However, now that Asians are beginning to adopt Western food preferences, chronic diseases are on the rise in these countries. In contrast, diets in Western industrialized cultures are high in meat, sugar, and fat.

Introducing the Case Study

James Thomas, a 76-year-old African American man, lives alone in a two-story urban row home. His wife passed away months ago. He has a history of hypertension (HTN), which is controlled with Lasix 40 mg od and Vasotec 5 mg bid. He has come for a routine blood pressure (BP) check but complains that he doesn't feel well and has lost 10 lb since his wife's death. He states, "I have no energy, and I'm always tired."



Performing the Nutritional Assessment

One of the first steps in assessing nutrition is screening the patient for possible nutritional risks. At the very least, you should evaluate the health history for symptoms and situations related to nutritional problems, perform a basic physical examination, and obtain laboratory data associated with malnutrition. A comprehensive nutritional assessment involves a detailed dietary history, focused **anthropometry**, and evaluation of laboratory values. It is recommended for people with any of the following nutritional risks:

- Weight less than 80 percent or more than 120 percent of ideal body weight (IBW).
- History of unintentional weight loss (>10 lb or 10 percent of usual weight).
- Serum albumin concentration lower than 3.5 g/dL.
- Total lymphocyte count lower than 1500 cells/mm³.
- History of illness, surgery, trauma, or stress.
- Symptoms associated with nutritional deficiency or depletion.
- Factors associated with inadequate nutritional intake or absorption.

Health History

Nutritional health risks and problems are not always obvious to the patient or the nurse. So during the health history, stay alert for clues. The following sections summarize health history data that may signal nutritional problems warranting further investigation.

Biographical Data

Scan the biographical data for clues that may affect the patient's nutritional status. Note age to determine normal dietary requirements. Nutritional needs also vary according to gender. Religion and cultural background may influence dietary preferences. Financial status may also affect the person's ability to maintain a healthy diet.

Current Health Status

Inadequate nutrition is often discovered indirectly during a routine health history and physical examination. For instance, diminished growth and delayed development related to inadequate nutrition may be identified during a routine well-child checkup. Patient concerns that result from malnutrition usually present as a specific symptom or functional problem rather than a focused nutritional problem. For example, a person with iron deficiency anemia may complain of a lack of energy and an inability to concentrate.

Ask your patient if his or her health status has changed. If so, consider the potential influence of the health change on nutrition. Acute and chronic illnesses, debilitating conditions, medications, surgery, and trauma

Cultural Dietary Preferences

CULTURE	DIETARY HABITS
African American	<ul style="list-style-type: none"> High-fat, high-sodium foods, fried foods, large amounts of animal fat, little fruit and vegetables.
Amish	<ul style="list-style-type: none"> High in fats and carbohydrates.
Appalachian	<ul style="list-style-type: none"> Believe that plenty of food is associated with wealth. Use large amounts of lard and preserve meats and other foods with salt.
Arab American	<ul style="list-style-type: none"> Cook with spices and herbs, eat lamb and chicken, but no pork. Believe it is unhealthy to drink and eat at same time or mix hot and cold foods. Main meal is at midday.
Chinese American	<ul style="list-style-type: none"> Food at meals is eaten in specific order. Food often used as medicine to cure disease and increase strength. Peanuts and soybeans popular.
Cuban American	<ul style="list-style-type: none"> Staples include root crops, plantains, and grains. Meat is usually chicken and pork. Typical diet is high in calories, starch, and saturated fats. Leisurely noontime meal, late dinner.
Egyptian American	<ul style="list-style-type: none"> Food is well done and well seasoned. Rice is main staple. Usually do not eat pork. Do not mix hot and cold, sweet and sour foods. Main meal is at midday.
Fillipino American	<ul style="list-style-type: none"> Meals are a time for socialization. Sharing food is important. Salt and vinegar used as preservatives, leading to high sodium intake. Plants and herbs also used for medicinal purposes.
Mexican American	<ul style="list-style-type: none"> Celebrate with food. Food rich in color, flavor, and spice. Rice, beans, and tortillas are staples. Hot and cold theory applies to food.
Navajo Native American	<ul style="list-style-type: none"> Food used to celebrate life events. Corn is a staple. Many have lactose intolerance and do not drink milk, leading to vitamin D deficiency.
Vietnamese American	<ul style="list-style-type: none"> Because of people's smaller size, normal caloric intake two-thirds less than that of other Americans. Rice is main staple. Hot and cold theory applies to food. High incidence of lactose intolerance.

Source: Adapted from Purnell, L., and Paulanka, B. (2003). *Transcultural Health Care: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.

all affect nutrition. Illness and trauma stimulate the stress response and increase nutritional requirements. Vomiting and diarrhea can cause fluid and electrolyte loss. Febrile illnesses accelerate metabolic processes and insensible fluid loss. Energy and fluid requirements are also greater during infections and febrile illnesses. Certain diseases, such as DM, cystic fibrosis, and celiac disease, are linked to specific nutritional deficits.

Ask about changes in diet or weight. Diet changes may be a result of physical, economic, or other factors that could contribute to malnutrition. Weight gains can occur with certain endocrine problems, and weight loss may accompany cancer, DM, and hyperthyroidism. Sudden weight changes are more likely to be related to fluctuations in hydration status caused by such conditions as congestive heart failure (CHF) or severe diarrhea.

Ask about prescription and over-the-counter (OTC) medications. Many of these drugs adversely affect nutrition. Note the patient's use of diuretics because they can cause non-nutritional weight changes. (See *Drugs That Adversely Affect Nutritional Status*.)

Past Health History

Ask the person if she or he has experienced a major illness, surgery, or trauma. These are usually associated with increased nutritional needs, and depending on how recently they occurred and the person's recovery status, they may continue to pose a nutritional risk. Also ask if she or he has any chronic conditions, such as cancer, that could affect utilization of nutrients. In addition, inquire about dental or oral problems. Loss of teeth or pain and

Drugs That Adversely Affect Nutritional Status

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Antacids	<ul style="list-style-type: none"> ■ Calcium carbonate ■ Magnesium hydroxide, magnesium oxide ■ Sodium bicarbonate 	<ul style="list-style-type: none"> ■ Hypercalcemia, milk-alkali syndrome ■ Hypermagnesemia ■ Metabolic alkalosis
Anti-infectives	<ul style="list-style-type: none"> ■ Amphotericin B ■ Aminoglycosides ■ Tetracyclines 	<ul style="list-style-type: none"> ■ Hypokalemia, hypomagnesemia, reversible normochromic, normocytic anemia ■ Hypocalcemia, hypokalemia, hypomagnesemia, hyponatremia ■ Impaired absorption of calcium, magnesium, and iron
Antilipemics	<ul style="list-style-type: none"> ■ Sulfasalazine ■ Cholestyramine resin ■ Niacin 	<ul style="list-style-type: none"> ■ Impaired absorption of folic acid ■ Impaired absorption of folic acid, calcium, iron, and fat-soluble vitamins (A, D, E, and K); steatorrhea ■ Hyperglycemia and abnormal glucose tolerance with long-term use of high doses
Antineoplastics	<ul style="list-style-type: none"> ■ Colestipol ■ Cisplatin ■ Cyclophosphamide 	<ul style="list-style-type: none"> ■ Impaired absorption of vitamins A, D, E, and K ■ Hypokalemia, hypomagnesemia ■ Hypokalemia, hyponatremia
Corticosteroids	<ul style="list-style-type: none"> ■ Prednisone, methylprednisolone, dexamethasone 	<ul style="list-style-type: none"> ■ Sodium retention, hypokalemia, hypocalcemia
Diuretics	<ul style="list-style-type: none"> ■ Thiazides ■ Loops 	<ul style="list-style-type: none"> ■ Hypokalemia, hypochloremic alkalosis, hypercalcemia, hypomagnesemia ■ Hypokalemia, hypomagnesemia, hypochloremic metabolic alkalosis, hyponatremia, hypocalcemia
Miscellaneous agents	<ul style="list-style-type: none"> ■ Mannitol ■ Amiloride, spironolactone, triamterene ■ Alcohol ■ Chlorpropamide, tolbutamide ■ Oral contraceptives ■ Orlistat 	<ul style="list-style-type: none"> ■ Fluid and electrolyte imbalances ■ Hyperkalemia ■ Decreased vitamin B₁₂ absorption; thiamine, magnesium, or folic acid deficiencies ■ Hyponatremia ■ Decreased glucose tolerance, altered carbohydrate and lipid metabolism, folic acid or B₆ deficiency ■ Impaired absorption of fat-soluble vitamins (A, D, E, and K)

discomfort associated with eating can significantly impair ingestion. Last, ask about recent camping trips or foreign travel because of the risk of food- and waterborne pathogens.

Family History

Certain genetic and hereditary conditions affect digestion, absorption, and metabolism. Ask the person if he or she has a history of food intolerance or allergies, such as lactose intolerance. Ask about hereditary or familial

health problems, such as Crohn's disease, DM, or cystic fibrosis; or anemias, such as thalassemia. Inquire about a family history of cardiovascular disease, atherosclerotic disease, or obesity.

Review of Systems

The review of systems (ROS) provides a focused screening for past and present problems related to or affecting each of the physical systems. It may also identify problems or symptoms that indicate a nutritional risk.

Review of Systems

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>General Health Survey</p> <ul style="list-style-type: none"> ■ Have you had unexplained weight loss, fatigue, activity intolerance, or inability to concentrate? 	<ul style="list-style-type: none"> ■ May indicate metabolic problems (e.g., DM, hyperthyroidism, or malignancy), iron deficiency anemia, or dehydration.
<p>Integumentary</p> <ul style="list-style-type: none"> ■ Have you noticed changes in skin texture, skin discolorations, poor wound healing, or bruising? 	<ul style="list-style-type: none"> ■ May signal vitamin and mineral deficiencies, protein malnutrition, or metabolic disorders.
<p>HEENT</p> <p>Eyes</p> <ul style="list-style-type: none"> ■ Do you have poor night vision or eye dryness? <p>Nose, Mouth, and Throat</p> <ul style="list-style-type: none"> ■ Do you have nosebleeds or bleeding gums? ■ Cavities or lost teeth? <p>Cardiovascular</p> <ul style="list-style-type: none"> ■ Do you have chest pain or pressure? 	<ul style="list-style-type: none"> ■ May indicate vitamin A deficiency or dehydration. ■ Possible vitamin, mineral, or protein deficiencies. ■ May indicate excess sugar intake. ■ <i>Cardiac disease:</i> May be caused by high cholesterol, elevated triglycerides, or high-calorie diet.
<p>Gastrointestinal</p> <ul style="list-style-type: none"> ■ Are you constipated? ■ Do you have diarrhea? 	<ul style="list-style-type: none"> ■ <i>Constipation:</i> Low-fiber diet, poor fluid intake. ■ <i>Diarrhea:</i> High-fiber diet, food allergies, hereditary and genetic disorders (e.g., cystic fibrosis).
<p>Reproductive</p> <p>Women</p> <ul style="list-style-type: none"> ■ Have you had frequent miscarriages or irregular menses? ■ How much caffeine do you consume a day? <p>■ What is your diet like?</p> <p>Men</p> <ul style="list-style-type: none"> ■ Do you suffer from impotence? <p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Do you have muscle weakness? <p>Neurological</p> <ul style="list-style-type: none"> ■ Are you nervous or irritable? Do you have headaches, numbness or tingling, or muscle tics? <p>Lymphatic</p> <ul style="list-style-type: none"> ■ Do you have frequent infections? ■ Allergies? 	<ul style="list-style-type: none"> ■ Maternal malnutrition can lead to fetal growth restriction. ■ Consumption of 300 mg of caffeine/d increases risk for spontaneous abortion. ■ Eating disorders can result in irregular menses. ■ Alcohol intake increases risk for impotence. ■ May be caused by potassium deficiency or dehydration. ■ Possible vitamin B deficiencies or hypocalcemia. ■ <i>Frequent infections:</i> Protein malnutrition. ■ <i>Allergies:</i> Food sensitivity (e.g., lactose).

Psychosocial Profile

Because physical problems are frequently assumed to have a physical origin, nurses may underestimate the relation-

ship of everyday life to health. Taking a psychosocial profile can yield valuable clues about a person's nutritional status. Helping the person change unhealthy lifestyle behaviors may also be the most viable avenue of intervention.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Health Practices and Beliefs/Self-Care Activities</p> <ul style="list-style-type: none"> ■ Who shops and prepares meals in your family? ■ Are you able to plan and cook meals yourself, or do you depend on others to do this? 	<ul style="list-style-type: none"> ■ Dependent people may not eat properly if caregivers are lax or uninformed about good nutrition. ■ Diminished mental capacity or physical limitations from aging or debilitation may impair ability to buy, prepare, or eat food.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ When was your last dental exam? ■ Do you go for routine physical exams? <p>Typical Day</p> <ul style="list-style-type: none"> ■ How do you usually spend your days? Do you go to restaurants frequently, and if so, where do you go and what types of foods do you eat? How much time do you spend shopping, preparing, and eating food? 	<ul style="list-style-type: none"> ■ Good dentition is required for proper mastication and digestion of food. ■ Identifies teaching needs. ■ Shows importance of food and good nutrition in person's life, what type of food person normally eats, and how often and where she or he eats out.
<p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ What is your typical daily diet? How much water do you drink daily? ■ Do you actively try to maintain good nutrition and healthy weight? How? ■ Have your eating habits and appetite changed recently? 	<ul style="list-style-type: none"> ■ Can immediately point to malnutrition and/or poor hydration. ■ Indicates conscious modifications of diet, although not necessarily correct information or awareness of health risks. ■ Appetite changes can accompany depression, medication therapy, and some physical illnesses.
<p>Activity and Exercise Patterns</p> <ul style="list-style-type: none"> ■ What is your usual activity and exercise level? ■ Do you feel that you have adequate energy? What activities do you not pursue because of lack of energy? 	<ul style="list-style-type: none"> ■ Validates that the person has a normal, active lifestyle. ■ Some changes may be a normal part of aging. Loss of energy or inability to pursue simple functional activities can accompany iron and protein/calorie deficiencies.
<p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ How many hours of sleep do you get a night? Do you wake up during the night? Are you taking more naps than usual? Do you feel rested? Do you use sleep aids? If so, what kinds? 	<ul style="list-style-type: none"> ■ Hunger or mild dehydration can interrupt sleep. Lack of energy or excessive sleep can be caused by iron or protein/calorie deficiency.
<p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you smoke? ■ Do you drink alcohol? If so, how much/day? Do you drink coffee, tea, or soft drinks every day? If so, how much? ■ Do you use OTC drugs? ■ Do you use illegal drugs? 	<ul style="list-style-type: none"> ■ Smoking interferes with absorption of vitamin C. ■ Alcohol and soft drinks are non-nutritive food sources. Excessive amounts impede adequate nutrition. Caffeine is mildly addictive. Withdrawal can cause nervousness and irritability. Excessive intake can contribute to calcium deficiency. ■ Some OTC drugs impair nutrient absorption and metabolism. ■ Artificial sweeteners help avert obesity but have potential cancer risks. ■ Purchase of food may be sacrificed to support an addictive drug habit.
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ What do you do for a living? ■ How does your job affect daily meal routines? 	<ul style="list-style-type: none"> ■ Exposure to chemical substances (e.g., lead) can interfere with absorption/metabolism of nutrients. ■ Rotating or night shifts can interfere with meal routines and planning. People may substitute fast or convenience foods for more appropriate food sources.
<p>Socioeconomic Status</p> <ul style="list-style-type: none"> ■ Is your income adequate to meet your food needs? 	<ul style="list-style-type: none"> ■ Good-quality food and food variety are expensive. Poverty and homelessness put people at serious risk for malnutrition. A significant number of the poor and homeless are women, children, and adolescents. Alcoholics and the chronically mentally ill are another large component of the homeless population.

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Environmental Health Patterns</p> <ul style="list-style-type: none"> Where do you buy your food? How do you store it? How do you prepare it? <p>Roles and Relationships</p> <ul style="list-style-type: none"> Do you have regular social interaction? Do you usually eat with other people or alone? <p>Cultural Influences</p> <ul style="list-style-type: none"> Do you have any cultural influences that may affect your dietary practices? If so, what are they? <p>Religious/Spiritual Influences</p> <ul style="list-style-type: none"> Do you have any religious influences that may affect your dietary practices? If so, what are they? <p>Stress and Coping</p> <ul style="list-style-type: none"> Are your social relationships satisfying or stressful? How much stress do you have in your life? How do you cope with it? 	<ul style="list-style-type: none"> Contamination and spoilage may be a concern. Poor hygiene and inadequate cooking increase risk of foodborne pathogens. Eating and preparing food have a strong basis in culture and serve important social functions. Changes in this structure (e.g., the death of a spouse) can affect nutrition. Different cultures have different dietary preferences that can affect overall nutrition. Some religions restrict certain foods, which can affect overall nutrition. Food can become a substitute for satisfying relationships. Emotional or psychological stress can weaken the immune system. Loss of appetite can occur with stress.

Comprehensive Nutritional History

If the person has one or more known nutritional risks, perform a comprehensive nutritional history. If there is no time to perform a comprehensive history, perform a focused history. Although nutritional patterns are learned early in life and are hard to break, weight and nutrition are major health issues in Western cultures. Information on dieting, exercise, and nutrition pervades the news media, along with the message that people should take responsibility for making healthy changes. Consequently, many people worry that they will be scolded for poor nutritional practices, so they withhold or embellish information. So remember to convey an attitude of acceptance and caring, and never be authoritarian or paternalistic.

Two dietary analysis techniques are discussed—24-hour recall and food intake records. You can use either technique as part of your comprehensive history.

24-Hour Recall

Ask the person to write down what he or she ate and drank during the previous 24 hours. Then use the Food Guide Pyramid to sort and categorize the foods and determine the general quality of his or her diet. People often have trouble accurately recalling what they ate, so prompt them by saying, “Start with the first thing you had when you got out of bed.” Also ask them to record between-meal drinks and snacks, desserts, bedtime snacks, condiments (e.g., mayonnaise, butter, sugar, or

Focused Nutritional History

If the situation or time prohibits a detailed nutritional history, focus on the person's weight, symptoms of malnutrition, and identification of nutritional risk factors. The person's response will help you appropriately focus your assessment. Key questions include:

- Have you lost or gained weight unintentionally in the past 6 months?
- What is the most you have ever weighed?
- How much did you weigh 6 months ago, and how much do you weigh now?
- What do you normally eat every day? Has your diet changed significantly? If so, how?
- Do you have any stomach or bowel symptoms (e.g., nausea, vomiting, diarrhea, or anorexia) that have lasted more than 2 weeks?
- Has anything happened in your life that has affected your ability to obtain or prepare food? If so, what?

cream), and food preparation items (e.g., cooking oil or lard). Water is critical to nutritional metabolic processes, so ask them to record water intake, too. Last, be sure to have them record the amount of each food or liquid they consumed, and translate these into standard servings according to the Food Pyramid. A person's typical serving may actually equal several servings.

The 24-hour recall is a valuable screening and assessment tool only if it represents the person's typical daily intake. If a person is ill or has a change in routine, her or his intake will vary from normal. So be sure to ask if what she or he recorded represents a typical day. If she or he says no, ask her or him to substitute foods typically eaten.

Use the Food Pyramid to categorize 24-hour-recall data. Then tabulate food items alongside each corresponding Pyramid group, and document them in the person's health record.

Food Intake Records

Food intake records are typically done on people who are debilitated, have severe burns, or are on chemotherapy. A food intake record is a quantitative listing of all food and fluid consumed within a designated time frame—usually 3 to 5 days. Because food intake patterns often change on weekends and holidays, records kept during outpatient or home-care situations should reflect one atypical day for a 3-day period and one weekend for a 5-day period. Again, be sure to clarify serving sizes.

To analyze the data, reduce the recorded food items into their constituent nutrients, using USDA food composition tables. To get a daily average intake for each nutrient, add up the total nutrients and divide by the number of days the record was kept. Evaluate the averages against the RDAs. Two-thirds of the RDAs is considered adequate for the general healthy population. However, acceptable levels may vary in disease, risk, or deficiency situations and will be reflected in therapeutic treatment decisions.

A less specific but more practical approach involves analyzing the patient's food intake record using food labels on packages. Once you have determined the total number of calories consumed, calculate the total grams of carbohydrate, protein, and fat and the percentage of caloric intake contributed by each of these. Protein should compose 10 percent of the diet; fat, 20 to 30 percent, and carbohydrates, the remaining calories.



Case Study Findings

Reviewing Mr. Thomas'

Biographical data:

- 78-year-old African American male.
- Wife died 6 months ago.
- Lives alone in urban area.
- Retired postal worker.

- Baptist religion.
- Medicare/Medicaid health insurance.
- Daughter is primary contact.

Current health status:

- Vague complaints of not feeling well, being very tired, and having no energy; 10-lb weight loss since wife's death.
- Current medications are Lasix 40 mg od and Vasotec 5 mg bid. No OTC medications except occasional milk of magnesia (MOM) for constipation. No illegal drug use.

Past health history:

- Positive history of HTN for 10 years.
- Hospitalized last year for uncontrolled HTN.
- Appendectomy at age 12.
- No allergies to food, drugs, or environmental factors.
- Medications listed previously.

Family history:

- Daughter, age 50, alive and well.
- Wife, deceased age 74, breast cancer.
- Mother, deceased age 75, HTN, stroke.
- Father, deceased age 80, HTN, myocardial infarction (MI).
- Sister, age 70, HTN.
- Sister, age 73, alive and well.
- Brother, age 75, DM, HTN, coronary artery disease (CAD).
- Maternal aunt, deceased age 80, CAD, MI.
- Maternal uncle, deceased age 78, stroke.
- Paternal aunt, deceased age 82, colon cancer.
- Paternal uncle, deceased age 81, stroke.

Review of systems:

- *General Health Survey*: 10-lb weight loss, fatigue.
- *Integumentary*: Dry, flaky skin.
- *Head, Eyes, Ears, Nose, and Throat (HEENT)*: Wears dentures.
- *Respiratory*: No changes.
- *Cardiovascular*: No chest pain; positive history of HTN.
- *Gastrointestinal*: Bowel movement every 2 to 3 days, brown and formed. Takes MOM as needed.
- *Genitourinary/Reproductive*: On Lasix; no problems with urination; voids about 8 times/day. Not sexually active since wife died.
- *Musculoskeletal*: Weakness and occasional cramping in legs.
- *Neurological*: No sensory changes. States that he is very lonely since wife's death.
- *Lymphatic*: No allergies or recent infections.

Psychosocial profile:

- Finds it hard to cook for one person; wife used to prepare meals.
- Says that his dentures are a little loose since he lost weight.
- Visits doctor every 6 months for BP check.
- Typical day: Awakens at 7 A.M., has breakfast, walks to corner for newspaper, takes midmorning nap, usually skips lunch, watches TV, has dinner around 5 P.M., watches TV after dinner, usually falls asleep while watching TV, goes to bed after the 10 P.M. news.
- Has three 8-oz glasses of water a day, not trying to lose weight. Eats out about once a week, likes fried foods. Not much appetite since wife died. Usually eats alone.
- Daily walk to corner for paper, no other scheduled exercise.
- Sleeps 7 to 8 hours a night. Awakens at least once to go to bathroom. Takes two 20- to 30-minute naps during day.
- Never smoked or used drugs. Occasional alcoholic drink about 2 times a month.
- Retired postal worker on fixed income with government pension; able to meet financial needs.
- Most close friends have passed away.
- No cultural or religious influence on dietary habits.
- Says life is “pretty routine without much stress.”
- Daughter is main source of support; lives 1 hour away, visits once a week, takes him shopping.

24-hour recall:

- **Breakfast:** 8 ounces black coffee, 8 ounces water, bowl of Cheerios with whole milk.
- **Lunch:** None.
- **Afternoon snack:** 4 pretzels.
- **Dinner:** Frozen prepared dinner of Salisbury steak, potatoes, and corn, piece of apple pie, 8 oz water, 8 oz black coffee.

**Case Study Evaluation**

Before you conduct an objective health assessment on Mr. Thomas, review his health history information from the comprehensive nutritional assessment.

**CRITICAL THINKING ACTIVITY #3**

What information from the health history indicates that Mr. Thomas is at risk for or has a nutritional deficiency?

Physical Assessment

Now proceed to the objective part of the assessment. Findings from the health history will determine the depth and scope of your physical examination. As you perform the examination, be alert to findings in various body systems that might signal malnutrition. Remember, determination of malnutrition cannot be made on physical findings alone. Many diseases and disorders mimic nutritional deficiencies. Always corroborate your physical findings with the health history and the results of laboratory assessments. The assessment includes performing a head-to-toe scan and taking various anthropometric measurements.

Approach

You will mainly use the techniques of inspection and palpation. Examining the skin and mucous membranes is crucial, but additional data are derived from assessment of other body systems.

Optimal nutrition cannot occur without adequate hydration. Therefore, evaluate the person's hydration status simultaneously. Red flags include reports of minimal fluid intake, excessive thirst or excessive fluid intake, increased urination, diarrhea, or diuretic use. Very young infants, very frail older adults, and chronically ill and debilitated people are less tolerant of fluid loss and are at particular risk for dehydration from vomiting and/or diarrhea.

**TOOLBOX**

The tools for a comprehensive nutritional assessment include a weight scale, flexible measuring tape, calipers, and stethoscope. You should also have access to growth charts, weight and height tables, anthropometric tables, and laboratory values. To help the person accurately estimate quantities of food, have a variety of containers on hand.

Performing a Head-to-Toe Physical Assessment

Look for changes in every system that might signal a nutritional problem. (See *Performing a Head-to-Toe Physical Assessment*.)

Anthropometry

Anthropometry literally means *human measurement*. It includes measuring overall body mass (particularly growth, fat reserves, and somatic protein stores) and evaluation of related laboratory values. Growth charts that plot height, weight, and head growth are used for children up to age 18. By adulthood, growth has stabilized and ratio measurements of body mass are used. The following sections describe common anthropometric techniques for assessing nutrition in children and adults.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL VARIATIONS	ABNORMAL FINDINGS/RATIONALE
<p>General Health Survey</p> <ul style="list-style-type: none"> ■ No unexplained weight changes. ■ Vital signs within normal limits for person's age. 	<ul style="list-style-type: none"> ■ Weight changes within a short period of time usually reflect fluid loss or gain (1 mL = 1 g). ■ Elevated BP may reflect fluid overload, high sodium intake, and obesity. Low BP may reflect dehydration.
<p>INSPECT/PALPATE</p>	
<p>Integumentary</p>	
<p>Skin</p>	
<ul style="list-style-type: none"> ■ Skin intact, warm, and dry. ■ Texture smooth, no lesions. ■ Color consistent with ethnicity. 	<ul style="list-style-type: none"> ■ <i>Scaling</i>: Low or high vitamin A, zinc, essential fatty acids. ■ <i>Transparent, cellophane appearance</i>: Protein deficit. ■ <i>Cracking (flaky paint, cracked-pavement appearance)</i>: Protein deficit. ■ <i>Follicular hyperkeratosis</i>: Vitamins A, C deficits. ■ <i>Petechiae (especially perifollicular)</i>: Vitamin C deficit. ■ <i>Purpura</i>: Vitamins C, K deficits. ■ <i>Pigmentation/desquamation of sun-exposed areas</i>: Niacin deficit. ■ <i>Edema</i>: Protein and thiamine deficits. ■ <i>Skin lesions, ulcers, wounds that do not heal</i>: Protein, vitamin C, zinc deficits. ■ <i>Yellow pigmentation (except sclerae)</i>: Excess carotene (benign). ■ <i>Poor skin turgor</i>: Dehydration.
<p>Hair</p>	
<ul style="list-style-type: none"> ■ Even hair distribution; no alopecia. 	<ul style="list-style-type: none"> ■ <i>Transverse pigmentation of hair shaft</i>: Protein deficit. ■ <i>Hair easy to pluck, breaks easily</i>: Protein deficit. ■ <i>Sparse hair distribution</i>: Protein, biotin, zinc deficits, excess vitamin A. ■ <i>Corkscrew hairs, unemerged hair coils</i>: Vitamin C deficit.
<p>Nails</p>	
<ul style="list-style-type: none"> ■ Nails smooth and pink. 	<ul style="list-style-type: none"> ■ <i>Transverse ridges in nails</i>: Protein deficit. ■ <i>Concave "spoon" nails</i>: Iron deficit.
<p>HEENT</p>	
<p>Eyes</p>	
<ul style="list-style-type: none"> ■ Eyes clear and bright, vision intact. 	<ul style="list-style-type: none"> ■ <i>Papilledema</i>: Vitamin A excess. ■ <i>Night blindness</i>: Vitamin A deficit. ■ <i>Sunken eyeballs, dark circles, decreased tears</i>: Dehydration. ■ <i>Sunken fontanels</i>: Dehydration.
<p>Nose</p>	
<ul style="list-style-type: none"> ■ Sense of smell intact. ■ Nasal mucosa pink, moist, and intact without lesions. 	<ul style="list-style-type: none"> ■ Anosmia can affect taste. ■ <i>Hyposmia (defect in sense of smell)</i>: Zinc deficit.
<p>Mouth</p>	
<ul style="list-style-type: none"> ■ Oral mucosa pink, moist, and intact without lesions. Tonsils pink; gums pink and intact with no bleeding. 	<ul style="list-style-type: none"> ■ <i>Angular stomatitis</i>: Riboflavin, niacin, pyridoxine deficits. ■ <i>Cheilosis (dry, cracked, ulcerated lips)</i>: Riboflavin, niacin, pyridoxine deficits. ■ <i>Atrophic lingual papillae (coated tongue)</i>: Riboflavin, niacin, folic acid, vitamin B₁₂, protein, iron deficits. ■ <i>Glossitis (raw, red tongue)</i>: Riboflavin, niacin, pyridoxine, folic acid, vitamin B₁₂ deficits. ■ <i>Hypogeusia (blunting of sense of taste)</i>: Zinc deficit.
<ul style="list-style-type: none"> ■ Taste sensation intact. 	

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL VARIATIONS

- Swallow and gag reflex.
- Full mobility of tongue.

Respiratory

- Lungs clear, respirations within normal limits.

Cardiovascular

- Regular heart rate/rhythm. PMI 1 cm at apex. No extra sounds.

Gastrointestinal

- Abdomen soft, nontender, no organomegaly.

Musculoskeletal

- No deformities, tenderness, or swelling.

Neurological

- No headache.
- AAO \times 3.
- Deep tendon reflexes (DTRs) + 2/4.
- Senses intact.

ABNORMAL FINDINGS/RATIONALE

- Absent swallow and gag reflexes may make eating difficult.
- *Dysphagia*: Impaired tongue mobility.
- *Swollen, retracted, bleeding gums (if teeth present)*: Vitamin C deficit.
- *Dry mucous membranes*: Dehydration.
- *Parotid gland enlargement*: Protein deficit (consider bulimia).
- *Increased respiratory rate*: Iron deficiency anemia.
- *Heart failure, S₃*: Thiamin, phosphorus, iron deficits.
- *Sudden heart failure, death*: Vitamin C deficit.
- *Tachycardia and systolic murmur*: Iron deficiency anemia.
- *Hepatomegaly, ascites*: Protein deficits, vitamin A excess.
- *Hyperactive bowel sounds*: May indicate hyperperistalsis associated with an absorption problem.
- *Beading of ribs, epiphyseal swelling, bowlegs*: Vitamin D deficit.
- *Tenderness, superperiosteal bleeding*: Vitamin C deficit.
- *Headache*: Vitamin A excess.
- *Drowsiness, lethargy, vomiting*: Vitamin A, D excess.
- *Dementia*: Niacin, vitamin B₁₂ deficit.
- *Confusion, irritability*: Dehydration.
- *Disorientation*: Thiamin (Korsakoff's psychosis) deficit.
- *Ophthalmoplegia*: Thiamin, phosphorus deficit.
- *Peripheral neuropathy (weakness, paresthesia, ataxia, and decreased DTRs), and diminished tactile, vibratory, and position sensation*: Niacin, pyridoxine, vitamin B₁₂ deficits.
- *Tetany, increased DTRs*: Calcium, magnesium deficits.

Growth Charts

Children

Growth charts for height, weight, and head circumference (see Appendix B) are excellent indicators of nutrition in children because they allow you to visualize the child's growth progress. Two sets of charts are commonly used: one for birth to 36 months (includes head circumference) and one for age 2 to 18 (does not include head circumference because cerebral growth is complete by age 2). New charts have also been developed to track the unique growth patterns of premature infants.

Until age 2, take growth measurements with the child nude or wearing only a diaper and lying supine. For children age 2 to 18, take a standing height without shoes, with the patient dressed in usual examination clothing.

Record measurements on the corresponding axes of the chart at the point that intersects with the child's current age. To visualize a growth pattern, take serial measurements. The important factor is the relative consistency of the child's growth within the norms of the curve.

As long as a child receives adequate nutrition, his or her growth largely reflects genetic heritage. If a child shows a consistent decline into a lower percentile or falls below the fifth percentile, suspect undernutrition. Suspect overnutrition in children who begin to deviate into higher percentiles or who fall above the 95th percentile. Growth chart abnormalities can also signify problems of a non-nutritional nature, such as endocrine disorders. If your history does not support a nutritionally related growth problem, a medical referral is warranted.

Adults

By age 18, growth is largely complete and weight-for-height tables replace growth charts. The Metropolitan Life Insurance Company's height and weight tables, revised in 1983 (see Appendix B), are the standard, although they do not reflect ideal weights. Instead, they represent the weights of people with the most longevity in each height category and recommend weight ranges based on height and skeletal frame size. Figures 8.2 and 8.3 illustrate height and weight measurement.



HELPFUL HINT

To estimate frame size, have the person wrap the thumb and middle finger of one hand around her or his opposite wrist. If the fingers overlap by 1 cm or more, she or he has a small frame; if the fingertips meet, a medium frame; and if the fingertips do not meet by 1 cm or more, a large frame.

Body Mass Index

Body mass index (BMI) is an accurate indicator of fat in adults. The most commonly used BMI is Quetelet's Index, which is obtained by dividing weight in kilograms by height in meters squared. BMIs between 20 and 25 kg/m² are associated with the least mortality; BMIs under 16 kg/m² are associated with eating disorders. The relatively larger proportion of muscle in athletes and body builders and the greater blood and tissue volume in pregnant and lactating women make BMI measurements inappropriate for these groups. It is also not recommended for growing children or frail and sedentary older adults.



FIGURE 8.2. Measuring height.



FIGURE 8.3. Measuring weight.

Arm Measurements

Triceps skin fatfold (TSF) estimates body fat using a double fold of skin and subcutaneous adipose tissue from the patient's dominant arm. With the patient's arm at his or her side and elbow flexed at 90 degrees, measure and mark the midpoint overlying the triceps muscle between the elbow and the shoulder. Tell the patient to relax the arm. Then use your thumb and index finger to compress a symmetrical fold of skin and adipose tissue ½ inch above the marked site. Use calipers to measure in the middle of the skin fold at the marked site, about ½ inch below your fingers. Release the calipers, and wait 4 seconds before reading the measurement. To ensure reliability, take two to three additional measurements at least 15 seconds apart. They should not vary by more than 1 cm.

Compare fatfold measurements with equivalent age- and gender-specific percentiles (see Appendix B). Values below the 10th percentile or above the 90th percentile indicate diminished or extensive fat reserves, respectively. Figure 8.4 illustrates TSF measurement.



FIGURE 8.4. Measuring triceps skin fatfold.

Body Mass Index

BMI	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35
<i>Height</i>	<i>Weight in Pounds</i>																
4'10" (58")	91	96	100	105	110	115	119	124	129	134	138	143	148	153	158	162	167
4'11" (59")	94	99	104	109	114	119	124	128	133	138	143	148	153	158	163	168	173
5' (60")	97	102	107	112	118	123	128	133	138	143	148	153	158	163	168	174	179
5'1" (61")	100	106	111	116	122	127	132	137	143	148	153	158	164	169	174	180	185
5'2" (62")	104	109	115	120	126	131	136	142	147	153	158	164	169	175	180	186	191
5'3" (63")	107	113	118	124	130	135	141	146	152	158	163	169	175	180	186	191	197
5'4" (64")	110	116	122	128	134	140	145	151	157	163	169	174	180	186	192	197	204
5'5" (65")	114	120	126	132	138	144	150	156	162	168	174	180	186	192	198	204	210
5'6" (66")	118	124	130	136	142	148	155	161	167	173	179	186	192	198	204	210	216
5'7" (67")	121	127	134	140	146	153	159	166	172	178	185	191	198	204	211	217	223
5'8" (68")	125	131	138	144	151	158	164	171	177	184	190	197	203	210	216	223	230
5'9" (69")	128	135	142	149	155	162	169	176	182	189	196	203	209	216	223	230	236
5'10" (70")	132	139	146	153	160	167	174	181	188	195	202	209	216	222	229	236	243
5'11" (71")	136	143	150	157	165	172	179	186	193	200	208	215	222	229	236	243	250
6' (72")	140	147	154	162	169	177	184	191	199	206	213	221	228	235	242	250	258
6'1" (73")	144	151	159	166	174	182	189	197	204	212	219	227	235	242	250	257	265
6'2" (74")	148	155	163	171	179	186	194	202	210	218	225	233	241	249	256	264	272
6'3" (75")	152	160	168	176	184	192	200	208	216	224	232	240	248	256	264	272	279

The National Institutes of Health (NIH) interpret BMI values for adults with one fixed number, regardless of age or sex, using the following guidelines: Underweight: <18.5 Overweight: 25.0–29.9 Obese: 30.0 or more
Source: Evidence Report of Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults, 1998.

NIH/National Heart, Lung, and Blood Institute (NHLBI). Centers for Disease Control and Prevention, United States Department of Health and Human Services.

Midarm circumference (MAC) provides a crude estimate of muscle mass and is most useful when combined with TSF. Measure the circumference of the patient's dominant arm at the same site where you obtained the TSF. Wrap the measuring tape firmly around the patient's arm without compressing the skin. Take two or three more measurements to ensure reliability. They should not vary by more than 1 cm. Compare midarm measurements with equivalent age- and gender-specific percentiles (see Appendix B). Figure 8.5 illustrates MAC measurement.

Midarm muscle circumference (MAMC) is mathematically derived using TSF and MAC values. As an indirect measurement of muscle mass, it provides an index of protein stores. To calculate MAMC, multiply the TSF (in centimeters) by 3.143 and subtract the result from the MAC. Because MAMC estimates skeletal muscle reserves, it should generally fall within 90 percent of standard. Values falling between 60 and 90 percent suggest moderate protein deficiency and those less than 60 percent indicate severe malnutrition. Compare MAMC measurements with equivalent age- and gender-specific percentiles (see Appendix B).



FIGURE 8.5. Measuring midarm circumference.

Waist-to-Hip Ratio

The **waist-to-hip ratio (WHR)** estimates obesity by evaluating the amount of abdominal fat. People with a greater proportion of upper body fat are at greater risk for HTN, DM, elevated triglycerides, and other atherosclerotic risk factors. WHR is calculated simply by dividing the waist circumference by the hip circumference. A WHR of 1.0 or greater in men and 0.8 or greater in women indicates upper body obesity.



Case Study Findings

Your general health survey and head-to-toe examination of Mr. Thomas have concluded. The following are your findings:

- **General Health Survey:** Awake, alert, and oriented (AAO) \times 3; cooperative, neatly dressed 76-year-old black male who appears stated age. Affect sad, poor eye contact, responds appropriately.
- **Vital Signs:** BP 130/70; pulse 100 and regular; respirations 20; temperature 97.5°F; height 5'11"; weight 175; 10-lb weight loss since last visit 6 months ago.
- **Integumentary:** Skin dry and flaky, hair evenly distributed, no alopecia.
- **HEENT:** Dry, pink mucous membranes, dentures poorly fitted.
- **Respiratory:** Breath sounds equal bilaterally, clear to auscultation but decreased at bases because of poor inspiratory effort.
- **Cardiovascular:** Normal heart rate and rhythm, $+S_4$, point of maximum impulse (PMI) 2 cm at apex.

- **Gastrointestinal:** Bowel sounds present, regular, slightly diminished; abdomen soft, nontender, with no masses or organomegaly.
- **Musculoskeletal:** +4 muscle strength on upper and lower extremities.
- **Neurological:** Cranial nerves intact; deep tendon reflexes +2, senses intact.



Case Study Analysis and Plan

Now that you have completed your comprehensive nutritional assessment of Mr. Thomas, list key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mr. Thomas, and then use this information to develop a list of nursing diagnoses. Some possible ones are listed. Cluster the supporting data.

1. Nutrition: imbalanced, less than body requirements, related to inadequate iron intake
2. Fluid Volume, deficient, related to medications and poor intake
3. Knowledge, deficient, related to nutrition

Identify any additional nursing diagnoses and any collaborative nursing diagnoses.



CRITICAL THINKING ACTIVITY #4

Now that you have identified some nursing diagnoses for Mr. Thomas, select one from above and develop a brief teaching plan for him. Include learning outcomes and teaching strategies.

RESEARCH TELLS US

Metabolic syndrome, insulin resistance syndrome or syndrome X, is characterized by dyslipidemia (elevated triglycerides and low HDLs), hypertension, impaired glucose intolerance and central obesity. Metabolic syndrome is a major risk factor for developing heart disease and is a predictor for Type II diabetes.

Obesity is a major health problem in the United States, with more than half of all adults being overweight or obese. Treatment is aimed at each component of the syndrome with dietary changes, exercise and medication. Dietary changes include:

- decrease caloric intake and portion size to achieve weight loss

- decrease saturated fat to decrease insulin resistance
- decrease sodium intake to control high blood pressure
- decrease triglycerides to lower glycemic index.

Assessment should be aimed at identifying those at risk and those with metabolic syndrome. Assessment includes:

- measuring height and weight to calculate BMI
- measuring waist circumference.

Nurses are involved in all aspect of healthcare and at all levels, placing them in a key position to screen and educate those who are at risk for developing Metabolic Syndrome.

Diagnostic Criteria for Metabolic Syndrome

WHO Diagnostic Criteria (insulin resistance plus two of the following):

- **Abdominal/central obesity:** Waist to hip ratio: >0.90 (men), >0.85 (women), or BMI >30 kg per m²
- **Hypertriglyceridemia:** ≥150 mg/dL (≥1.7 mmol/L)
- **Low HDL cholesterol:** <35 mg/dL (<0.9 mmol/L) for men, <39 mg/dL (<1.0 mmol/L) in women
- **High blood pressure:** ≥140/90 Hg or documented use of antihypertensive therapy
- **High fasting glucose:** Impaired glucose tolerance, impaired fasting glucose, insulin resistance, diabetes
- **Microalbuminuria:** Urinary albumin to creatinine ratio: 30 mg/g, or albumin excretion rate: 20 mcg/minute

Adult Treatment Panel Diagnostic Criteria (three of the following):

- **Abdominal/central obesity:** Waist circumference: >102 cm (40 in.) in men, >88 cm (35 in) in women
- **Hypertriglyceridemia:** ≥150 mg/dL
- **Low HDL cholesterol:** <40 mg/dL (1.036 mmol/L) for men, <50 mg/dL (<1.295 mmol/L) for women
- **High blood pressure:** ≥130/85 Hg or documented use of antihypertensive therapy
- **High fasting glucose:** ≥110 mg/dL (≥6.1 mmol/L)

When reviewing the above diagnostic criteria, please note the following:

- Insulin resistance is type 2 DM or impaired fasting glucose.
- American Diabetic Association suggest lowering fasting threshold to 100.

Source: Deen, D. (2004). Metabolic syndrome: Time for action. *American Family Physician* 69 (12) June 15, pp. 2875–2882, 2887–2888.

HEALTH CONCERNS**F**acts about obesity and disease

- Obesity affects roughly 35 percent of women and 31 percent of men in the United States. This is an increase of one-third over the past decade, and the rate continues to rise.
- Genetics account for only 33 percent of variations in body weight and has a greater overall influence on fat distribution than on total body fat.
- Moderate obesity, particularly abdominal obesity, can increase the risk of non-insulin-dependent DM (adult onset) by tenfold.
- The risk of cardiovascular disease increases with increasing levels of obesity, independent of other risk factors.
- Overweight men have a significantly higher mortality rate for colorectal and prostate cancer.
- Overweight women are three times as likely to have breast, uterine, cervical, and ovarian cancer and are seven times more likely to have endometrial cancer.
- Obese girls begin to menstruate at a younger age than normal-weight girls.
- Obese women have more irregular menstrual cycles and a greater frequency of other menstrual abnormalities than normal-weight women.
- Overweight women have more problems during pregnancy, including toxemia and HTN.
- Overweight people have an increased risk of developing gallbladder disease.
- Increased weight causes more wear and tear on the joints, leading to osteoarthritis.
- As weight increases, breathing at night becomes progressively more difficult, leading to sleep apnea or Pickwickian syndrome (obesity hypoventilation syndrome).
- In the United States, obesity-related illness is second only to smoking-related illness as a leading cause of death.

Common Abnormalities**ABNORMALITY****AREA****ASSESSMENT FINDINGS****Anorexia Nervosa**

A preoccupation with being thin and dieting, leading to excessive weight loss.

- General Health Survey
- Behavioral

- Continued dieting despite weight loss.
- Feeling fat despite weight loss
- Intense fear of weight loss.
- Preoccupation with food, especially calories and fat content.
- Eating in isolation, lying about food.

(continued)

Common Abnormalities (continued)

ABNORMALITY	AREA	ASSESSMENT FINDINGS
Bulimia Nervosa Uncontrolled binge eating (excessive quantities of food consumed in one sitting) alternated with purging (ridding the body of food) in an attempt to lose weight. It is characterized by intense feelings of guilt and shame.	<ul style="list-style-type: none"> ■ Integumentary 	<ul style="list-style-type: none"> ■ Exercising compulsively. ■ Cold intolerance, cold hands and feet. ■ Weakness and exhaustion. ■ Dry, brittle skin and hair; hair loss. ■ Growth of fine body hair on extremities. ■ Blotchy or yellow skin.
	<ul style="list-style-type: none"> ■ Respiratory ■ Cardiovascular 	<ul style="list-style-type: none"> ■ Shortness of breath. ■ Low BP. ■ Dysrhythmias.
	<ul style="list-style-type: none"> ■ Gastrointestinal ■ Musculoskeletal 	<ul style="list-style-type: none"> ■ Constipation. ■ Skeletal muscle atrophy. ■ Loss of fatty tissue.
	<ul style="list-style-type: none"> ■ Neurological ■ Genitourinary 	<ul style="list-style-type: none"> ■ Depression, anxiety. ■ Amenorrhea.
	<ul style="list-style-type: none"> ■ Behavioral 	<ul style="list-style-type: none"> ■ Preoccupation with body weight. ■ Uncontrolled eating alternating with periods of strict dieting and exercise. ■ Inducing vomiting or taking laxatives to lose weight. ■ Frequent use of bathroom after a meal.
Diabetes Mellitus A disorder in the metabolism of carbohydrates caused by decreased production of insulin or decreased ability to use insulin.	<ul style="list-style-type: none"> ■ HEENT 	<ul style="list-style-type: none"> ■ Swollen glands in neck and face. ■ Deterioration of dental enamel related to gastric acidity of vomitus. ■ Sore throat.
	<ul style="list-style-type: none"> ■ Integumentary ■ Cardiovascular 	<ul style="list-style-type: none"> ■ Poor skin turgor; dehydration. ■ Arrhythmias related to electrolyte imbalances; cardiac arrest possible.
	<ul style="list-style-type: none"> ■ Gastrointestinal 	<ul style="list-style-type: none"> ■ Hemoptysis from damage to throat when inducing vomiting. ■ Heartburn, bloating, indigestion.
	<ul style="list-style-type: none"> ■ Neurological ■ Genitourinary 	<ul style="list-style-type: none"> ■ Depression, mood swings, feeling out of control. ■ Amenorrhea or irregular menses.
	<ul style="list-style-type: none"> ■ General Health Survey ■ Integumentary 	<ul style="list-style-type: none"> ■ Increased thirst. ■ Fatigue. ■ Poor skin turgor. ■ Frequent skin infections.
Iron Deficiency Anemia A decrease in the number of red cells caused by too little iron.	<ul style="list-style-type: none"> ■ HEENT ■ Respiratory ■ Gastrointestinal 	<ul style="list-style-type: none"> ■ Blurred vision. ■ Acetone breath odor. ■ Weight loss in spite of increased appetite. ■ Nausea, vomiting.
	<ul style="list-style-type: none"> ■ Genitourinary 	<ul style="list-style-type: none"> ■ Impotence in men. ■ Amenorrhea. ■ Frequent bladder and vaginal infections.
	<ul style="list-style-type: none"> ■ Neurological 	<ul style="list-style-type: none"> ■ Depression. ■ Confusion.
	<ul style="list-style-type: none"> ■ Integumentary 	<ul style="list-style-type: none"> ■ Pale skin color (pallor). ■ Sore tongue. ■ Brittle nails.

ABNORMALITY	AREA	ASSESSMENT FINDINGS
	<ul style="list-style-type: none"> ■ HEENT ■ Respiratory ■ Cardiovascular ■ Gastrointestinal 	<ul style="list-style-type: none"> ■ Frontal headache. ■ Shortness of breath. ■ Orthostatic hypotension. ■ Unusual food cravings (pica). ■ Decreased appetite (especially children). ■ Fatigue.
<p>Kwashiorkor A protein deficiency caused by inadequate dietary protein, malabsorption, cancer, or acquired immunodeficiency syndrome (AIDS).</p>	<ul style="list-style-type: none"> ■ Neurological ■ Integumentary ■ Gastrointestinal ■ Musculoskeletal ■ Neurological 	<ul style="list-style-type: none"> ■ Darkened, scaly skin. ■ Hair depigmentation. ■ Pitting edema. ■ Extreme abdominal swelling/bloating. ■ Hepatomegaly. ■ Muscle atrophy. ■ Mental apathy. ■ Developmental delays, retardation.
<p>Marasmus A severe protein and calorie deficiency that causes tissue to be sacrificed to meet energy needs. It is also called <i>protein-calorie malnutrition</i>. Other causes include severe infections, burns, eating disorders, chronic liver disease, cancer, and AIDS.</p>	<ul style="list-style-type: none"> ■ HEENT ■ Integumentary ■ Gastrointestinal ■ Musculoskeletal ■ Neurological 	<ul style="list-style-type: none"> ■ Facial appearance of an aged person. ■ Wasting of subcutaneous fat and muscle. ■ Severe weight loss and emaciation. ■ Potbellied appearance caused by swollen abdomen. ■ Growth retardation. ■ Mental apathy. ■ Developmental delays, retardation.


S U M M A R Y

- Throughout the health assessment process, the nurse should be attuned to data related to nutrition.
- Nutrition is influenced by a myriad of factors. Basic information in the health history can identify the need to more thoroughly investigate nutritional status.
- Assessment of nutritional status involves eliciting data that are both directly and indirectly related to nutrition.
- Comprehensive nutritional assessment will provide specific data to enable the nurse to determine potential or actual nutritional health problems, to devise an appropriate plan of intervention, and to determine criteria for evaluation.

R E V I E W Q U E S T I O N S

1. You are performing a nutritional assessment. Which of the following conditions might result in primary malnutrition?
 - a. Anorexia nervosa
 - b. Diarrhea
 - c. Ulcerative colitis
 - d. Bulimia
2. When performing a nutritional assessment, which technique would best screen for nutritional deficits?
 - a. BMI measurement
 - b. Triceps skinfold measurements
 - c. 24-hour recall
 - d. Food composition
3. Part of your nutritional assessment includes calculating your client's BMI. What measurements are needed to calculate a BMI?
 - a. Caloric intake and weight
 - b. Height and weight
 - c. Midarm and waist circumference
 - d. Triceps skinfold and weight
4. When measuring midarm circumferences, the difference between arm measurements should not exceed:
 - a. 1 cm.
 - b. 2 cm.
 - c. 3 cm.
 - d. 4 cm.

5. The BMI is a valuable nutritional assessment tool. What does the client's BMI reflect?
- Body fat
 - Body weight
 - Muscle mass
 - Body frame
6. Which measurement is useful in estimating body fat?
- Weight
 - Height
 - Body frame
 - Triceps skinfolds



Spiritual Assessment

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify spiritual components of health
- Discuss spirituality for various age groups
- Discuss spirituality for various cultural groups
- Perform a spiritual assessment
- Identify influence of spirituality on healthcare

Historically, the practice of nursing has been based on holistic care of the patient. This **holism** respects the interrelated nature of body, mind, and spirit. The body has been well represented by research and innovations in the care of physiological processes. Psychology and sociology have contributed to the understanding of the emotional processes.

More recently, an increased nursing emphasis on holism is reflected in the increased importance given to the spiritual dimension. What is the spiritual dimension? **Spirituality** may be defined as the inherent quality of humans to believe in something greater than the self and in a faith that affirms life. Spirituality also involves interconnectedness between the self, nature, and others. Spirituality is larger than religion, although religion is a forum for expressing our spiritual nature. For example, the spiritual dimension encompasses how we feel when we see a new baby or a beautiful sunset. In essence, spirituality is all behavior that gives meaning to life and gives us strength. As a component of health, spirituality is characterized by meaning, hope, purpose, love, trust, forgiveness, and creativity.

Are you equipped to assist the patient and family with issues that relate to spirituality? As with any physical system, the goal of a spiritual assessment is to detect strengths, potential problems, or dysfunction. You need adequate information in order to plan appropriate interventions. Assessing your patient's spirituality is unique because the only "tools" needed are you and time. Your ability to listen actively is essential to obtaining a spiritual assessment.

Even so, your assessment will still follow a systematic approach. Although physical examination is not actually a part of the assessment, people often have physical symptoms that are related to underlying spiritual concerns. So spiritual assessment is a part of a holistic assessment, conducted along with the assessment of physiological and psychological systems.

Nursing is a humanistic science that focuses on caring for the whole person. Several authors have chosen to depict the nature of spirituality in different ways, but in each, the spiritual dimension is seen as all-encompassing or pervasive throughout a person's being. Nurses appreciate the interrelatedness and interconnectedness of all the dimensions of a person. As a result, we are especially suited to treat the patient in a holistic manner. You may examine each system separately, but to achieve complete care, you must consider the patient as a whole, using the mind-body-spirit concept.

Brief Review of Spiritual Health

To adequately assess the spiritual dimension, you need to understand its general nature. Spiritual health and wellness reflect the spiritual nature of humans and are related to

the patient's overall health. Because humans are composed of multiple related aspects, you must assess and intervene in all these aspects to provide holistic and humanistic care. Many researchers believe that all human beings have three needs in common: the need for meaning and purpose, the need for love and belonging, and the need for forgiveness. When patients are having difficulty in these areas, the nurse is often the one who identifies the need for spiritual assistance.

When taking a patient's history, your first questions about spirituality should focus on the basics. Ask the patient the following questions:

- Do you identify with any organized religion? If so, what religion?
- If you do not identify with a particular religion, do you have a belief system that provides comfort and strength?
- Are you an **agnostic** or an **atheist**? If so, do you have any belief system that gives meaning to your life?

While asking these questions, keep in the back of your mind:

- What is the primary nature of the patient's religion?
- Are people of this religion **monotheistic** (worshipping one God) or **polytheistic** (worshipping many gods)?

As you can see, not all these questions have easy answers. To best serve your patient's spiritual needs, you must remain open and nonjudgmental when assessing him or her.



HELPFUL HINT

Recognizing your own spiritual beliefs and biases will help you be as open as possible when assessing your patient's spirituality.

Developmental, Cultural, and Ethnic Variations

The development of the spiritual dimension is closely linked to intellectual, psychological, and sociological factors. In addition, you must also consider your patient's age and culture. Intellectual factors often depend on physiological factors such as brain development. Psychological and sociological factors are greatly influenced by family dynamics and environment.

Infants

Infants are dependent on their parents and other caregivers. They are capable of feeling trust or mistrust but are incapable of the abstract thought necessary for spiritual faith. So when assessing and caring for infants, you

must consider the spiritual needs of the family and other caregivers and listen actively to their concerns.

When the patient is an infant, ask the parents or caregivers the following questions:

- What concerns do you have about your child's illness?
- How can I help support your use of religious practices (e.g., by referral to the hospital chaplain or hospital meditation room)?

Ask yourself the following questions:

- Do parents ask why their child is ill?
- Do they see the infant's illness as a religious punishment?
- Are they practicing religious rituals?

Toddlers and Preschoolers

Toddlers learn values by mimicking others and by receiving positive or negative feedback in response to their behavior. Toddlers who participate in their family's celebrations of religious ceremonies or rites will begin to incorporate those aspects of spiritual development. They also learn values, including the importance of respect for life and others. These interactions set the stage for further spiritual development. Toddlers and preschoolers are just beginning to develop language and intellectual skills and an appreciation for cause and effect. A true understanding of the nature of spiritual or religious values is not possible at this level; however, the young child is aware of the interconnectedness of family.

When the patient is a preschooler, ask the following questions:

- How do you feel about what is happening to you?
- Whom do you talk to when you are in trouble, sad, lonely, or scared?
- What makes you feel better when you are scared, sad, or lonely?

Ask yourself the following questions:

- Does the child express concerns or show anxiety about illness and dying?
- Does she or he speak of being punished by a deity for "being bad"?
- Is he or she practicing religious rituals, such as saying bedtime prayers?

School-Age Children

Once a child begins school, she or he is influenced by peers as well as by family. Development of a spiritual awareness parallels cognitive and emotional development. The younger school-age child often believes in the magic value of religious ritual, whereas the older school-age child may begin to understand the abstractness of faith. To have faith in a power bigger than yourself requires abstract thought—which is usually not common in children before age 11 or 12.

Children begin to question spiritual issues as they explore their world. Although the loss of a loved one at this time is difficult for a child to grasp because he or she can see no benefit to the death, nurses who work with dying children often find that they are mature beyond their years regarding spiritual concerns. If the conditions are right, spiritual development and awareness can appear at almost any age. Therefore, do not dismiss the need for assessment of spiritual needs in children.

When the patient is a school-age child, ask the same questions as were used with toddlers and preschoolers.

Adolescents

Adolescents are searching, trying to form a unique identity for themselves. Consequently, they begin examining their parents' beliefs in all aspects of life. In the area of spiritual development and awareness, they may review, critique, rebel, search for alternatives, and sometimes embrace their parents' values and beliefs as their own. This phase may continue into young adulthood. Do not assume that the adolescent is without spiritual concerns because of her or his behavior.

When the patient is an adolescent, ask the following questions:

- How do you feel about what is happening to you?
- Whom do you usually go to for support?
- What gives your life meaning? Has this changed since you got sick?

Also ask yourself the following questions:

- Does the adolescent express concerns about dying or the seriousness of his or her illness?
- Does she or he practice any religious rituals?
- Does he or she verbalize his or her own beliefs and values?

Young Adults

Young adults are a diverse group, and so each person may be in a different stage of spiritual development. Some young adults may be very spiritual, with a well-developed sense of religious commitment or a commitment to something more personal, such as music or the arts. Others appear to have no sense of spirituality and may live just for the moment, focusing on survival. Significant life events, such as the death of a friend, a close brush with death, or the beginning of a new relationship, may be the nucleus for the start of a spiritual awakening.

The experience of pregnancy and birth may have profound effects on spiritual growth and development. Impending parenthood is often the impetus for a deepening of existing spirituality or an awakening of new feelings. The responsibility of parenting and the desire to transmit beliefs and values to a new generation often stimulates a search to clarify those beliefs and values and

find answers to spiritual questions. As the family grows and matures, parents' lives usually revolve around their children. First, they strive to impart values to young children; then eventually they deal with adolescents beginning to question these values.

When the patient is a young adult, ask the same questions as you would an adolescent (see previous section).

Older Adults

As people grow older, they may cling more tightly to their connection with spirituality. Older adults are confronted with resolving the developmental task of integrity versus despair. Those who find a sense of meaning and purpose to their lives and enjoy the interconnectedness with others that may be found in a place of worship are successfully working through this stage of development.

When the patient is an older adult, ask the same questions as you would an adolescent or a young adult. However, older patients often suffer many losses and therefore have fewer support systems, and so you may need to make referrals to community services.

People of Different Cultures or Ethnic Groups

Culture is a multidimensional factor in human development. It includes shared values, beliefs, and practices of a group that are transmitted from one generation to another. Culture often guides the way in which a group of people thinks and acts. Values, beliefs, and practices related to spiritual issues are also part of culture and are passed on to each generation. Just as it is important to be aware of physiological variations when assessing a patient, it is important to be aware of variations among people's spiritual and/or religious values, beliefs, and practices.

Spirituality develops from a person's worldview, which is often strongly influenced by cultural background. Before you can assess your patient's spirituality, familiarize yourself with her or his cultural domain. This will help you choose the right questions to ask, depending on the patient's age. The following are important areas to consider:

- *Use of prayer:* Learn about the use of prayer, meditation, and other activities or symbols that help people of this cultural or ethnic group reach fulfillment.
- *Meaning of life and individual sources of strength:* Know what gives meaning to life for the patient's cultural group. Identify the patient's individual sources of strength.
- *Spiritual beliefs and healthcare practices:* Become familiar with the way in which spiritual beliefs affect this cultural group's healthcare practices.



CRITICAL THINKING ACTIVITY # 1

Consider the different cultures described in table on page 191. How might your patient's culture influence your approach to planning his or her healthcare?



CRITICAL THINKING ACTIVITY # 2

What questions should you ask your patient to help identify the impact of spirituality on her or his healthcare?



CRITICAL THINKING ACTIVITY # 3

For each of the cultures described in table on page 191, identify interventions that would incorporate the patient's spiritual belief into his or her healthcare.



ALERT

Although many ethnic groups are linked with a particular spiritual viewpoint or religion, do not assume a patient's spiritual perspective by her or his ethnic background alone. Assess each patient as a unique being.



Introducing the Case Study

January Mays is a 68-year-old retired schoolteacher. She lives in a single, two-story home in a small town in Pennsylvania. Her husband died last year, and her only child, a daughter, lives in California with her husband and two sons. Her sister lives across the street. Mrs. Mays was admitted to the hospital after apparently suffering a "flare-up" of congestive heart failure (CHF). As you begin your interview, she states, "I don't know why people are making so much fuss."

Performing the Assessment

Data for a spiritual assessment are obtained during the health history. Before conducting the history, you can also make preliminary observations related to the patient's behavior, relationships, and environment that may result in a more thorough spiritual assessment. You will also take a spiritual history that specifically relates to the patient's spiritual needs and identifies potential or actual problems.

How Culture Affects Spirituality

CULTURE	DOMINANT RELIGIONS/USE OF PRAYER	MEANING OF LIFE/SOURCES OF STRENGTH	SPIRITUAL BELIEFS/HEALTHCARE PRACTICES
African American	<ul style="list-style-type: none"> Majority are Baptists and Methodists. Strong belief in prayer. 	<ul style="list-style-type: none"> Sickness and pain are weaknesses coming from Satan, whereas God is a source of inner strength. 	<ul style="list-style-type: none"> Sickness is a separation from God. God is the Supreme Healer.
Amish	<ul style="list-style-type: none"> No national church hierarchy; local leaders are chosen from community. Silent prayer before and after meals is important. 	<ul style="list-style-type: none"> Share view of most Christians, but emphasis is on community, not individual. 	<ul style="list-style-type: none"> May seek counsel from religious leaders on health issues, but final decision is made by immediate family.
Appalachian	<ul style="list-style-type: none"> Variety of religions, but most stress fundamentalism. Prayer is a primary source of strength. 	<ul style="list-style-type: none"> Family and living right provide meaning. 	<ul style="list-style-type: none"> Fatalistic view—illness is God's will.
Arab American	<ul style="list-style-type: none"> Most are Muslims. Prayer five times a day is a duty. 	<ul style="list-style-type: none"> Faith means submission to Allah and is a part of everyday life. 	<ul style="list-style-type: none"> Religious practices are combined with medicine. Cleanliness is essential before praying. Illness is seen as a punishment or a trial that is endured.
Chinese American	<ul style="list-style-type: none"> Variety of religions, including Buddhism. Formal religious services are minimal. Pray alone rather than as a community. Prayer is a source of comfort. 	<ul style="list-style-type: none"> Life viewed in terms of cycles and interrelationships. Life forces are a source of strength. 	<ul style="list-style-type: none"> Cyclic explanation of life is connected to health.
Cuban American	<ul style="list-style-type: none"> Most are Catholic, but practice is more personal than institutional. 	<ul style="list-style-type: none"> Family is important source of strength. 	<ul style="list-style-type: none"> Fatalistic, lack control, use of folk remedies.
Egyptian American	<ul style="list-style-type: none"> Islam or Christian. Pray during illness. 	<ul style="list-style-type: none"> Inner peace through practice of religious rituals. 	<ul style="list-style-type: none"> When ill, you are at the mercy of God (Christian) or Allah (Islam).
Filipino American	<ul style="list-style-type: none"> Most are Catholic. Prayer is important. 	<ul style="list-style-type: none"> Strength comes from relationship with God. 	<ul style="list-style-type: none"> Fatalistic—illness is seen as God's will or induced by evil spirits.
French Canadian	<ul style="list-style-type: none"> Most are Catholic. Prayer is important. 	<ul style="list-style-type: none"> Family is important source of strength, but divorce has affected this. 	<ul style="list-style-type: none"> Little connection made between spirituality and healthcare. Spirituality is still seen as religion.
Greek American	<ul style="list-style-type: none"> Greek Orthodox faith and daily prayer very important. 	<ul style="list-style-type: none"> Life has meaning through the different roles one performs. 	<ul style="list-style-type: none"> Icons of family saint or Virgin Mary are important when ill.
Iranian American	<ul style="list-style-type: none"> Islam influence. Prayer is important. 	<ul style="list-style-type: none"> Accepting Allah's will is source of strength, as are family and friends. 	<ul style="list-style-type: none"> Fate and Allah's will explain illness.

(continued)

How Culture Affects Spirituality (continued)

CULTURE	DOMINANT RELIGIONS/USE OF PRAYER	MEANING OF LIFE/SOURCES OF STRENGTH	SPIRITUAL BELIEFS/HEALTHCARE PRACTICES
Irish American	<ul style="list-style-type: none"> Most are Catholic. Prayer is individual and private. 	<ul style="list-style-type: none"> Church is a source of strength. 	<ul style="list-style-type: none"> Somewhat fatalistic. Religion is important when ill. Sacraments and religious medals may provide solace.
Jewish American	<ul style="list-style-type: none"> Monotheistic belief in God. Pray alone or in groups. 	<ul style="list-style-type: none"> Preservation of life is most important; therefore, good health is very important. 	<ul style="list-style-type: none"> Caring for the sick takes precedence over keeping Sabbath.
Mexican American	<ul style="list-style-type: none"> Most are Catholic. Prayer is important and individualized. 	<ul style="list-style-type: none"> Family is most important. 	<ul style="list-style-type: none"> When ill, it is important to talk about soul or spirit.
Navajo Native American	<ul style="list-style-type: none"> Spirituality based on harmony with nature. 	<ul style="list-style-type: none"> Strength comes from inner self and being in harmony with surroundings. 	<ul style="list-style-type: none"> Illness results from not being in harmony. Healing ceremonies help restore mental, physical, and spiritual harmony.
Vietnamese American	<ul style="list-style-type: none"> Most are Buddhists. Prayer and offerings may be made at altar in home or temple. 	<ul style="list-style-type: none"> Family and religion are important. 	<ul style="list-style-type: none"> Harmony and order influence perspective of illness that is characterized by self-control, acceptance of one's destiny, and fatalism.

Source: Data from Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Health Care: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.

Health History

First, obtain the patient's biographical data; then ask questions about his or her current health history, past health history, family history, and psychosocial profile.

Biographical Data

A review of the patient's admission sheet may reveal basic religious information such as church or temple affiliation. Be sure to ask questions to clarify this information because religious affiliation may range from being in name only to being a very involved relationship.



ALERT

Be aware that the absence of religious identification does not mean that the patient has no spiritual needs.

Other helpful information includes marital status and contact person. This provides clues about the state of connectedness of the patient to others. A patient who has not listed relatives may have spiritual needs that a patient with a strong family network does not. Age is

another indicator of spiritual needs. Older adults with an identified connection to organized religion may find that connection to be a source of comfort and support.

Current Health Status

Your patient's current physical health and spiritual health are intertwined. A sense of spirituality is influential for the patient or a significant other during periods of physical and/or mental illness. For some, acceptance of losses is easier because life as a whole has been full of meaning; for others, losses are seen as punishment for a life that could have been lived differently.

Although anyone may have spiritual needs, patients with chronic or terminal diseases are especially prone to spiritual distress. Exploring the connection between medical diagnosis and spiritual needs is an ongoing process, because as time passes, your patient's experience and situation change.

Past Health History

Health problems that occurred in your patient's past can have long-lasting physical and spiritual effects. The past health history will give you clues about your patient's

overall health up to this point in her or his life. If she or he has been relatively healthy until now, an acute illness may precipitate spiritual distress. Or a series of past health problems may have tested and strengthened the patient's spirituality. Whatever the case, you need to explore the connection between your patient's health and spiritual needs.

Family History

The family history is invaluable in identifying familial problems that pose a threat to your patient's health. This in turn may affect his or her spiritual needs. The family history can also be helpful in identifying supports and relationships.

Psychosocial Profile

The psychosocial profile reveals patterns in the patient's lifestyle that affect spiritual health. Begin by examining your patient's typical day. Are there any activities that reflect spirituality, such as praying or attending religious services? Dietary preferences may also be influenced by religious practices. For example, some religions have dietary restrictions that need to be considered when caring for your patient, such as **kosher** diet for patients of Jewish faith.

The psychosocial history also identifies relationships and support systems. These supports can be within the family or the community. Organized religion can also be a source of support for your patient. The amount of stress in your patient's life and her or his ability to deal with it can be greatly influenced by spirituality. Spirituality can be a source of comfort and coping.



Case Study Findings

Reviewing Mrs. Mays'

Biographical data:

- 68-year-old African American woman.
- Husband died 1 year ago.
- One grown daughter who lives in California with her husband and two sons, ages 4 and 6.
- Lives in two-story home; sister lives across the street in her own home.
- Born in Grady Falls, Pennsylvania.
- Methodist Episcopal religion.
- M.S. in Education, retired from teaching elementary school.
- Has Medicare with supplemental HMO from teacher's union. Hospital admission for CHF.
- Good historian; alert and seems reliable.

Current health status:

- Difficulty breathing at rest, lower extremity edema, change in sleep patterns, uses two pillows.
- Currently hospitalized for exacerbation of CHF.
- Also has hypertension.

- Current medication lisinopril (Prinivil), 10 mg/d, for hypertension.

Past health history:

- CHF, treated for 3 to 5 years.
- Hypertension, treated for 15 years.

Family history:

- Father died of heart failure at age 89.
- Mother died at age 70 from complications of stroke.

Psychosocial profile:

- *Typical Day:* Arises at 7 A.M., makes her bed, eats breakfast, and reads the paper. Usually does light housework, such as dusting and washing breakfast dishes. She then rests because the activity leaves her short of breath. She generally calls her sister to come over for lunch and watch soap operas. Sister leaves by 4 P.M. to get dinner for her husband. Mrs. Mays usually eats a light supper and then does needlework until 10 P.M., when she watches the news. She goes to bed at 10:30 P.M.
- *Nutrition:* Eats three meals a day. Lunch is her main meal, and she enjoys it most because she usually has her sister for company. Doesn't like to cook just for herself, so she eats a light supper.
- *Activity and Exercise:* Until last week, she walked to the corner store for her daily paper. She now has it delivered because the walk is too difficult. She is also unable to walk to services at her church, which is three blocks away. She does not drive.
- *Sleep and Rest:* Goes to bed at 10:30 P.M. and arises at 7 A.M. Not sleeping well because of breathing problems caused by CHF.
- *Self-Care Activities:* Sees her doctor regularly for hypertension and CHF.
- *Roles/Relationships:* Sister lives across the street and spends most weekday afternoons with patient. Patient's only child and two grandchildren live in California. They talk on the phone weekly, but she sees them rarely. Attended church and taught Sunday school and vacation Bible school until recently.
- *Occupational Activities:* Retired teacher.
- *Hobbies:* Needlepoint, reading, watching television.
- *Environmental Risks:* Lives in two-story house with bedroom on second floor.
- *Personal Habits:* Has never smoked; doesn't drink alcohol.
- *Stress and Coping:* Says she doesn't like to make a "big fuss" over her health problems or "burden anyone" with them.



CRITICAL THINKING ACTIVITY #4

List all the strengths and areas of concern that have a bearing on Mrs. Mays' health status.



Case Study Evaluation

Before moving on, pause a moment and consider what you have learned about Mrs. Mays. Document the history related to Mrs. Mays.

Spiritual Assessment

A person's spirituality cannot be "measured," so you must rely on other evidence to make nursing diagnoses in the spiritual dimension. For example, emotional and physical behaviors often signal underlying spiritual disturbances. To assess your patient's spiritual health, focus your attention in four particular areas—behavior, verbal communication, relationships, and environment—and ask yourself questions in the following areas.

Assessing Behavior

Is the patient:

- Praying or meditating?
- Shutting others out?
- Constantly complaining?
- Having sleep difficulties?
- Pacing?
- Requesting frequent pain or sleep medication without apparent need?

Assessing Verbal Communication

Is the patient talking about:

- God or another deity?
- Church, temple, mosque, or other place or worship?
- Prayer, hope, faith, or the meaning of life?
- Effect of the diagnosis on his or her quality of life?

Assessing Relationships

Does the patient:

- Have many visitors? Who are her or his visitors (family, friends, clergy, or other spiritual support people)?
- Interact with them well?

Assessing Environment

Does the patient have:

- A **Bible**, **Koran (Qur'an)**, or other religious reading material?
- Religious jewelry or symbols, such as a **cross**, **Star of David**, prayer cap or shawl, flowers from a church altar, or religious greeting cards?

Additional Questions

Assessing the previous four areas will help you pinpoint your patient's spiritual resources. In addition, ask these questions:

- Who are your support people?
- What provides you with strength and hope?
- What gives your life meaning and purpose?
- How has your life changed since you became ill?

Phrase these questions carefully so that you do not offend the patient or seem to be unduly prying. If a specific religious affiliation was listed on the admission record, be sure you have a general understanding of its perspective. For example, is the religion monotheistic, such as **Christianity** or **Judaism**; polytheistic, such as **Hinduism**; or **nontheistic**, such as **animism** or **New Age**? You need to know basic similarities and differences to be able to assist the patient in the best possible way. Answers to the broad questions can then be used to further refine questions that are in line with the patient's belief system. Examples might be:

- What is important to you?
- How is your religion, deity, or faith important to you?
- Is prayer important?
- What accommodations can be made to assist you in continuing any spiritual practices (e.g., religious symbols and/or dietary needs)?




Spiritual Assessment Instruments

Several assessment surveys have been developed to help in the identification of spiritual needs. Some of the surveys are biased toward a Judeo-Christian viewpoint and are not suitable for all patients. Some of the surveys are long or designed for self-report; they may be difficult to use with an acutely ill patient. Some healthcare facilities have developed their own spiritual assessment tools and included them on their overall nursing assessment instrument. You may find the spiritual assessment questions in the previous section and other questions in this chapter more helpful than using any one survey. The Jarel Spiritual Well-Being Scale provides a scale for assessing spiritual well-being. (See *Jarel Spiritual Well-Being Scale*.)

Keep in mind that people need to understand the meaning of their existence and need to be connected to other human beings regardless of the presence or absence of organized religious affiliations.



(Text continued on page 200)



How Belief Systems Influence Healthcare

BELIEF SYSTEM	SIGNIFICANT BELIEFS AND SYMBOLS	EFFECTS ON HEALTHCARE
<p>Christianity Roman Catholic</p> 	<ul style="list-style-type: none"> ■ Important religious rites: baptism, holy communion, and anointing of the sick. ■ Crucifix and rosary (beaded chain with small crucifix) are important symbols. Missal and Bible are important reading materials. ■ Abstain from meat during Lent and other holy days. 	<ul style="list-style-type: none"> ■ Modern healthcare accepted, but many rely on a strong sense of faith for healing. Healing services may be held. ■ Rites may be requested in the hospital as needed. Stillborn or critically ill infants are baptized as soon as possible after delivery. ■ Prayers for and by ill person are encouraged. ■ Presence of symbols often a source of comfort. ■ Opposed to abortion and most forms of birth control.
<p>Crucifix</p> <p>Eastern Orthodox</p> 	<ul style="list-style-type: none"> ■ Practices same rites as Roman Catholic and Anglican Church: baptism, holy communion and anointing of the sick. ■ Orthodox cross has a diagonal cross bar near the bottom of the upright bar and two cross bars at the top. Wearing this cross is important to the religious person. ■ Parishes are often established along ethnic/national lines (e.g., Greek, Russian, Ukrainian). ■ Church follows different liturgical calendar, so most high holy days are celebrated at different times than rest of Christian world. 	<ul style="list-style-type: none"> ■ Generally accepts modern healthcare. ■ Disapproves of birth control and strongly opposes abortion. ■ Orthodox priest may be called on for blessings, exorcisms, and direct healing. ■ Having religious symbols near is a source of faith and comfort. ■ Organ donation is acceptable. ■ Similar to Roman Catholics.
<p>Orthodox cross</p> <p>Episcopalian or Anglican</p> 	<ul style="list-style-type: none"> ■ Important religious rites: baptism, holy communion, and anointing of the sick. ■ Both crucifix and plain cross are used. Book of Common Prayer and Bible are important reading materials. ■ Fast on high holy days and abstain from meat on Fridays in Lent. 	<ul style="list-style-type: none"> ■ Some Episcopal parishes practice healing by “laying on of hands” and prayers of special intention. ■ Rites may be requested in the hospital as needed. Stillborn or critically ill infants are baptized as soon as possible after delivery. ■ Prayers for and by ill person are encouraged.

(continued)

How Belief Systems Influence Healthcare (continued)

BELIEF SYSTEM	SIGNIFICANT BELIEFS AND SYMBOLS	EFFECTS ON HEALTHCARE
		<ul style="list-style-type: none"> ■ Responsible family planning and free choice are important. No sanctioned restrictions on birth control or abortion.
Plain cross	<p>Other Protestant Denominations</p> 	<ul style="list-style-type: none"> ■ Wide range of practices found among a diversity of denominations (e.g., Baptist, Pentecostal, Lutheran, Methodist). ■ A plain cross and Bible are symbols common to almost all Protestant denominations. ■ When practiced, baptism is usually reserved for older children or adults. Holy communion is practiced infrequently, if at all. ■ Special prayers for sick are a more common practice than anointing. <ul style="list-style-type: none"> ■ Because there are many different denominations, nurse must ask questions that address specific religious practices that may affect patient's health and healthcare (e.g., dietary restrictions). ■ Use of prayers and healing by "laying on of hands" varies. Strict Christian Scientists reject all medical treatment and rely on prayer for healing. ■ Interpretation of biblical scripture within each denomination determines acceptance of such practices as blood transfusions, abortion, or organ transplantation.
Plain cross	<p>Jehovah's Witness</p>	<ul style="list-style-type: none"> ■ Believe that blood transfusions and blood products are forbidden.
Christian Scientist	<ul style="list-style-type: none"> ■ Believe that God is all good and nothing that is real exists outside. ■ God, sin, sickness, and death seen as mortal error or illusion. ■ Learning about body harms them spiritually. 	<ul style="list-style-type: none"> ■ Organic disease does not exist, disease is mental. ■ No understanding of signs or symptoms. ■ Refuse or delays medical treatment. ■ Suffer needlessly, neglect life-threatening conditions.
Mormon	<ul style="list-style-type: none"> ■ Believe physical body and spirit make up soul. ■ Purpose on earth is to gain experience and prove self worthy to return and live with God. 	<ul style="list-style-type: none"> ■ Caution against use of tobacco, alcohol, tea, coffee. ■ Emphasize positive benefits of good eating habits, physical and spiritual fitness. ■ Misuse of drugs (illegal, legal, prescription, or controlled) seen as a violation of health code. ■ May affect compliance with medical treatment plan.

BELIEF SYSTEM	SIGNIFICANT BELIEFS AND SYMBOLS	EFFECTS ON HEALTHCARE
<p>Judaism</p>  <p>Star of David</p>	<ul style="list-style-type: none"> ■ Encompasses three groups of believers with different practices: Orthodox (most traditional), Reform (most liberal) and Conservative (has elements of both traditional and liberal practices). ■ Common symbols are the Star of David and the <i>Torah</i>, the first five books of the Bible written on scrolls. It contains guidelines for life. A prayer book may be carried for reading of special prayers. ■ Services held after sundown on Friday and/or before sundown on Saturday. ■ Male infants usually circumcised at 8 days old in a religious ceremony known as a <i>bris</i>. ■ Orthodox and some Conservative Jews keep kosher dietary laws. ■ Men may wear a yarmulke, a small skull cap. Cap is often worn in temple by all men but may be worn at all times. 	<ul style="list-style-type: none"> ■ Be sure to ask questions about diet and religious practices, such as, "Do you follow any dietary rules that we should know about while you are here in the hospital?" ■ Prayer is important to all groups. ■ Observation of Sabbath and other holy days is important. Some patients may refuse treatment on those days or refuse to eat on traditional days of fasting. Kosher food is available in most hospitals. ■ Religious law provides for the individual's responsibility to stay healthy, both physically and mentally. Traditional medicine is valued as well as prayer for relief of suffering. ■ Organ transplants are allowed, but donation of organs from dying people is not permitted without the donor's consent.
<p>Islam</p>  <p>Crescent moon</p>	<ul style="list-style-type: none"> ■ Based on teachings of Mohammed. ■ In North America, adherents of Islam may be either traditional ethnic Muslims or members of the Nation of Islam, a primarily African American group. ■ The Qur'an (Koran) is the sacred book of Islam. Many have a special prayer rug that they carry. Symbols vary, but generally include a crescent moon in some position. ■ Ritual prayer, five times a day while kneeling and facing toward Mecca (in Middle East) and specific dietary restrictions are common to all members. ■ Alcohol and pork are forbidden. ■ Many practice fasting during the month of Ramadan. ■ Women dress in clothing that hides the body. 	<ul style="list-style-type: none"> ■ Caregivers must be of the same gender to protect modesty. ■ Women may continue to wear traditional head scarf with or without face veil while hospitalized. Traditional hospital gowns may be unacceptable. ■ Some have a fatalistic view of health and may refuse treatment. ■ Islam has rituals related to all aspects of life and death. Be aware of these needs and show sensitivity for the patient and family. ■ Medications that have an alcohol base, such as elixirs and cough syrups, may be unacceptable.


(continued)

How Belief Systems Influence Healthcare (continued)

BELIEF SYSTEM	SIGNIFICANT BELIEFS AND SYMBOLS	EFFECTS ON HEALTHCARE
Hinduism	<ul style="list-style-type: none"> ■ Ancient religion originating in India. Many Indians now living in North America continue to follow their religion. ■ Universal symbol is the om symbol. ■ A major tenet is achieving a state of oneness with the universe. Prayer is important to achieve oneness. ■ Many gods have power over the individual. Religious shrines are often erected in the home. ■ Killing any living creature is forbidden, so many sects are vegetarians. 	<ul style="list-style-type: none"> ■ Patients may want to set up a small shrine in the hospital room. ■ Although the traditional sari is not religion based, some women may wish to wear it while in the hospital. It is a long length of soft cloth that is wrapped around the body like a long skirt, with the loose end draped over the head like a scarf. ■ The belief in <i>karma</i> (unchangeable fate) may influence the approach to self-care and the seeking of healthcare.
New Age	<ul style="list-style-type: none"> ■ Although not a religion, this belief system is growing in popularity. ■ Beliefs are borrowed from Buddhism, Native Americans, humanism, physics, and occultism, among others. ■ Objects such as pyramids and crystals are thought to impart strength and healing energy. ■ There is great diversity of thinking, but some common beliefs are that all people are divine, right and wrong are determined by the individual, and reality can be controlled. 	<ul style="list-style-type: none"> ■ The <i>crystal</i> is a common symbol. It is thought to impart healing energy and may be rubbed over the skin in close proximity to diseased tissue. Nurses need to recognize the crystal as a possible belief object and not as jewelry. ■ No clergy, but several believers may meditate with patient to increase healing power. These beliefs do not usually prevent the patient from seeking traditional therapies.
Nonreligious Agnostic	<ul style="list-style-type: none"> ■ Believes that it is impossible to know if God or other Supreme Being exists. ■ Does not believe in any God but may have beliefs that are of the spiritual dimension. 	<ul style="list-style-type: none"> ■ Diverse and individual influences on healthcare. ■ Assess each patient individually. ■ Diverse and individual influences on healthcare. ■ Must assess each patient individually.
Atheist	<ul style="list-style-type: none"> ■ People who believe in a Supreme Being that is unidentified may avoid an organized religion but still have spiritual beliefs and needs. 	<ul style="list-style-type: none"> ■ Diverse and individual influences on healthcare. ■ Must assess each patient individually.
No Organized Religion	<ul style="list-style-type: none"> ■ People who believe in a Supreme Being that is unidentified may avoid an organized religion but still have spiritual beliefs and needs. 	<ul style="list-style-type: none"> ■ Diverse and individual influences on healthcare. ■ Must assess each patient individually.



Om symbol



Crystal

Jarel Spiritual Well-Being Scale©

Directions: Please circle the choice that *best* describes how much you agree with each statement. Circle only one answer for each statement. There are no right or wrong answers or good or bad scores. The purpose of this scale is to help the patient talk about his or her spirituality in a nonthreatening way and to give you a sense of the importance of spirituality in your patient's life so that you can plan appropriate interventions.

	STRONGLY AGREE	MODERATELY AGREE	AGREE	DISAGREE	MODERATELY DISAGREE	STRONGLY DISAGREE
1. Prayer is an important part of my life.	SA	MA	A	D	MD	SD
2. I believe I have spiritual well-being.	SA	MA	A	D	MD	SD
3. As I grow older, I find myself more tolerant of others' beliefs.	SA	MA	A	D	MD	SD
4. I find meaning and purpose in my life.	SA	MA	A	D	MD	SD
5. I feel there is a close relationship between my spiritual beliefs and what I do.	SA	MA	A	D	MD	SD
6. I believe in an afterlife.	SA	MA	A	D	MD	SD
7. When I am sick, I have less spiritual well-being.	SA	MA	A	D	MD	SD
8. I believe in a Supreme Power.	SA	MA	A	D	MD	SD
9. I am able to receive love from and give love to others.	SA	MA	A	D	MD	SD
10. I am satisfied with my life.	SA	MA	A	D	MD	SD
11. I set goals for myself.	SA	MA	A	D	MD	SD
12. God has little meaning in my life.	SA	MA	A	D	MD	SD
13. I am satisfied with the way I am using my abilities.	SA	MA	A	D	MD	SD

(continued)

Jarel Spiritual Well-Being Scale[©] (continued)

14. Prayer does not help me in making decisions.	SA	MA	A	D	MD	SD
15. I am able to appreciate differences in others.	SA	MA	A	D	MD	SD
16. I am pretty well put together.	SA	MA	A	D	MD	SD
17. I prefer that others make decisions for me.	SA	MA	A	D	MD	SD
18. I find it hard to forgive others.	SA	MA	A	D	MD	DS
19. I accept my life situations.	SA	MA	A	D	MD	DS
20. Belief in a Supreme Being has no part in my life.	SA	MA	A	D	MD	SD
21. I cannot accept changes in my life.	SA	MA	A	D	MD	SD

Source: Adapted from Hungelmann, J., Kenkel-Rossi, E., Klassen, L., and Stollenwerk, R. (1987). Marquette University College of Nursing, Milwaukee, Wisconsin.



Case Study Findings

Mrs. Mays' focused spiritual assessment findings include:

- **Behavior**
 - Lies in bed with her face turned away from the door.
 - Blood pressure remains elevated despite medication.
 - Eyes are closed but her hands are folded and her lips move silently.
- **Verbal Communication**
 - Asks if anyone from church has called for her when she was off the floor for a test.
 - Asks why she is sick. States that when she was teaching she felt alive and useful. "What use am I now?"
- **Relationships**
 - No visitors have been noted for Mrs. Mays since her admission 2 days ago.
 - Patient wonders why her sister hasn't visited or even called.
- **Environment**
 - A well-worn Bible is on Mrs. Mays' nightstand

- She wears a small gold cross around her neck. States that it was a confirmation gift from her godparents many years ago.

■ Additional questions:

- Who are your support people?

Patient states: "I usually rely on my sister, but I don't want to burden her. My daughter is so far away, and it's expensive for her to come back here. Besides, she has her own family."

- What provides you with strength and hope?

Patient states: "I usually find strength in my church, the services are so beautiful. The people are great, too. We're just like a family. I haven't been able to get there recently, and I really miss it. I guess they don't know I'm here yet. Our pastor recently left, and we have a temporary right now. I don't know him very well."

- What gives your life meaning and purpose?

Patient states: "Teaching the children was my first love after my husband and daughter, of course. Those little ones were so eager to learn, you know. I retired 3 years ago. My husband and I were only able to travel to California one time to see our daughter

and her family before he got so sick. My, how he suffered. I miss him terribly.”

- How has your life changed since you became ill?

Patient says: “Since I retired, I have taught Sunday school and vacation Bible school. It keeps me young to be around the children. But since I’ve been bothered by this pesky heart, I don’t even have the energy to walk to church let alone handle teaching. I guess I’ll have to find something else to do. Why has this happened to me now? What is God thinking?”



Case Study Analysis and Plan

Now that your assessment of Mrs. Mays is complete, document your key history and key spiritual assessment findings. List key history findings and spiritual assessment findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

The next step is to analyze the data from Mrs. Mays’ spiritual assessment and develop nursing diagnoses. The following are possible nursing diagnoses for this patient. Cluster the supporting data for each.

1. Spiritual Distress, related to current life situation
2. Hopelessness, related to chronic illness
3. Social Isolation

Identify any additional nursing diagnoses.



CRITICAL THINKING ACTIVITY #5

Now that you have identified some nursing diagnoses for Mrs. Mays, select one from the previous list and develop a nursing care plan and a brief teaching plan.

RESEARCH TELLS US

A qualitative study using grounded theory was designed to gain a better understanding of spirituality as seen by patients, nurses, and various religious representatives in the United Kingdom. Phase I of this study included interviews with 12 nurses, 15 patients from acute care and hospice, and 5 representatives from four major world religions. Constant comparative method was used for data analysis with line-by-line analysis, identification of categories, and central themes.

Two categories were identified: definitions of spirituality and diverse perceptions of spirituality. The nurses had similar definitions of spirituality. Nurses identified descriptors of spirituality as essence, individual, and whole, affecting every aspect of life. For the nurses, spirituality was seen as a universal concept.

For the 15 patients, their description of spirituality was not as clear. Often, spirituality was seen as syn-

onymous with religion. The assumption that spirituality comes into play when ill was not expressed by these patients.

The 5 religious representatives’ perceptions of spirituality varied. The Islamic representative identified elements similar to those identified by nursing. Two representatives struggled to define spirituality. Most of the religious representatives’ perceptions were different from those of nursing.

The findings suggest that spirituality is viewed differently by different groups and that the concept of spirituality may not be universal. The implications for nursing affect both practice and education. Nurses need to be aware of the diversity surrounding the concept of spirituality. Students need to be educated about this diversity so that they may develop an individualistic, nonjudgmental approach in assessing the spiritual needs of their patients.

Source: McSherry, W., Cash, K., and Ross, L. (2004). Meaning of spirituality: Implications for nursing practice. *Journal of Clinical Nursing* 13(8):934–941.

S U M M A R Y


- Because the body, mind, and spirit are interrelated, illness can have an impact on your patient’s spiritual well-being.
- The opposite is also true—your patient’s spirituality can have a positive or a negative effect on his or her illness.
- Performing a spiritual assessment involves listening closely and compassionately and knowing what ques-

tions to ask. It also involves performing a systematic assessment.

- Armed with this information, you can then formulate appropriate nursing diagnoses to help your patient cope with her or his illness.

R E V I E W Q U E S T I O N S


1. Mary Braman, age 65, is hospitalized for a total knee replacement for degenerative joint disease. While performing the admission nursing assessment, you note that she is of Jewish faith. Because she is of Jewish faith, what aspect of her care should be considered while planning her care?
 - a. Pain management
 - b. Dietary restrictions
 - c. Anesthesia preference
2. John Schmidt, age 35, was involved in an automobile accident and has blunt trauma to the abdomen. You see on his admission record that he is a Jehovah's Witness. Which treatment may be prohibited owing to his religious beliefs?
 - a. Surgery
 - b. Pain medication
 - c. Blood transfusions
 - d. Anesthesia
3. Theresa McCauley, age 75, has metastatic breast cancer. You see that she is Catholic. Which intervention may be helpful in meeting her spiritual needs?
 - a. Give her a Bible
 - b. Have priest visit and administer sacrament of the sick
 - c. Give her a crucifix
4. John Wilcox, age 50, was involved in brought to the ED with head injuries from a fall. You note he is a Christian Scientist. What implications may his religious beliefs have on his decisions for healthcare?
 - a. May refuse treatment
 - b. May refuse blood transfusions
 - c. May want to see priest
5. Tara Ali, age 25, has just delivered a healthy baby boy. She is of Islamic faith. Considering her religious belief, how might this affect her care?
 - a. Caregiver must be a woman
 - b. May refuse blood transfusion
 - c. May refuse pain medication



UNIT

3

Systems Physical Assessment



Assessing the Integumentary System

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent integumentary history questions
- Obtain an integumentary history
- Perform an integumentary physical assessment
- Document integumentary findings
- Identify actual/potential health problems of the integumentary system
- Recognize developmental and cultural variations of the integumentary system
- Differentiate between normal and abnormal integumentary system findings

The **integumentary system**, consisting of the skin, hair, and nails, is the largest organ of the body and the easiest of all systems to assess. It provides invaluable information about all other bodily systems. The skin, hair, and nails provide clues about general health, reflect changes in environment, and signal internal ailments stemming from other organs. Because integumentary system cells reproduce rapidly, changes in the skin, hair, and nails may be an early warning of a developing health problem. Yet, the importance of carefully assessing the integumentary system for subtle changes is often overlooked. A thorough assessment of this system may help you detect actual or potential problems, not only in the skin but also in underlying systems.

Anatomy and Physiology Review

Before you begin your assessment, you need a basic understanding of the integumentary system, including its general function and purpose. A knowledge of normal functions and structures will enable you to detect and interpret any abnormalities.

Structures and Functions of the Integumentary System

The structures of the integumentary system are the skin, hair, nails, sweat glands, and sebaceous glands. Their functions are described in the following paragraphs.

The Skin

The **skin** is a layer of tissue that covers all exposed body surfaces. Although similar to the mucous membranes, the skin also includes appendages such as hair follicles and sebaceous glands. Its thickness varies according to location or site. The **epidermis**, the outer visible layer, contains keratin, an extremely tough, protective protein substance that can cause tissue to become hard or horny. The deeper **dermis** is made up of proteins and **mucopolysaccharides**, thick, gelatinous material that provides a supporting matrix for nerve tissue, blood vessels, sweat and sebum glands, and hair follicles. Beneath the dermis lies the **subcutaneous layer**, made up of fatty connective tissue. Together, the layers of the skin protect underlying structures from physical trauma and ultraviolet (UV) radiation. The skin is essential to maintaining body temperature, fluid balance, and sensation. It is involved in absorption and excretion, immunity, and the synthesis of vitamin D from the sun (Fig. 10.1).

The Hair

The **hair** is also made up of keratinized cells. Hair is found over most of the body. It grows from hair follicles supplied by blood vessels located in the dermis. **Vellus**, which is short, pale, and fine hair, is located over all of the body. **Terminal hairs**, which are dark and coarse, are found on the scalp, brows, and, after puberty, on the legs, axillae, and perineum. The texture and color of hair are highly variable. Hair provides protection by covering the




FIGURE 10.1. Structures of the skin.

Structures and Functions of the Skin

STRUCTURE	FUNCTION
Epidermis	<ul style="list-style-type: none"> ■ Covers, protects, and waterproofs. ■ Contains four main layers: stratum corneum, stratum lucidum, stratum granulosum, and stratum germinativum.
<i>Stratum Corneum</i>	<ul style="list-style-type: none"> ■ Keratinized layer. Prevents loss or entry of water; protects against pathogens and chemicals.
<i>Stratum Lucidum</i>	<ul style="list-style-type: none"> ■ Translucent layer of dead cells with eleidin, precursor of soft keratin. ■ Found only on palms of hands and soles of feet; protects against UV sunrays to prevent sunburn.
<i>Stratum Granulosum</i>	<ul style="list-style-type: none"> ■ Keratinization begins at this layer. ■ Contains keratohyaline, keratinocytes, and Langerhans' cells. ■ Precursor of soft keratin.
<ul style="list-style-type: none"> ■ Keratohyaline ■ Keratinocytes ■ Langerhans' Cells 	<ul style="list-style-type: none"> ■ Secrete interleukin-1, which affects skin T-cell maturation. ■ Identify foreign substances (antigens), initiating the immune response.
<i>Stratum Germinativum</i>	<ul style="list-style-type: none"> ■ Continually produces new cells to replace worn-off surface cells. ■ Includes stratum spinosum and stratum basale.
<i>Stratum Spinosum</i>	<ul style="list-style-type: none"> ■ Has polyhedral cells (spinelike projections) that block entry to bacteria.
<i>Stratum Basale</i>	<ul style="list-style-type: none"> ■ Has single layer of columnar or cuboidal cells and melanocytes, which produce melanin. ■ Production of new cells occurs at this layer. ■ Site of basal cell carcinoma. ■ Melanin production increases with UV light exposure; melanin determines pigmentation and protects from UV light.
Dermis	<ul style="list-style-type: none"> ■ Contains collagen, reticular, and elastic fibers. ■ Adds strength and elasticity to skin. Contains papillary layer, reticular layer, sweat glands, sebaceous glands, cholesterol, and arterioles.
<i>Papillary Layer</i>	<ul style="list-style-type: none"> ■ Contains capillaries that supply the stratum germinativum; also contains nerve endings, touch receptors, and fingerprint pattern; double layer on hands and feet.
<i>Reticular Layer</i>	<ul style="list-style-type: none"> ■ Contains connective tissue with collagen and elastic fibers, blood and lymphatic vessels, nerves, free nerve endings, fat cells, sebaceous glands and hair roots, deep pressure receptors, and smooth muscle fibers.
<i>Sweat Glands (Sudoriferous)</i>	<ul style="list-style-type: none"> ■ Most numerous on palms of hands and soles of feet. ■ Two types are eccrine and apocrine glands.
<ul style="list-style-type: none"> ■ Eccrine Glands 	<ul style="list-style-type: none"> ■ Respond to external temperature and psychological stress. ■ Found over most of body but most numerous on palms of hands and soles of feet; secrete sweat, which helps regulate body temperature and, to a lesser degree, excretes wastes such as urea.
<ul style="list-style-type: none"> ■ Apocrine or Odoriferous Glands 	<ul style="list-style-type: none"> ■ Found in axilla and genital area. ■ Respond to stress; secrete pheromones, a substance with a barely perceptible odor; when apocrine secretions react with bacteria, body odor results. ■ Ceruminous glands are a type of apocrine gland found in the external ear canal. They secrete cerumen, which prevents drying of the ear drum and traps foreign substances.
<i>Sebaceous Glands</i>	<ul style="list-style-type: none"> ■ Produce sebum, which lubricates and protects skin and hair.
<i>Cholesterol</i>	<ul style="list-style-type: none"> ■ Converts to vitamin D when exposed to UV lights.
<i>Arterioles</i>	<ul style="list-style-type: none"> ■ Dilate when hot to increase heat loss and constrict when cold to conserve heat. ■ Constrict in response to stressful situations to shunt blood to vital organs.
Hypodermis/Subcutaneous	<ul style="list-style-type: none"> ■ Contains connective tissue and adipose tissue.
<i>Connective Tissue</i>	<ul style="list-style-type: none"> ■ Connects skin to muscles; contains white blood cells.
<i>Adipose Tissue</i>	<ul style="list-style-type: none"> ■ Contains stored energy, cushions bony prominences, provides insulation.

Structures and Functions of the Hair

STRUCTURE	FUNCTION
<p>Shaft: Portion that protrudes</p> <ul style="list-style-type: none"> Medulla has soft keratin, central core. Cortex has hard keratin. Cuticle has hard keratin, outer layer. <p>Root: Portion embedded in skin</p> <ul style="list-style-type: none"> Bulb is composed of a matrix of epithelial cells enclosed by a follicle. Arrector pili muscle. 	<ul style="list-style-type: none"> Scalp hair provides insulation, protection from heat and cold. Eyebrows cushion and protect eyes from glare and perspiration. Eyelashes protect eyes from foreign substances; nasal hair traps foreign particles. Pigment found in cortex gives hair color. Cellular mitosis occurs here. Causes hair to stand up (as in “goosebumps”) and squeezes sebum from sebaceous glands.

scalp and filtering dust and debris away from the nose, ears, and eyes.

The Nails

Nails are made up of hard, keratinized cells and grow from a nail root under the cuticle. Other nail structures include the free edge, which overhangs the tip of the finger or toe; the nailbed, or epithelial layer of skin; and the **lunula**, the proximal part of the nail. The nailbed's vascular supply gives the nail a pink color, although the nail itself is generally transparent. The purpose of the nails is to protect the distal portions of the digits and aid in picking up objects (Fig. 10.2).

Other Structures

Other appendages to the integument include the **sweat glands** and **sebaceous glands**. There are two types of sweat glands: **eccrine glands**, which are distributed over much of the body, and **apocrine glands**, which are limited to the genitalia, axillae, and areolae. Sebaceous glands are located near hair follicles, over




FIGURE 10.2. Structures of the nails.

most of the body. They secrete **sebum**, which lubricates the hair shaft.

Interaction With Other Body Systems

Changes in the integumentary system may reflect a problem in any of the systems described in the following paragraphs.

The Respiratory System

The respiratory system is responsible for obtaining the oxygen necessary for cellular metabolism, as well as for eliminating the carbon dioxide produced through the metabolic processes. If the process of respiration is impaired, alterations in the skin are most often evident through the development of **cyanosis**, a bluish discoloration, as hemoglobin becomes unsaturated with oxygen. **Central cyanosis** occurs when oxygen saturation is less than 80 percent and results in diffuse changes in the skin and mucous membranes. In contrast, **peripheral cyanosis**, which occurs in response to decreased cardiac output, is evident in areas of the body such as the nailbeds and lips, which are cooler than other regions; it may also be evident when an individual is chilled. In severe and chronic cardiopulmonary diseases, clubbing of the nails occurs owing to hypoxia (Fig. 10.3).




FIGURE 10.3. Nail clubbing.

The Cardiovascular System

The skin layer also contains a network of blood vessels, which contribute to its ability to regulate temperature and obtain nourishment. Alterations in the cardiovascular system can lead to circulatory impairment and changes in skin coloring and temperature, as well as to the development of **lesions**, **ulcerations**, and **necrosis**. When cardiac output is decreased, cyanosis may develop.

The Gastrointestinal System

The primary roles of the gastrointestinal (GI) system are the conversion of food to absorbable nutrients and the elimination of wastes. When GI disorders impair the body's ability to excrete toxins, the accumulation of toxins may become evident in the skin. For instance, when bile excretion is impaired so that bile builds up, **jaundice**, a yellow discoloration, often results. When dietary lipids accumulate, **xanthomas**, which are lipid deposits, or **papules** may develop. Nutritional deficits, which may stem from GI problems, are often evident in the skin. For example, deficits in vitamin A, riboflavin, vitamin C, iron and protein may all result in skin, hair, or nail alterations.

The Urinary System

The urinary system is primarily responsible for filtering the blood, but it is also involved in the production of red blood cells and the regulation of electrolyte and fluid status. When renal function is altered and filtration decreases, toxins and fluids build up in the body. The toxins often include **pigmented metabolites**, which alter the skin coloring. For example, an increased concentration of urea may lead to a residue of urea on the skin, which is called uremic frost. Toxins are also responsible for the development of **pruritus**, or itching. Calcium deposits may lead to **excoriations** of the skin. Altered hematologic status may be evident through **ecchymoses** and **hematomas**. Increased fluid volume associated with diminishing renal function may result in **edema**.

The Neurological System

The skin contains an intricate system of sensory and autonomic nerve fibers and serves as the body's largest sensory organ. Not only does this network of fibers permit the sensations of touch, temperature, pressure, vibration, and pain, the autonomic nervous system fibers control the skin's blood vessels and glands, regulating the skin's temperature, moisture, and oiliness. Alterations in the nervous system can have profound effects on the skin, placing it at risk for injury or discomfort. For instance, if sensation is decreased, a person is more likely to experience trauma to the skin because he or she is less likely to detect the need to withdraw from potentially dangerous objects or activities. Irritated nerves can

produce disagreeable sensations in the skin, such as burning. Alterations in the autonomic nervous system can result in drying of the skin.

The Endocrine System

Alterations of the endocrine system may affect the skin in myriad ways. Diabetes leads to alterations in skin integrity through complex processes involving changes in the immune, vascular, and neurological systems. Diabetic foot ulcers are examples of altered skin associated with diabetes. When thyroid disease occurs, the skin is often affected. In hypothyroidism, the skin is often dry and cool and becomes puffy, with nonpitting edema. It may develop a yellow hue as **carotene** accumulates. The hair is affected, becoming dull, brittle, and sparse. In contrast, hyperthyroidism causes the skin to be warmer, sweatier, and smoother than usual. The nails are thin and brittle and may separate from the nail plate. The hair is fine and silky, with patchy hair loss. Adrenal diseases affect the skin, hair, and nails. **Hypofunction** of the adrenals can result in **hyperpigmentation** of the skin (a bronze color) and **alopecia** (baldness). **Hyperfunction** results in thin and fragile skin, petechiae, plethora, bruises, and poor wound healing.


The Lymphatic/Immune System

The immune system is involved in protecting the body from both external and endogenous factors. Impairments in the immune system are reflected in the skin when infectious diseases result in their typical rashes or lesions. Abnormalities of the immune system can also result in hypersensitivity and the development of **atopic**, or **allergic**, skin changes, including pruritus or rashes such as **atopic dermatitis** or **psoriasis**. Skin changes are also common in some systemic autoimmune disorders such as **lupus erythematosus**.

Developmental, Cultural, and Ethnic Variations

Infants

Various differences can be noted in the newborn skin. Infants have very smooth skin, partly because of their lack of exposure to the environment, but also because there is less subcutaneous tissue. Color changes can be readily seen. Newborns often appear pinker or redder because of the lack of subcutaneous tissue. Physiological jaundice may occur 2 to 3 days after birth as a result of the breakdown of excessive red blood cells at birth and immature functioning of the liver. Newborns have little or no coarse terminal hair. They shed their hair at



Relationship of the Integumentary System to Other Systems

approximately 3 months and it is soon replaced. **Eccrine** sweat glands begin to function within a month after birth. The immature sweat glands lead to poor thermoregulation. With no functioning **apocrine** sweat glands, babies' skin is less oily than adults' and lacks offensive odor. The secretion of **sebum** by the sebaceous glands can result in cradle cap, seborrheic dermatitis, thick yellow-crusting lesions. Numerous skin lesions may also be seen on the newborn, such as **mongolian spots**, **nevus flammeus** (port-wine stains), **capillary hemangiomas** (stork bites), **hemangioma simplex** (strawberry marks), **milium**, and **erythema toxicum neonatorum**.

Adolescents

Adolescence is a time of rapid hormonal change that may affect the integumentary system. During adolescence, the apocrine glands begin to enlarge and function. At this time, young people develop increased axillary sweating and the potential for a more pronounced body odor. The sebaceous glands increase sebum production and the skin becomes more oily, leading to the onset of **acne**. During adolescence, pubic and axillary hair and male and female body hair patterns become apparent.

Pregnant Women

During pregnancy, there is increased blood flow to the skin, particularly to the hands and feet, as peripheral vessels dilate and the number of capillaries increases to dissipate heat. Along with this increased flow, there is an increase in sweating and sebaceous activity. The skin thickens and separates with stretching, with the appearance of **striae**. Hormonal changes result in hyperpigmentation. The pigmentary changes occur on the face resulting in **chloasma**, on the abdominal midline (the **linea alba** becomes the **linea nigra**) and on the nipples, areolae, axillae, and vulva.

Menopausal Women

During menopause, hormonal fluctuations result in hot flashes, often accompanied by flushing of the skin and increased pigmentation. There may be an increase in facial hair and some degree of scalp hair loss. Chloasma may occur. The incidence of skin tags increases at menopause.

Older Adults

With age, the skin atrophies. There is a decrease in production of sebum and sweat. The skin becomes drier and flattens, often becoming paperlike. The elasticity decreases and wrinkles develop. There is a decreased melanocyte

function, so that the hair grays and the skin becomes more pale. Target areas of increased melanocyte function result in "age spots." There is a decrease in axillary, pubic, and scalp hair. Women may experience increased facial hair as estrogen function is lost; men experience an increase in nasal and ear hair growth. The nails grow more slowly and become thicker and more brittle. Furthermore, specific skin lesions are more common in elderly persons, including: **actinic keratoses**, **basal cell carcinomas**, **seborrheic keratoses**, **stasis ulcers**, **senile pruritus**, and **keratotic horns**.

People of Different Cultures and Ethnic Groups

Cultural and ethnic variations can readily be seen in the integumentary system. Genetic factors determine the skin color. The greater the amount of melanin, the darker the skin color. Assessing for subtle changes in the skin becomes more of a challenge the darker the skin color. The oral mucosa is best for assessing color changes in dark-skinned people. Also, assessing the sclera for jaundice is more accurate than assessing the skin in an Asian person. Fair-skinned persons of Irish, German, or Polish descent have an increased risk for skin cancer with prolonged sun exposure. African Americans have a higher incidence of **keloids**, **pseudofolliculitis**, and **mongolian spots**.

Differences in hair can also be readily seen with different ethnic groups. Asians often have black, straight, silky hair, and Chinese men have very little facial hair. The hair texture of African Americans is often thick and kinky. Cultural variations may also be noted in the amount of sweat production. For example, Asians produce less sweat and therefore have less body odor. Be alert to your patient's ethnicity when assessing the integumentary system; changes may be culturally related rather than an indication of pathology.

Introducing the Case Study



As a nurse in a community health setting, you have volunteered to participate as an examiner in a community skin cancer screening program. Eva Green, a 48-year-old woman with fair, freckled skin, presents for a total skin assessment. She states that several of her relatives have been diagnosed with skin cancer and she is very much afraid that she might have it. She goes on to say that because she has always had very freckled skin that "easily blemishes," she has difficulty determining what is normal and what should be a concern. As part of the screening program, you will complete a health history and physical examination that focuses on the integumentary system.

Performing the Integumentary Assessment

Assessment of the integumentary system includes a comprehensive health history and physical examination. The history identifies any symptoms related to the integumentary system, risk factors for skin problems, and the presence of diseases in other systems that could contribute to skin problems. The physical examination identifies the current condition of the integument, including any abnormal function. Throughout the assessment, be attentive to signs or symptoms of both actual and potential problems of the integument.

Health History

The health history includes obtaining biographical data and asking questions about the patient's current health, past health, and family and psychosocial history. It also involves a review of systems. If you don't have the time to perform a complete health history, make sure to at least perform a focused health history of the integumentary system.

Focused Integumentary History

If your patient's condition or time prohibits a detailed integumentary history, ask the following questions:

- Do you have any changes in your skin, hair, or nails? (If yes, ask questions following the PQRST format.)
- Do you have any food, drug, or environmental allergies?
- Do you have any medical problems?
- Are you on any medications, prescribed or over-the-counter (OTC)?

Biographical Data

Briefly review all biographical data. Identify your patient's age; skin function varies by age, and certain skin diseases are more likely to develop at particular ages. For example, **papules**, **vesicles**, and **pustules** associated with **impetigo** are mostly seen in children; **acne** frequently occurs during puberty, and **plaques** and **malignancies** are more common in older adults. Your patient's occupation and recreational activities can provide clues regarding the potential for exposures to harmful chemicals, trauma, or environmental hazards. Exposure to chemicals may cause a **contact dermatitis**. Excessive sun exposure from either work or play may increase the risk for skin cancer. Similarly, the patient's living situation can suggest environmental exposures to factors that might be harmful to the skin.

Current Health Status

If the patient has a specific skin complaint, analyze it as you begin your history, using the PQRST system. The major complaints to be alert for include: changes in moles or other lesions, nonhealing sore or ulcer or chronic irritation, pruritus/itching, and rash, a very common complaint. Because the integumentary system also includes the hair and nails, a symptom analysis should focus on changes in these areas. Generally, the problems that trigger integumentary complaints are of a stable nature and not life-threatening, and so you will be able to proceed with the full history and physical. If the patient is overly distressed by the symptoms (e.g., severe itching or fear of malignancy), focus first on the presenting problem and perform a comprehensive history at a later time.



HELPFUL HINT

- Remember, skin, hair, and nails cells reproduce rapidly, and so changes are often seen when a serious acute health problem occurs or with use of chemotherapeutic agents.
- Changes in skin, hair, and nails are often seen with underlying endocrine problems.

Symptom Analysis

Symptom analysis tables for the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Change in Mole or Lesion

Skin cancer is the most common type of cancer, and changes in a mole (**nevus**) or skin lesion can often evoke fear in the patient. Although there are many types of lesions, most of which are benign, it is important to be able to detect skin cancer at its earliest stages, when treatment yields the best results. There are three types of skin cancer: basal cell and squamous cell carcinomas, which affect the epidermal keratinocytes, and melanoma, which affects the melanocytes of the basal layer of the epidermis. Sun exposure is a risk factor in all types.

The majority of skin cancers are basal cell. Basal cell carcinomas are directly related to sun exposure, with 90 percent of lesions occurring on the head and neck. Basal cell carcinomas are easily treated and relatively benign. Squamous cell carcinoma is often preceded by **actinic keratosis** (pre-malignant macule or papule of rough, sandpaper texture, caused by excessive UV exposure). Sixty-six percent of these lesions occur on sun-exposed areas and respond well to treatment. Even though melanoma occurs less frequently than basal cell and squamous cell carcinomas, it is the most deadly type of skin cancer. **Congenital** nevi and **dysplastic** nevi may be precursor lesions to melanoma.

Age-Related Skin Disorders

AGE	DISORDERS
Children	<ul style="list-style-type: none"> ■ Impetigo ■ Atopic dermatitis or eczema ■ Pityriasis rosea ■ Juvenile plantar dermatosis ■ Rashes secondary to bacterial or viral infections ■ Pediculosis capitis ■ Varicella
Adolescents/ Young Adults	<ul style="list-style-type: none"> ■ Acne ■ Pityriasis rosea ■ Tinea versicolor
Adults	<ul style="list-style-type: none"> ■ Psoriasis ■ Seborrheic dermatitis ■ Malignant melanoma ■ Herpes simplex virus type 2 (HSV-2) ■ Tinea cruris ■ Seborrheic interigo ■ Rosacea
Older Adults	<ul style="list-style-type: none"> ■ Actinic keratosis ■ Seborrheic keratosis ■ Basal cell carcinoma ■ Squamous cell carcinoma ■ Xerosis ■ Herpes zoster
Children and Adults	<ul style="list-style-type: none"> ■ Nummular eczema/dermatitis ■ Scabies ■ Insect bites, poisonous plants ■ Contact dermatitis ■ Herpes zoster ■ Tinea pedis

Unfortunately, there are usually no symptoms associated with skin cancers unless the lesion has metastasized regionally or distantly. In these cases, various symptoms might be present related to whether a lesion, for instance a malignant melanoma, had metastasized to another organ such as the bowel, lung, liver, or brain.

Nonhealing Sore or Chronic Ulceration

When your patient's history includes a sore that won't heal or a chronic ulceration, the routine symptom analysis questions will provide a good picture of the lesion and any previous self-treatment applied. Keep in mind that a nonhealing wound or chronic irritation is often associated with an underlying disease. The most common types of nonhealing wounds or chronic skin ulcerations are caused by vascular disease or pressure or by diabetes.

Pruritus

Pruritus is severe itching. It may be localized or generalized and caused by a dermatologic problem or underlying systemic problem. Pruritus is often accompanied by a rash. Itching, when not associated with a rash, may be indicative of significant systemic disease or simply dry skin.

Itching arises from free nerve endings (nonmyelinated), which are especially abundant in the flexor aspects of the wrist and ankles. It occurs as a result of a spinal reflex and external stimuli, such as heat, dryness, inflammation, and vasodilation. Psychological factors, such as depression, can influence the perception of itching, which explains the varied responses to it. A thorough symptom analysis will help you to pinpoint the underlying cause.

Rashes

Rashes, like itching, may be localized or generalized, acute or chronic, and caused by an obvious dermatologic problem or an underlying systemic problem. A thorough symptom analysis will help you to pinpoint the problem and direct your physical assessment.

Seasonal Skin Disorders

Certain skin disorders are more common during one time of year than others. Seasonal skin problems include those caused by temperature fluctuations, air humidity, and exposure to contaminants. It is important to remember, however, that although these problems may be more common at certain times of the year, they may appear at any time.

Seasonal Skin Disorders

SEASON	SKIN DISORDERS
Spring	<ul style="list-style-type: none"> ■ Pityriasis rosea ■ Chickenpox ■ Acne flare-ups
Summer	<ul style="list-style-type: none"> ■ Contact dermatitis ■ Tinea ■ <i>Candida</i> ■ Impetigo ■ Insect bites
Fall	<ul style="list-style-type: none"> ■ Senile pruritus/winter itch ■ Pityriasis rosea ■ Urticaria ■ Acne flare-ups
Winter	<ul style="list-style-type: none"> ■ Contact dermatitis of hands ■ Senile pruritus/winter itch ■ Psoriasis ■ Eczema

Hair Changes

Hair loss (**alopecia**) is probably the most distressing change in hair that can occur because of its cosmetic effect. Alopecia not only refers to scalp hair but also to body hair. Normally hair growth is cyclical, with 85 to 90 percent of scalp hair in the growth phase (**anagen**), and the remaining 10 to 15 percent in the resting phase (**telogen**). Scalp hair grows about 0.25 mm/d, and about 100 strands of hair are lost per day. Hair loss can occur for many reasons. Alopecia can be classified as scarring or **cicatricial** (resulting from injury such as burns, radiation, or traction with irreversible damage to the hair follicles) and nonscarring or **noncicatricial** (resulting from hormonal changes, medications, infectious diseases, or thyroid disease, in

which the follicles remain intact with a potential to reverse the process).

Nail Changes

Changes in the nails also often reflect an underlying systemic problem. Changes in color and texture are frequent complaints. A symptom analysis will help you identify any underlying problems.

Past Health History

The past health history allows you to determine what illnesses or problems the patient has had in the past, including those related to the skin, hair, or nails. It additionally permits you to determine episodes of illnesses involving other systems that might have an impact on the integument.

Past Health History

RISK FACTORS/QUESTIONS TO ASK

Childhood Illnesses

- Do you have a history of infectious diseases such as chickenpox or rubella?
- Did you have asthma or allergies as a child?

- Did you have frequent skin problems or chronic skin illness?

Surgery/Injuries

- Have you ever had surgery, biopsies, or a major traumatic injury?

Hospitalizations/Diagnostic Procedures

- Have you have been hospitalized or had procedures such as electrophysiological studies or scans?

Allergies

- Are you allergic to foods, drugs, chemicals, or environmental agents? If yes, what type of reaction?

Other Medical Problems

- Do you have a history of skin cancer, eczema, or psoriasis?
- Do you have a history of asthma?
- Do you have a history of diabetes, thyroid disease, respiratory disorders, liver disease, or heart disease?

RATIONALE/SIGNIFICANCE

- Explains scarring from rashes and degree of immunity for diseases that have skin eruptions.
- History of asthma/allergic disorders commonly associated with skin problems such as eczema or psoriasis.
- Explains scarring or chronic changes to skin and potential for recurrence.

- History of surgery/trauma helps explain scars. Keloids may also develop on scar tissue.
- Biopsies may signal history of skin cancer or other malignancies associated with skin changes.

- Helps identify previous disorders that might be associated with skin condition.
- History of diagnostic procedures provides information on major problems that might influence integumentary system.

- Skin changes are often related to substances that are ingested, applied, or come into contact with skin.

- Consider potential for recurrence.
- Often associated with atopic problems including atopic dermatitis, psoriasis.
- May be reflected in skin's appearance. Diabetes causes poor wound healing; hyperthyroidism, moist skin and skin necrosis; hypothyroidism, dry, scaly skin, hair loss, brittle nails and transverse ridges; chronic lung disease, cyanotic color changes; liver disease, jaundice and itching; and cardiovascular disease, color changes, edema, leg ulcers.

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Exposures to Infectious or Contagious Illnesses</p> <ul style="list-style-type: none"> ■ Have you recently been exposed to any infectious or contagious disease? ■ Have you been exposed to anyone with lice or scabies? <p>Immunizations</p> <ul style="list-style-type: none"> ■ What immunizations have you had; for example, mumps, measles, rubella (MMR), varicella, hepatitis A and B? <p>Medication</p> <ul style="list-style-type: none"> ■ Are you taking any prescribed or OTC medications? ■ Do you use recreational drugs? 	<ul style="list-style-type: none"> ■ Impetigo, chickenpox, herpes, and other contagious disease cause skin rashes. ■ Exposures to infestations can cause skin and hair problems. ■ Identifies likelihood of immunity against vaccine-preventable diseases that have associated skin changes. ■ Any medications can cause adverse skin reactions and allergic dermatitis. See <i>Drugs That Adversely Affect the Integumentary System</i>. ■ Antimalarials may cause pigmentary changes; many others result in photophobia, rashes, or other skin effects. ■ Use of specific medications points to diseases of other organs that may be associated with skin changes. For instance, insulin indicates a diagnosis of diabetes. ■ Provides information on patient's understanding of currently prescribed treatments.

Drugs That Adversely Affect the Integumentary System

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Corticosteroids	<ul style="list-style-type: none"> ■ Methylprednisolone, prednisone, dexamethasone 	<ul style="list-style-type: none"> ■ Urticaria, skin atrophy and thinning, acne, facial erythema, striae, allergic dermatitis, petechiae, ecchymoses.
	<ul style="list-style-type: none"> ■ Corticotropin (ACTH) 	<ul style="list-style-type: none"> ■ Urticaria, pruritus, scarlatiniform exanthema, skin atrophy and thinning, acne, facial erythema, hyperpigmentation.
Anticonvulsants	<ul style="list-style-type: none"> ■ Carbamazepine 	<ul style="list-style-type: none"> ■ Pruritic rash, toxic epidermal necrolysis, Stevens-Johnson syndrome.
	<ul style="list-style-type: none"> ■ Lamotrigine 	<ul style="list-style-type: none"> ■ Same as Carbamazepine.
	<ul style="list-style-type: none"> ■ Valproic acid 	<ul style="list-style-type: none"> ■ Alopecia.
	<ul style="list-style-type: none"> ■ Phenytoin ■ Ethosuximide 	<ul style="list-style-type: none"> ■ Morbilliform (measles-like) rash, hirsutism. ■ Urticaria, pruritic (itchy) and erythematous (reddened) rashes
Antimalarial	<ul style="list-style-type: none"> ■ Chloroquine 	<ul style="list-style-type: none"> ■ Pruritus; pigmentary skin changes, eruptions resembling lichen planus (with prolonged therapy).
Antineoplastic Agents	<ul style="list-style-type: none"> ■ Bleomycin 	<ul style="list-style-type: none"> ■ Skin toxicity may be accompanied by hypoesthesia that may progress to hyperesthesia, urticaria, erythematous swelling, hyperpigmentation, patchy hyperkeratosis, alopecia.
	<ul style="list-style-type: none"> ■ Busulfan 	<ul style="list-style-type: none"> ■ Cheilosis, hyperpigmentation, urticaria, dry skin, alopecia, anhidrosis (absent or deficient sweating).
	<ul style="list-style-type: none"> ■ Cyclophosphamide 	<ul style="list-style-type: none"> ■ Skin and fingernail pigmentation, alopecia

(continued)

Drugs That Adversely Affect the Integumentary System (continued)

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Barbiturates	<ul style="list-style-type: none"> ■ Pentobarbital, phenobarbital 	<ul style="list-style-type: none"> ■ Urticaria; maculopapular, morbilliform, or scarlatiniform rash.
Cephalosporins	<ul style="list-style-type: none"> ■ Cefazolin, cefoxitin, cefuroxime, ceftriaxone, cefotaxime, and others in this class 	<ul style="list-style-type: none"> ■ Rash, pruritus, urticaria, erythema multiforme.
Gold Salts	<ul style="list-style-type: none"> ■ Auranofin, gold sodium thiomalate 	<ul style="list-style-type: none"> ■ Rash, pruritus, alopecia, photosensitivity, urticaria.
Nonsteroidal Anti-Inflammatory Agents	<ul style="list-style-type: none"> ■ Diflunisal ■ Ibuprofen ■ Sulindac 	<ul style="list-style-type: none"> ■ Rash, pruritus, erythema multiforme, Stevens-Johnson syndrome. ■ Rash, erythema multiforme, Stevens-Johnson syndrome. ■ Rash, pruritus, photosensitivity, erythema multiforme, Stevens-Johnson syndrome.
Oral Antidiabetic Agents	<ul style="list-style-type: none"> ■ All types 	<ul style="list-style-type: none"> ■ Photosensitivity, various skin eruptions.
Penicillins	<ul style="list-style-type: none"> ■ Amoxicillin, ampicillin, dicloxacillin, penicillin G potassium, penicillin V potassium, piperacillin, ticarcillin, nafcillin 	<ul style="list-style-type: none"> ■ Urticaria, erythema, maculopapular rash, pruritus.
Phenothiazines	<ul style="list-style-type: none"> ■ Chlorpromazine, thioridazine, trifluoperazine, and others in this class 	<ul style="list-style-type: none"> ■ Dermatoses, pruritus, marked photosensitivity, urticaria, erythema, eczema, exfoliative dermatitis.
Sulfonamides	<ul style="list-style-type: none"> ■ Sulfamethoxazole, sulfasalazine, sulfisoxazole 	<ul style="list-style-type: none"> ■ Rash, pruritus, erythema nodosum, erythema multiforme, Stevens-Johnson syndrome, exfoliative dermatitis, photosensitivity.
Tetracyclines	<ul style="list-style-type: none"> ■ Demeclocycline, doxycycline, minocycline, tetracycline 	<ul style="list-style-type: none"> ■ Photosensitivity.
Miscellaneous Agents	<ul style="list-style-type: none"> ■ Allopurinol ■ Captopril ■ Griseofulvin ■ Isotretinoin ■ Oral contraceptives (estrogen) ■ Thiazide and loop diuretics ■ Lithium ■ Warfarin 	<ul style="list-style-type: none"> ■ Pruritic maculopapular rash, exfoliative dermatitis, urticaria, erythematous dermatitis. ■ Maculopapular rash, pruritus, erythema. ■ Rash, urticaria, photosensitivity, erythema multiforme or lupus-like syndrome. ■ Alopecia, photosensitivity, hirsutism, pruritus, urticaria. ■ Chloasma or melasma, rash, urticaria, erythema. ■ Photosensitivity. ■ Acne, alopecia. ■ Skin necrosis.

Family History

The family history allows you to determine what, if any, integumentary problems are common to the patient's family members. Diseases of the integument are as likely as others to have a familial predisposition. The history

also helps to identify familial diseases that directly affect other systems and that might have some affect on the skin. This portion of the history also begins to explore the potential for problems stemming from the patient's living environment.

Family History

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Skin Disorders

- Do you have a family history of skin cancer, psoriasis, or eczema?

- Identifies familial predisposition for skin disease that can be inherited.

Cardiovascular Disorders

- Do you have a family history of heart anomaly, output disorders, or peripheral vascular disease?

- Identifies familial disorders that might affect skin.

Pulmonary Disorders

- Do you have a family history of asthma or other chronic lung diseases?

Gastrointestinal Disorders

- Do you have a family history of liver or gallbladder disease?

Endocrine Disorders

- Do you have a family history of diabetes or thyroid disorders?

Immune Disorders

- Do you have a family history of allergies or autoimmune diseases such as lupus erythematosus?

- Identifies familial tendencies and environmental triggers.

Infectious/Contagious Diseases in Family

- Has anyone in your family recently had an infectious or a contagious disorder?

- Identifies contagious diseases patient has been exposed to.

Review of Systems

The review of systems is extremely important when exploring a complaint related to the integumentary system. It helps you identify problems in other systems that directly affect the skin, hair, and nails. Health problems

directly affecting many systems can have profound effects on the integument. The review of systems is also useful in prompting the patient to identify problems that she or he previously felt were not related or worth mentioning.

Review of Systems

SYSTEM/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- Have you noticed a change in your energy level?

- Helps identify problems that are relatively nonspecific to any one system. Many endocrine and immune diseases present with this type of complaint.

- Do you have fevers, weight changes, or sweats?

Head, Eyes, Ears, Nose, and Throat (HEENT)

Head/Neck/Throat

- Do you have swollen glands/nodes, congestion, nasal discharge, sore throat, or sneezing?

- Swollen glands or nodes may be associated with infectious or malignant cause of skin problems.
- Congestion and sore throat are common symptoms of several viral illnesses that typically cause skin changes.

Eyes

- Do you have watery or itchy eyes?

- Congestion with watery/itchy eyes is common in atopic disorders that may be associated with urticaria, eczema, and other skin problems.

Respiratory

- Do you have a cough, wheezing, tightness in your chest, or difficulty breathing?

- Chronic lung disorders such as emphysema and bronchitis may limit oxygenation to the point of cyanosis.
- Reactive airway disease (asthma) may be associated with atopic disorders of the skin.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Cardiovascular</p> <ul style="list-style-type: none"> ■ Do you have pain in your legs when walking (claudication), shortness of breath, difficulty breathing (dyspnea), chest pain, or palpitations? 	<ul style="list-style-type: none"> ■ Dyspnea, palpitations, and chest pain are seen with cardiac disease and result in poor output, which contributes to cyanosis. ■ Claudication is common in vascular disorders and results in discoloration, hair loss, and ulceration of skin over lower extremities.
<p>Gastrointestinal</p> <ul style="list-style-type: none"> ■ Do you have abdominal pain, loss of appetite, nausea/vomiting, jaundice, changes in bowel habits or in color of stool? 	<ul style="list-style-type: none"> ■ Abdominal pain, anorexia, nausea, changes in bowel habits and clay-colored stools occur with liver diseases. ■ Liver disease can result in jaundice and pruritus.
<p>Genitourinary</p> <ul style="list-style-type: none"> ■ Have you noticed a change in the color of your urine? ■ Do you have increased urination or burning? 	<ul style="list-style-type: none"> ■ Dark orange urine is often seen with obstructive jaundice. ■ Urinary tract infections (UTIs) are frequently seen in type 2 diabetes, which can lead to skin changes. ■ Can cause dermatitis.
<p>Female Reproductive</p> <ul style="list-style-type: none"> ■ When was your last menstrual period? Are they regular periods? ■ Are you pregnant? ■ Do you practice safe sex? ■ Do you have any vaginal discharge? 	<ul style="list-style-type: none"> ■ Menstrual irregularities are often found in endocrine disorders, including hypothyroidism and glucose intolerance. ■ Pregnancy may explain localized changes in skin pigmentation. ■ Unprotected sex increases the risk for sexually transmitted diseases (STDs). ■ Discharge may be associated with STD and may also cause skin lesions of genitalia or other skin.
<p>Male Reproductive</p> <ul style="list-style-type: none"> ■ Do you practice safe sex? ■ Do you have any penile discharge or rashes? 	<ul style="list-style-type: none"> ■ Unprotected sex increases the risk for STDs. ■ Discharge may be associated with STD, which may also cause lesions on genitalia or other skin.
<p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Do you have joint pain, swelling, or redness? 	<ul style="list-style-type: none"> ■ Common in some autoimmune disorders such as lupus and in infectious syndromes, such as Reiter's syndrome, that affect the skin. ■ Gout is associated with development of tophi, uric acid deposits beneath the skin
<p>Neurological</p> <ul style="list-style-type: none"> ■ Do you have numbness, tingling, or burning? 	<ul style="list-style-type: none"> ■ The altered sensation found in neuropathic disorders can predispose to skin injuries. Altered sensation is also found in some neurological disorders that result in skin dryness.
<p>Immune/Hematologic</p> <ul style="list-style-type: none"> ■ Do you have any bleeding disorders, or are you on any anticoagulants? ■ Are you on any immunosuppressive drugs? ■ Do you have any immune disorders? 	<ul style="list-style-type: none"> ■ Bleeding disorders or use of anticoagulants may result in vascular lesions such as petechiae, purpura, and ecchymosis. ■ Immunosuppressive drugs can cause skin changes. For example, steroids can cause acne, decreased wound healing, ecchymosis, thinning of skin, hirsutism and petechiae. ■ Immune diseases such as lupus or human immunodeficiency virus (HIV) often have skin manifestations, such as Kaposi's sarcoma.

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Endocrine</p> <ul style="list-style-type: none"> ■ Do you have increased thirst, frequent urination, fatigue, or weight loss? ■ Have you gained weight? ■ Do you have energy loss or dry skin? 	<ul style="list-style-type: none"> ■ Polyuria, polydipsia, polyphagia, fatigue and weight loss are common symptoms in diabetes, which alters the healing ability of the skin and promotes other changes through alterations in circulation and sensation. ■ Fatigue and weight gain are commonly found in hypothyroid disorders, which result in dry skin and pruritus.

Psychosocial Profile

The psychosocial profile serves several purposes. It provides important information regarding dietary and other habits, as well as occupational, social, and recreational

activities that could influence the condition or health of the skin, hair, and nails. It provides an opportunity to explore the patient's self-care and social activities, which may identify his or her response to an integumentary system problem.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Health Practices and Beliefs</p> <ul style="list-style-type: none"> ■ Do you get yearly physical exams? ■ Do you examine your skin regularly for any unusual lesions, or changes in moles? 	<ul style="list-style-type: none"> ■ Determines if your patient takes preventive measures concerning skin care. ■ Health promotion activities such as annual exams, self-examinations, and use of sunblock are proactive measures. ■ Helps determine health promotion and protective patterns. ■ May identify teaching needs.
<p>Self-Care Activities</p> <ul style="list-style-type: none"> ■ Do you use sunblock? If yes, number? ■ Do you go to tanning salons? If, yes how often? ■ Do you use protective eyewear? Do you wear sunglasses? ■ How often do you bathe and shampoo your hair? What products do you normally use? Do you use detergents or hair dyes? ■ How do you care for your nails? ■ How often do you have manicures or pedicures? 	<ul style="list-style-type: none"> ■ Certain products or overuse of products can result in altered integrity or appearance of the skin, hair, and nails. ■ Poor hygiene can also alter skin, hair, and nail integrity.
<p>Typical Day</p> <ul style="list-style-type: none"> ■ How would you describe your typical day? 	<ul style="list-style-type: none"> ■ Identifies frequent exposure to chemicals, light, or temperature variations that affect skin.
<p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ What did you eat and drink in the past 24 hours? ■ What is your usual fluid intake? ■ What dietary supplements do you use? 	<ul style="list-style-type: none"> ■ Nutritional deficits and inadequate hydration can affect the skin. ■ Anorexia nervosa can cause loss of hair and blotchy, sallow skin. ■ Diets high in yellow vegetables can cause carotemia (yellow-orange color of skin). ■ Iron deficiency anemia can cause spoon nails and pallor. ■ Dehydration can cause dry skin.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Activity/Exercise Patterns <ul style="list-style-type: none"> ■ Do you exercise routinely? ■ What type of exercise do you do? 	<ul style="list-style-type: none"> ■ Exercise patterns may explain skin calluses, abrasions, potential for injuries and the need for education regarding safety and use of protection.
Recreation, Pets, Hobbies <ul style="list-style-type: none"> ■ What type of recreational activities do you do? ■ Do you have any pets? What kind? ■ Do you have any hobbies? 	<ul style="list-style-type: none"> ■ Outdoor activities increase sun exposure and risk for skin cancers. ■ Assess use of protection, such as sunscreens, hats. ■ Allergies are frequently associated with animals. Pets may also be carriers of fleas and ticks, which can lead to infestation and skin irritations and disease. ■ Chemicals associated with some hobbies may result in contact dermatitis.
Sleep/Rest Patterns <ul style="list-style-type: none"> ■ How many hours of sleep do you get daily? ■ Are you having any sleep problems? 	<ul style="list-style-type: none"> ■ Itching and other skin symptoms may result in altered sleep habits. ■ Difficulty sleeping may also help identify other problems that might be associated with skin changes, including thyroid or renal diseases.
Personal Habits <ul style="list-style-type: none"> ■ Do you smoke cigarettes? If so, how many/day and for how many years? ■ Do you use recreational drugs or alcohol? 	<ul style="list-style-type: none"> ■ Tobacco causes vascular impairment with related skin changes. Cigarette smoking increases wrinkling of skin, and nicotine stains fingers and nails yellow. ■ Alcohol abuse is a contributing factor for liver disease and related skin changes. ■ Illegal drug use increases risk for hepatitis B, HIV, and other infectious diseases with associated skin changes.
Occupational Health Patterns <ul style="list-style-type: none"> ■ What type of work do you do? ■ Are you exposed to chemical irritants? 	<ul style="list-style-type: none"> ■ Identifies potential risks associated with patient's occupation, as well as her or his understanding of proper use of protective devices and equipment.
Socioeconomic Status <ul style="list-style-type: none"> ■ Do you have health insurance? 	<ul style="list-style-type: none"> ■ Determine available resources for patient and determine if it has impact on his or her health care.
Environmental Health Patterns <ul style="list-style-type: none"> ■ How would you describe your home environment? ■ What type of heating and cooling do you use? 	<ul style="list-style-type: none"> ■ Important in determining potential for exposures, including infestations of mites, lice, etc. ■ Ability to properly heat and cool a home or work setting influences skin integrity.
Roles/Relationships/Self-Concept <ul style="list-style-type: none"> ■ Do you participate in group activities? ■ Have any changes in your skin, hair, or nails affected your personal relationships? 	<ul style="list-style-type: none"> ■ Groups increase risk for contracting contagious diseases. ■ Because skin, hair, and nails are so evident, alterations may cause some people to withdraw from or limit interaction with others.
Stress and Coping <ul style="list-style-type: none"> ■ How much stress do you routinely have? ■ How do you cope with stress? 	<ul style="list-style-type: none"> ■ Stress can trigger changes in skin, hair, and nails. ■ Persons with alterations in the skin, hair, or nails may be embarrassed about their appearance, and this can add to usual stressors.



Case Study Findings

Reviewing Mrs. Green's

Biographical data:

- 48-year-old woman, married, with two adult children (27 and 22 years old).
- Works part-time in retail sales.
- Born in the United States, of English-Scottish descent.
- Protestant religion.
- Husband is accountant and provides health insurance through his employer.

Current health status:

- Feels she is in very good health. No current problems.
- Has recently started taking estrogen and progesterone (Prempro) daily for menopause symptoms. Is tolerating therapy well.
- Occasionally takes aspirin or acetaminophen for headache or back and shoulder discomfort, but not monthly.

Past health history:

- Usual childhood diseases, without sequelae.
- One miscarriage at 10 weeks' pregnancy, age 22.
- Hospitalized for two childbirths, with uncomplicated vaginal deliveries and a cholecystectomy 2 years ago.
- No history of allergies, skin disorders.
- No recent exposures to persons with infections.
- Unsure about immunization status. Received all immunizations in youth, but cannot recall later tetanus, others.

Family history:

- Mother died of breast cancer at age 65, had previously had two basal cell lesions removed.
- Brother (45) has hypertension and a history of childhood asthma.
- Father (73) has "borderline high blood pressure" and adult-onset diabetes but is in generally good health. He also had two basal cell cancers on his face and has several "sun spots."
- Two older siblings (49 and 50) are in generally good health, although one had a small malignant melanoma between two toes, which was surgically removed 3 weeks ago, with no lesions or spread detected.
- Children are both in good health.

Review of systems:

- *General Health Survey:* Feels well overall.
- *Integumentary:* Skin is somewhat dry; itches occasionally. No sores or rashes. Reports no changes in hair and nails.

- *Head, Eyes, Ears, Nose, and Throat (HEENT):* Runny nose/itchy eyes.

- *Respiratory:* No shortness of breath, congestion, cough.

- *Cardiovascular:* No chest pain.

- *Gastrointestinal:* No abdominal discomfort, nausea, change in bowel pattern.

- *Genitourinary:* No urinary problems.

- *Reproductive:* Last menstrual period was about 1.5 years ago. Before that they were regular until the final year, when they fluctuated a little.

Musculoskeletal: Joint pain (except mild after extensive gardening).

Neurological: No dizziness.

Psychosocial profile:

- She bathes or showers daily, usually with very warm water. She shampoos her hair every other day and rarely uses any hair spray, mousse, or other products. She keeps her nails closely filed and wears clear polish. She wears cosmetics daily and has never had problems tolerating any brand. She colored her hair in the past, but does not do so at this time.
- She is often out in the Florida sun as she walks and gardens, but only occasionally wears sunscreen. She does not reapply the product when she has been out for an extended time, which often occurs. She thought that skin cancer was something she would worry about in old age; however, now that her sister has had skin cancer she has been worried about any damage she may have done to her skin.
- Typical day starts at 7 A.M., when her alarm awakens her. On days when she works (2 to 3 days a week), she showers, dresses, eats breakfast, then works from 9 A.M. to 5 P.M. On her off days, she cleans her house, works in the yard/garden when weather permits, and just "keeps busy" until about 5 P.M. She then fixes dinner and watches TV or does needlework in the evening. She generally goes to bed at 10:30 to 11:00 P.M.
- She eats three meals daily, including foods from all categories. She feels that she should lose 10 to 15 lb, which she has gradually gained over the past 10 years. She tries to remember to drink several glasses of water daily, but sometimes only drinks one or two 8-oz glasses.
- Gardening is one of her main forms of exercise. She also walks 2 or 3 miles a day, three or four times a week.
- She sleeps very well, with rare interruptions to void (two to three times per week).

(continued on page 222)



Case Study Findings (continued)

- Her only medications are those mentioned earlier. She drinks 1 cup of coffee and 4 to 5 glasses of iced tea daily. She rarely drinks colas. She has never smoked but used to be exposed to cigarette smoke until her husband quit 5 years ago. She does not drink alcohol or use recreational drugs.
- She is not exposed to chemicals or unusual environmental situations at work. She uses various chemicals in her gardening (fertilizers, pesticides, weed killers) and doesn't always wear gloves, masks, or other protective equipment.
- She has no pets. She has lived in her current home, a ranch style, for 12 years. The home is very adequate for her needs, and has central heat and air conditioning.
- She enjoys meeting customers at work and has several close friends with whom she often has lunch or shops. She and her husband have a very close and caring relationship. She has good relationships with her two children, who live nearby. Her father and siblings live 600 miles away, but she sees them two or three times a year for an enjoyable visit. She is not currently involved in any specific community group and rarely attends church.



Case Study Evaluation

Before you proceed with the physical examination of the skin, hair, and nails, document the key information you have learned from Mrs. Green's health history and the integumentary history related to her.



CRITICAL THINKING ACTIVITY #1

What history findings put Mrs. Green at risk for skin problems?



CRITICAL THINKING ACTIVITY #2

What strengths can you identify that will help Mrs. Green adapt to or prevent skin disorders?

Physical Assessment

Once you have taken the history, proceed to collect objective data through your physical examination. Even though the skin, hair, and nails are easily accessible and we look at them every day, you still need to be very objective and attentive to details that could easily be overlooked.

Approach

The techniques used in the examination of the integument are inspection and palpation. As you conduct the assessment, along with your sense of sight and touch, use your sense of smell to note any unusual odors. It is important to inspect all areas of skin, including **intertriginous** areas, which lie between or under folds of skin. Throughout the examination, compare symmetrical parts. Also be aware of the "feel" of the skin, hair, and nails. You can inspect the skin in one of three ways:

1. Using a head-to-toe approach.
2. Observing all skin on the anterior, posterior, and lateral surfaces of the body.
3. Inspecting the skin by regions, as you examine the cardiovascular, respiratory, and other systems.

Regardless of your approach, a complete examination is necessary, and a systematic approach will help you avoid omissions. During the examination, keep in mind the underlying structures or organs because they may explain changes in the overlying skin. Also compare exposed with unexposed areas. Variations might be signs of "wear and tear," poor alignment, or injury, or they may indicate the need for further history.



TOOLBOX

The tools necessary for this assessment include gloves, a flexible transparent ruler, a marker, a penlight, a glass slide, and a magnifier. Good overall lighting is important. The examination area must be adequately screened to provide privacy, and the person must be adequately covered or draped with only the area being immediately examined exposed at any particular time. Also make sure the room temperature is comfortable. Examination in an overly warm or cold environment may alter the skin's condition. For example, if the room is cold, it can change the surface texture of the skin or lead to peripheral cyanosis; if the room is too warm, the skin may become flushed.

Performing a General Survey

A general survey of the integumentary system is typically done while obtaining the health history. Be attentive to any signs that suggest alterations in the integrity of the skin, hair, and nails. During the early observation phase, note general skin coloring, as well as any obvious variations by region. Coloring is a highly variable feature, even among people of the same race or ethnic background. Variations can indicate sun or chemical exposure, emotional responses, illness, or just a personal characteristic. Also observe the general distribution of the hair, including its condition, color, and sheen. Note the general condition of the nails. The examination also relies on your

sense of smell and touch. Be attentive for any unusual odors. As you shake the patient's hand or take vital signs, check the temperature and moisture of the skin.

In addition, consider the patient's overall skin condition in comparison with her or his stated age. If the skin appears chronologically older, it may have been aged by chronic illness, substance use, or environmental exposures. Exposure to sunlight increases the risk of malignancy; exposure to chemicals increases the risk for various forms of dermatitis.

Also do the following:

- Determine the patient's overall nutritional status. Signs of nutritional deficiencies are often evident on inspection of the skin, hair, or nails.
- Note the patient's apparent emotional status. This provides insight into his or her ability to cope with any real or perceived disfigurement associated with alterations in the integument. It may also suggest the likelihood that lesions might be self-inflicted.
- Be aware of the patient's overall body habitus—weight distribution, posture, and muscle mass. Besides providing information on nutritional status, this suggests other health problems that influence the skin. For instance, a person who has truncal obesity and a tripod posture (sitting leaning forward on elbows) may have chronic lung disease requiring systemic corticosteroid treatment. Lung disease may be reflected by nail clubbing and cyanosis; corticosteroid therapy may cause the skin to thin and become friable.
- Obtain the patient's vital signs. Elevations in temperature may indicate an infection that might be accompanied by a rash or other skin lesion.

Performing a Head-to-Toe Physical Assessment

Next, do a head-to-toe survey, checking for more specific signs of diseases affecting other organ systems that might alter the skin, hair, or nails.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

General Health Survey

INSPECT

- Awake, alert, and oriented.
- Memory intact. Able to provide adequate history.

HEENT

Head

INSPECT/PALPATE

- Facial movements symmetrical

Neck

- No neck vein distention
- No unexplained hypertrophy of neck accessory muscles
- No palpable or visible cervical nodes
- Thyroid not palpable

ABNORMAL FINDINGS/RATIONALE

- *Confusion and irritability:* May indicate hypoxia due to cyanosis.
- *Inability to recall information/provide history:* May indicate neurological disorders that cause impaired judgment and altered sensation, placing skin at risk for injury.
- *Lethargy/somnolence:* May indicate a variety of disorders including disease of thyroid, liver, kidney, or cardiovascular or neurological systems, which may be reflected in skin.
- Facial expression and movements often reflect neurological or psychological diseases. For example, pruritus may be associated with depression and neurotic excoriations. Depression may also show in the client's facial expression.
- *Neck vein distension:* Altered cardiac function that could be reflected with cyanosis of skin.
- *Hypertrophy of neck muscles:* Normal in a person who lifts weights. Otherwise, suggests chronic pulmonary disease, with potential for cyanosis or ruddy skin.
- *Palpable/visible nodes:* May indicate infectious process or malignancy that could cause skin rashes, pruritus.
- *Palpable thyroid:* Hyper- or hypothyroidism, both of which cause integumentary changes.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p>Eyes</p> <ul style="list-style-type: none"> ■ Clear and bright; no lesions or edema 	<ul style="list-style-type: none"> ■ <i>Sclera icterus (yellow sclera)</i>: Jaundice. ■ <i>Exophthalmos</i>: Hyperthyroidism with associated skin changes.
<p>Nose/Throat</p> <ul style="list-style-type: none"> ■ Nares patent, nasal mucosa pink, no nasal drainage, oral mucous membranes pink 	<ul style="list-style-type: none"> ■ <i>Red, swollen mucous membranes</i>: Allergy or infection. Infections such as streptococcal infection or mononucleosis can cause a macular rash.
<p>Respiratory <i>INSPECT/PALPATE/PERCUSS/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Respiratory rate within normal limits. Respiratory movements symmetrical and unlabored. ■ Breath sounds clear without adventitious sounds. ■ Percussion tones resonant and palpable. Fremitus within normal range. 	<ul style="list-style-type: none"> ■ <i>Tachypnea</i>: Respiratory distress. ■ <i>Asymmetrical or labored respiratory movements</i>: Pneumothorax or chronic restrictive lung diseases. ■ <i>Crackles, wheezes, rhonchi, or bronchial breath sounds</i>: Altered pulmonary function. ■ <i>Dull percussion tones/increased fremitus</i>: Consolidation within pulmonary tissue. ■ <i>Hyperresonant percussion tones and decreased/absent fremitus</i>: Emphysemic changes. ■ All of these respiratory alterations have the potential to impair oxygenation and cause cyanosis. Chronic cardiopulmonary disease with hypoxia can result in clubbing.
<p>Cardiovascular <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ Extremities pink, warm, and dry, +2 pulses, no edema. 	<ul style="list-style-type: none"> ■ <i>Thin, shiny, hairless, cool skin with decreased pulses</i>: Arterial insufficiency. ■ <i>Peripheral edema, leathery skin with cyanosis and brownish discoloration</i>: Venous insufficiency.
<p><i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Radial and apical pulses within normal limits. S₁ and S₂ present and regular. No extra or adventitious heart sounds. 	<ul style="list-style-type: none"> ■ <i>Tachycardia</i>: Respiratory with impaired oxygenation, causing cyanosis. ■ <i>Irregular heart rate/extra or adventitious heart sounds</i>: Cardiovascular disease impairs cardiac output and/or peripheral oxygenation, causing altered skin appearance and integrity, such as diaphoresis and cyanosis.
<p>Gastrointestinal <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ Abdomen round, soft, nondistended, nontender, no organomegaly, positive bowel sounds. 	<ul style="list-style-type: none"> ■ <i>Hepatomegaly</i>: May cause jaundice and pruritus.
<p>Reproductive <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ External genitalia pink, intact, no lesions. 	<ul style="list-style-type: none"> ■ <i>Genital warts (acuminatum condyloma)</i>: human papillomavirus. ■ <i>Burning, itching, vesicles</i>: Herpes simplex II.
<p>Musculoskeletal/Neurological</p> <ul style="list-style-type: none"> ■ Grade 5 muscle strength, no atrophy, sensations intact, +2 deep tendon reflexes (DTR). 	<ul style="list-style-type: none"> ■ <i>Decreased sensation/DTR</i>: Associated with neuropathy, which increases risk for injury to skin.

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> Changes in energy level Weight changes Fevers	General	<i>Inspect for:</i> Signs of distress <i>Measure:</i> Vital signs Height and weight
HEENT <i>Head and Neck</i>	<i>Ask about:</i> Lumps or swelling in neck Difficulty swallowing History of endocrine problems	HEENT <i>Head and Neck</i>	<i>Inspect for:</i> Facial expression Neck vein distention Enlarged accessory muscles <i>Palpate for:</i> Lymph node enlargement Thyroid gland enlargement
<i>Eyes</i>	<i>Ask about:</i> Watery eyes and allergies Changes in eye color	<i>Eyes</i>	<i>Inspect for:</i> Red eyes Icteric eyes
<i>Ears, Nose and Throat</i>	<i>Ask about:</i> Ear, throat or sinus infections Sore throats Nasal discharge	<i>Ears, Nose and Throat</i>	<i>Inspect for:</i> Red, swollen nasal and oral mucous membranes
Respiratory	<i>Ask about:</i> Cough, breathing difficulty History of respiratory disease	Respiratory	<i>Inspect for:</i> Signs of hypoxia Asymmetrical chest movement <i>Auscultate for:</i> Abnormal/adventitious breath sounds
Cardiovascular	<i>Ask about:</i> History of cardiovascular disease Leg pain	Cardiovascular	<i>Inspect for:</i> Signs of impaired circulation Skin changes in extremities <i>Palpate for:</i> Pedal pulses Edema <i>Auscultate for:</i> Irregular rhythms, rates and extra sounds
Gastrointestinal	<i>Ask about:</i> History of liver disease Nausea/vomiting, loss of appetite Change in stool to clay color	Gastrointestinal	<i>Inspect for:</i> Ascites <i>Palpate for:</i> Liver enlargement, tenderness <i>Percuss for:</i> Liver size
Genitourinary/Reproductive	<i>Ask about:</i> Changes in urine color Urinary tract infections Incontinence History of STDs Safe-sex practices	Genitourinary/Reproductive	<i>Inspect for:</i> Lesions on external genitalia skin
Musculoskeletal	<i>Ask about:</i> History of joint disease, rheumatoid arthritis	Musculoskeletal	<i>Inspect for:</i> Joint deformity Decreased ROM Skin changes over joints
Neurological	<i>Ask about:</i> Loss of sensation	Neurological	<i>Test for:</i> Sensory perception changes, both superficial and deep sensations Deep tendon reflexes
Endocrine	<i>Ask about:</i> History of thyroid disease, diabetes	Immune/Hematologic	<i>Inspect for:</i> Ecchymoses or petechiae
Immune/Hematologic	<i>Ask about:</i> Immune disorders Use of immunosuppressive drugs Bleeding Use of anticoagulants or aspirin	Immune/Hematologic	

Performing an Integumentary Physical Assessment

Once the general survey and head-to-toe assessment are completed, begin the focused examination of the skin, hair, and nails, using inspection and palpation. For purposes of simplicity, inspection and palpation are discussed separately below. However, rather than inspecting all areas of skin, hair, and nails, and then palpating all areas and suspicious lesions, you are more likely to inspect and palpate specific areas almost simultaneously. As you read the following information, keep in mind that areas that vary from normal should be explored using palpation.

Assessing the Skin

Use inspection and palpation to examine the skin.

Inspection

Examine the patient's skin, noting color, odor, and the presence of lesions. Once you have determined the patient's overall skin coloring, take a moment to decide if the coloring suggests something other than a normal variation.

Assessing Color

In addition to alterations in general coloring, it is normal for various *regions* of a person's skin to differ in color, depending primarily on the amount of exposure to light. These variations are generally symmetrical. If you notice that one area—for instance, the shoulders or arms—is darker than other areas—such as the anterior chest or buttocks—make sure that the difference is symmetrical or explained. For instance, a long-distance truck driver's left arm might be darker than the right because that arm receives greater sun exposure during daylight driving hours. (See *Skin Color Variations*.)

INSPECTION OF THE SKIN

1. Color

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect both exposed/unexposed areas for color.
- Differentiate central cyanosis from peripheral cyanosis by inspecting oral mucosa and conjunctiva.
- Gently pull lower eyelids down to examine conjunctiva.



INSPECTION OF THE SKIN

Color (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Uniform skin color with slightly darker exposed areas. Ethnic/racial differences account for many variations in skin color. Mucous membranes and conjunctiva pink.*



HELPFUL HINT

When assessing for color changes in dark-skinned patients, check oral mucous membranes.

ABNORMAL FINDINGS/RATIONALE

- Color changes may be benign or indicate underlying pathology. (See *Skin Color Variations*.)

2. Odor

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note unusual odors.
- *No unusual odor.*

ABNORMAL FINDINGS/RATIONALE

- *Unusual body odor:* Poor hygiene or underlying disease. If from poor hygiene, may be related to self-care deficit that warrants nursing intervention.
- *Odors from excessive sweating (hyperhidrosis):* Possible thyrotoxicosis.
- *Odors from night sweats:* Possible tuberculosis.
- *Urine odor:* Incontinence problem. Stale urine odor may be associated with uremia.
- *Mousy odor:* Liver disease.

3. Lesions

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Differentiate primary and secondary lesions.
- Identify vascular lesions.
- If vascular lesions present, gently palpate and note blanching and pulsation.
 - Describe ABCD of suspicious lesions. A is for asymmetry, B is for border irregularity, C is for color variations, and D is for diameter > 0.5 cm.
 - *No skin lesions.*
 - *No vascular lesions.*

(continued)

INSPECTION OF THE SKIN

Lesions (continued)

ABNORMAL FINDINGS/RATIONALE

- Classify lesion as primary or secondary.
- Primary lesion is an initial alteration in the skin. Secondary lesion arises from a change in a primary lesion.
- A thorough description of lesions should include:
 - Morphological (clinical) description:
 - Size.
 - Shape.
 - Color.
 - Texture.
 - Surface relationship.
 - Exudate.
 - Tenderness.
 - Configuration.
 - Location and distribution.
- For vascular lesions, also note:
 - Pulsations.
 - Blanching.

Skin Color Variations

COLOR/CAUSE/DESCRIPTION

EXAMPLE

Bronzing/Tanning

- *Addison's disease/adrenal insufficiency*: Generalized, most evident over exposed areas.
- *Hemochromatosis*: Generalized, may be gray-brown coloring.



Addison's disease

Tan

- *Chloasma*: "Mask of pregnancy" (on face).
- *Lupus*: Butterfly rash on face.
- *Scleroderma*: Generalized tanning/yellowing of skin, associated with loss of elasticity.
- *Ichthyosis*: With coarse scaliness.
- *Sprue*: Tan/brown patches of any area.
- *Tinea versicolor*: Fawn color or yellow patchy.



Chloasma



Tinea versicolor

**COLOR/CAUSE/
DESCRIPTION**
EXAMPLE
Yellow

- *Uremia*: Generalized.
- *Liver disease, such as hepatitis, cirrhosis, liver cancer, gallbladder with obstructive jaundice*: Generalized.
- *Carotemia*: Not found in conjunctiva or sclera.

Dusky Blue

- *Arsenic poisoning*: Paler spots on trunk and extremities.
- *Central cyanosis with hypoxia; peripheral cyanosis from vasoconstriction*: Caused by cold exposure or vascular disease.



Carotemia



Cyanosis


HELPFUL HINTS

- Jaundice from liver disease is seen in the sclera and conjunctiva, whereas pseudojaundice—yellow color variations associated with carotemia—is seen on the skin but not in the eyes.
- When differentiating peripheral cyanosis (caused by vasoconstriction or decreased circulation) from central cyanosis (caused by hypoxia), check the oral mucous membranes and conjunctiva. Cyanotic mucous membranes and conjunctiva indicate a central process.

Pallor

- *Anemia*: Also on conjunctiva and mucous membranes.
- *Vitiligo*: Patchy.
- *Albinism*: Generalized.

Red

- *Polycythemia*.
- *Erythema*: Dilated superficial capillaries, such as rosacea.

Cherry

- Carbon monoxide poisoning.



Jaundice



Vitiligo



Albinism



Erythema

Assessing Lesions

The skin should be a continuous tissue, and so note breaks, erosions, or lesions. Document localized and/or pigmented variations, including moles, freckles, or vascular lesions, and examine them closely.

Use a flashlight or penlight and a magnifier to determine the surface, pigmentary, or border characteristics of many lesions, particularly when they are small. Besides providing brighter light to a specific skin area, the penlight or flashlight can also be used to shed tangential or oblique light to a lesion. **Tangential lighting** will cause the distal edge of the lesion to cast a shadow if the lesion is raised. Another use of the light is to transilluminate a lesion. For very small lesions, you will need either a small-beam penlight or a transilluminator attachment for an otoscope or ophthalmoscope. Transillumination of a raised lesion helps determine whether the lesion is solid or fluid filled. Fluid-filled lesions have a yellow or pink glow, whereas solid lesions do not.

Use a transparent ruler with centimeter markings to measure any lesions you detect. Clean the ruler after each use, using the method recommended by your facility. Either the transparent ruler or a glass slide can be used as a diascopy to determine whether or not a vascular lesion blanches. Press the ruler or slide gently against the lesion, noting whether it blanches or pales with the pressure. Vascular lesions are red to purple in color. They may be caused by an **extravasation** of blood into the skin tissue or by visible superficial vascular irregularities.

Always be attentive to the signs of malignant melanoma when assessing a skin lesion. The warning signs are easily recalled using the mnemonic “ABCD.” Any time a patient indicates that a pigmented area has newly developed or changed significantly from its original appearance, you must be alert to the potential of malignancy. In addition to malignant melanomas, several other types of skin malignancies are less aggressive and less likely to be fatal (Figs. 10.4 and 10.5).

If you detect a lesion, inspect it closely and palpate it to determine its characteristics. Decide whether it represents a primary or secondary skin lesion. A **primary lesion** is one that appears in response to some change in the internal or external environment of the skin and is not altered by trauma. Primary lesions are categorized by whether or not they are raised and by their overall dimensions. Different sources use different dimensions (0.5 or 1.0 cm) to determine the “cutoff” at which a lesion is given one label or another. This text uses 1.0 cm as the dimension at which lesions are differentiated.

Secondary lesions result from changes in primary lesions. They either add to or take away from an existing primary lesion.

Pressure Ulcers

Pressure ulcers are a type of secondary lesion caused by unrelieved pressure. Healthy People 2010 has established a goal of reducing pressure ulcers in nursing homes by



FIGURE 10.4. Moles (nevi). Moles are generally uniformly tan or brown, round or oval in shape, and have well-defined borders. Moles tend to look very similar to one another. They usually start out flat and, with time, become raised. They should be less than 0.5 cm in diameter. A person may normally have 10 to 40 scattered moles, which generally appear above the waist. When a mole changes in appearance, it needs evaluation, including biopsy, to rule out malignant melanoma.

50 percent to 8 per 1000 residents from 16 per 1000 residents reported in 1997. Assessment begins with identifying those at risk for pressure ulcer development and developing a plan to prevent pressure ulcer formation. If a pressure ulcer develops, assessment focuses on staging pressure ulcers and developing and evaluating pressure ulcer treatment plans.

Lesions may also be categorized according to their pattern, configuration, and distribution.



HELPFUL HINT

Pressure ulcers often develop over bony prominences, such as the sacrum and heels, so inspect these areas carefully.



FIGURE 10.5. Warning signs of malignant melanoma. When assessing for malignant melanoma, think of the acronym *ABCD*. A is for asymmetry, B is for border irregularity, C is for color variations, and D is for diameter > 0.5 cm.

(text continues on page 240)

Clinical Description of Lesions

CHARACTERISTIC	SIGNIFICANCE
Size	<ul style="list-style-type: none"> Major determinant of correct category for primary lesions. Pigmented lesions are typically <0.5 cm. If larger, consider potential for malignancy. Depth of pressure ulcers is major determinant of assigned grade (see Staging Criteria for Pressure Ulcers, page 239).
Shape	<ul style="list-style-type: none"> Macules, wheals, and vesicles are circumscribed. Fissures are linear. Irregular borders are associated with melanoma.
Color	<ul style="list-style-type: none"> Varies widely, and many changes are diagnostic of specific skin diseases. Variegated-colored lesions may signal melanoma. Pustules are usually yellow-white. New scars are red and raised; old scars, white or silver. Petechiae are red. Purpura are red to purplish. Vitiligo is white.
Texture	<ul style="list-style-type: none"> Macules are smooth. Warts are rough. Psoriasis is scaly.
Surface Relationship	<ul style="list-style-type: none"> Surface characteristics help differentiate potential causes of a change and between various primary and secondary lesions: <ul style="list-style-type: none"> <i>Flat (nonpalpable)</i>: Macules, patches, purpura, ecchymoses, spider angioma, venous spider. <i>Raised (palpable) solid</i>: Papules, plaques, nodules, tumors, wheals, scale, crust. <i>Raised (palpable) cystic</i>: Vesicles, pustules, bullae, cysts. <i>Depressed</i>: Atrophy, erosion, ulcer, fissures. <i>Pedunculated</i>: Skin tags, cutaneous horns.
Exudate	<ul style="list-style-type: none"> <i>Clear or pale, straw-yellow exudate</i>: Serous oozing/weeping from noninfected lesion. <i>Thicker, purulent discharge</i>: Infected lesion. <i>Clear serous exudates</i>: Vesicles, as seen with herpes simplex; or bullae, larger than vesicles, as seen with second-degree burns. <i>Yellow pus exudates</i>: Pustules, as seen with impetigo or acne.
Tenderness or Pain	<ul style="list-style-type: none"> Tenderness or pain associated with a lesion depends on the underlying cause. May be associated with bullae from a burn or ecchymoses (bruise).

Vascular Lesions

LESION/CAUSES/DESCRIPTION

EXAMPLE

Ecchymosis


- Extravasation of blood into skin layer.
- Caused by trauma/injury.
- Does not blanch.

Petechiae or Purpura

- Extravasations of blood into skin.
- Caused by steroids, vasculitis, systemic diseases.
- Does not blanch.



Ecchymosis



Petechiae

(continued)

Vascular Lesions (continued)

LESION/CAUSES/ DESCRIPTION

EXAMPLE

Venous Star

- Blue color.
- Irregular-shaped, linear, spider.
- Does not blanch.
- Caused by increased pressure on superficial veins.



Venous star



Spider angioma

Telangiectasia

- Red color.
- Very fine and irregular vessels.
- Blanches.
- Seen with dilation of capillaries.



Telangiectasia

Spider Angioma

- Red color, type of telangiectasia.
- Looks like a spider, with central body and fine radiating legs.
- Blanches; seen in liver disease, vitamin B deficiencies, idiopathic origin.

Capillary Hemangioma

- Red color.
- Irregular-shaped macular patch.



Capillary hemangioma

Port-Wine Stain

- Red color.
- Does not blanch.
- Seen with dilation of dermal capillaries.



Port-wine stain

Primary Lesions

SURFACE CHARACTERISTICS


LESION

EXAMPLE

Flat, Nonpalpable

Examples:


- Cherry angioma
- Freckle
- Flat mole
- Lentiginos
- Melanoma
- Petechiae
- Scarlet fever




Macule

- Macule: <1 cm
- Patch: >1 cm

- Birthmark
- Café-au-lait spot
- Chloasma
- Mongolian spot
- Port-wine stain
- Tinea versicolor
- Vitiligo



Cherry angioma




Vitiligo

Palpable, Raised, but Superficial

Examples:

- Basal cell carcinoma
- Kaposi's sarcoma
- Lichen planus
- Psoriasis
- Raised mole
- Seborrheic keratosis



Papule

- Papule: <1 cm



Mole



Seborrheic keratosis, psoriasis

(continued)

Primary Lesions (continued)

SURFACE CHARACTERISTICS


LESION

EXAMPLE

Raised, Superficial, Temporary

Examples:

- Allergic reaction
- Hives (urticaria)
- Insect bite



Wheal/Hive




Transient lesion (hive)

Palpable, Solid With Depth Into Dermis

Examples:

- Bartholin's cyst
- Erythema nodosum
- Keratogenous cyst
- Lipoma
- Xanthoma



Cyst

- Tumor: >2 cm



Keratogenous cyst




Neoplasm (lipoma)

Palpable, Fluid Filled

Examples:


- Blister
- Contact dermatitis
- Herpes simplex



- Vesicle (serous): <1 cm



Herpes simplex


SURFACE CHARACTERISTICS	LESION	EXAMPLE
<p>Examples:</p> <ul style="list-style-type: none"> ■ Blister ■ Burn ■ Contact dermatitis 	<ul style="list-style-type: none"> ■ Bulla (serous): >1 cm 	
<p>Examples:</p> <ul style="list-style-type: none"> ■ Acne vulgaris ■ Folliculitis ■ Impetigo 		

Blisters

Pustule (pus filled)



Acne vulgaris

Secondary Lesions

SURFACE CHARACTERISTICS	LESION	EXAMPLE
<p>Thickening and Scaling With Increased Skin Markings</p> <p>Examples:</p> <ul style="list-style-type: none"> ■ Contact dermatitis ■ Eczema ■ Lipoma ■ Psoriasis 	<p>Secondary lesions that add to:</p>	
	<p>Lichenification</p>	<p>Contact dermatitis</p>

(continued)

Secondary Lesions (continued)

SURFACE CHARACTERISTICS	LESION	EXAMPLE
<p>Shedding, Dead Skin Cells; Scales Can Be Either Dry or Oily, Adherent or Loose, Variable in Color</p> <p><i>Examples:</i></p> <ul style="list-style-type: none"> ■ Psoriasis 	 <p>A 3D cross-section diagram of skin showing a layer of dead, flaking skin cells (scales) on the surface. The underlying skin layers are shown in shades of pink and yellow.</p>	 <p>A photograph showing the knees of a person with psoriasis. There are large, raised, silvery-white scales on the skin, particularly on the patellar region.</p>
<p>Dried Exudates</p> <p><i>Examples:</i></p> <ul style="list-style-type: none"> ■ Dried herpes simplex ■ Impetigo 	 <p>A 3D cross-section diagram of skin showing a thick, dark red, crusted area on the surface. The underlying skin layers are shown in shades of pink and yellow.</p>	 <p>A close-up photograph of a person's mouth showing dried herpes simplex lesions. The lips and surrounding skin are covered in red, crusted exudates.</p>
<p>Replacement Connective Tissue Formations</p> <p><i>Examples:</i></p> <ul style="list-style-type: none"> ■ Surgical site ■ Trauma site 	 <p>A 3D cross-section diagram of skin showing a raised, fibrous scar on the surface. The underlying skin layers are shown in shades of pink and yellow.</p>	 <p>A photograph showing the knee of a person with a surgical site. The skin is raised and fibrous, characteristic of a scar.</p>

SURFACE CHARACTERISTICS

LESION

EXAMPLE

Hypertrophic Scarring
 Because of Excess Collagen Formation;
 Raised and Irregular
Examples:

- Ear piercing site
- Keloids
- Surgical site
- Tattoo



Keloid




Keloids

Abrasion or Other Loss That Does Not Extend Beyond the Superficial Epidermis
Examples:

- Atopic dermatitis
- Excoriations (scratch marks)
- Insect bite
- Scabies
- Stasis dermatitis
- Vascular rupture site

Secondary lesions that take away from:




Excoriation



Excoriation from uremic pruritus

Loss of Superficial Epidermis
Examples:

- Abrasion
- Candidiasis erosion
- Dermatophyte infection
- Fragile skin
- Impetigo
- Intertrigo






Erosion



Candidiasis

(continued)

Secondary Lesions (continued)

SURFACE CHARACTERISTICS	LESION	EXAMPLE
<p>Linear Breaks in the Skin With Well-Defined Borders, May Extend to the Dermis</p> <p><i>Examples:</i></p> <ul style="list-style-type: none"> ■ Athlete's foot ■ Cheilitis ■ Hand dermatitis (chapped hands) ■ Syphilis 		 <p data-bbox="983 636 1054 657">Cheilitis</p>
<p>Irregularly Shaped Loss Extending to or Through the Dermis; May Be Necrotic</p> <p><i>Examples:</i></p> <ul style="list-style-type: none"> ■ Pressure ulcer ■ Stasis ulcer 		 <p data-bbox="983 1010 1078 1031">Stasis ulcer</p>
<p>Thinning of Skin With Transparent Appearance and Loss of Markings</p> <p><i>Examples:</i></p> <ul style="list-style-type: none"> ■ Aging ■ Arterial insufficiency ■ Topical corticosteroids 		 <p data-bbox="983 1409 1031 1430">Aging</p>



ALERT

Extensive undermining often occurs, extending through the dermal layer to the bone. The visible pressure ulcer may be only the tip of the iceberg.

Risk Factors for Pressure Ulcers

- Impaired mental status
- Impaired nutritional status
- Sensory deficits
- Immobility
- Mechanical forces
- Shearing and friction
- Increased by temperature
- Excessive exposure to moisture from bodily secretions, such as urinary and fecal incontinence

Staging Criteria for Pressure Ulcers




STAGE	APPEARANCE	CHARACTERISTICS
I		Nonblanchable erythema of intact skin; indicates potential for ulceration.
II		Partial-thickness loss involving both epidermis and dermis. Ulcer is still superficial and appears as a blister, abrasion or very shallow crater.
III		Full-thickness loss involving subcutaneous tissue. Ulcer may extend to but not through fascia. A deep crater that may undermine adjacent tissues.
IV		Full-thickness loss with extensive involvement of muscle, bone, or supporting structures. This deep ulcer may involve undermining and sinus tracts of adjacent tissues.
		Ulcers that are covered with eschar cannot be staged without débridement.




Palpation

After inspecting the skin, explore any findings through palpation. Palpation is used to determine the skin's temperature, moisture, texture, and turgor. It can also help to determine whether a localized lesion is raised, indented, or pedunculated and its surface characteristics.


As you palpate for temperature, you will find that the dorsal part of your hands and your fingers are most sensitive to temperature variations. Remember to wear gloves when palpating any potentially open areas of the skin.

Pattern and Configuration of Lesions


PATTERN	DESCRIPTION	EXAMPLE
ROUND/OVAL 	<ul style="list-style-type: none"> ■ Coin or oval-shaped, as in nummular eczema. 	
DISCRETE 	<ul style="list-style-type: none"> ■ Lesions that remain separate and apart are common in many skin disorders. Moles (nevi) are an example. 	
GROUPED 	<ul style="list-style-type: none"> ■ Lesions that are grouped, or clustered, such as herpes simplex. 	
CONFLUENT 	<ul style="list-style-type: none"> ■ Lesions that run together or are confluent are common in childhood diseases such as rubella. 	

PATTERN	DESCRIPTION	EXAMPLE
<p>LINEAR</p> 	<ul style="list-style-type: none"> Lesions arranged in lines are common in contact dermatitis due to poison ivy or herpes zoster. 	
<p>ANNULAR/CIRCULAR</p> 	<ul style="list-style-type: none"> Ring-shaped lesion may be ringworm. 	
<p>ARCIFORM</p> 	<ul style="list-style-type: none"> Lesions arranged in partial rings, or arcs, occur in syphilis. 	
<p>IRIS</p> 	<ul style="list-style-type: none"> A bull's-eye lesion, or round lesion with central clearing, is typical in erythema multiforme and Lyme disease. 	


(continued)

Pattern and Configuration of Lesions *(continued)***PATTERN****DESCRIPTION****EXAMPLE****RETICULAR**

- Meshlike pattern as in lichen planus.

**GYRATE**

- Lesions have serpentine configuration as in gyrate erythema.

**POLYCYCLIC**

- Coalesced, concentric circles such as urticaria.




Distribution of Skin Lesions

AREA

DESCRIPTION

EXAMPLE


DIFFUSE/GENERALIZED




- Lesions distributed over entire body, as in urticaria from allergic reactions.



SCATTERED






- Lesions that are sparsely distributed, as in seborrheic keratosis.



(continued)

Distribution of Skin Lesions (continued)

AREA	DESCRIPTION	EXAMPLE
<p>LOCALIZED</p> 	<ul style="list-style-type: none"> ■ Lesions in a very limited, discrete area ■ Location may indicate contact with an allergen or a wheal from insect bite. 	
<p>REGIONAL Head</p> 	<ul style="list-style-type: none"> ■ Confined to a specific body area. ■ Tinea capitis. 	
<p>Torso</p> 	<ul style="list-style-type: none"> ■ Pityriasis rosea 	

Distribution of Skin Lesions (continued)


AREA

DESCRIPTION

EXAMPLE


Extensor Surfaces

■ Psoriasis




Flexor Surfaces

■ Intertrigo






Dermatome

■ Herpes zoster



(continued)

Distribution of Skin Lesions (continued)

AREA	DESCRIPTION	EXAMPLE
<p>Hairy Areas</p> 	<ul style="list-style-type: none"> ■ Herpes 2, pediculosis pubis 	
<p>Intertriginous Areas (Folds of Skin)</p> 	<ul style="list-style-type: none"> ■ Contact dermatitis, diaper rash, intertrigo (erythema and scaling of body folds) 	
<p>Sun-Exposed Areas</p> 	<ul style="list-style-type: none"> ■ Actinic keratosis (precancerous skin lesion), and skin cancers, basal, squamous and melanoma 	

The skin's moisture varies among body parts, as well as with changes in the environmental temperature, physical activity, or body temperature. Perspiration is produced to cool the body. In the winter, the skin tends to be drier because of the lower ambient temperature and decreased humidity in the environment.

Turgor is assessed as an indication of elasticity. To determine turgor, pinch a fold of skin over an unexposed area, such as below the clavicle, or on the abdomen or sternum. You may also use the forearm. Do not test turgor on the dorsal hand or other areas where the skin is noticeably loose or thin. As you pinch the skinfold, it

PALPATION OF THE SKIN

1. Temperature

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS



Temperature

- Compare side to side using the dorsal aspect of your hand.
- *Skin warm.*
- *Temperature varies depending on area being assessed; for example, exposed areas may be cooler than unexposed areas.*

ABNORMAL FINDINGS/RATIONALE

- *Local area with increased temperature:* Inflammatory process, infection, or burn, caused by increased circulation to area.
- *Generalized increase in temperature:* Fever.
- *Local area with decreased temperature:* Decreased circulation to area, as with arterial occlusion.
- *Generalized decrease in skin temperature:* Exposure or shock.

2. Moisture

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use light palpation to assess skin moisture.
- *Depends on environmental conditions and patient's age. Elderly people have drier skin because of decreased sweat production.*
- *Exposed areas are usually drier than unexposed areas. Also, moisture varies according to body area; for example, the axillae are usually more moist than other areas.*

ABNORMAL FINDINGS/RATIONALE

- *Increased moisture:* Fever, thyrotoxicosis.
- *Decreased moisture:* Dehydration, myxedema, chronic nephritis.

(continued)

PALPATION OF THE SKIN

3. Texture

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use light palpation to assess texture.
- *Varies from soft and fine to coarse and thick, depending on area assessed and patient's age.*
- *Exposed skin usually not as soft as unexposed.*
- *Extensor surfaces, such as elbows, have coarser skin.*
- *Usually, the younger the patient, the softer the skin, so infants have very soft skin.*

ABNORMAL FINDINGS/RATIONALE

- *Coarse, thick, dry skin:* Hypothyroidism.
- *Skin that becomes more fine-textured:* Hyperthyroidism.
- *Smooth, thin, shiny skin:* Arterial insufficiency.
- *Thick, rough skin:* Venous insufficiency.

4. Turgor

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS



Turgor

- Test turgor by gently pinching a fold of skin on an unexposed area (such as below the clavicle) and note any “tenting.”
- *Elasticity decreases with age.*
- *Exposed areas may have less turgor.*



HELPFUL HINT

Turgor is often used to assess hydration status. But because turgor decreases with age, it is not a useful tool for assessing hydration status in elderly people.

ABNORMAL FINDINGS/RATIONALE

- *Decreased turgor or tenting:* Dehydration or normal aging.
- With scleroderma, the turgor is actually increased and the tension does not allow the skin to be pinched upward. This may also be seen with edema.

should feel resilient, move easily, and return to place quickly when released.

Assessing the Nails

Assess the nails through inspection and palpation.

Inspection

The condition of the nails often provides important clues about the patient's overall health status. Inspect the color

and shape of the nails. The color beneath the nails should be similar to the overall skin coloring, although somewhat rosier. There should be no hemorrhage. Nail texture should be uniform and not brittle. Note any grooves or lines in the nail or nailbed.

Also assess for **clubbing**, or loss of the normal angle (Lovibond's angle) between the nail base and the finger. When no clubbing is present, the nailbed is firm. You can further assess for clubbing by having the patient

INSPECTION OF THE NAILS

1. Color

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS



Color

- Inspect nails for color.
- *Normal nails vary from pink in light-skinned patients to light brown in darker-skinned patients.*

ABNORMAL FINDINGS/RATIONALE

- Color changes in nails may indicate a local or systemic problem.
- *Very distal band of reddish-pink or brown covering <20% of nail (Terry's nails):* Cirrhosis, disorders causing hypoalbuminemia.

- *Yellow nails:* Cigarette smoking, fungal infections, psoriasis.



Fungal infection

- *Distal band of reddish-pink brown covering 20% to 60% of nail (Lindsay's nails or half-and-half nails):* Renal disease, hypoalbuminemia.

- *Blue (cyanotic) nails with clubbing:* Peripheral disease or hypoxia.

- *White nails (leukonychia):* Trauma; cardiovascular, liver, or renal disease.

- *Black nails:* Trauma.



Half-and-half nails



Blue nails



Leukonychia


(continued)

INSPECTION OF THE NAILS

2. Shape

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect nail shape.
- Have patient place fingers together and note space (opening) between nails.
- *Angle of nail attachment 160 degrees; nails convex*
- *Clubbing is present if nails meet and angle of attachment is 180 degrees or greater.*



ABNORMAL FINDINGS/RATIONALE

- *Splinter hemorrhages*: Bacterial endocarditis or trauma.
- *Angle of nail attachment 180 degrees or more*: Clubbing associated with diseases that affect level of oxygenation, such as congenital heart disorders, cystic fibrosis, and chronic pulmonary diseases.
- *Spooning or concave nail (koilonychia)*: Severe iron deficiency anemia, hemochromatosis, thyroid and circulatory diseases, in response to some skin diseases and local trauma.
- *Onycholysis, separation of the nail from nailbed*: Fungal infections, psoriasis, thyrotoxicosis, eczema, systemic diseases, following trauma, or as allergic response to nail products/contactants



Splinter hemorrhages

- *Red and inflamed perionychium (paronychia)*: Infection or ingrown nail tuberculosis.

- *Pitting*: Psoriasis.



Onycholysis



Psoriasis




Paronychia

PALPATION OF THE NAILS

1. Texture

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use light palpation to assess texture.



- *Firm*
- *Longitudinal ridges usually benign*



ABNORMAL FINDINGS/RATIONALE

- *Soft, boggy nails*: Clubbing caused by poor oxygenation.
- *Brittle nails*: Hyperthyroidism, malnutrition, calcium and iron deficiency, repeated use of harsh nail contactants or products.
- *Pitting*: Psoriasis, eczema, alopecia areata.



- *Beau's lines (transverse ridges)*: Serious illness that causes nail growth to slow or halt.



(continued)

PALPATION OF THE NAILS

2. Capillary Refill

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Gently press on nail and note blanching, then release and note speed of refill (color return).



Pressing nail



Releasing pressure



HELPFUL HINT

Positive capillary refill may be affected by cold temperatures.

ABNORMAL FINDINGS/RATIONALE

- *Poor capillary refill:* Cardiopulmonary problems or anemia.

place the dorsal aspect of two opposite distal fingers together, so that the nails rest against one another. In the absence of clubbing, you should be able to detect a window of light caused by the space created by Lovibond's angle.

Palpation

Palpate the nail for texture and refill. Nail texture should be uniform and not brittle. Note any grooves or lines or pitting in the nail or nailbed. To check for capillary refill, press on the tip of the nail. It should blanch, and upon release the color should return within 3 seconds.

Assessing the Hair

Assessing the hair is done by inspection and palpation.

Inspection

Inspect the hair for distribution, color, and condition of the scalp. Note any increased hair growth or areas of thinning or **alopecia**. Also, assess the body for normal distribution of hair. The color of the hair can be very difficult to assess, primarily because so many people color their hair. Inspect the scalp as you would any area of skin, assessing any lesions for size, relationship to the overall scalp plane, color, and surface integrity. A morphological description of individual lesions often provides clues to their cause. Almost any of the common skin disorders can affect the scalp.

Note whether there is any adherent material on the hair. Small 1- to 2-mm white eggs are found with lice or **pediculosis**, which occurs on the hairs of the scalp, beard, axillae, or pubic areas. Although head lice can be

INSPECTION OF THE HAIR

1. Scalp-Hair Distribution

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect hair quantity and distribution. Assess areas for the pattern. Note whether there is actual hair loss, with smooth skin beneath, or whether hair has been broken off near the scalp, with palpable stubble over the skin. True hair loss occurs in many conditions.
- *Gender, age, and genetics affect hair distribution. Hair should be evenly distributed; exceptions are normal balding patterns common to men or persons of advanced age. Hair thins with age.*



Scalp-hair distribution



Male pattern-baldness

ABNORMAL FINDINGS/RATIONALE

- *Generalized hair loss:* Nutritional deficiencies, hypothyroidism, lupus erythematosus, thyroid disease, and in response to disorders or situations that stress the integumentary system, such as serious illnesses or side effects of medications.
- *Patchy alopecia* associated with alopecia areata, trichotillomania, and fungal infections such as tinea capitis.



Alopecia



Tinea capitis



Alopecia areata

(continued)

INSPECTION OF THE HAIR

2. Body Hair

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Fine body hair (vellus) noted over body. Gender, age, and genetics influence amount of body hair.*
- *Men usually have more hair on chest.*
- *Puberty marks the onset of pubic hair growth and increased growth on legs and axillae.*

ABNORMAL FINDINGS/RATIONALE



Hirsutism

- **Hirsutism**, usually caused by endocrine disorders or medications such as steroids, is hair in male patterns in a female; for instance, excess facial or trunk hair.

3. Color

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color.
- *Wide range of normal color variations. Gray coloring occurs with aging.*

ABNORMAL FINDINGS/RATIONALE

- **Localized areas of white or gray hair:** In patients recovering from alopecia areata and in those with vitiligo and piebaldism.
- **Diffuse white hair:** *Albinism.*
- **Green hair:** Copper exposure and pernicious anemia.

4. Condition of Scalp

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect scalp.
- *Scalp intact and free of lesions and pediculosis.*

ABNORMAL FINDINGS/RATIONALE

- **Scaling of scalp:** Dandruff, seborrhea, psoriasis, certain tineas, and eczema (atopic dermatitis).

seen with the naked eye, they are quite small and mobile and their eggs, called **nits**, are easier to see. Nits are deposited near the base of the hair shaft, so that fresh nits are usually found within $\frac{1}{8}$ inch of the scalp or skin. When an infestation of lice has persisted for some time, or if nits were not removed from an earlier infestation, they will be found along a greater portion of the hair

shaft because the hair will have grown during the period. Nits found $\frac{1}{4}$ inch or more from the skin have probably already hatched.

Palpation

Palpate the texture of the hair. If it is unusually coarse or fine, consider a thyroid disorder.

PALPATION OF THE HAIR AND SCALP

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use light palpation to assess hair.



- Use light palpation to assess scalp.
- *Scalp mobile, nontender*



HELPFUL HINT

Genetics influence hair texture.

ABNORMAL FINDINGS/RATIONALE

- *Dry, coarse hair:* Hypothyroidism.
- *Fine, silky hair:* Hyperthyroidism.
- *Tenderness:* May indicate a localized infection.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Green have concluded. The following are your findings:

■ General Health Survey

- Well-developed, 48-year-old, fair-skinned woman.
- Appearance consistent with stated age.
- Awake, alert, and oriented (AAO) \times 3, memory intact.
- No signs of distress, appears comfortable, and provides history with no difficulty.
- Head-to-toe scan reveals no signs of problems in other systems/areas that would affect integumentary system.

■ Vital Signs

- Height, 5'4".
- Weight, 138 pounds.
- Temperature, 97.4°F
- Pulse, 82 and regular.
- Respirations, 14 and unlabored.
- Blood pressure (BP), 134/82, left arm.

■ Skin

- Warm, with good turgor indicated by brisk recoil.
- Generally intact, with smooth texture.
- Coloring symmetrical: pale pink on unexposed areas, moderately tanned on exposed areas with diffusely scattered freckles, consistently light brown in color and all 0.2 to 0.4 cm in diameter.

(continued on page 256)



Case Study Findings (continued)

- Approximately 30 moles, 0.2 to 0.4 cm in diameter, with well-circumscribed, regular borders; all very dark brown and consistent in color.
- Three patches of slightly pink, scaly, flat skin, approximately 0.8 cm each (two right forearm, one left forearm); no telangiectasia or ulcerations of the sites; no changes in surrounding skin.
- **Nails**
 - Short, with smooth edges.
 - Coat of clear, glossy polish present.
 - Nail bed pink, firm to palpation.
 - No hemorrhages or discoloration of nails or surrounding tissues.
 - No clubbing.
- **Hair**
 - Generally brown with scant amount of gray interspersed and distributed evenly over scalp.
 - Shortly cut and well-groomed.
 - Shiny, with no signs of damage.
 - No areas of excess hair growth on body.



Case Study Analysis and Plan

Now that your examination of Mrs. Green is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.



CRITICAL THINKING ACTIVITY #3

List the strengths and areas of concern that you have identified related to Mrs. Green's current health status.

Nursing Diagnoses

Next, consider all of the data you have collected during your assessment of Mrs. Green. Use this information to identify a list of nursing diagnoses. Some possible nursing diagnoses are provided below. Cluster the supporting data.

1. Health-Seeking Behaviors, related to the fear of skin cancer
2. Risk for Impaired Skin Integrity
3. Fear, related to family history of skin cancers

Identify any additional nursing diagnoses.



CRITICAL THINKING ACTIVITY #4

Now that you have identified some nursing diagnoses for Mrs. Green, select one diagnosis and create a nursing care plan and a related teaching plan including learning outcomes and teaching strategies.

RESEARCH TELLS US

The risk of UV light exposure and the associated risk for skin cancer are well documented. In 2000, approximately one million Americans were newly diagnosed with skin cancer, and malignant melanoma was on the rise. Of concern is the increasing use of tanning salons in spite of public educational efforts concerning unprotected sun exposure.

To determine the effects of tanning salon exposure on skin, Whitmore, Morison, Potten, and Chadwick (2001) studied 11 subjects who had full body exposure in tanning salons over 2 weeks. The researchers examined the molecular changes associated with the tanning salon UV exposure.

Pretreatment peripheral blood lymphocytes and epidermal biopsy specimens were compared with post-exposure specimens. The findings revealed that cyclobutane pyrimidine dimers in DNA and p53 protein expression were present in the epidermal keratinocytes, but not in the lymphocytes reflecting molecular changes associated with UV exposure. The same molecular changes are seen with sunlight exposure and are associated with skin cancer.

As nurses, we are in a prime position to educate the public of the dangers of unprotected sun exposure and the risk of skin cancer while at the same time informing them that the same risk applies to tanning salons.

HEALTH CONCERNS

Facts About Melanoma

Skin cancer is the most common type of all cancers. Although melanoma accounts for only 4 percent of skin cancer cases, it has the highest mortality rate among skin cancers—and its incidence is on the rise in the United States. In 2005, the American Cancer Society estimated that there would be nearly 60,000 new cases of melanoma for that year and that 7,770 people would die from the disease.

Risk Factors for Melanoma

- Too much exposure to UV radiation (sunlight, tanning lamps or booths).
- Moles.
- Fair skin, freckling, or red or blond hair.
- Positive family history of melanoma (10 percent of people with melanoma have a positive family history of melanoma).
- History of immunosuppressive treatment.
- Older age.
- Male gender.
- History of xeroderma pigmentosum, a rare inherited skin condition.
- Past history of melanoma.

Preventative Measures

- Avoid excessive UV exposure.
- Avoid being outdoors in the middle of the day, when UV light is most intense.
- Wear a hat and long-sleeve shirt when outdoors.
- Use sunscreen with a sun-protection factor of 15 or greater and lip balm. Apply sunscreen and lip balm about 20 to 30 minutes before going outdoors, and then reapply every 2 hours.
- Wear wraparound sunglasses with 99 percent UV absorption.
- Avoid tanning salons and sun lamps.
- Protect children from the sun because severe, blistering sunburns during childhood and adolescence increase the risk for melanoma.
- Have suspicious moles checked by physician and removed if needed.
- Consider genetic counseling if you have:
 - Positive history of melanoma.
 - Positive family history of melanoma.
 - Had a melanoma at a young age.
 - History of dysplastic nevi.

A mutated gene has been identified in some families that have a high incidence of melanoma. Research is being done to test for this gene.

Source: American Cancer Society. *All About Skin Cancer: Melanoma*. (2005). Available at: www.cancer.org. Last accessed January 23, 2005.

Common Abnormalities**ABNORMALITY****ASSESSMENT FINDINGS****Acne Vulgaris**

- Caused by sebaceous gland overactivity with plugging of hair follicles and retention of sebum, resulting in comedones, papules, and pustules. Onset is typically at puberty, but acne may last into advanced age. Greater incidence in males.
- Pimples present as papules or pustules.
- Cysts may develop and leave extensive scarring.
- Most common on face, back, and shoulders.
- *Bacillus* is cause.
- Lesions may be sore and painful.
- Aggravated by emotional distress, greasy topical applications (cosmetics), and certain medications (oral contraceptives, isoniazide, rifampin, lithium, phenobarbital).



(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Actinic Keratosis</p> <ul style="list-style-type: none"> Causes reddish, irregular, slightly raised lesions that have a rough, gritty surface. Sign of sun-damaged skin. Precancerous lesion, may progress to squamous cell cancer. 	<ul style="list-style-type: none"> Typically less than 1 cm in diameter. Generally on sun-exposed areas of face, head, neck, and hands. 
<p>Basal Cell Carcinoma</p> <ul style="list-style-type: none"> An epidermoid cancer, one of the most common malignant skin diseases, but rarely metastatic. 	<ul style="list-style-type: none"> Typically has pearly, flesh-colored or transparent “rolled” border. Central area develops telangiectasia and may ulcerate. Variations can present with nodular, sclerotic, and/or pigmented appearance. Usually occurs on sun-exposed surfaces, especially the face. 
<p>Contact Dermatitis</p> <ul style="list-style-type: none"> Localized skin irritation, inflammation, and pruritus from contact with an irritating substance. Can occur as an additive effect of multiple irritants (soaps, detergents, or chemicals) or allergy to a specific agent (topical to a specific agent, topical medication, plant oils, or metals). Secondary infections may occur at the site. 	<ul style="list-style-type: none"> Edema may occur, with development of vesicles and bullae. Vesicles or bullae may rupture, causing crusting. Edema may be very significant, particularly when face or genitalia are involved. Person may have history of previous reaction to agent and recent exposure. 
<p>Eczema/Atopic Dermatitis</p> <ul style="list-style-type: none"> Causes redness, pruritus, scratching, and skin lesions in a person with a predisposition to skin irritations 	<ul style="list-style-type: none"> Red to red-brown, slightly scaly lesions. Lichenification with increased skin markings common. Exudative As sites resolve, skin pigmentation is often permanently altered. Common sites include face, neck, upper trunk, wrists, hands, and flexor surfaces (folds) of knees and elbows. Lesions on face, neck, and upper trunk are called <i>monk's cowl</i>. Person also often has asthma or allergic rhinitis; family history is often positive for asthma, rhinitis, eczema, or other allergy problems. Itching can be quite severe. Sites may develop secondary infection. May be triggered by changes in temperature, emotional stress, or food allergies. 

ABNORMALITY

ASSESSMENT FINDINGS

Herpes Simplex

- A common, contagious disease caused by the herpes simplex virus type 1. More prevalent in women than in men.

- Recurrent clusters of small vesicles on erythematous base.
- Sites burn and sting; neuralgia often occurs.
- Typically found on perioral and genital areas.
- May initially follow a minor infection.
- Later recurrences may be triggered by trauma, stress, or sun exposure.
- Often associated with lymphadenopathy of regional nodes.

**Herpes Zoster**

- Also called *shingles*; an acute, infectious disease caused by the varicella zoster virus. Postzoster neuralgia discomfort can last for months. Ocular involvement can lead to blindness.

- Pain along a nerve dermatome is often the first symptom.
- Discomfort followed in 2 to 4 days by erythematous area that develops papules or plaques followed by painful grouped vesicles unilaterally along the dermatome.
- Vesicles or bullae rupture with crusting.
- Most common sites are face and trunk.
- Most common in people over age 60 and those with impaired immunity.

**Intertrigo**

- A superficial dermatitis in the skinfolds. It is caused by heat, moisture, and friction, and is most common in obese people.

- Pink to reddened skin in body folds (between and beneath buttocks, beneath fatty abdominal pad, or beneath pendulous breasts).
- Areas in folds develop erythema, fissures, and denudation.
- Lesions may itch, burn, or sting.



(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Kaposi's Sarcoma</p> <ul style="list-style-type: none"> ■ A type of malignant skin cancer seen most often in people who are HIV positive. 	<ul style="list-style-type: none"> ■ Purple-blue to red papules. ■ Mild scaling of surface that progresses to ulcerate and bleed. ■ Dissemination common, with widespread involvement of skin and mucosa. 
<p>Malignant Melanoma</p> <ul style="list-style-type: none"> ■ An invasive, cancerous skin tumor with strong potential for metastasis to both regional and distant sites and organs. 	<ul style="list-style-type: none"> ■ Commonly presents as a black or purple nodule. ■ Other color variations include pink, tan, brown, red, or even "normal" tones. ■ May also be flat or pedunculated. ■ Erythema or halo of coloring may surround lesion. ■ May ulcerate or become friable. ■ May be found on any location, including sun-exposed areas, palms, or soles. 
<p>Pityriasis Rosea</p> <ul style="list-style-type: none"> ■ A common, mild, acute inflammatory skin disease occurring most often in the spring and fall. 	<ul style="list-style-type: none"> ■ Herald (or mother) patch usually precedes onset of smaller fawn-colored, oval, scaly eruptions 1 to 2 weeks later. ■ Later rash has individual oval lesions with a diagonal orientation. Often described as "Christmas tree" rash because of shape. ■ Rash may last 4 to 8 weeks. ■ Can be intensely pruritic. ■ As lesions exfoliate, they develop a "crinkly" scale and clear centrally so that they may mimic tinea. ■ Believed to be caused by viral infection. 

ABNORMALITY

ASSESSMENT FINDINGS

Psoriasis

- A common dermatitis that has genetic causes and may begin at any age.

- Silvery scales on bright red papules.
- Scales generally thick; area beneath bleeds if scale is removed.
- Usually occurs on extensor surfaces of knees, elbows, and scalp.
- Can occur elsewhere, including between buttocks.
- Nails may develop a stippled, “pitted” appearance and separations.
- Itching may be mild or severe.
- A genetic predisposition is suggested by family history.
- May occur with arthritis.

**Rosacea**

- A chronic disorder of unknown cause that occurs mainly on the face.

- Vascular component with erythema and telangiectasias.
- Acne component with papules, pustules, and seborrhea.
- Glandular component with hyperplasia of soft tissue of nose (rhinophyma).
- Usually rosy hue is diagnostic.
- Onset usually in 40s and 50s.
- May be aggravated by alcohol, caffeine, chocolate, heat, and spicy foods, as well as by situations that promote flushing.

**Seborrhea**

- A disorder of the sebaceous glands that causes an increase in the amount of sebaceous secretion and may also alter its quality.

- Greasy scales may have underlying, flat erythema site.
- Some degree of papules and pustules possible.
- Typical sites include scalp, face (between brows, along sides of nose, at mustache/beard areas) and on preternal, interscapular, and umbilical regions.
- Genetic tendency, with family history possible.
- Itching may be present.
- Fissuring is possible with secondary infections.
- “Super dandruff” is common term because it occurs along sites with greater hair distribution.



(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS	
<p>Seborrheic Keratosis</p> <ul style="list-style-type: none"> ■ A benign skin lesion that may be pigmented. 	<ul style="list-style-type: none"> ■ Sharply demarcated lesions. ■ Brown to black pigmentation. ■ Rough, dry surface. ■ Elevation, with pasted or stuck-on appearance. ■ Surrounding skin generally normal. ■ Incidence increases with age. ■ Generally found on trunk, although potentially can occur anywhere. 	
<p>Squamous Cell Carcinoma</p> <ul style="list-style-type: none"> ■ A form of skin cancer occurring mainly in the squamous cells. 	<ul style="list-style-type: none"> ■ Pink, scaly, elevated lesions ■ Base of lesions may be inflamed. ■ Scablike appearance is common. ■ Typically on sun-exposed surfaces, including scalp, hands, lips, and ears. 	
<p>Stasis Dermatitis</p> <ul style="list-style-type: none"> ■ Eczema of the legs with pigmentation, edema and, at times, chronic inflammation resulting from venous insufficiency. Stasis ulcer is associated with stasis dermatitis and develops from venous insufficiency. 	<ul style="list-style-type: none"> ■ Red, scaly patch often initial sign. ■ Site develops vesicles and crusts. ■ Ulcer may develop as a result of trauma, edema, or infection. ■ When site does not progress to ulceration, reddish-brown hyperpigmentation may develop. ■ Caused by poor circulation, which can be related to peripheral vascular diseases, obesity, or poor nutrition. 	
<p>Tinea Capitis</p> <ul style="list-style-type: none"> ■ A fungal infection of the scalp. 	<ul style="list-style-type: none"> ■ Well-demarcated, reddened area. ■ Scaling, itching. ■ Dry, brittle hair. 	

ABNORMALITY

ASSESSMENT FINDINGS

Tinea Corporis

- Ringworm, a fungal skin disease occurring anywhere on the body.

Tinea Cruris

- *Jock itch*, a fungal skin disease occurring in the genital and anal areas in males.

Tinea Pedis

- *Athlete's foot*, a fungal skin disease occurring in the foot. *Tinea manum* occurs on the palms.

Vitiligo

- Characterized by white patches of skin surrounded by areas of normal pigmentation. Progresses slowly and is more common in dark-skinned people.

- Ring-shaped erythematous lesions on body.
- Central clearing.
- Advancing border with small vesicles.
- Pruritic.
- Most often on exposed surfaces.

- Sharply demarcated, reddened areas.
- Central clearing.
- Severe pruritus.
- Intertriginous area in groin.
- When it occurs on scalp, proper term is *tinea capitis*.

- Exfoliating, fissuring, macerated area of erythema.
- Sites itch, burn, and/or sting.
- Tinea manum occurs in interdigital folds of fingers or on palms.
- Tinea pedis occurs in interdigital folds between toes or on soles of feet.

- Irregular areas of depigmentation.
- May have hyperpigmented border.
- Flat, nonraised, with smooth surface.
- Most common sites are face, hands, and feet.
- Probably autoimmune cause; also associated with various endocrine disorders.



ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the integumentary system.

Ask the patient the following:

- Do you have...
 - Changes in moles or other lesions?
 - Nonhealing sore or chronic ulceration?
 - Pruritus/itching?
 - Rashes?
 - Changes in skin, hair, or nails?
- Do you have any food, drug, or environmental allergies?
- Do you have any medical problems?
- Are you on any medications, prescribed or over-the-counter?

PHYSICAL ASSESSMENT**Equipment**

Gather equipment needed for the exam:

- Gloves
- Penlight
- Ruler
- Marker
- Magnifier

Position

Assess all areas, change position as needed.

Technique

First, scan your patient checking for specific signs of diseases affecting other organ systems that might alter the skin, hair, or nails.

ASSESSMENT CHECKLIST (continued)

Inspection/Palpation

Skin

Inspect

- Color, odor, integrity
- Lesions, if any (if found, describe morphology, distribution, pattern, and location).

Palpate

- Temperature.
- Turgor.
- Texture.
- Moisture.
- If lesions found on inspection, palpate them for texture, tenderness, pulsations, blanching.

Hair

Inspect

- Color, quantity, distribution, condition of scalp, lesions, pediculosis.

Palpate

- Texture.
- Scalp for tenderness, mobility.

Nails

Inspect

- Color, condition, shape, angle of attachment.

Palpate

- Texture, capillary refill.

DOCUMENT

Document your findings.


S U M M A R Y

- The integumentary system provides invaluable information about your patient's overall health status. Therefore it is important that you learn to objectively assess the skin, hair, and nails and be aware of the wide range of normal variations, which further differ according to age, race, and ethnic background.
- During your assessment, use a consistent approach and make careful observations about the overall integrity of the tissues, as well as any specific areas of abnormality.

R E V I E W Q U E S T I O N S

1. You are assessing an 80-year-old male patient admitted to your unit with the diagnosis of congestive heart failure (CHF). When assessing his skin, which finding would be considered normal, taking his age into account?
 - a. Increased skin temperature
 - b. Increased moisture
 - c. Decreased erythema
 - d. Decreased turgor
2. Which area of the skin is best for assessing skin turgor?
 - a. Hand
 - b. Face
 - c. Neck
 - d. Below the clavicle
3. Which part of your hand is best for assessing skin temperature changes?
 - a. Dorsal aspect
 - b. Fingerpads
 - c. Palmar surface
 - d. Ulnar surface
4. A nursing assistant reports that a patient is "blue." What area is best for assessing central cyanosis?
 - a. Nails
 - b. Lips
 - c. Ear lobes
 - d. Oral mucosa
5. What area is best for detecting color changes in patients with dark skin?
 - a. Nails
 - b. Oral mucosa
 - c. Palms of hands
 - d. Lips
6. Mr. Heenan, age 81, has a pressure ulcer that is covered with eschar. How would you stage this pressure ulcer?
 - a. Stage 2
 - b. Stage 3
 - c. Stage 4
 - d. Can't be staged

7. When assessing the nails, what is the normal angle of attachment?
- 45 degrees
 - 90 degrees
 - 160 degrees
 - 180 degrees
8. Mrs. Green, age 75 years, has a stroke with left side weakness. As you reposition her, you notice a red-denied sacral area about 3 cm in size. How would you stage this pressure ulcer?
- Stage 1
 - Stage 2
 - Stage 3
 - Stage 4
9. You are working as a school nurse and are assessing the children for pediculosis. You know that the nits have probably hatched if the nits are located:
- On the scalp
 - At the base of the hair shaft
 - $\frac{1}{8}$ inch from the scalp on the hair shaft
 - 1 inch from the scalp on the hair shaft
10. You are working in a pre-natal clinic. As you assess your patients, which skin changes would you normally expect to find? Circle all that apply.
- Striae
 - Linea alba
 - Chloasma
 - Palmar erythema



Assessing the Head, Face, and Neck

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent head, face, and neck history questions
- Obtain a history specific for head, face, and neck
- Perform a physical assessment of the head, face, and neck
- Document head, face, and neck assessment findings
- Identify actual or potential health problems stated as a nursing diagnosis
- Differentiate between normal and abnormal head, face, and neck assessment findings

The head, face, and neck form a large portion of what is often referred to as the *head, eyes, ears, nose, and throat (HEENT) system*. This is actually a complex set of varied organs, combined during assessment because of their proximity to one another and the integration among the components of the system. The HEENT encompasses almost all of the systems: integumentary, respiratory, cardiovascular, gastrointestinal, musculoskeletal, neurological, endocrine, and lymphatic. The vascular, neurological, and musculoskeletal components of the HEENT, as well as the eyes and ears, are covered in separate chapters. The components addressed in this chapter include the head, face, nose, sinuses, neck, mouth, and pharynx. These components are complex in their actions and are involved in expression, communication, nourishment, respiration, and sensation,

among other functions. Furthermore, disorders involving the head and face can be devastating to patients because they can greatly affect appearance. Even minor disorders involving the head, face, or neck can be perceived as disfiguring by patients.

Anatomy and Physiology Review

Before you begin your assessment, you need an understanding of these complex structures, their basic anatomy (Figs. 11.1 through 11.10) and function, the ways in which they relate to other systems, and expected normal findings. (See *Head, Face, and Neck Structures and Functions*.)




FIGURE 11.1. Skull.




FIGURE 11.2. External nose.




FIGURE 11.3. Paranasal sinuses: Frontal, ethmoid, sphenoid, and maxillary.




FIGURE 11.4. Internal nose.




FIGURE 11.5. Structures of the mouth.

Interaction With Other Body Systems

The head, face, and neck include many structures with highly varied functions. Disruption or disease of several other systems can affect the organs of the head, face, and neck. These other systems include those described in the following paragraphs.

The Respiratory System

The nasal and oral cavities are entry points to the respiratory system. Injuries or diseases of these two structures can result in impaired ability to breathe. The ears, nose, and pharynx form the upper respiratory system. These structures communicate with the lower respiratory




FIGURE 11.6. Deciduous teeth.




FIGURE 11.7. Permanent teeth.




FIGURE 11.8. Salivary and parotid glands.




FIGURE 11.9. Thyroid gland.




FIGURE 11.10. Cervical lymph nodes.

system through the trachea, so that infections in one area can be transmitted to the other.

The Cardiovascular System

Many structures of the head, face, and neck system receive rich vascular supply. The mucosa of the nasal cavity includes a plexus of vessels that bleed easily. The temporal arteries are assessed during the examination of the face, as are the carotids during the examination of the neck. Disorders of the cardiovascular system may be reflected in the structures of the head, face, and neck. Infarct or ischemic pain may radiate to the jaw or throat. Facial edema can reflect fluid retention.

The Musculoskeletal System

The muscles of the face (Fig. 11.11) are highly involved in expression, communication, and nourishment. Diseases of the musculoskeletal system can have profound effects on these actions. Inflammatory changes of the tempo-

mandibular joint (TMJ) can cause limited jaw motion and jaw pain and can be unilateral or bilateral. Disorders of the cervical vertebrae or strain or inflammation of supporting structures can have profound effects on neck comfort and motion.

The Neurological System

The head, face, and neck are highly involved in many sensory processes. The neurological system includes the structures responsible for olfaction and taste as well as for sensation in the face and related structures. Dysfunction of the neurological system can have intense effects on these senses. Furthermore, the neurological system is involved in the complex movements necessary for speech, feeding, and expression. Altered neurological function is often evident in these actions. For example, Bell's palsy, an inflammatory paralysis of the trigeminal nerve, has a profound effect on the function of the motor components on the affected side.

(Text continued on page 274)


Head, Face, and Neck Structures and Functions

STRUCTURE	FUNCTION
Skull	
<i>Cranial Skull</i>	
Parietal, temporal, occipital, frontal, sphenoid, and ethmoid	Protects and supports the brain.
<i>Facial Skull</i>	
Maxillary, zygomatic, lacrimal, nasal, inferior nasal concha, palatine, mandible, and vomer	Forms facial framework.
<i>Joints</i>	
The sutures: coronal, lambdoid, sagittal, and squamous	Immovable joints that connect cranial bones. (Bones are joined at birth by pliable membranes called fontanel, which permit delivery through the birth canal and allow the brain to grow and develop.)
<i>Temporomandibular Joint (TMJ)</i>	Joins mandible and temporal bone, allowing movement of the mandible.
<i>Paranasal Sinuses</i>	Air cavities that add resonance to the voice and lighten the weight of the head. Lined with mucous membranes, the sinuses drain through the meatus, located beneath the turbinates on the lateral aspects of the nasal cavity.
Nose	
<i>Nares (Nostrils) and Vestibule</i>	Nares provide access to the nasopharynx through structures lined with mucous membranes, which are rich with blood vessels and covered with cilia (see below).
<i>Nasal Mucosa</i>	Ciliated epithelium cells with goblet cells produce mucus, which traps bacteria and air pollutants.
<i>Nasal Septum</i>	Composed of the ethmoid bone and vomer, divides the nasal cavities.
<i>Cilia</i>	Small hairs that filter air.
<i>Turbinates</i>	
Superior, middle, and inferior	Bony conchae (folds) of the internal nasal walls that increase the surface area for air to be filtered, warmed, and humidified before entering the lungs.
<i>Olfactory Nerve (CN I)</i>	Controls sense of smell. Stimulated by olfactory receptor cells located in upper nasal cavities.
Mouth	
<i>Lips</i>	Muscular folds that surround the mouth; contain sensory receptors and are very sensitive. Lips assist with eating, expression, and speech.
<i>Superior and Inferior Labial Frenulum</i>	Connect the lips to the gums.
<i>Buccal Mucosa</i>	Moist mucous membrane of the cheeks that assists with eating.
<i>Teeth</i>	
20 deciduous, 32 permanent	Needed for chewing; also have esthetic value.
<i>Gingiva (Gums)</i>	Help to anchor teeth, part of oral mucosa, cover alveolar bone.
<i>Tongue</i>	Muscle controlled by CN XII; needed for chewing, swallowing, and speech.
	Sensory (taste) CN VII and CN IX.
<i>Lingual Frenulum</i>	Attaches the tongue to the floor of the mouth.

(continued)

Head, Face, and Neck Structures and Functions (continued)



STRUCTURE	FUNCTION
<p>Pharynx Nasopharynx, oropharynx, and laryngopharynx</p>	<p>The pharynx is a muscular tube. Contraction of the oropharynx and laryngopharynx is part of swallowing reflex.</p> <p>The nasopharynx is the uppermost portion behind the nasal cavity. The adenoids are located on the posterior nasopharynx. The eustachian tubes open into the nasopharynx.</p> <p>The oropharynx is a part of the oral cavity and the palatine tonsils located on the lateral walls.</p> <p>The laryngopharynx is the most posterior portion of the pharynx. The anterior portion opens into the larynx; the posterior portion opens into the esophagus.</p>
<p>Tonsils Palatine, lingual, and pharyngeal</p>	<p>Lymphatic tissue that prevents infection. The palatine tonsils lie laterally on each side of the soft palate between the palatoglossal and the palatopharyngeal arches. The lingual tonsils are at the base of the tongue, and the pharyngeal tonsils (adenoids) are behind the nose in the posterior pharyngeal wall.</p>
<p>Hard Palate</p>	<p>Forms the floor of the nasal cavity.</p>
<p>Soft Palate</p>	<p>Posterior to hard palate, elevates during swallowing to prevent food and saliva from entering the nasopharynx.</p>
<p>Uvula</p>	<p>Fleshy conelike structure in the center of the soft palate that prevents food from entering nasal passages.</p>
<p>Salivary Glands</p>	<p>Secrete saliva, which contains amylase to convert starch to maltose.</p>
<p>Parotid Glands</p>	<p>Salivary glands anterior to the ear. Secretions help lubricate food to facilitate chewing and swallowing.</p>
<p>Stensen's Duct</p>	<p>Drains the parotid gland, enters the oral cavity through the buccal mucosa, opposite the second upper molar.</p>
<p>Submandibular Glands</p>	<p>Located at the posterior corner of the mandible.</p>
<p>Sublingual Glands</p>	<p>Located below the floor of the mouth.</p>
<p>Wharton's Duct</p>	<p>Drains submandibular and sublingual salivary glands, enters the oral cavity under the tongue on the floor of the mouth.</p>
<p>Neck</p>	
<p>Thyroid Gland</p>	<p>Controls metabolism and helps regulate calcium. The thyroid is the largest endocrine organ and produces thyroxine (T_4) and triiodothyronine (T_3), which are largely involved in the body's metabolism as well as cardiovascular, gastrointestinal, and neuromuscular functions. The thyroid also produces calcitonin, which lowers calcium and phosphate blood levels.</p>
<p>Parathyroid Glands</p>	<p>Located on the posterior side of the thyroid; increase blood calcium levels.</p>
<p>Cervical Lymph Nodes</p>	<p>Part of the lymphatic system. Drain the structures of the head and neck.</p>



NEUROLOGICAL

Cranial nerves located in head, face and neck.

Cranial nerves influence ability to communicate both verbally and nonverbally and to eat.



ENDOCRINE

Thyroid and parathyroid glands located in neck.

RESPIRATORY

Respiratory tract begins at nasal and oral cavities.

Injuries to head and face can affect breathing.

Respiratory infections often begin in upper airways of nose and throat.



INTEGUMENTARY

Skin-color changes on face (e.g., cyanosis, pallor, jaundice) may indicate systemic problems.



CARDIOVASCULAR

Temporal and carotid arteries located in head and neck.

Neck and jaw pain may indicate cardiovascular disease.


DIGESTIVE

Mouth is beginning of digestive tract.



MUSCULAR

Facial muscles needed for expression, communication and nutrition.



URINARY

Changes in face (e.g., edema or uremic frost) may reflect renal problems.



LYMPHATIC

Cervical lymph nodes located in neck.

Tonsils located in pharynx.

Mast cells located in pharynx.



SKELETAL

Skull protects brain.



REPRODUCTIVE

Pregnancy can cause changes in facial color (chloasma).

Lips and mouth are erogenous areas.

Relationship of Head/Face/Neck to Other Systems




FIGURE 11.11. Facial muscles.

The Endocrine System

The neck houses the thyroid gland, a major endocrine organ. Dysfunction of the thyroid is often accompanied by organ hypertrophy, or enlargement, regardless of whether the dysfunction results in hyperactivity or hypoactivity of the thyroid gland. Enlargements, called **goiters**, are evident during the examination of the neck. Many endocrine disorders have typical facies. For example, when hypocalcemia develops from parathyroid disease, Chvostek's sign, a facial spasm, may be an early symptom.

The Lymphatic/Hematologic System

The nasopharynx and oropharynx are rich with mast cells, responsible for allergic control symptoms. After exposure to allergens, the mast cell and basophil mediators are triggered, with resulting inflammatory responses responsible for the typical sneezing, itching, and secretions of the nasal, nasopharyngeal, oropharyngeal, and other linings. The ultimate result can include congestion, drainage, and secondary infections involving additional structures including the sinuses. The cervical nodes are sensitive to infection or inflammatory changes in the regions they drain, with resulting enlargement of nodes often signaling a disorder.

Because of the rich vascular supply of the nasal and oral mucosa, hematologic changes may first be reflected here. For instance, thrombocytopenia may first cause petechiae of the mucous membranes or bleeding of the gums or nose. Anemia may be accompanied by pallor of the mucosa or glossitis.

Developmental, Cultural, and Ethnic Variations

Infants and Children

Head

At birth, the head is often molded into a less than round shape during vaginal delivery, yet it assumes a symmetrical, rounded shape in several days. The fontanels (anterior, posterior, anterolateral, and posterolateral), which are the openings between the skull bones, remain open for growth, but close over the first year of life. The head circumference is an important measurement because a large head can indicate hydrocephaly, whereas a small head can indicate a developmental delay. In comparison to the head of an adult, the head of an infant or toddler is large in proportion to its body.

Nose and Sinuses

Only the ethmoid and maxillary sinuses are fully developed at birth. Others develop over the first 7 years. Children have a tendency to place small objects in their noses. Unilateral purulent drainage often results.

Teeth

The number of teeth varies, depending on a child's age. Infants get their first incisors at approximately 6 months of age. At age 6 or 7 years, they gradually begin to lose the 20 deciduous teeth, which are replaced with 32 permanent ones.

Throat and Neck

The tonsils, which are small in infancy, may become larger during childhood. They often become smaller again by

approximately age 12. The neck muscles of an infant are weak, and the infant must develop control of them to control the head. The thyroid of infants and children is not palpable. It is not unusual for young children to have palpable, small, “shoddy” cervical nodes.

Pregnant Women

Neck

Pregnant women may have palpable thyroids because the gland is stimulated by estrogen and develops increased vascular supply.

Face

A brownish pigmentation (chloasma or *the mask of pregnancy*) may occur.

Mouth

Hypertrophy of the gums may occur during pregnancy.

Older Adults

Teeth

With proper care and use of improved products, teeth should now last through advanced age. However, it is not unusual for older people to lose at least some of their teeth. They often have dental prostheses. Dentition plays a significant role in the nutrition of elderly people.

Mouth and Nose

Older people tend to have decreased production of saliva. With age, the senses of olfaction and taste may be diminished, which can affect nutritional status.

People of Different Cultures and Ethnic Groups

You need to consider your patient’s cultural background when assessing the head, face, and neck. What might be considered an abnormal finding in one cultural group may be a normal cultural variation for another.

Face

Facial skin color can vary greatly depending on cultural background. Remember, cyanosis or jaundice may be difficult to assess in patients with dark skin tones. Cultural variations can also be seen in skin textures and hair growth patterns. For example, African Americans may develop facial pseudofolliculitis (razor bumps) and Chinese Americans normally have little facial hair. Irish Americans have a greater incidence of skin cancer, so be sure to thoroughly assess sun-exposed areas, especially the face and ears. The shape and position of the eyes and nose may also have cultural variations. Chinese Americans have Mongolian traits, and Filipino-Americans have almond-shaped eyes, mildly flared nostrils, and a low, flat, nose bridge.

Introducing the Case Study



John Henck, a 17-year-old high school senior, is required to have a physical examination before participating in a deep-sea diving course and receiving certification. He is very excited about spending his spring vacation diving. You plan to complete both a health history and a physical examination. Because of the requirement that John be able to tolerate use of the diving equipment, a major emphasis of the examination is on the head, face, and neck.

Performing the Head, Face, and Neck Assessment

Assessment of the head, face, and neck involves obtaining a complete health history and performing a physical examination. As you perform the assessment, be alert for signs and symptoms of actual and potential problems of the various components of the head, face, and neck.

Health History

The health history identifies any related symptoms or risk factors and the presence of diseases involving the head, face, and neck. It must also detect any other disorders that may affect these structures. Your history will include obtaining biographical data and asking questions about the patient’s current health, past health, and family and psychosocial history. It also includes a review of systems (ROS). If you don’t have the time to perform a complete health history, make sure to at least perform a focused health history of the head, face, and neck.

Biographical Data

Review the patient’s biographical information. Note your patient’s age—certain diseases are more prevalent in specific age groups. For example, children tend to have more upper respiratory problems and pharyngitis than older adults. Also ask about your patient’s occupation. Does he or she have a job that puts him or her at risk for head injury? Does he or she spend long hours at a computer terminal (may result in tension headaches)? Questions like these will help to identify the potential for exposures to physical and environmental situations that could harm the head, face, and neck structures.

Current Health Status

Determine whether the patient has any specific presenting complaints related to the head, face, or neck. Some

Focused Health History for the Head, Face, and Neck

If a detailed history is either not feasible or inappropriate, be sure to ask the following basic questions. Remember that you must pursue a symptom analysis (PQRST) for any complaint identified during this focused history.

- Do you have problems or complaints related to your head, face, nose, mouth, throat, or neck? Some examples are head pain, nasal congestion, nosebleeds, nasal discharge, mouth sores or pain, sore throat, postnasal drip, difficulty swallowing, and neck pain.
- Do you have allergies to any medications, foods, or environmental factors?
- What, if any, health problems do you have?
- What, if any, OTC or prescribed medications do you take?
- Is there anything specific that you think I should know related to your overall health or this specific complaint?

common complaints include headaches, jaw pain, neck pain or stiffness, masses, nasal congestion, epistaxis, mouth or dental pain, mouth lesions, sore throat, and hoarseness. Any such complaints, as well as others that you might identify later, should be explored and developed using an organized system of symptom analysis, such as the PQRST format. Because disorders of several systems can influence the head, face, and neck, it is important to determine the patient's overall health status. There will be times, however, when the patient's presenting problems are particularly acute or distressful, and the initial history and physical will have to be very focused.

Symptom Analysis

Symptom analysis tables for the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Head Pain

Head pain can be associated with a variety of problems, including migraines, tension, systemic infections, and trauma.

Jaw Tightness and Pain

When a patient presents with jaw tightness and/or pain, the cause may be TMJ syndrome, but it could also be trauma or infection/inflammation in the structures near the jaw. An important consideration for jaw discomfort is whether it might be caused by cardiovascular disease. Always ask patients if they have a personal or family history of heart disease.

Neck Pain and Stiffness

Neck pain and stiffness can stem from musculoskeletal problems as well as from infections. Symptom analysis can help identify any forgotten trauma or physical exertion that might explain the complaint.

Neck Mass

When a patient complains of a neck mass, it might be a goiter of the thyroid gland or enlarged lymph nodes. Enlarged nodes may signal either an infectious or a malignant disorder.

Nasal Congestion

Nasal congestion is usually caused by an upper respiratory infection or allergy.

Nosebleed

Epistaxis, or nosebleed, is usually self-limited and has relatively benign causes. However, it can be caused by coagulopathies or other hematologic disturbances, malignancies, hypertension, or trauma.

Mouth Lesions

A mouth lesion can be caused by a malignancy, trauma, nutritional deficit, or poorly fitted dentures or orthodontic appliances.

Mouth and Dental Pain

Mouth pain can be caused by ischemic heart disease, musculoskeletal disorders, or dental problems.

Sore Throat

When a patient complains of a sore throat, the most common cause is a bacterial or viral illness. However, throat discomfort can be associated with throat masses, including thyroid hypertrophy or malignancies, foreign objects in the throat, and other causes.

Hoarseness

Another common complaint is hoarseness. Hoarseness may be caused by overuse of the voice, for example, prolonged periods of shouting or loud speech. It can also be an indication of gastroesophageal reflux, malignancies, neuromuscular disorders, or other health problems.

Past Health History

Once you have investigated the patient's chief complaint, explore the past health history. This portion of the history includes childhood and adulthood illnesses, surgeries, or major injuries; hospitalizations; major diagnostic procedures; exposures to infectious diseases; and allergies; as well as an immunization and medication history.

Past Health History

CATEGORY/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Childhood Illnesses

- As a child, did you have frequent respiratory infections, such as strep throat, sinus infections, or tonsillitis?
- Did you have allergies or asthma?

- Asking these questions identifies the potential for recurrence of problems that have existed since youth and for surgical or diagnostic procedures that should be explored. It also identifies the potential for rhinitis, which is commonly associated with other atopic disorders.

Hospitalizations

- Have you ever been hospitalized? What for?
- Have you had radiation to the neck; biopsies of the head, face, and neck; or other procedures?

- This further explores the existence of health problems that may be related to the head, face, and neck or other problems that might have an influence on these structures. It may identify disorders or previous syndromes that are still unidentified.

Surgeries/Serious Injuries

- Have you ever had surgery? If so, what kind? For example: Tonsillectomy, adenoidectomy, thyroidectomy, or rhinoplasty.

- This may explain physical findings, such as scars or the presence or absence of palatine tonsils.
- Both past surgeries and traumatic injuries may suggest the potential for chronic problems.

Serious/Chronic Illnesses

- Have you ever been diagnosed with or experienced thyroid disease, a malignancy, headaches, Bell's palsy, rhinitis, sinusitis, frequent upper respiratory infections, or TMJ syndrome?

- These questions point out diseases of the head, face, or neck that have a potential for recurrence or sequelae or that should be evaluated during current examination.
- These questions also inform you about treatment at the time of incidence and the patient's understanding of the diagnosis and treatment.

Immunizations

- Are your immunizations up to date?
- Have you ever had a flu shot or pneumovax? If so, when?
- When was your last tetanus/diphtheria vaccination?

- This allows you to document immunization history, especially influenza and diphtheria vaccines, which are of special significance to the upper respiratory tract.

Allergies

- Have you ever had an adverse reaction to medications, foods, chemicals, or environmental agents?

- This allows you to record previous allergic reactions so that they can be avoided in the future. A history of one atopic disorder, such as asthma or eczema, causes higher incidence of others, such as allergic rhinitis.
- A common allergic reaction is angioedema (benign swelling of skin, mucous membranes, or viscera).

Medications

- Are you currently taking any prescribed or OTC medications? What are they?

- Medications can cause reactive symptoms, including a sense of throat tightness from angioedema or adverse reactions such as ototoxicity. Knowing all agents the patient is using alerts you to potential for interactions with new agents.
- Chemotherapy often causes stomatitis, and superimposed fungal infections often occur with antibiotic therapy. Other medications, such as steroids, may have side effects that affect the head, face, and neck (e.g., moon face or hirsutism). See *Drugs That Adversely Affect the Head, Face, and Neck*.

(continued)

Past Health History (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Recent Travel/Military Service/Exposure to Infections or Contagious Diseases</p> <ul style="list-style-type: none"> ■ Are you aware of any recent exposures to infectious or contagious illnesses? ■ Have you traveled to foreign countries? ■ Have you shared a glass or eating utensils with anyone? 	<ul style="list-style-type: none"> ■ Identifies the potential for pending problem and presents an opportunity to determine the patient's understanding of protection from infections. ■ Drinking water and eating in foreign countries can expose patient to parasitic infections. ■ Knowing previous exposure to infectious disease may initiate further investigation of symptoms and prompt diagnostic testing to assist with a differential diagnosis. Note dates of possible exposure. ■ Teens on sports teams may share water bottles and pass on infectious disease.

Drugs That Adversely Affect the Head, Face, and Neck

When obtaining a health history to assess a patient's head and neck, the nurse must ask about current drug

use. Many of the commonly used drugs listed below can produce adverse reactions in the head and neck, including gingivitis (gum inflammation), alopecia (hair loss), and epistaxis (nosebleed).

Drugs That Adversely Affect the Head, Face, and Neck

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Anticholinergics	<ul style="list-style-type: none"> ■ Atropine, scopolamine, glycopyrrolate, propantheline, belladonna alkaloids, dicyclomine, hyoscyamine 	<ul style="list-style-type: none"> ■ Decreased salivation, dry mouth
Anticonvulsants	<ul style="list-style-type: none"> ■ Phenytoin ■ Valproic acid 	<ul style="list-style-type: none"> ■ Gingival hyperplasia ■ Alopecia
Antidepressants	<ul style="list-style-type: none"> ■ Tricyclics, including amitriptyline and nortriptyline, paroxetine 	<ul style="list-style-type: none"> ■ Dry mouth
Antihistamines	<ul style="list-style-type: none"> ■ Diphenhydramine, brompheniramine, chlorpheniramine 	<ul style="list-style-type: none"> ■ Dry mouth
Antihypertensives	<ul style="list-style-type: none"> ■ Guanabenz, clonidine, methyldopa ■ All calcium channel blockers ■ Propranolol 	<ul style="list-style-type: none"> ■ Dry mouth ■ Gingival hyperplasia ■ Hyperkeratosis and psoriasis of the scalp, alopecia
Antineoplastics	<ul style="list-style-type: none"> ■ Bleomycin ■ Dactinomycin ■ Melphalan ■ Mitomycin ■ Methotrexate ■ Cyclophosphamide ■ Vincristine ■ Chlorambucil ■ Uracil mustard ■ Cisplatin 	<ul style="list-style-type: none"> ■ Ulcerated tongue and lips, alopecia ■ Mouth lesions, alopecia ■ Mouth lesions, alopecia ■ Mouth lesions, alopecia ■ Gingivitis, mouth lesions, alopecia ■ Mouth lesions, alopecia ■ Mouth lesions, alopecia ■ Mouth lesions ■ Mouth lesions ■ Gingival platinum line, alopecia

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Cardiac agents Genitourinary smooth muscle relaxants Gold salts Miscellaneous agents	<ul style="list-style-type: none"> ■ Hydroxyurea ■ Fluorouracil ■ Doxorubicin ■ Cytarabine ■ Daunorubicin ■ Etoposide ■ Disopyramide ■ Flavoxate, oxybutynin, propantheline ■ Gold sodium thiomalate ■ Auranofin 	<ul style="list-style-type: none"> ■ Mouth lesions ■ Alopecia, epistaxis ■ Mouth lesions, alopecia ■ Mouth lesions, alopecia ■ Mouth lesions, alopecia ■ Alopecia ■ Dry mouth ■ Dry mouth
	<ul style="list-style-type: none"> ■ Lithium ■ Metoclopramide ■ Penicillamine ■ Isotretinoin ■ Edrophonium ■ Pyridostigmine 	<ul style="list-style-type: none"> ■ Gingivitis, mouth lesions, alopecia ■ Mouth lesions, alopecia ■ Dry mouth, dry hair, alopecia ■ Dry mouth, glossal or periorbital edema ■ Mouth lesions ■ Inflamed and bleeding gums, epistaxis, hirsutism ■ Increased salivation ■ Increased salivation, increased tracheobronchial secretions
	<ul style="list-style-type: none"> ■ Fluorides ■ Warfarin ■ Amphetamines 	<ul style="list-style-type: none"> ■ Staining or mottling of teeth ■ Epistaxis ■ Dry mouth, continuous chewing or bruxism (tooth grinding) with prolonged use
	<ul style="list-style-type: none"> ■ Tetracycline 	<ul style="list-style-type: none"> ■ Enamel hypoplasia and permanent yellow-gray to brown tooth discoloration in children under age 8 and in offspring of pregnant patients
	<ul style="list-style-type: none"> ■ Cyclosporine 	<ul style="list-style-type: none"> ■ Gingival hyperplasia

Family History

The purpose of the family history is to identify health problems that are familial or genetic. The history should include information on close relatives, both living and dead. The focus should be on problems that either have

a genetic component or are attributed to environmental/living situations shared with the patient. The genogram described in earlier chapters is a helpful way to organize the information obtained through the family history.

Family History

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Head and Neck Disorders</p> <ul style="list-style-type: none"> ■ Do you have a family history of migraine, cluster, or other headaches? ■ Do you have a family history of neck masses, thyroid disease, or malignancies? <p>Nose/Sinus Disorders</p> <ul style="list-style-type: none"> ■ Do you have a family history of rhinitis or nasal polyps? 	<ul style="list-style-type: none"> ■ Headache syndromes are often found in several family members. ■ Identify the history of neck masses or enlargements, including thyroid goiter and lymphadenopathy associated with lymphomas or other malignancies. ■ Rhinitis has a familial component. A family history of rhinitis, like other allergic problems, may have environmental triggers.

(continued)

Family History (continued)

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Mouth/Throat Disorders</p> <ul style="list-style-type: none"> Do you have a family history of cancers of the mouth or throat? 	<ul style="list-style-type: none"> This may indicate an increased likelihood of similar problems.
<p>Respiratory Disorders</p> <ul style="list-style-type: none"> Is there a history of asthma in your family? 	<ul style="list-style-type: none"> A family history of asthma increases the patient's chance of having rhinitis or other atopic diseases.
<p>Cardiovascular Disorders</p> <ul style="list-style-type: none"> Do you have a family history of heart disease or hypertension? 	<ul style="list-style-type: none"> A family history of coronary artery disease or cardiovascular disease is particularly important if the patient presents with neck or jaw pain unrelated to trauma, chewing, strain, or epistaxis.
<p>Hematologic Disorders</p> <ul style="list-style-type: none"> Is there any history of clotting or bleeding disorders among your close relatives? 	<ul style="list-style-type: none"> Familial bleeding disorders are significant when the patient has a history of bleeding gums or nosebleeds.
<p>Neurological Disorders</p> <ul style="list-style-type: none"> Do you have a family history of strokes, neuritis, or other neurological disorders? 	<ul style="list-style-type: none"> A family history of neurological disorders might help explain facial weakness or sensory disturbances, such as alteration of olfaction or taste.
<p>Endocrine Disorders</p> <ul style="list-style-type: none"> Do you have a family history of thyroid disease, acromegaly, or other endocrine diseases? 	<ul style="list-style-type: none"> Thyroid disease and other endocrine disorders have familial components. Acromegaly, a pituitary disorder, may be responsible for altered bone structure or soft tissue hypertrophy causing appearance changes. Many other endocrine disorders, such as Cushing's syndrome, present with typical facial changes.
<p>Musculoskeletal Disorders</p> <ul style="list-style-type: none"> Do you have a family history of arthritis or osteoporosis? 	<ul style="list-style-type: none"> A predisposition to arthritic disorders or osteoporosis is important when assessing pain of the neck or jaw.
<p>Lymphatic Disorders</p> <ul style="list-style-type: none"> Is there a family history of allergies or autoimmune disorders of the thyroid or other organs? 	<ul style="list-style-type: none"> Allergic disorders, which can be inherited, are a crucial consideration when assessing the nose and throat. Autoimmune disorders, such as rheumatoid arthritis or Sjögren's syndrome, are reflected in changes of the head, face, and neck.
<p>Infectious or Contagious Disorders</p> <ul style="list-style-type: none"> Have you recently been exposed to an upper respiratory infection or any other potentially contagious infection from family members or close friends? 	<ul style="list-style-type: none"> Identify the potential that any current symptoms might be related to contagious illness.

Review of Systems

The ROS allows you to explore each body system to determine whether the patient has a complaint that might affect the head, face, or neck. Ask the patient about problems that commonly affect the head, face, and neck, as well as all other systems. Although exploring the

patient's presenting complaint will already have alerted you to the potential for many symptoms, this review has the potential to trigger the patient to recall symptoms that she or he might otherwise have forgotten or to relate symptoms that she or he previously thought were unimportant.

Review of Systems

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- Has your energy level or weight changed?
- Have you had fevers or sweating?

- Endocrine and lymphatic system disorders often result in nonspecific symptoms, which are best elicited through the general ROS. Examples include thyroid disorders and allergies, both of which are important to the assessment of the head, face, and neck.

Integumentary

- Have you had any itching, rashes, sores, or other skin changes?
- Has the texture of your skin, hair, or nails changed?
- Have you noticed any changes in your skin, hair, or nails, such as color changes or an increase in facial hair?

- People with allergic rhinitis often have allergies affecting other systems, including the skin, and could have atopic rashes or pruritus.
- Sores or lesions might cause enlarged lymph nodes.
- Changes in the texture of skin, hair, and nails commonly occur with thyroid disorders.
- Color changes may be associated with endocrine disorders, such as the bronze color change of Addison's disease.
- Hirsutism (increase in facial or body hair growth) may be a result of Cushing's disease or a side effect of steroids.

HEENT

Head/Neck

- Have you had head or neck pain, neck masses, or swollen nodes?

- Although the symptom analysis and other history addresses symptoms of the head and neck, the ROS should again focus on that system in order to identify additional problems.

Eyes/Ears

- Have you noticed any changes in your vision or hearing?
- Have you had any drainage from your eyes or ears?
- Have you had any itching in your eyes or ears?
- Have you had any eye or ear pain?

- Because of their proximity, disorders that affect the eyes and ears may also affect other organs of the head, face, and neck. The ears communicate with the rest of the upper airways, so infections involving one structure may move to the other. Allergic disorders causing rhinitis often involve the ears and/or eyes as well.

Nose/Mouth

- Have you had nasal congestion, nasal drainage, sinus pain, or nosebleeds?
- Have you had mouth sores, dental pain, or jaw pain?

- The ROS should also include the nose and mouth to ensure that all potential problems have been explored.

Respiratory

- Have you had a cough or sputum or phlegm production?
- Do you ever wheeze or feel as though your chest is tight?
- Do you ever have trouble breathing?

- The upper and lower respiratory tracks are closely related, so infections in one often influence the other.
- People with asthma have a high incidence of allergic rhinitis.

Cardiovascular

- Have you had shortness of breath, chest pain, or palpitations?
- Do you have high blood pressure?
- How well do you tolerate activity?

- Cardiovascular disorders can result in fluid imbalances that cause facial edema. Veins may become distended and arterial pulses may be bounding. Extreme elevations in blood pressure can cause nosebleeds. Other symptoms of cardiac ischemia may help to explain jaw, tooth, or neck discomfort.
- Thyroid disorders have profound cardiovascular effects, including tachycardia and/or arrhythmias.

Gastrointestinal

- Have you had abdominal pain, appetite changes, nausea, vomiting, diarrhea, or constipation?

- Thyroid disorders can alter bowel habits.
- Infections that involve the upper respiratory tract, particularly viral ones, are often associated with nausea, abdominal discomfort, vomiting, and/or diarrhea.

(continued)

Review of Systems (continued)

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Genitourinary

- Do you have any history of kidney disease?

- Chronic renal failure may cause periorbital edema.
- Uremic frost may be seen on the face in end-stage renal disease.

Female Reproductive

- When was your last normal menstrual period?
- Have your periods been irregular?
- Have you ever been pregnant? When?
- Do you have vaginal discharge or sores?
- Have you had changes in sexual drive?

- Thyroid disorders frequently alter the menstrual pattern.
- Pregnancy may cause nasal congestion.
- Oral lesions and pharyngitis may be found in people with sexually transmitted diseases (STDs).
- Decreased libido is associated with both hypothyroid and hyperthyroid disease.

Male Reproductive

- Do you have penile discharge or sores?
- Do you have changes in sexual drive?

- Oral lesions and pharyngitis may be found in people with STDs.
- Decreased libido is associated with both hypothyroid and hyperthyroid disease.

Musculoskeletal

- Have you had joint pain or swelling, muscle aches, or limitations in motion?

- Joint and muscle symptoms suggest arthritic or neuromuscular disorders that would cause head, face, and neck discomfort or limit motion.
- Infectious diseases affecting the head, face, and neck may also cause generalized aches and pains.
- Hypothyroidism can result in an overall sense of fatigue or weakness; hyperthyroidism causes weakness of the proximal muscle groups of the extremities.

- Do you ever have weakness?

Neurological

- Do you have numbness, tingling, or muscle weakness?
- Do you have nervousness, changes in level of consciousness, trouble staying focused, forgetfulness, headaches, or history of head trauma?

- Neurological disorders can cause head pain and neck pain/stiffness. Paresthesias, paralysis, or weakness from neurological disorders may have a profound influence on facial expressions and ability to swallow.
- Thyroid disease may manifest as neurological symptoms.
- Nervousness, fine tremors, restlessness, and labile moods are associated with hyperthyroidism, apathy, lethargy; forgetfulness, slowed speech, and slowed mental processes are associated with hypothyroidism.
- A history of head trauma or headaches may also explain physical findings.

Endocrine

- Has your shoe size or ring size changed recently?
- Has your appearance changed significantly?
- Have your features changed significantly?
- Do you tire easily?
- Has your weight changed recently?
- Do you feel nervous or have trouble sleeping?

- Acromegaly, a pituitary hypersecretory disorder, causes enlargement and hypertrophy of bone and connective tissues, with profound alterations in facial structures and headaches. Fatigue, nervousness, altered sleep patterns, and weight changes are associated with thyroid disorders. Including these questions in the ROS ensures that issues missed by earlier history will be detected.

Lymphatic/Hematologic

- Do you have unusual bruising or bleeding?

- Disorders of the hematologic system may become evident through bleeding of other structures and help to explain nasal or oral bleeding.

Psychosocial Profile

The psychosocial profile provides information about the patient's occupation, social involvement, recreational interests, and daily activities and habits in order to identify factors that can influence the health of the head, face,

and neck. It determines risks associated with exposure to hazards, provides information about the patient's support system, and helps identify the patient's ability to perform self-care activities and obtain and carry out recommended treatments.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK

SIGNIFICANCE/RATIONALE

Health Practices and Beliefs/Self-Care Activities

- What is your usual bathing and toiletry routine?
- What toiletries or cosmetics do you usually use?

Typical Day

- How do you usually spend the day?

Nutritional Patterns

- What have you eaten in the past 24 hours?
- What types of foods do you usually eat?
- How much water and other fluids do you usually drink?
- What, if any, dietary supplements or vitamins do you take?

Activity and Exercise Patterns

- Do you exercise? What type of exercise do you do?

Recreation/Pets/Hobbies

- What do you do for recreation and hobbies? Do any of these activities involve exposure to extreme environmental conditions, chemicals, noise, or other harmful situations? What, if any, protective equipment do you use in your recreational activity?

Sleep and Rest Patterns

- How many hours of sleep do you usually get?
- Do you have problems with rest or sleep?
- Do you usually awaken feeling rested?

- Makeup and other toiletries may alter the skin integrity over the face.

- Knowledge of typical daily activities identifies potential exposures to hazards and habits that might adversely affect the head, face, and neck.

- Many nutritional deficits can affect the mucous membranes; for example, a "beefy" tongue is associated with pernicious anemia.
- Foods and additives may be sources of allergens.
- Nutrition influences many of the diseases that affect the head, face, and neck.
- Changes in appetite and weight are often associated with thyroid disease.
- Decreased appetite, nausea and vomiting, and weight gain are associated with hypothyroidism; increased appetite and thirst and weight loss are associated with hyperthyroidism.

- Information on the ability to exercise and exercise tolerance helps to provide a picture of any limitations presented by diseases such as rhinitis, headache, and neck pain. On the other hand, the type of exercises and activities performed may explain complaints such as headache or neck pain and stiffness.
- Thyroid disorders affect activity tolerance.

- Chemicals, environmental extremes, and noise can affect the head, face, and neck.

- Nasal congestion; sore throat; and other head, face, and neck complaints may limit or impair sleep.
- People with sleep apnea have altered sleep patterns and feel sleepy throughout the day.
- The need to nap during the daytime may suggest a poor night's sleep.
- Hypothyroidism often causes an increase in amount of sleep required; hyperthyroidism often interferes with ability to rest or sleep.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	SIGNIFICANCE/RATIONALE
<p>Personal Habits</p> <ul style="list-style-type: none"> Do you smoke cigarettes or use any other form of tobacco? If so, what type and quantity do you usually use daily? Do you drink alcohol? If so, what do you usually drink and how much? Do you use recreational drugs? If so, which ones, how often, and in what amount? <p>Occupational Health Patterns</p> <ul style="list-style-type: none"> What type of work do you do? Does your daily work involve exposures to loud noises, environmental extremes, or chemicals? What, if any, protection do you use in your work? <p>Environmental Health Patterns</p> <ul style="list-style-type: none"> Are you able to keep your home clean? Do you have problems with pests or dust? Is your home adequately heated and cooled? <p>Roles/Relationships/Self-Concept</p> <ul style="list-style-type: none"> Do you participate in any group activities? Have changes in your health recently affected your personal relationships? <p>Stress and Coping</p> <ul style="list-style-type: none"> How much stress do you usually experience? How do you usually deal with it? 	<ul style="list-style-type: none"> Tobacco affects the nose and sinuses. Smokeless tobacco is associated with oral cancers. Excessive alcohol use is associated with throat cancer. Drugs such as cocaine that are inhaled affect the nasal mucosa and erode the nasal septum. Many occupations expose people to injury from either sound or trauma. These questions help to determine the patient's access to, understanding of, and use of protective equipment. The home environment presents potential allergens. Roach excrement is an important allergen for people with asthma. Humidity and temperature affect the mucous membranes. The face, head, nose, mouth, and neck are very visible features. Any alteration may result in withdrawal from social activities because of depression or embarrassment. Stress and tension can trigger tension neck aches and headaches. Coping mechanisms are an important consideration for all patients.



Case Study Findings

Reviewing John's

Biographical data:

- 17-year-old unmarried male.
- Works part-time in fast-food restaurant.
- Born in the United States, was adopted immediately after birth.
- Protestant religion.
- Father is computer salesman and mother is radiology technician.
- Source:* Self, reliable.

Current health status:

- Describes current health as excellent. Says he is "health directed" and wishes to maximize health.
- No current complaints, although he does have "hay fever" in spring and fall. He is not currently experiencing any symptoms, but generally has several weeks of sneezing, runny nose, itching eyes, and head congestion. Antihistamines generally control the symptoms. On rare occasions at times other than in the fall or spring, he has symptoms

that last several hours, but he is not sure what triggers them.

Past health history:

- Usual childhood illnesses, without complications.
- Tonsils and adenoids removed at age 4, after frequent throat infections.
- Tympanoplasty at age 3, after frequent bouts of ear infections.
- Immunizations up to date; tetanus updated last year when he stepped on a piece of glass while wading on a shoreline.
- Seasonal hay fever since childhood.
- Takes OTC antihistamines when he has no prescribed antihistamines on hand, but usually takes Claritin during seasonal allergy attacks; has always taken one multivitamin daily; takes no other medications
- No recent exposures to known infections, but notes that he works in a busy restaurant and attends public school, so could be exposed to "whatever's going around."

Family history:

- Biological family history unknown; would like to know his family health history, but has been told this is not available.
- Both adoptive parents are in good health.

Review of systems:

- *General Health Survey:* States that health is usually good. No changes in weight or energy level.
- *Integumentary:* No changes in skin, hair, and nails. No skin rashes.
- *HEENT:* No swollen nodes or masses in head or neck. “Allergy attack” causes ear fullness; nasal congestion and clear drainage; postnasal drip; sneezing; and scratchy/itchy throat, eyes, and nose. Vision 20/20, no problems reported.
- *Respiratory:* No cough, shortness of breath (SOB), or wheezing.
- *Cardiovascular:* No chest pain or history of cardiovascular disease.
- *Gastrointestinal:* Appetite good, no dietary restrictions, no nausea or vomiting, has daily bowel movement.
- *Genitourinary:* No history of renal disease.
- *Reproductive:* Heterosexual, not currently sexually active, is aware of safe sex practices.
- *Musculoskeletal:* No weakness reported.
- *Neurological:* No headaches, dizziness, tremors, or paresthesia.
- *Endocrine:* No changes in weight or shoe size (11).
- *Lymphatic/Hematologic:* Positive seasonal allergies, no bleeding disorders.

Psychosocial profile:

- Bathes daily; uses no cosmetics on face other than occasional OTC acne medication; sees doctor and dentist for checkup once a year.
- Typical day starts at 6:45 A.M., when his mother awakens him to get ready for school. Attends school from 8 A.M. until 3:15 P.M. on weekdays, and works 5 P.M. until 11 P.M. on Monday and Friday nights and noon until 6 P.M. on Saturday. Goes to bed at 11:30 P.M. on school nights and 1 A.M. on weekends.
- Eats three meals plus one or two snacks daily. Breakfast consists of cereal, milk, and juice. Lunch is fast food, obtained between classes. Dinner is usually a cooked meal, shared with his parents. No specific dietary restrictions or food intolerances.
- Plays baseball and competes on swim team; practices 1 to 2 hours a day, depending on time of year. Wears protective gear while playing baseball.

- Tolerates swimming well, wearing goggles. Does some amount of physical activity daily, and this is well tolerated.
- Normally sleeps well and awakens rested. During “allergy season” sometimes has sleep difficulty. When he takes OTC antihistamines, they make it difficult to sleep and he wakes frequently through the night. If he takes no antihistamines or decongestants, he often awakens with congestion and/or sinus pressure.
- No medications other than seasonal use of antihistamines and daily vitamin. Uses no recreational drugs. Admits to rare use of alcohol (beer).
- Works part-time at restaurant.
 - Exposed to second-hand smoke in the restaurant where he works; otherwise not generally around smokers or exposed to any known chemicals. Tries to avoid exposure to smoke, as he is concerned about the potential health hazards.
- Lives in a single-family, two-story home described as “more than adequate” for his family.
- Has many friends and enjoys spending time with groups. Dates, but has no steady girlfriend.
- Does well in school and has been accepted to the state university. No major yet chosen.
- Not currently sexually active.

**Case Study Evaluation**

Before you proceed with the physical examination of the head, face, and neck, document the key history information you have learned from John’s health history and his related head, face, and neck history. The information from your patient history will be important to consider before and during your physical examination.

**CRITICAL THINKING ACTIVITY # 1**

Based on his history, is John at risk for any problems of the head, face, or neck?

**CRITICAL THINKING ACTIVITY # 2**

What strengths can you identify that will help John prevent or adapt to any problems of the head, face, or neck?

Anatomical Landmarks

Before you begin your physical assessment of the head, face, and neck, you need to visualize the underlying structures and identify landmarks. Two landmarks on the face that are useful in determining symmetry of facial features are the palpebral fissures and the nasolabial folds (Fig. 11.12). The palpebral fissure is the distance between the upper and the lower eyelid. The nasolabial fold is the distance from the corner of the nose to the edge of the lip. This is the facial crease that is often seen when someone smiles.

The anterior and posterior triangles (Fig. 11.13) are important landmarks of the neck. The **sternocleidomastoid** and **trapezius** muscles form the triangles. Both triangles are helpful in locating the underlying structures of the neck.

Physical Assessment

During the history, you probably developed a sense of the patient's concerns and may have begun to cluster the data obtained to help guide your physical examination. You should have an awareness of any physical limitations or discomfort that will influence the physical examination. Throughout the history, you observed the patient's body posture, fluidity of movements, facial expressions, and speech—all of which are important observations for the head, face, and neck. Now, as you approach the physical examination, you must be very objective in your observations as you inspect the internal structures of the nose, mouth, and throat.




FIGURE 11.12. Palpebral fissures and nasolabial folds.




FIGURE 11.13. Anterior and posterior triangles.

Approach

All four techniques of physical assessment—inspection, palpation, percussion, and auscultation—are used in the examination of the head, face, and neck. Some structures, like the throat and internal nose, can only be inspected; generally only the sinuses are percussed, and only the vessels of the neck and thyroid are auscultated. The cranial nerve (CN) assessment is generally incorporated in the examination of the face, mouth, nose, throat, and neck. The assessment of the arteries and veins is also incorporated into the examination of the neck and face. Although there is no “right” sequence to follow for the examination of these structures and organs, you should develop, practice, and adhere to a set routine in order to avoid omitting a test. One common sequence is the head-to-toe approach that



TOOLBOX

The tools that will be necessary to examine the head, face, nose, mouth, throat, and neck are a penlight or otoscope for focused light, tongue blades, gauze, stethoscope, transilluminator, cup of water, and gloves. If you are using an otoscope as a light source, you will want a wide-tipped speculum. A nasal speculum is another useful piece of equipment. Lighting is very important, and some examiners prefer also using a gooseneck lamp or headlamp when examining the mouth and throat.

begins with inspection of the shape and general placement of the head and facial structures, followed by a thorough inspection of the facial muscles and then the neck. Some examiners prefer to examine the nose, mouth, and throat along with the face, whereas others do this only after they have completed the examination of the neck. No matter what sequence you use, always take into consideration the structures' symmetry during your examination.

Performing a General Survey

The first step of the physical examination is a general survey, although in many cases this is accomplished during history taking. Besides providing early information regarding speech and movements, the general survey allows you to detect clues about the patient's emotional status, nutritional status, and overall posture. During the general survey, obtain vital signs. Altered pulse can be associated with thyroid disease. Respiratory rate changes can be related to an altered airway, including the nose, mouth, or pharynx. The temperature is an important consideration for infection. Blood pressure elevations may explain epistaxis. Aside from the vital signs, be alert for other signs that may indicate underlying problems with the head, face, and neck. For example:

- Note facial expression. Is it appropriate? Nervousness or a flat expression may be associated with thyroid disease. A masklike expression is seen with Parkinson's disease.
- Note any gross abnormalities, such as exophthalmus, which is seen with thyroid disease.
- Consider dress and grooming. Are they appropriate? Temperature intolerance associated with thyroid disease may cause people to overdress or underdress.
- Note speech and thought processes. Are responses appropriate? Are thought processes intact? Problems with focusing may be related to thyroid disorders.
- Look for changes in weight or weight distribution. A buffalo hump (fat pads on the lower midcervical and upper thoracic areas) is associated with Cushing's disease or steroid use.

Performing a Head-to-Toe Physical Assessment

The head, face, and neck reflect many different systems, so look for changes that may indicate underlying pathology. Next, perform a head-to-toe physical assessment, checking for specific signs of disease in other organ systems that might be reflected in the head, face, and neck.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

ABNORMAL FINDINGS/RATIONALE

General Health Survey

INSPECT

- Awake, alert, and oriented.
- Memory intact, able to provide adequate history.

- Altered mental status may indicate meningeal infection, which could be associated with head or neck pain or stiffness.
- Forgetfulness or slow mental processes are associated with hypothyroidism; mood lability, personality changes, and restlessness are associated with hyperthyroidism.

Integumentary

Skin

INSPECT/PALPATE

- Skin color varies according to racial and genetic makeup. Oral mucosa pink, skin warm, moist, intact, with good turgor. Should have stippled appearance, resembling orange peel.

- Color changes can readily be seen in mucous membranes. For example, cyanosis related to hypoxia, jaundice related to liver dysfunction, pallor related to anemia, or bronze/tan discoloration related to Addison's disease. Dry, thick, cool skin with poor turgor is associated with hypothyroidism; warm, smooth, moist skin is associated with hyperthyroidism.

Hair

- Hair evenly distributed. Texture dependent on racial and genetic makeup.

- Dry, sparse, coarse hair seen with hypothyroidism; patchy alopecia, and fine, silky hair, seen with hyperthyroidism. Increased facial hair in women is associated with Cushing's disease or steroid use.

Nails

- Nails pink and firm.

- Thick, brittle nails seen with hypothyroidism; thin, brittle nails seen with hyperthyroidism.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p>HEENT</p> <p>Eyes <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ Conjunctiva moist and pink. No drainage. ■ Palpebral fissures equal. No lid lag. <p>Ears</p> <ul style="list-style-type: none"> ■ Hearing intact. Tympanic membrane gray; light reflex intact. ■ Landmarks visible. No drainage or bleeding. <p>Respiratory <i>INSPECT/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Normal respiratory rate. Respirations regular and unlabored. ■ No cough. Clear breath sounds; no adventitious sounds. <p>Cardiovascular <i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Apical and radial pulses within normal limits. S₁, S₂ present and regular, with no adventitious or extra heart sounds. Pulses +2/4. ■ No bruits. <p>Gastrointestinal <i>INSPECT/PALPATE/PERCUSS/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Abdomen round, soft. Positive bowel sounds. Nontender, no organomegaly. <p>Genitourinary <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ External genitalia pink, intact, no lesions. <p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Full range of motion (ROM). No tenderness. Strength 5/5 (see Chapter 20, The Motor-Musculoskeletal System). No joint deformities. <p>Neurological</p> <ul style="list-style-type: none"> ■ Intact cranial nerves and sensation. Coordinated movements. 	<ul style="list-style-type: none"> ■ Conjunctival redness and drainage suggest infection or allergy, both of which may involve the nose and sinuses. ■ Exophthalmos and lid lag are associated with hyperthyroidism. <ul style="list-style-type: none"> ■ Hearing changes and altered appearance of tympanic membrane and landmarks associated with fluid accumulations behind the tympanic membrane (serous otitis) or infection (otitis media), both of which may affect the nose or throat. ■ Bleeding from the ear and nose signals possible skull fracture. <ul style="list-style-type: none"> ■ Tachypnea, cough, and wheezing may indicate asthma, which is highly related to other atopic diseases, such as allergic rhinitis or a respiratory infection that may also involve the upper respiratory tract. Masses or lesions in the nose, mouth, or throat can influence respiratory status. <ul style="list-style-type: none"> ■ Bounding temporal arteries paired with temporal pain suggests temporal arteritis, an explanation for head pain. ■ Bruits of the carotid arteries are detected during the examination of the neck and may indicate carotid stenosis. Proper examination should differentiate between bruits of the carotids and vascular sounds over the thyroid. <ul style="list-style-type: none"> ■ The mouth is the beginning of the GI tract, so oral problems may affect the GI system. Jaundice can be seen on the palate. <ul style="list-style-type: none"> ■ Oral lesions may be caused by STDs. <ul style="list-style-type: none"> ■ Limitations of motion, pain, and/or joint deformities are found with various forms of arthritis and may affect the TMJ and cervical joints. Weakness can be a sign of thyroid disorder as well as of various neuromuscular disorders that may affect expression, speech, and eating. Limited ROM of the neck and nuchal rigidity (resistance to neck flexion) are associated with meningitis. <ul style="list-style-type: none"> ■ Cranial nerves innervate most structures of the head, face, and neck and affect their function. Changes in sensation and coordination indicate neurological disorders that influence function of the nose (olfaction) and muscles of facial expression, eating, and speech.

Performing the Physical Assessment for the Head, Face, and Neck

After the general survey and head-to-toe assessment, perform a physical examination that focuses on the head, face, and neck. Although inspection and palpation are discussed separately below, they are not distinct, sequential activities. They are actually performed almost in concert. Although you inspect an area or structure before touching or moving it, this takes only a moment and is usually followed immediately by touching or palpating the area. The only area of the head, face, and neck to be percussed is the sinus area, and this generally occurs after

you have applied pressure over the sites during palpation. The only areas to be auscultated are the carotids and jugulars and, if it is enlarged, the thyroid.

Assessing the Head and Face

Examination of the head and face involves inspection and palpation.

Inspection

Have patients remove hats, wigs, or hair ornaments if present. Put on gloves in case there are open lesions under the hair. Begin with inspection. Identify the prominences of the brows, cheeks, mastoids, and occiput.

INSPECTION OF THE HEAD AND FACE

1. Head Size

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect head size and shape and symmetry of facial features.
- *Variation is wide, between and within gender and racial/ethnic group.*



Inspecting head size and shape and symmetry of facial features

ABNORMAL FINDINGS/RATIONALE

- *Abnormal increase in head size in young child:* May indicate hydrocephalus.
- *Inconsistently large head size in adolescent or adult:* May indicate acromegaly.



Acromegaly

(continued)

INSPECTION OF THE HEAD AND FACE

2. Head Shape

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Variation is wide, although shape should be symmetrical and contour rounded.*



HELPFUL HINT

Abnormal shapes or flat occiputs in infants and small children may result from positioning during the first year of life.

ABNORMAL FINDINGS/RATIONALE

- *Asymmetry of shape and contour:* Previous trauma, surgery, or congenital deformity.

3. Facial Appearance

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Facial appearance varies by gender, age, and racial/ethnic group. However, there should be symmetry of features and movement.*

ABNORMAL FINDINGS/RATIONALE

- *Facial appearance inconsistent with gender, age, or racial/ethnic group:* May indicate an inherited or chronic disorder with typical facies, such as Graves' disease, hypothyroidism with myxedema, Cushing's syndrome, or acromegaly.
- *Asymmetry of features:* Previous trauma, surgical alterations, congenital deformity, paralysis, or edema. Asymmetry is also seen with Bell's palsy and stroke.



Cushing's syndrome



Bell's palsy.

- *Asymmetry of movement:* Suggests neuromuscular disorder or paralysis. Tics, or spastic muscular contractions, usually occur in the head and face.



HELPFUL HINT

Two good places to inspect for symmetry of facial features are the palpebral fissures and the nasolabial folds.

Palpation

Next, palpate the head and face. There should be no tenderness and, except in infants, no soft areas in the head. As you palpate the TMJ, ask the patient to open and close his or her mouth and deviate his or her jaw from side to side. Determine sensation, motion, and strength of the face, as described in Chapter 20, Assessing the Motor-Musculoskeletal System, and assess the temporal artery,

as described in Chapter 15, Assessing the Peripheral Vascular and Lymphatic Systems.

Assessing the Sinuses

Assessment of the sinuses includes inspection (with transillumination), palpation, and percussion. Only the frontal and maxillary sinuses are readily accessible for assessment. Envision the areas of the face that overlay the sinuses.

PALPATION OF THE HEAD AND FACE

1. Head Contour/Facial Structures

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use light palpation to note head size, shape, symmetry, masses or areas of tenderness.
- Use light palpation to palpate the scalp for mobility and tenderness.
- *No tenderness or lesions.*
- *Relatively smooth with no unexpected contours or bulges.*



Palpating the head

ABNORMAL FINDINGS/RATIONALE

- *Contour abnormalities, including bulges or projections:* Previous trauma, surgery, or congenital deformity.
- *Tenderness:* Trauma, TMJ syndrome, temporal arteritis, or inflammatory process.

2. TMJ

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Palpate the TMJ by placing fingers over the TMJ and palpating the joint as the patient opens and closes his or her mouth.
- *Smooth, symmetrical motion, with no pain, crepitus, or clicking.*



Palpating the TMJ

ABNORMAL FINDINGS/RATIONALE

- *Irregular or uneven movement, pain with motion, or crepitus/popping:* TMJ syndrome.

Remember, the frontal sinuses are located above the eyebrows and the maxillary sinuses are located below the eyes.

Inspection

The sinus areas are inspected for edema and discoloration. If you suspect a sinus problem after regular inspection, palpation, and percussion, you can also transilluminate the sinuses. A transilluminator should be used; however, either a penlight or an otoscope with a speculum attached are good alternatives. Transillumination requires a darkened room. To transilluminate the frontal sinuses, hold the light source so that the light is directed upward from just below the brows. A glow of light may be

detected over the brow. To transilluminate the maxillary sinuses, have the patient open her or his mouth and position her or his head so that you can observe the roof of the mouth. Place the light source below the eyes and above the cheek, with the patient's mouth opened, and look for a glow on the roof of the mouth. Absence of transillumination suggests sinus fullness or thickening. Any glow noted with transillumination of either the frontal or the maxillary sinus should be symmetrical. However, absence of transillumination may not always indicate pathology. It may simply be a normal variant caused by the thickness of the bones overlying the sinuses or underdevelopment of the sinuses.

INSPECTION OF THE SINUSES

1. Frontal and Maxillary Sinuses

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect frontal sinuses above the eyes and maxillary sinuses below the eyes.
- *No periorbital edema or discoloration.*

ABNORMAL FINDINGS/RATIONALE

- *Periorbital edema and dark undereye circles: Sinusitis.*

2. Frontal and Maxillary Sinuses by Transillumination

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Transilluminate frontal sinuses by shining light upward under eyebrow.
- Transilluminate maxillary sinuses by shining light below eyes while looking for a red glow on the roof (palate) of the mouth.



Transilluminating frontal sinuses



Transilluminating maxillary sinuses

- **Frontal sinus:** *Normally, red glow noted above eyebrow.*
- **Maxillary sinus:** *Normally, red glow noted on roof of mouth.*
- *Expected variations include absence of transillumination because the ability to transilluminate is dependent on the thickness of the bones overlying the structure examined.*

ABNORMAL FINDINGS/RATIONALE

- *Absence of transillumination over one sinus when opposite structure transilluminates: Mucosal thickening or sinus fullness with sinusitis.*
- *Absence of transillumination must be considered with other findings.*

Palpation

Palpate the sinuses for tenderness. To palpate the frontal sinuses, press upward just below the medial third of each eyebrow. To palpate the maxillary sinuses, apply pressure to the lower portion of the cheeks, below the eyes.

Percussion

Percussion is performed to further assess for sinus discomfort. If tenderness is elicited with palpation, omit percussion over that area. Otherwise, you should percuss the

sinuses by tapping over these same areas. Direct percussion, using the tapping finger to strike directly over the bony prominence, is most frequently used. Because the sinuses are normally filled with air, percussion should elicit somewhat of a resonant tone.

Assessing the Nose

To examine the nose, inspect the external structures, palpate the external structures, and then inspect the internal structures (nasal cavity).

PALPATION OF THE SINUSES

1. Frontal and Maxillary Sinuses

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Palpate frontal sinuses by pressing upward just below eyebrows; note tenderness.



Palpating the frontal sinuses

- Palpate maxillary sinuses by pressing below eyes; note tenderness.



Palpating the maxillary sinuses

- *No tenderness.*

ABNORMAL FINDINGS/RATIONALE

- *Tenderness:* May indicate infectious or allergic sinusitis.

PERCUSSION OF THE SINUSES

1. Frontal and Maxillary Sinuses

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Percuss frontal sinuses with direct or immediate percussion above eyebrows.



Percussing the frontal sinus

- Percuss maxillary sinuses with direct or immediate percussion below eyes.



Percussing the maxillary sinus

- *No tenderness. Resonant tone.*

ABNORMAL FINDINGS/RATIONALE

- *Tenderness:* Suggests sinusitis.
- *Dull tone:* Indicates thickening or fullness of sinus cavity or cavities, associated with chronic or acute sinusitis.

Inspection

Inspection of the internal structures includes the septum, nasal mucosa, and medial and inferior turbinates. If you are using an otoscope with a wide-tipped speculum, stabilize the patient's head with one hand and then slowly and gently insert the speculum into the nares. If

you are using a penlight and nasal speculum, insert the closed speculum and then gently open it once it is in the nose, being careful not to open it too much. Take care not to scrape or press on the central septum because this area is sensitive. During your assessment, take note of any sounds the patient is making with his or her nose, such as sniffing or snorting.

INSPECTION OF THE NOSE

1. External Nose

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note size, shape, and symmetry.
- *Midline placement. Shape symmetrical and consistent with age, gender, and race/ethnic group.*
- *No nasal flaring.*
- *No drainage.*

ABNORMAL FINDINGS/RATIONALE

- *Misalignment of nose or shape inconsistent with patient's biographical information:* Previous trauma, congenital deformity, surgical alteration, or mass. Abnormal shape also associated with typical facies, including acromegaly or Down syndrome.
- *Nasal flaring:* Suggests respiratory distress, especially in infants, who are obligatory nose breathers.
- *Clear, bilateral drainage:* Allergic rhinitis.
- *Clear, unilateral drainage:* May be spinal fluid as a result of head trauma or fracture.
- *Clear, mucoid drainage:* Viral rhinitis.
- *Yellow or green drainage:* Upper respiratory infection.
- *Bloody drainage:* Trauma, hypertension, or bleeding disorders.

2. Internal Nasal Mucosa

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Tilt head back and use nasoscope or penlight to inspect nasal mucosa.



Inspecting the nasal mucosa

- *Pink, variations consistent with ethnic group/race and with oral mucosa.*
- *Moist, with only clear, scant mucus present.*
- *Intact, with no lesions or perforations.*
- *No crusting or polyps.*
- *Septum located midline.*

INSPECTION OF THE NOSE

2. Internal Nasal Mucosa (continued)

ABNORMAL FINDINGS/RATIONALE

- *Bright red mucosa*: Inflammation from rhinitis or sinusitis; also suggests cocaine abuse.
- *Pale or gray mucosa*: Allergic rhinitis.
- *Copious or colored discharge*: Allergic or infectious disorder, epistaxis, head or nose trauma.
- *Clustered vesicles*: Herpes infection.
- *Ulcers or perforations*: Chronic infection, trauma, or cocaine use.
- *Dried crusted blood*: Previous epistaxis.
- *Polyps (elongated, rounded projections)*: Allergies, irritation or chronic infections.
- *Deviated septum*: Normal variant or following trauma.



Polyps



ALERT

A deviated septum is cause for concern if breathing is obstructed.



Deviated septum

3. Turbinates

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect the turbinates. The middle turbinate is located more medially, the inferior turbinate is more lateral, and the superior is not visible.
- *Medial and inferior turbinates visible, symmetrical and shape/size consistent with general features of patient. Overlying mucosa coloring consistent with other mucous membranes.*

RATIONALE/SIGNIFICANCE

- *Enlarged, boggy turbinates*: Allergic disorder.
- *Pale or gray mucosa overlying turbinates*: Allergic disorder.

Palpation

Palpate the bony ridge and soft tissues of the external nose. The cartilaginous, distal two-thirds of the nose should be mobile, without pain. Gently occlude one nostril at a time and have the patient inhale through the nose to determine patency. (See *Palpation of the Nose*.)

Assessing the Mouth and Throat

The mouth and throat are components of both the respiratory and the digestive tracts. Assessment involves inspection and palpation. The assessment begins with examination of the lips, then the structures of the mouth and throat. Remember to wear gloves when

PALPATION OF THE NOSE

1. External Nose

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Occlude each nostril and note patency.



Palpating the nostril

- *Cartilaginous portion is slightly mobile. Nontender, no masses. Nares patent.*

ABNORMAL FINDINGS/RATIONALE

- *Deviations or masses:* Previous trauma or infection.

examining the internal structures of the mouth. You will also need a penlight and tongue depressor to perform the examination.



HELPFUL HINT

Because the mucous membranes reproduce cells rapidly, mouth lesions tend to heal quickly with treatment. Therefore any persistent lesion requires medical attention. Be aware of the possibility of oral cancer.

Common Findings With Oral Cancer

- Nonhealing lip or mouth sore.
- Swelling or mass in the mouth or throat; for denture wearers, swelling of jaw can affect denture fit and be uncomfortable.
- Erythroplakia/leukoplakia.
- Unusual bleeding in the mouth.
- Unusual sensation, such as pain or numbness, in the mouth.
- Pain with chewing and swallowing.
- Chronic sore throat.
- Foreign body sensation in the throat.
- Dysphagia.
- Voice change.
- Ear pain.

Inspection

Inspect the lips, gingiva, buccal mucosa, tongue, and pharynx for color, lesions, and exudates. Note the color, number, condition, and occlusion of the teeth. The upper and lower molars should approximate with the jaw closed. The front teeth should slightly override the lower ones. Observation of the teeth with a physical assessment does not replace a dental examination, so remind the patient of the importance of maintaining routine dental care.

Using the tongue blade to displace the cheeks and lips, first inspect the **buccal mucosa**. The **Stensen's ducts**, openings for the parotid glands, are located on the buccal mucosa at the point of the second upper molars. The **Wharton's ducts**, openings for the submandibular glands, are located on either side of the **frenulum** under the tongue.

Inspect all aspects of the tongue: dorsal, ventral, and lateral edges. Note the color, moisture, and surface texture and observe for any swelling. Observe the frenulum and the mobility of the tongue.

Palpation

Palpate the tongue for nodules or areas of thickening. Palpate the floor of the mouth for nodules or masses. To palpate the floor of the mouth, use your nondominant hand to press upward beneath the patient's chin to provide support while palpating downward with a gloved hand inside the mouth. The support provided externally helps ensure that the examining hand actually palpates a mass and does not merely push it away. The use of two hands will help to assess any mass, examining the dimensions, consistency, tenderness, and texture.

(Text continued on page 305)

INSPECTION OF THE MOUTH AND THROAT

1. Lips

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color, condition, lesions, odor.



Inspecting the lips

- *Midline, symmetrical, skin intact, pink, and moist.*
- *Coloring consistent with ethnic group/race.*
- *No unusual odors.*

ABNORMAL FINDINGS/RATIONALE

- *Asymmetry of placement:* Congenital deformity, trauma, paralysis, or surgical alteration.
- *Pallor:* Anemia.
- *Redness:* Inflammatory or infectious disorder.
- *Cyanosis:* Vasoconstriction or hypoxia.
- *Lesions:* Infectious or inflammatory disorder.



Cancer on the lip

- *Cheilitis (inflammation of lips), drying, and cracking:* Dehydration, allergy, lip licking.
- *Cheilosis (fissures at corners of lips):* Deficiency of B vitamins or maceration related to overclosure.



Cheilitis and cheilosis

- *Chancre:* Single, painless ulcer of primary syphilis.



Chancre

- *Angioedema:* Allergic response.



Angioedema

- *Herpes simplex (clustered area of fullness/nodularity that forms vesicles, then ulceration):* Herpes viral infection.



Herpes simplex

- *Halitosis:* Infections or gastrointestinal problems.

(continued)

INSPECTION OF THE MOUTH AND THROAT

2. Teeth and Bite

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Have patient open and close mouth. Note occlusion and number, color, condition of teeth.
- *Most adults have 28 teeth, or 32 if the four third molars, or wisdom teeth, are erupted. (However, they are usually impacted or extracted.)*
- *Teeth should be white, not loose, with good occlusion, and in good repair.*



Inspecting the teeth and bite

ABNORMAL FINDINGS/RATIONALE

- Various abnormalities include loose, poorly anchored teeth, malalignment, dental caries.



Malocclusion



Dental caries

- *Discoloration of teeth:* Chemicals or medications (tetracycline may discolor teeth gray if administered before puberty).

- *Mottled enamel:* Fluorosis (excessive fluoride).



Tetracycline staining



Fluorosis

INSPECTION OF THE MOUTH AND THROAT

3. Oral Mucosa and Gums

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color, condition, lesions of mucosa.
- Note condition of gingiva, bleeding, retraction, or hypertrophy.



Inspecting the oral mucosa and gums

- *Pink, moist, intact mucosa. Color variants acceptable if consistent with patient's ethnic group/race for instance, dark stippling in dark-skinned patients.*
- *Gums consistent in color with other mucosa and intact, with no bleeding.*

ABNORMAL FINDINGS/RATIONALE

- **Gum hyperplasia:** Side effect of medications, such as dilantin or calcium channel blockers.
- **Gum recession or inflammatory gum changes (gingivitis/periodontal disease):** Poor dental hygiene or vitamin deficiency.
- **Pale or gray gingivae:** Chronic gingivitis.



Gum hyperplasia



Gingival recession



Chronic gingivitis

- **Abrasions, erosion of underlying mucosa:** In denture wearers, poorly fitted dentures. Inflamed, bleeding gingivae may also be seen with leukemia and human immunodeficiency virus (HIV).



Leukemia



Early HIV periodontitis



Advanced HIV periodontitis

(continued)

INSPECTION OF THE MOUTH AND THROAT

3. Oral Mucosa and Gums (continued)

ABNORMAL FINDINGS/RATIONALE

- *Painful, reddened mucosa, often with mildly adherent white patches:* Candida albicans.
- *Reddened, inflamed oral mucosa, sometimes accompanied by ulcerations:* Allergic stomatitis.
- *Small, painful vesicles that often have a reddened periphery and a white or pale yellowish base:* Aphthous ulcer, believed to be caused by viral infection, stress, or trauma.
- *Nodular, macular, or papular lesions widely involving the integument and often evident on the oral mucosa:* Kaposi's sarcoma. Incidence has increased with the development of acquired immunodeficiency syndrome (AIDS).
- *Inflammatory changes of the integument, often found on oral mucosa as chronic gray, lacy patches with or without ulceration:* Lichen planus. May progress to neoplasm.



Aphthous ulcer



Lichen planus

- *Reddened mucosal change that may progress to form cancer:* Erythroplakia
- *White, adherent mucosal thickening:* Leukoplakia. May progress to cancer.
- *Cancer on lip or oral mucosa:* Cancers can be found on the lips, gums, oral mucosa or other areas of the mouth and are associated with tobacco use and alcohol abuse.
- *Enlarged sebaceous glands on buccal mucosa, white/yellow raised lesions:* Fordyce granules (very common).



Leukoplakia



Cancer on the oral mucosa



Fordyce granules

- Medication reactions can cause a variety of changes to oral mucosa, as can infections.

INSPECTION OF THE MOUTH AND THROAT

4. Hard and Soft Palate

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color and condition of hard and soft palate.
- *Palate intact, smooth, pink.*
- *Bony, mucosa-covered projection on the hard palate (torus palatinus) or on floor of mouth (torus mandible) are normal variations.*



Inspecting the palate



Torus palatinus



Torus mandible

ABNORMAL FINDINGS/RATIONALE

- *Perforation:* Congenital or from trauma or drug use.



Cocaine use



HIV palatal candidiasis

5. Salivary Ducts

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Stensen's duct:* Inspect inner aspect of cheek (buccal mucosa) opposite the second upper molar.
- *Wharton's duct:* Have patient lift tongue and inspect the floor of mouth.



Inspecting Stensen's ducts



Inspecting Wharton's ducts

(continued)

INSPECTION OF THE MOUTH AND THROAT

5. Salivary Ducts (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Stensen's duct intact at buccal mucosa at level of second molars.*
- *Wharton's duct intact at either side of frenulum.*
- *Both ducts with moist and pink mucosa; no lesions, swelling, or nodules.*

ABNORMAL FINDINGS/RATIONALE

- *Fullness or inflammatory changes of glands:*
Blockage of duct by calculi, infection, malignancy.
Parotitis is inflammation of parotid glands.



Parotitis

6. Tongue

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color, texture, moisture, and mobility.



Inspecting the tongue



Inspecting the underside of the tongue

- *Pink and moist.*
- *Coloring may vary consistent with ethnic group/race.*
- *Mucosa intact with no lesions or discolorations.*
- *Papillae intact. Tongue is freely and symmetrically mobile (CN XII intact).*
- *Geographic tongue is a normal variation.*



Geographic tongue

INSPECTION OF THE MOUTH AND THROAT

6. Tongue (continued)

ABNORMAL FINDINGS/RATIONALE

- *Absence of papillae, reddened mucosa, ulcerations:* Allergic, inflammatory, or infectious cause.
- *Color changes:* May indicate underlying problems; for example, red “beefy” tongue is seen with pernicious anemia.
- *Black, hairy tongue:* Fungal infections.



Red, beefy tongue



Black, hairy tongue

- *Hypertrophy and discoloration of papillae:* Antibiotic use.
- *Reddened, smooth, painful tongue, with or without ulcerations (glossitis):* Anemia, chemical irritants, medications.



Glossitis

- Cancers may form on the tongue and on other oral mucosa.



Cancer of the tongue

7. Oropharynx

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect oropharynx for color, lesions, and drainage.
- *Mucosa is pink, moist, intact. The lymphoid-rich posterior wall may have a slightly irregular surface.*
- *No lesions, erythema, swellings, exudate, or discharge.*



Inspecting the oropharynx

(continued)

INSPECTION OF THE MOUTH AND THROAT

7. Oropharynx (continued)

ABNORMAL FINDINGS/RATIONALE

- *Yellowish or green streaks of drainage on the posterior wall:* Postnasal drainage.
- *Gray membrane/adherent material:* Diphtheria.
- *White or pale patches of exudates with erythemic mucosa:* Infection, including streptococcal bacterial infection or mononucleosis viral infection. Gonorrhea and chlamydia are also associated with exudative pharyngitis.
- *Erythema:* Inflammatory response, typically associated with infectious pharyngitis; also common in smokers.
- *Scattered vesicles/ulcerations:* Herpangioma.



Herpangioma

8. Tonsils

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Locate tonsils posterior to arches on sides of throat.
- Note color, size, and exudate.
- *Symmetrical, pink, clean crypts. Crypts may have normal variation of small food particles (tonsillar pearls) or scant amounts of white cellular debris.*

ABNORMAL FINDINGS/RATIONALE

- *Bulges adjacent to the tonsillar pillars:* Potential peritonsillar abscess.
- *Reddened, hypertrophic tonsil, with or without exudates:* Acute infection or tonsillitis.
- Lymphoid cobblestoning.
- Enlarged tonsils with exudates.



Enlarged tonsils with exudates. See *Tonsil Grading Scale* for specific tonsil grades.

9. Uvula

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Have patient say “AH!” and note symmetrical rise of the uvula.
- *Midline, pink, moist, without lesions.*
- *Symmetrical rise of the uvula (CN IX, glossopharyngeal, and CN X, vagus, intact).*

ABNORMAL FINDINGS/RATIONALE

- *Erythema, exudate, lesions:* Infectious process.
- *Asymmetrical rise of the uvula:* Problem with CN IX and CN X.

Tonsil Grading Scale

Grade	Description
1+	Tonsils extend to arches.
2+	Tonsils extend just beyond arches.
3+	Tonsils approximate the uvula.
4+	Tonsils meet midline ("kiss").

The parotid, submandibular, and sublingual glands should also be palpated for enlargement and tenderness. The parotid glands are located anterior to the ear, and the submandibular and sublingual glands are located under the mandible.



HELPFUL HINT

If the salivary glands are enlarged, palpate very gently because they may be very tender.

PALPATION OF THE MOUTH AND THROAT

1. Lips

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Lightly palpate lips for consistency and tenderness.
- *Soft, nontender, no masses.*



Palpating the lips

(continued)

PALPATION OF THE MOUTH AND THROAT

1. Lips (continued)

ABNORMAL FINDINGS/RATIONALE

- *Areas of induration, thickening, nodularity, or masses:* Neoplasm.
- *Tender induration that soon develops vesicles:* Herpes simplex.

2. Tongue

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Lightly palpate tongue for consistency and tenderness.
- Tissue is soft, without masses, nodules, thickenings, or tenderness.
- Tissue is soft, supple, without nodules, thickenings, masses, or tenderness. Sublingual glands may be palpable under the tongue but should be nontender, soft, and supple.

ABNORMAL FINDINGS/RATIONALE

- *Areas of induration, thickening, nodularity:* Potential malignancy.
- *Areas of unexpected induration, thickening, nodularity or other mass:* Malignancy.

3. Glands (Parotid, Submandibular, and Sublingual)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Parotid:* Palpate in front of ears.
- *Submandibular and sublingual:* Palpate under the mandible.



Palpating the parotid gland



Palpating the sublingual gland

- *Parotid glands are nonpalpable and nontender. Submandibular and sublingual glands may be palpable but should be nontender, supple, and soft.*

ABNORMAL FINDINGS/RATIONALE

- *Enlarged, tender parotid glands:* Parotitis, blocked ducts, infection, or malignancy.

Assessing the Neck

Examination of the neck integrates components of the vascular, respiratory, musculoskeletal, neurological, lymphatic, and endocrine systems. This chapter focuses on the lymphatic and endocrine systems, primarily on assessment of the cervical lymph nodes and the thyroid through inspection, palpation, and auscultation.

Inspection

You will be inspecting the cervical lymph nodes and thyroid gland. Remember, lymph nodes drain toward the

center of the body. When examining the thyroid gland, focus your attention on the middle to lower third of the anterior neck, checking for enlargements.

Palpation

The order in which you palpate the cervical lymph nodes is not important, although it is best to develop a sequence and be consistent to ensure that you do not omit a group. One common sequence is to start with the preauricular nodes, followed by the postauricular nodes, then move to the tonsillar, submandibular, and submental

INSPECTION OF THE NECK

1. Neck, Lymph Nodes, and Thyroid

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect neck in neutral and hyperextended positions and as patient swallows.



Inspecting the neck from the neutral position



Inspecting the neck when hyperextended



Inspecting the neck when the client swallows water

- *Neck erect, midline, no lumps, bulges, or masses.*
- *Thyroid not visible. No masses, swelling, or hypertrophy in mid to lower half of anterior neck.*

ABNORMAL FINDINGS/RATIONALE

- **Enlargements:** Lymphadenopathy, lymphoma, or other malignancy.
- **Torticollis** (*deviation of neck to one side caused by spasmodic contraction of neck muscles*): Scars, tonsillitis, adenitis, disease of cervical vertebrae, enlarged cervical glands, cerebellar tumor, rheumatism retropharyngeal abscess.
- **Enlarged, visible thyroid:** Goiter or malignant mass.



Cervical adenitis

nodes along the mandible. Next, palpate the occipital area, followed by the superficial and deep cervical, posterior cervical, and supraclavicular nodes. Palpate node groups gently with one or two fingers, applying alternate pressure. Palpate any identified nodes between two fingers to establish their dimensions, texture, consistency, and shape.

Although the lymph nodes are generally not palpable, it is not unusual to identify them at 1 cm or less in

size. Palpable lymph nodes should be described according to their location, size, shape, consistency, mobility, and tenderness. Palpable small nodes should be soft to rubbery in consistency and be freely mobile, distinct, round, and nontender. The term used to describe enlarged nodes (>1 cm in diameter) is **lymphadenopathy**. Lymphadenopathy can be regional (involving one or two groups) or more generalized (involving three or more groups).



HELPFUL HINTS

- Lymphatic tissue is largest in childhood and decreases in size with age. Normal palpable nodes are more likely to be found in children than in adults.
- Patients who present with a sore throat often complain about “swollen glands.” They are actually feeling their submandibular salivary glands. To distinguish between salivary glands and lymph nodes, remember: A normal lymph node is either small (< 1 cm), round, soft to rubbery, movable, and nontender or tender and enlarged with infection. Submandibular glands are larger, soft, glandular, and not freely movable.
- A palpable normal node is more likely to be a superficial node than a deep cervical one. Deep cervical nodes are normally nonpalpable.

To palpate the thyroid, use an anterior or posterior approach. Some examiners combine both approaches when assessing a thyroid nodule or enlargement. Both approaches are depicted here, so that you can determine which works best for you. Begin by locating the thyroid gland.

Although the thyroid is usually nonpalpable, you may be able to feel the isthmus, which connects the two lobes and lies below the cricoid cartilage. The lobes are located behind the sternocleidomastoid muscle. The likelihood of palpating the isthmus increases with very thin or pregnant

patients. You may be able to feel the edge of the gland, especially in women, who have larger thyroid glands than men.

The thyroid gland moves as the individual swallows. Therefore, have the patient drink water while you palpate the gland, to facilitate detection. Instruct her or him to take a sip from the cup and hold it in the mouth until you ask her or him to swallow. Do this at least twice, as you examine both the left and the right thyroid lobes.

To use the posterior approach, stand behind the patient and ask him or her to flex his or her neck slightly forward and to the left. This relaxes the muscles and the skin overlying the left side of the neck, making it easier to detect the tissue of the left thyroid lobe. Using the fingers of your left hand, locate the cricoid process. Push the trachea slightly to the left with your right hand as you palpate just below the cricoid process and between the trachea and the sternocleidomastoid muscle with your left hand. As you palpate, ask the patient to swallow the water he or she is holding in the mouth. Now, repeat the steps with the patient’s head flexed slightly forward and to the right. This time, displace the trachea slightly to the right with your left hand and palpate the right lobe of the thyroid with your right hand.

To use the anterior approach, stand in front of the patient and ask her or him to flex the neck slightly forward and in the direction you intend to palpate. Place your hands on the neck and apply gentle pressure to one side of the trachea while palpating the opposite side of the neck for the thyroid. The patient should take a sip of water during this approach as well.

Differentiating Lymph Nodes

CHARACTERISTICS	NORMAL	INFECTION	MALIGNANCY
Size	<1 cm	Enlarged	Enlarged
Shape	Round	Round	Irregular
Consistency	Soft to rubbery	“Boggy”	Hard
Delimitation	Well defined	Well defined	Irregular borders
Mobility	Mobile	Mobile	Immobile
Tenderness	Nontender	Tender	Nontender

PALPATION OF THE NECK

1. Neck

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use light palpation and check for masses or areas of tenderness.
- *Supple, nontender, no masses.*

ABNORMAL FINDINGS/RATIONALE

- *Masses:* Lymphadenopathy, malignancies, thyroid masses.

2. Cervical Nodes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- **Occipital node:** Lightly palpate at the back of the head at the base of the skull.



Palpating the occipital node

- **Postauricular node:** Lightly palpate behind the ears.



Palpating the postauricular node

- **Preauricular node:** Lightly palpate in front of the ears.



Palpating the preauricular node

- **Tonsillar node:** Lightly palpate at the angle of the jaw.



Palpating the tonsillar node

- **Submandibular:** Lightly palpate under the mandible.



Palpating the submandibular node

- **Submental:** Lightly palpate under the tip of the chin.



Palpating the submental node

- **Superficial cervical node:** Lightly palpate the upper portion of the sternocleidomastoid muscle.



Palpating the superficial cervical node

- **Posterior cervical node:** Lightly palpate in the posterior triangle behind the sternocleidomastoid muscle.



Palpating the posterior cervical node

- **Deep cervical node:** Palpate the sternocleidomastoid muscle.



Palpating the deep cervical node

(continued)

PALPATION OF THE NECK

2. Cervical Nodes (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Supraclavicular node:* Lightly palpate above the clavicle.



Palpating the supraclavicular node

- *Infraclavicular node:* Lightly palpate below the clavicle.



Palpating the infraclavicular node

- *Lymph nodes normally not palpable especially the deep and clavicular nodes. If a node is palpable, normal characteristics include small, 1 cm, mobile, soft, nontender, and usually superficial.*

ABNORMAL FINDINGS/RATIONALE

- *Palpable nodes (1 cm or greater):* Malignancy, inflammatory, or infectious process of glands or area they drain.

3. Thyroid

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Locate the thyroid isthmus below the cricoid cartilage.



Landmarking the thyroid

- *Anterior approach:* To palpate the right lobe, slide fingers to the right, gently displace trachea to right, and palpate gland as patient swallows. To palpate left lobe, slide fingers to the left, gently displace trachea to left, and palpate gland as patient swallows.



Palpating the thyroid:
Anterior approach

PALPATION OF THE NECK

3. Thyroid (continued)

- *Posterior approach:* Have patient tilt head to right, and then gently displace trachea to right, slide fingers to right, and palpate right thyroid lobe as patient swallows (the gland moves up with the cartilage as the patient swallows). To palpate the left lobe, repeat the same technique but have patient tilt head to left, displace trachea to left, and palpate the left lobe.



Palpating the thyroid: Right posterior approach



Palpating the thyroid: Left posterior approach

- *Generally nonpalpable.* If some tissue is palpable (e.g., in very thin or pregnant patients), consistency is firm, smooth, and meaty, with no nodularity, enlargement, or tenderness.

ABNORMAL FINDINGS/RATIONALE

- *Enlarged thyroid:* Tumor, goiter.
- Nodular thyroid tissue.
- *Tender thyroid:* Inflammatory process such as acute thyroiditis.

Auscultation

The final portion of the neck examination, auscultation of the thyroid gland, is generally reserved for situations in which the thyroid is enlarged or a mass is palpated. To auscultate the gland, place the bell of your stethoscope over one lobe, then the other. Ask the patient to briefly stop breathing as you auscultate, to optimize your ability to hear without the distraction of the tracheal breath sounds. There should be no thyroid sounds. Because the thyroid is a very vascular organ,

vascular sounds are sometimes present in hyperthyroidism. (See *Auscultation of the Thyroid*.)



HELPFUL HINT

An enlarged thyroid gland may occur with either hyperthyroidism or hypothyroidism. If the gland is enlarged, further assessment of presenting signs and symptoms and thyroid function studies are warranted to determine the cause.

AUSCULTATION OF THE THYROID

1. Thyroid

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- If thyroid gland is palpable, have patient hold breath and then listen over the thyroid gland with the bell portion of the stethoscope for bruits.
- *No sounds detected.*



Auscultating the thyroid

ABNORMAL FINDINGS/RATIONALE

- *Bruit:* Increased vascularity of hyperthyroidism.

Signs and Symptoms of Hyperthyroidism and Hypothyroidism

Hyperthyroidism

- Muscle weakness/tremors
- Irritability or emotional lability, manic behavior
- Decreased weight
- Sleep disturbance
- Thyroid enlargement, goiter
- Heat intolerance
- Irregular menses, amenorrhea
- Flushed skin, fine hair

Hypothyroidism

- Muscle weakness, aches and pain
- Apathy, agitation, paranoia, depression
- Increased weight
- Increased sleep
- Thyroid enlargement, goiter
- Cold intolerance
- Changes in menses
- Coarse, dry skin and hair



Case Study Findings

Your general health survey and head-to-toe assessment of John have concluded. The following are your findings:

■ General Health Survey

- Well-developed 17-year-old male.
- Tanned skin, healthy appearance consistent with stated age.
- No apparent distress.
- Head-to-toe scan reveals no signs of head, face, and neck problems.
- Projects a confident and comfortable affect.
- Provides history with ease.
- Head-to-toe scan reveals no signs of problems in other systems/areas that would affect head, face, and neck.

■ Vital Signs

- Temperature 97.8°F
- Pulse 62.
- Respirations 14.
- Blood pressure (BP) 100/72, left arm.
- Height 5'10".
- Weight 160 lb.

■ Head and Face

- Head is normocephalic, with no abnormal contour, bulges, lesions, or tenderness.
- Face is symmetrical at rest and at motion, with no lesions.

■ Nose and Sinuses

- Sinuses are nontender, resonant to percussion, and maxillary/frontal sinuses transilluminate bilaterally.

(continued on page 314)

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	Ask about: Changes in energy level Weight changes Fever Night sweats	General	Inspect: Features for symmetry and proportion Measure: Vital signs Height and weight
Integumentary	Ask about: Changes in skin, hair and nails Rashes, itching	Integumentary	Inspect: Skin for color changes (e.g., cyanosis, pallor), lesions, hair distribution Palpate: Skin for temperature, turgor, texture Nails and hair texture
HEENT <i>Head and Neck</i>	Ask about: Head or neck pain Masses or swollen nodes	HEENT <i>Head and Neck</i>	Inspect: Size, shape, symmetry of head and facial features Facial expression Palpate: Thyroid Lymph nodes
Eyes	Ask about: Vision changes, drainage, itching, pain	Eyes	Inspect: Conjunctiva Palpebral fissures Lid lag, exophthalmos Test: Visual acuity
Ears, Nose and Throat	Ask about: Changes in hearing Ear drainage, itching, pain Nasal congestion, drainage, sinus pain Nosebleeds Mouth sores or dental pain Jaw pain	Ears, Nose and Throat	Inspect: Gross hearing Structures of the mouth and throat External ear and tympanic membrane
Respiratory	Ask about: Cough, congestion, wheezing, mucus production	Respiratory	Measure: Respiratory rate Auscultate: Breath sounds
Cardiovascular	Ask about: History of cardiovascular disease (e.g., CAD, HTN) Chest pain, palpitations	Cardiovascular	Measure: Heart rate Pulse amplitude Palpate: Pulse Auscultate: Heart sounds, bruits
Gastrointestinal	Ask about: Changes in appetite Bowel changes Nausea, vomiting	Gastrointestinal	Inspect: Abdomen for size, shape and symmetry Palpate: Organomegaly Auscultate: Bowel sounds
Genitourinary/ Reproductive	Ask about: History of STD Changes in libido Women: Last menstrual period Menstrual changes Vaginal discharge or sores Men: Penile discharge, sores	Genitourinary/ Reproductive	Inspect: External genitals for lesions or discharge
Musculoskeletal	Ask about: Joint pain Muscle weakness Limited movement	Musculoskeletal	Inspect: ROM of head and neck Palpate: Muscle strength, joint deformities
Neurological	Ask about: Nervousness Difficulty staying focused Forgetfulness Changes in level of consciousness Numbness and tingling Headaches, head injury	Neurological	Test: Level of orientation Thought process and memory Cranial nerves Sensation Inspect: Abnormal movements
Endocrine	Ask about: Changes in shoe size or ring size Changes in facial features Changes in energy level Weight changes Sleep problems	Endocrine	Inspect: Masses, goiters Palpate: Thyroid enlargement, nodules Auscultate: Thyroid, if palpable
Lymphatic/ Hematologic	Ask about: Unusual bleeding, bruising Current/recent infection History of cancer	Lymphatic/ Hematologic	Inspect: Echymoses or petechiae Palpate: Lymph node enlargement

Assessment of the Head/Face/Neck's Relationship to Other Systems



Case Study Findings (continued)

- External nose is midline, symmetrical, with no drainage or lesion.
- Nasal mucosa is intact, slightly paler than oral mucosa, with no lesions or polyps.
- Nares are patent, no nasal flaring.
- Septum is midline, mucosa intact and pink, with no deviation or perforation.
- Turbinates are symmetrical, slightly enlarged.
- Scant clear, mucoid secretions.
- **Mouth and Teeth**
 - Lips are midline, symmetrical, close easily.
 - 28 white teeth in good repair, good occlusion.
 - Lips are pink, supple, with no lesions or indurations.
 - Oral mucosa is uniformly deep pink, glistening, moist; no lesions or indurations.
 - Tongue is supple, freely mobile, pink mucosa with papillae intact, no lesions.
 - Stensen's and Wharton's ducts with mucosa intact, no induration or swelling.
- **Pharynx**
 - Tonsils are symmetrical, small, pink, with clean crypts; no lesions or exudates.
 - Uvula is midline with symmetrical rise and pink mucosa.
 - Posterior wall is moist, smooth, pink, with no exudates or lesions.
- **Neck**
 - Head and neck are erect; neck is supple, with full range of motion, no bulges or masses.
 - Cervical nodes are nonpalpable.
 - Trachea is midline and mobile.
 - Thyroid is nonpalpable, nontender.



Case Study Analysis and Plan

Now that your examination of John is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.



CRITICAL THINKING ACTIVITY # 3

What strengths and areas of concern have you identified related to John's current health status?

Nursing Diagnoses

Consider all of the data you have collected during your assessment of John Henck. Use this information to identify a list of nursing diagnoses. Some possible nursing diagnoses are provided below. Cluster the supporting data.

1. Health-Seeking Behaviors, related to the desire to participate in recreational activities and to maintain overall health.
2. Risk for Trauma, related to frequent athletic activities, teenage driving, occasional alcohol intake.
3. Disturbed Sleep Pattern, related to adverse reaction to medication and/or sinus/nasal discomfort and congestion.

Identify any additional nursing diagnoses.



CRITICAL THINKING ACTIVITY # 4

Now that you have identified appropriate nursing diagnoses for John, design a nursing care plan and a brief teaching plan for one of the diagnoses identified above, including learning outcomes and teaching strategies.

RESEARCH TELLS US

In 2005, the American Cancer Society reported that an estimated 29,370 Americans (19,100 men and 10,270 women) would be diagnosed with oral or pharyngeal cancer, and nearly 7320 Americans would die from these disorders. In fact, oral cancer claims more lives than melanoma and cervical cancer, and squamous cell carcinoma accounts for 90 percent of oral cancers. Although the oral cavity is readily accessible and easily examined, most oral malignancies are detected in the advanced stages, and the survival rate for oral cancer has not changed over the past 3 decades.

In 2001, the American Dental Society (ADA) conducted a public service program to increase public awareness of oral cancer. The ADA performed two surveys, a telephone survey to consumers (n = 1270) and a mailed survey to dentists (n = 2500). The

results of the consumer survey revealed that the majority did not recognize the severity of the disease and the dentist's role in detecting oral cancer. The majority of the dentists surveyed were aware of the ADA initiative and became more active in screening for oral cancer. Stahl, Meskin, and Brown (2004), who studied the initiative's impact on consumer and dentist awareness of oral cancer, concluded that continued efforts to provide health education and health promotion interventions to patients may result in earlier detection and better prognosis.

Because nurses are present in all aspects of health-care, you are in an ideal position to provide health education and screenings for your patients. Remember that screening for oral cancer begins with the patient history and the identification of risk factors.

HEALTH CONCERNS

Risk factors for oral cancer include:

- Use of tobacco, either through smoking, chewing, or the use of “snuff.”
- Exposure to secondhand smoke.
- Alcohol consumption (75 to 80 percent of patients with oral cancer have history of alcohol heavy use, and oral cancer is more common among alcohol drinkers than nondrinkers).
- Ultraviolet light exposure.
- Poor nutrition (diets low in fruits and vegetables increase risk).
- Plummer-Vinson syndrome, a rare iron deficiency condition with associated tongue, nail, esophagus, and red blood cell abnormalities.
- Human papillomavirus (HPV) infection.
- Immunosuppression.
- Older age (half of all patients with oral cancer are older than age 65).
- Male gender.
- Chronic irritation of the mouth, such as with poorly fitted dentures, has been suggested as a risk factor for oral cancers since causative factors, such as alcohol and tobacco particles, can become trapped. Denture wearers should remove dentures at night and clean them every day. Denture fit should be evaluated at least every 5 years.

Source: American Cancer Society. (2005). What are Risk Factors for Oral Cavity and Oropharyngeal Cancer? Available at: www.cancer.org. Last accessed January 23, 2005.

Common Abnormalities

ABNORMALITY

ASSESSMENT FINDINGS

Acute Cervical Adenitis

Localized or regional enlarged and tender lymph nodes

- Affected area warm and erythematous



Acute Parotitis

Inflammation of the parotid glands

- Fever
- History of cat scratch, dental infection, upper respiratory infection, or other exposure
- Pain over salivary gland site
- Swelling with gland visible above jaw angle and anterior to earlobe
- Swollen and tender over the duct
- Firm consistency (hard growth that develops gradually may indicate malignancy)
- Purulent discharge at duct orifice possible
- Fever
- Generalized overlying erythema possibly present

(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Acute Rhinitis, Viral Acute nasal congestion with increased mucus secretion</p>	<ul style="list-style-type: none"> ■ Clear or mucoid drainage ■ Congestion ■ Fever ■ Mild scratchy sore throat ■ Mild dry cough ■ Sneezing
<p>Acute Sinusitis Inflammation of the sinuses</p>	<ul style="list-style-type: none"> ■ Mucopurulent nasal discharge ■ Postnasal drainage ■ Cough ■ Fever ■ Inflamed mucosa ■ Tenderness over sinuses ■ Periorbital swelling ■ Facial pain
<p>Allergic Rhinitis Hay fever</p>	<ul style="list-style-type: none"> ■ Sneezing ■ Rubbing nose and eyes ■ “Allergic salute” ■ Transverse nasal crease ■ Watery clear discharge ■ Allergic “black eye” ■ Periorbital puffiness
<p>Candidal Stomatitis Fungal inflammation of the mouth</p>	<ul style="list-style-type: none"> ■ History of underlying condition (HIV, diabetes mellitus) or medication (corticosteroid, antibiotic) ■ Tongue usually involved ■ Erythematous mucosa ■ Thin, white plaques often present
<p>Chronic Sinusitis Chronic inflammation of the sinuses</p>	<ul style="list-style-type: none"> ■ Nasal discharge ■ Congestion ■ Cough ■ Dull headache ■ Thick postnasal drainage ■ Ear pressure ■ Halitosis ■ Fatigue
<p>Epistaxis Nosebleed</p>	<ul style="list-style-type: none"> ■ History of trauma, local irritation, inflammation of nasal mucosa ■ Scant to significant bleeding ■ 90% of anterior bleeding at Kesselbach’s plexus



ABNORMALITY	ASSESSMENT FINDINGS	
<p>Herpes Simplex Vesicular eruption caused by a virus</p>	<ul style="list-style-type: none"> ■ Tender, enlarged lymph nodes ■ Clustered, small (3–4 mm) vesicles on mucosa ■ Vesicles rupture and form ulcers ■ Severe pain ■ Fever ■ History of previous episodes 	
<p>Ludwig's Angina A suppurative inflammation of subcutaneous tissue adjacent to a submaxillary gland</p>	<ul style="list-style-type: none"> ■ History of dental disease ■ Cellulitis on floor of mouth (tender, red, swollen) ■ Fever ■ Systemic symptoms 	
<p>Nasal Polyp A tumor with a pedicle, found in the nose</p>	<ul style="list-style-type: none"> ■ Mobile (unlike fixed turbinates) ■ Pale, nontender rounded masses ■ Pale coloring ■ Variable size ■ Usually bilateral ■ History of atopic disease, chronic rhinitis, or chronic sinusitis 	
<p>Periodontal Disease Disease of the supporting structures of the teeth</p>	<ul style="list-style-type: none"> ■ Loose teeth ■ Inflamed gingiva ■ Bleeding ■ Edema of gums ■ Possible tooth loss 	
<p>Pharyngitis Inflammation of the pharynx</p>	<ul style="list-style-type: none"> ■ Throat pain ■ Upper respiratory infection symptoms ■ Tender anterior cervical nodes ■ Posterior pharyngeal erythema ■ Fever ■ Scant white or yellow exudate possibly present 	
<p>Pharyngitis, Mononucleosis Inflammation of the pharynx caused by mononucleosis</p>	<p>Same symptoms as pharyngitis, plus:</p> <ul style="list-style-type: none"> ■ Increased fever ■ Fatigue ■ Enlarged lymph nodes ■ Palatal petechiae 	
<p>Pharyngitis, B Streptococcal Inflammation of the pharynx caused by streptococci</p>	<p>Same symptoms as pharyngitis, plus:</p> <ul style="list-style-type: none"> ■ Increased fever ■ Dysphagia ■ Abdominal pain, nausea and/or vomiting ■ Headache ■ No upper respiratory symptoms ■ Increased exudate ■ "Sandpaper" rash 	

(continued)

Common Abnormalities *(continued)*

ABNORMALITY	ASSESSMENT FINDINGS
<p>Tonsillitis Inflammation of the tonsils</p>	<p>Same symptoms as pharyngitis, plus:</p> <ul style="list-style-type: none"> ■ Exudative tonsils ■ Tonsils enlarged 2+ to 4+ 
<p>Torus Palatinus A benign exostosis located in the midline of the hard palate</p>	<ul style="list-style-type: none"> ■ Dysphagia ■ Midline bony growth projecting outward from the hard palate ■ Covered with normal mucosa ■ May have multiple “lobules” 

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the head, face, and neck.

Ask the patient the following:

- Do you have . . .
 - Head pain/headaches?
 - Jaw tightness and pain?
 - Neck pain and stiffness?
 - Neck mass?
 - Nasal congestion?
 - Nosebleed?
 - Mouth and dental pain?
 - Sore throat?
 - Hoarseness?
 - Do you have problems or complaints related to your head, face, nose, mouth, throat, or neck?
 - Do you have allergies to any medications, foods, or environmental factors?
 - Do you have any health problems?

- Are you taking any medications, prescribed or over-the-counter (OTC)?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam:

- | | |
|--------------|------------------|
| Gloves | Stethoscope |
| Penlight | Otoscope |
| Tongue blade | Transilluminator |
| Gauze | Cup of water |

Position

Sitting

Technique

First, scan your patient head to toe checking for specific signs of diseases affecting other organ systems that might alter the head, face and neck.

(continued)

ASSESSMENT CHECKLIST (continued)

Inspection, Palpation, Percussion, Auscultation

Head

Inspect

- Size, shape, and symmetry
- Facial appearance, symmetry of facial features, lesions, hair distribution, and abnormal movements

Palpate

- Tenderness, masses
- TMJ joint

Sinuses

Inspect

- Edema
- Transilluminate sinuses

Palpate

- Tenderness

Percuss

- Tenderness

Nose

Inspect

- Position, septal deviation, discharge, flaring
- Nasal mucosa color, intact septum, turbinates, polyps

Palpate

- Patency

Mouth and Throat

Inspect

- Lips:** Color, lesions, purse-lip, odor
- Oral mucosa:** Color, lesions
- Gingivae:** Color, bleeding, retraction, hypertrophy
- Teeth:** Occlusion, number, condition, color
- Tongue:** Color, lesions, texture, position, mobility
- Oropharynx/soft and hard palate/tonsils:** Color, condition, lesions, drainage, exudates

Palpate

- Lips and tongue:** Tenderness, muscle tone, lesions
- Oropharynx:** Test gag reflex

Neck

Inspect

- Neutral position, hyperextended, and as patient swallows
- Note ROM, symmetry, condition of skin

Palpate

- Cervical lymph nodes:** Occipital, postauricular, preauricular, tonsillar, submandibular, submental, superficial, deep, posterior, supraclavicular, infraclavicular
- Thyroid gland:** Anterior or posterior approach

Auscultate

- Thyroid:** If palpable, listen for bruits with the bell over the thyroid.

DOCUMENT

Document your findings and develop your plan of care.


SUMMARY

- Disorders of the head, face, and neck have the potential to exert profound influence over the morbidity and mortality of patients.
- The physical changes associated with these disorders can have devastating effects.
- The structures and function of the head, face, and neck are greatly influenced by changes in other systems.
- It is essential that nurses be skilled in performing examinations of this system and differentiating between normal and abnormal findings.

REVIEW QUESTIONS

1. Sean Mooney, 3 years old, is being seen for an upper respiratory infection. Part of the assessment includes assessing his sinuses. Which of his sinuses is not yet fully developed?
 - a. Frontal
 - b. Maxillary
 - c. Ethmoid
2. As you assess Sean's throat and mouth, considering his age, how many teeth would you expect to find?
 - a. 20
 - b. 28
 - c. 30
 - d. 32

3. When assessing the face, which area is best for assessing symmetry of facial features?
 - a. Nasolabial folds
 - b. Inner canthus
 - c. Anterior triangle
 - d. Posterior triangle
4. The anterior and posterior triangles are two landmarks used to assess the neck. What two muscles make up these triangles?
 - a. Omohyoid and trapezius
 - b. Masseter and sternocleidomastoid
 - c. Trapezius and sternocleidomastoid
 - d. Temporalis and omohyoid
5. You are assessing the lymph nodes of the head and neck. Where are the occipital nodes located?
 - a. In front of the ears
 - b. Behind the sternocleidomastoid muscle
 - c. Behind the ears
 - d. At the base of the skull
6. You are assessing the lymph nodes of the head and neck. Where are the submental nodes located?
 - a. Under the jaw
 - b. Behind the ears
 - c. Under the chin
 - d. At the angle of the jaw
7. You are assessing the lymph nodes of the head and neck. Where are the tonsillar nodes located?
 - a. Under the jaw
 - b. At the angle of the jaw
 - c. Under the chin
 - d. In front of the ears
8. You are assessing the lymph nodes of the head and neck. Where are the posterior cervical nodes located?
 - a. On the sternocleidomastoid muscle
 - b. In the anterior triangle
 - c. In the posterior triangle
 - d. Above the clavicle
9. When assessing the thyroid gland, you locate the isthmus. Where is the isthmus located?
 - a. Below the hyoid bone
 - b. Below the thyroid cartilage
 - c. Below the cricoid cartilage
 - d. Below the tracheal rings
10. To facilitate palpation of the thyroid gland, you instruct your patient to:
 - a. Flex neck.
 - b. Hyperextend neck.
 - c. Swallow.
 - d. Hold breath.
11. You are palpating your patient's parotid glands. Where are the parotid glands located?
 - a. Anterior to the ears
 - b. Posterior to the ears
 - c. At the angle of the jaw
 - d. Under the jaw



Assessing the Eye and the Ear

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent eye and ear history questions
- Obtain an eye and an ear history
- Perform an eye and an ear physical assessment
- Document eye and ear assessment findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal eye and ear findings

The eye and the ear are sensory structures that connect us with the environment. They allow us to perceive our surroundings through sight and sound. Disorders of the eye and the ear can range from minor annoyances to life-threatening problems. Most problems do not result in acute illness; however, they may be associated with more serious neurological conditions such as brain tumor, stroke, or head injury.

No matter what the cause, visual and hearing problems can have a major impact on physiological functioning as well as psychological and social well-being. Early detection reduces the likelihood of problems related to social interaction.

Determining whether a patient has adequate vision and hearing is crucial before assessing mental status or providing instructions. The eyes and the ears are common sites of injury; they also exhibit structural variations as a result of age, cultural background, and genetic influences. Although, for the sake of clarity, the eyes and the ears are covered separately here, they are usually examined along with the head and neck because of their location.

A thorough assessment of the eyes and the ears includes vision and hearing screenings and examination of the external and internal structures. The assessment provides not only specific data about the eyes and the ears but also vital information on the health status of other systems.

Before you begin your assessment, an understanding of the anatomy and physiology of the eyes and the ears is essential. You need to be able to identify normal structures before you can identify abnormal findings, accurately perform the assessment, and correctly interpret your findings. This chapter covers assessment of the eyes first and then assessment of the ears.

Anatomy and Physiology

Review: The Eye

The primary function of the eye is vision, including central and peripheral vision, near and distance vision, and differentiation of colors. To accomplish these tasks, the external and internal structures of the eye work together to receive and transmit images to the occipital lobe of the brain for interpretation. Visual difficulties can result from disease or injury to any of the structures involved in the visual pathway.

Structures and Functions of the Eye

The eye consists of internal and external structures that support or protect it (Figs. 12.1 through 12.5).

How We See

The ability to see objects in the environment depends on light waves that reflect off images. Natural lighting produces gradations of shading that help to determine an object's shape and position in the environment. The light rays pass through the cornea, anterior chamber, pupil, lens, and posterior chamber to the back of the eye (retina). The **pupil**, which is actually created by the aperture of a muscular layer of tissue called the **iris**, dilates or constricts to allow more or less light onto the retina. The **lens**, an elastic biconvex disc, bends the light waves entering the eye by either flattening or increasing the lens curvature. The precise functioning of the pupil and iris together allows a clear image to focus on the retina.

(Text continued on page 326)




FIGURE 12.1. Frontal view of eye and eyelid.




FIGURE 12.2. Lacrimal glands and ducts.




FIGURE 12.3. Eye globe with attachment of extraocular muscles.




FIGURE 12.4A. Cross-section of the eye, showing anterior and posterior chambers.




FIGURE 12.4B. Cross-section of the eye, showing choroid layer.




FIGURE 12.5. Right eye, showing optic disc.

Structures and Functions of the Eye

STRUCTURE	FUNCTION
Eyelid	<ul style="list-style-type: none"> Protects eye from injury and distributes lubrication over the eye globe. Lid margins contain meibomian glands, which lubricate the lid margins.
Orbicularis oculi muscle	<ul style="list-style-type: none"> Closes the eyelid. Is innervated by cranial nerve (CN) VII (facial).
Levator palpebrae	<ul style="list-style-type: none"> Raises the eyelid. Is innervated by CN III (oculomotor).
Eyelashes	<ul style="list-style-type: none"> Protect eyes from injury from foreign bodies.
Medial canthus	<ul style="list-style-type: none"> Contains the punctum, which drains the lacrimal gland fluids (tears) into the nasolacrimal sac.
Nasolacrimal sac	<ul style="list-style-type: none"> Located alongside the nasal bridge. Receives excessive tears from the lacrimal gland and drains them into the inferior meatus in the nose.
Lacrimal gland	<ul style="list-style-type: none"> Located in the upper lid just below the orbital rim. Produces tears to moisturize the globe and wash away foreign bodies.
Conjunctiva	<ul style="list-style-type: none"> A mucous membrane that lines the eyelids (palpebral conjunctiva) and the outermost portion of the globe (bulbar conjunctiva). Contains goblet cells that secrete fluids to lubricate the eyes.
Cornea	<ul style="list-style-type: none"> The anterior outermost layer of the eyeball, which covers the pupil and iris and extends to the limbus. Contains sensory innervation for pain, thus providing an early warning system for eye injury. Avascular and transparent.
Sclera	<ul style="list-style-type: none"> Underlies the bulbar conjunctiva and gives the eyeball its white color. A tough avascular layer that gives the eye its structure and shape, which is important for transmission of visual images to the back of the eye for interpretation.
Iris	<ul style="list-style-type: none"> A circular, muscular tissue that forms the colored part of the eye. The aperture of the iris forms the pupil and controls the amount of light entering the retina.
Extraocular muscles	<ul style="list-style-type: none"> A set of six muscles, innervated by cranial nerves, that move the eyes in a conjugate (parallel) manner (see Fig. 12.3). <i>Superior rectus</i>: Rolls eyeball upward, innervated by CN III. <i>Inferior rectus</i>: Rolls eyeball downward, innervated by CN III.

STRUCTURE	FUNCTION
Anterior chamber	<ul style="list-style-type: none"> ■ <i>Lateral rectus</i>: Rolls eyeball laterally, innervated by CN VI. ■ <i>Medial rectus</i>: Rolls eyeball medially, innervated by CN III. ■ <i>Superior oblique</i>: Rolls eyeball downward and laterally, innervated by CN IV. ■ <i>Inferior oblique</i>: Rolls eyeball upward and laterally, innervated by CN III.
Posterior chamber	<ul style="list-style-type: none"> ■ The area between the cornea and the iris. ■ Filled with a clear fluid called aqueous humor.
Aqueous humor	<ul style="list-style-type: none"> ■ Space between the iris and the lens. ■ Contains aqueous humor. ■ Clear fluid produced continuously by the ciliary processes. ■ Circulates between the posterior and the anterior chambers and is drained by a collecting vessel called Schlemm's canal.
Ciliary body	<ul style="list-style-type: none"> ■ Supplies the lens and cornea with nutrition. ■ A muscle attached around the lens via zonules that change the shape of the lens when constricted. ■ Produces aqueous humor and some vitreous humor.
Schlemm's canal	<ul style="list-style-type: none"> ■ Pathway in the sclerocorneal region that provides drainage for aqueous humor. ■ Opening for canal is located in the anterior chamber. ■ Pressure in the eye is regulated by filtration of aqueous humor through Schlemm's canal, where it is recirculated into the blood supply of the conjunctiva.
Lens	<ul style="list-style-type: none"> ■ A biconvex disc through which light and images pass. ■ Lies behind the pupil and is partly covered by the iris. ■ The shape of the lens flattens when focusing on far objects and returns to its natural convex shape when focusing on near objects (accommodation).
Zonules	<ul style="list-style-type: none"> ■ Tough fibers that hold the lens in place and facilitate adjustment of the lens shape for seeing near and far objects.
Vitreous humor	<ul style="list-style-type: none"> ■ The area behind the lens of the eye. ■ Filled with a clear, jelly-like substance called vitreous humor that serves to cushion the retina and maintain the eye's shape and pressure.
Choroid layer	<ul style="list-style-type: none"> ■ Highly vascular middle layer of the eye, which is continuous with the posterior wall of the posterior chamber. ■ Provides nourishment for the eye and dissipates heat produced by the light energy that enters the eye.
Retina	<ul style="list-style-type: none"> ■ The sensory network of the eye located on the inner surface of the posterior chamber. ■ Retina contains photoreceptors (rods and cones) that transform light and color impulses into electrical impulses and transmit them to the brain.
Optic disc	<ul style="list-style-type: none"> ■ Yellow-orange disc where nerves and blood vessels enter and leave the eye. ■ Serves as a reference point for measuring and locating other structures on the retina.
Physiological cup	<ul style="list-style-type: none"> ■ Lighter whitish/yellow area of optic disc area located near center of the disc. ■ Diameter of cup is not more than half of the optic disc's diameter.
Macula	<ul style="list-style-type: none"> ■ An indistinct, darker, avascular area on the retina responsible for night, color, and central vision and motion detection.
Fovea centralis	<ul style="list-style-type: none"> ■ Located 2 DD temporally from the optic disc. ■ A slight depression in the center of the macula that appears as a light reflection during ophthalmoscope examination.
Arteries	<ul style="list-style-type: none"> ■ Contains a high concentration of cones for color and bright light vision. ■ Emerge from the center of the optic disc. ■ Supply retina with oxygen and nutrients. ■ Arteries are about one-third smaller than veins, are brighter red, and display a light reflex as a thin streak of light reflecting off each artery.
Veins	<ul style="list-style-type: none"> ■ Emerge from the center of the optic disc. ■ Remove waste products from retina. ■ Are larger, darker, and duller in color than arteries and show no light reflex.




FIGURE 12.6. A. Normal eye. B. Myopia. C. Hyperopia. D. Astigmatism.

Several conditions result from variations in how or where the light rays entering the eye converge and focus. People with **myopia** (nearsightedness) need to hold objects close to the eye to see them clearly. In the myopic eye, the globe is elongated, causing light rays to focus in front of the retina. When the globe is shorter than normal, light rays focus at a point beyond the retina, producing a condition called **hyperopia** (farsightedness). People with hyperopia must move close objects farther away to see them clearly. **Astigmatism** is an irregular curvature of the lens or cornea that causes light rays to scatter, blurring images on the retina (Fig. 12.6).

The **retina** is rich in sensory neurons, which are necessary for reception and transmission of accurate images. The retina contains specialized nerve cells called **rods**, which are sensitive to dim light, and **cone cells**, which are sensitive to bright light and color. From the rods and cones of the retina, the visual image is transmitted to the **optic disc**, where the nerve fibers of the retina converge to enter the optic nerve.

Nerve fibers from the optic disc join to form the **optic nerve**. The neural impulses are then transmitted by the optic nerve to the **optic tract** and **optic radiations**, where they are interpreted by the visual cortex. Nerve fibers from the nasal portion of each eye cross over to the opposite side of the brain at the **optic chiasm**. Fibers from the temporal portion of the retina of each eye do not cross over before being received by the visual cortex in the occipital lobe of the brain (Fig. 12.7).

Interaction With Other Body Systems

The functions of the eyes are interconnected with those of the cardiovascular, musculoskeletal, and neurological systems.





FIGURE 12.7. Optic pathways and visual fields.

**NEUROLOGICAL**

CN II, III, IV and VI are responsible for vision and eye movement.

Changes in pupils can reflect changes in intracranial pressure.

Vision changes can reflect underlying neurological problems such as MS.

ENDOCRINE

Hyperthyroid disease may be seen in the eyes as exophthalmos.

Diabetes can cause visual changes that can be seen on fundoscopic examination.

**LYMPHATIC/
HEMATOLOGICAL**

Anemia can be seen as a pale conjunctiva.

RESPIRATORY

Pulmonary problems that result in hypoxia can be seen in the conjunctiva.

CARDIOVASCULAR

Xanthelasma (fat deposits) can be seen on the eyelids.

Vascular changes can be seen on fundoscopic examination.

DIGESTIVE

Jaundice from liver disease can be seen in the sclera.

MUSCULOSKELETAL

Eye muscles are needed for movement.

Rheumatoid arthritis can cause iritis and scleritis.

INTEGUMENTARY

Skin covers the lids.

Lashes protect the eyes.

URINARY

Renal disease can result in periorbital edema.

Relationship of the Eyes to Other Systems

The Cardiovascular System

The **optic fundus** is the only area in the body where blood vessels can be directly observed without using invasive techniques. Use of an ophthalmoscope provides direct visualization of the **optic disc**, where the vessels that supply the retina emerge. Changes in the optic disc, blood vessels, macula, and general background of the retina can reveal systemic problems with circulation as a result of chronic hypertension (HTN) or diabetes. Or they can reveal localized circulatory problems that occur with glaucoma, increased intracranial pressure, and other neurological problems.

The Musculoskeletal System

Movement of the eyes in a parallel or conjugate manner is made possible by the coordinated movement of the **extraocular muscles**. Each of the six extraocular muscles is responsible for rotating the eyes in a specific direction and maintaining the eyes' conjugate movement. Each extraocular muscle is innervated by a specific cranial nerve (CN). The superior rectus, inferior rectus, medial rectus, and inferior oblique muscles are innervated by CN III. Damage to CN III can therefore result in limited range of movement in the upward, downward, nasal, and upper diagonal fields of vision. The remaining extraocular muscles, the lateral rectus and superior oblique, provide movement of the eye in the temporal lateral and nasal inferior direction, respectively. Damage to the nerves that innervate the lateral rectus muscle (CN VI) and the superior oblique muscle (CN IV) can result in limited eye movement in the corresponding directions.

Movement of the eyelid is controlled by another set of muscles, the orbicularis oculi and the levator palpebrae. The orbicularis oculi encircles the eyelids and is innervated by CN VII. Damage to the orbicularis oculi muscle or the part of the cranial nerve that innervates it results in an inability to close the eyelid completely. The ability to raise or open the eyelid depends on an intact CN III, which innervates the levator palpebrae muscle.

Constriction and relaxation of the muscular tissue of the iris and ciliary body allows **visual adaptation**. In a darkened environment, contraction of the smooth muscle of the iris causes the aperture of the iris or pupil to dilate. As a result, more light enters the retina and night vision is enhanced. In brightly lighted environments, the retina does not need to receive additional light; as a result, the pupil is constricted.

The ciliary body, located posterior to the outer edge of the iris, alters the shape of the lens and allows the eye to adjust to near or far objects, an occurrence referred to as **accommodation**. Constriction of the ciliary body results in flattening of the normal convex shape of the lens. Flattening of the lens facilitates the eye's ability to focus on objects in the distance. Relaxation of the ciliary body allows the lens to assume its normal convex shape and fa-

cilitates focusing on near objects. Both the iris and the ciliary body can be affected by damage to CN III, resulting in pupil dilatation (**mydriasis**) and loss of accommodation.

The Neurological System

The ability to see images depends on an intact visual pathway. From the time when an image is received on the retina to the time when it is interpreted by the visual cortex, the neurological system plays a key role. Damage to the retina can result in diminished visual acuity or diminished color and night vision. Damage to the optic nerve, a source of retinal nervous tissue, can result in similar visual changes. Damage to specific points along the optic track or to the visual cortex can produce deficits in corresponding visual fields (Fig. 12.8).

The neurological system also innervates the extraocular eye muscles, which control the movement of the eyes, and the muscles of the eyelids, which control the opening and closing of the eyes. The cranial nerves responsible for innervation of each of the six extraocular muscles include CN III (oculomotor), CN IV (trochlear), and CN VI (abducens) (Fig. 12.9). Inability of these muscles to function properly is largely caused by damage to the nerves that innervate the muscles.

Increased intracranial pressure from intracranial tumors, head injuries, or severe intracranial hemorrhage may impinge on CN II (optic), CN III, CN IV, or CN VI, causing specific eye changes. The optic nerve innervates the retina and is surrounded by a meningeal sheath that is continuous with the meninges of the brain. When intracranial pressure increases, the pressure is transmitted from the brain to the optic disc, where swelling occurs. Pressure on a specific part of the optic nerve tract can produce visual loss (**hemianopia**) on the ipsilateral (same side) or contralateral (opposite side) visual field, depending on the location of the injury or lesion.

Developmental, Cultural, and Ethnic Variations

Infants

Several variations may be noted in the eyes of infants. The shape, slope, spacing, and color of the eyes should be noted. Normal shape is oval. Slope is determined by drawing an imaginary line through the inner canthus to the occiput. Except in people of Asian descent, the slope line transects the outer canthus. Measurement of the distance between various structures of the eye can be plotted on a growth chart. Normal spacing measurements are plotted between the 10th and the 90th percentiles.

Infants usually open their eyes when held upright, permitting inspection of the iris, pupil, and sclera. The color of the iris after birth is normally blue-gray in light-skinned infants and brown in darker-skinned infants. Permanent eye color is usually established by 9 months of age. White




FIGURE 12.8. Optic nerve pathways, visual fields, and selected associated visual defects.




FIGURE 12.9. Cranial nerves and eye movements (6 Cardinal Fields of Gaze).

specks in the iris called **Brushfield spots** can be a normal variant or a sign of Down syndrome. Edema of the eyelids and irritation of the conjunctiva may be caused by birth trauma or silver nitrate prophylaxis. The sclera is very thin at birth, so it may have a slightly blue undertone.

A gross assessment of visual acuity is made by testing for pupillary light reflexes and also by noting the infant's behavior. The pupils should normally constrict to light. After 3 weeks, if no pupillary light reflex is present, blindness is indicated. However, the presence of pupillary reaction alone does not confirm an infant's ability to see. A blink reflex in response to bright light and observing the infant for ability to follow objects or light with the eyes confirm that some degree of vision is intact. By 2 to 4 weeks, an infant should be able to fixate on an object and, by 1 month, to fixate and follow an object. An infant's visual acuity is usually about 20/200; 20/20 vision is usually achieved by school age.

During the first 1 to 2 months, infants' eye movements are often disconjugate (not working in unison), making screening for strabismus difficult. Persistence of disconjugate eye movements after this time may indicate strabismus and warrants referral to a specialist. In infants, the fundoscopic examination is difficult but still important. The internal structures of the eye should be examined regularly during the first few years of life. One of the first things to note is the presence of a **red reflex**, which is a normal finding. Absence of a red reflex may indicate congenital cataracts or retinal detachment. The general background in infants is typically paler than that in adults because the blood vessels to the area are not fully developed. The macula is also not fully developed until about 1 year of age.

Toddlers

Visual acuity in toddlers is determined by the **Allen test**, which uses picture cards of seven common objects. The child should successfully identify three of the seven objects at a distance of 15 feet. If the child cannot do so, move closer until he or she is able to do so.

The corneal light reflex can provide an initial, rapid screening test for strabismus that can be followed by additional measures as the child grows. Untreated strabismus

can lead to permanent visual damage. Eventually, the brain suppresses information from the affected eye, and visual acuity in that eye deteriorates.

Preschool Children

Between the ages 3 and 5, the **Snellen E chart**, which uses various sizes of Es facing in different directions, can usually be used to determine visual acuity. Normal visual acuity for a 3-year-old is approximately 20/40 or better. By the time the child is 4 years old, visual acuity should be about 20/30 or better.

School-Age Children

By the time a child is about 5 to 6 years old, normal visual acuity approximates that of the adult—20/20 in both eyes. You should continue using the Snellen E chart until the child has acquired reading skills and can easily verbalize the letters seen on the Snellen chart. Some degree of myopia (nearsightedness) often occurs during adolescence.

Children should be screened for defects in color perception (colorblindness) between 4 and 8 years of age. Though many forms of color blindness exist, most cases involve inherited recessive X-linked traits in males that affect the ability to distinguish red and green.

Older Adults

Many changes in the structure and function of the eye occur with aging. Both central and peripheral visual acuity may be diminished with advanced age. Changes in near vision occur around the fourth and fifth decades, often resulting in decreased ability to focus clearly on near objects (**presbyopia**). The adult may compensate for these changes by holding near objects farther away.

External structures of the eye also undergo significant changes with advanced age. Tissues of the eyelids lose elasticity and fatty deposits, causing the eyes to appear sunken. The lower eyelid may sag away from the globe. The latter condition, called **ectropion**, is significant because the **punctum**, which drains the tears, is no longer in contact with the globe, resulting in constant tearing. The laxity that develops in the eyelids may also lead to an inward turning of the eyelids, referred to as **entropion**. With entropion, the punctum also may not be able to drain tears. In addition, the eyelashes may rub the conjunctiva and cornea, causing pain and injury to the cornea. Older adults also may experience dry eyes because of a decrease in tear production.

Changes in the internal structures of the eye are also common with aging. The lens becomes more opaque and yellowish, obscuring the transfer of light rays to the retina. This clouding of the lens is referred to as **senile cataract**. **Arcus senilis**, a white opaque ring around the edge of the cornea resulting from fat deposits, is a common benign finding. The older adult's pupil size at rest is generally smaller than that of younger adults. Pupillary reaction to

light and accommodation slow because of decreased ability to constrict and relax. The general background is paler, and the blood vessels of the eye may show signs of the same atherosclerotic processes that are occurring elsewhere throughout the body. Visual fields may be less than normal. Other visual changes that occur with aging reflect degeneration of the rods and cones. Color vision may be less vivid as a result of degeneration of the cones, and night vision may be impaired because of degeneration of the rods.

Several eye diseases also occur more commonly in older adults. Macular degeneration and glaucoma, the two leading causes of blindness in older adults, show a significant increase with aging.

People of Different Cultures and Ethnic Groups

Differences in physical characteristics of the eye and differences in the risk of certain eye diseases are found in various ethnic groups. People of Asian origin typically have an epicanthal fold at the medial canthus, giving the eyes an almond-shaped appearance. The outer canthus also may slant in an upward direction. In African Americans and others with normally dark skin, brown-pigmented spots on the sclera, referred to as **muddy sclera**, are common. In dark-skinned people, the color of the optic disc is also typically darker orange, and the retinal background is darker red than in fair-skinned people. An African American person's sclera also may have a blue-gray appearance or a yellowish cast at the peripheral margins.

The incidence and severity of glaucoma is greater in African Americans than in people of other races. Cataracts also occur with greater frequency in people living in sunny climates.

Introducing the Case Study



Laura Scammel is a 55-year-old secretary whom you are seeing in the clinic for the first time. She has a history of hypertension (HTN) for 10 years and was diagnosed with type 2 (non-insulin-dependent) diabetes mellitus (NIDDM) 3 years ago. She has experienced blurred vision, which has gradually worsened over the past several months. She takes the oral hypoglycemic agent Glucatorol as well as Lasix and Captopril for HTN. She admits to having difficulty sticking to a diabetic, low-sodium diet and taking her medications regularly.

Performing the Eye Assessment

Assessment of the eye includes taking a thorough health history and performing a physical examination. Data obtained are combined and analyzed to determine the patient's existing health status and to identify potential health risks and disorders of the eye.

Health History

The health history addresses the patient's personal and family history of eye diseases and diseases that affect the eye. A comprehensive health history also allows the nurse to identify areas of the physical examination that require more or less depth. The health history will include biographical data, current health status, past health history, family history, a review of systems, a psychosocial profile, and a detailed eye history. If time is an issue and you are unable to perform a complete health history, perform a focused eye history.

Focused Eye History

If the severity or acuity of the person's condition prohibits obtaining a detailed eye history, take a focused eye history. The person's responses will allow you to obtain the most relevant and crucial information. Follow up any key symptoms with additional questioning, using the PQRST format. Ask these questions:

- Have you noticed any changes in your vision?
- Do you wear glasses or contact lenses?
- Have you ever had an eye injury?
- Have you ever had eye surgery?
- Have you ever had blurred vision?
- Have you ever seen spots or floaters, flashes of light, or halos around lights?
- Do you have a history of frequent or recurring eye infections, styes, tearing, or dryness?
- When was your last eye examination?
- Do you have a history of diabetes or high BP?
- What medications are you currently taking?
- Do you use any prescription or OTC eye drops?

Biographical Data

First, review the patient's medical records, intake surveys, and other sources of data to identify her or his age, occupation, gender, and ethnic background. This information will help you decide what questions to ask and how to interpret subsequent history and examination findings. For example, the way you measure visual acuity and interpret the findings differs greatly with the patient's age. The patient's occupation can be a source of environmental risk for eye injury. Gender may be a factor in certain disorders such as colorblindness, which is more common in males. Knowledge of the patient's race or ethnic group is useful when interpreting many physical examination findings.



CRITICAL THINKING ACTIVITY #1

Considering Mrs. Scammel's age, what eye changes might you expect to see?

Current Health Status

Begin by asking about the person's chief complaint, asking him or her to describe the problem in his or her own words. Use the PQRST format to probe further about any symptoms reported. If the person has an eye problem, focus your questions on the eye symptoms prioritized later.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Vision Loss

Vision loss refers to the inability to see the shape, size, position, or color of objects. Vision loss may be **complete**, resulting in the inability to perceive light from dark, or **incomplete**, causing the person to see objects with varying degrees of detail.



ALERT

Acute primary angle closure (closed) glaucoma is a medical emergency.

Eye Pain

Eye pain is a subjective sensation of discomfort in the eye that may be caused by trauma, irritation, infection, or neurological conditions.

Double Vision

Double vision (diplopia) refers to seeing two overlapping images because of the inability of the eyes to focus on an object and to move in a conjugate manner. Double vision can be caused by a variety of conditions, including diseases of the cerebellum, cranial nerves, and extraocular muscles.

Eye Tearing

Tears are normally produced by the lacrimal gland, located along the upper outer orbit of the eye, and are distributed over the eye by blinking. Tearing is a discharge of clear, watery fluid as a result of the inability of the tears to drain through the punctum and into the nasolacrimal duct. Tearing occurs in a variety of conditions such as infections; irritation; and exposure to chemicals, irritants, or allergens.

Dry Eyes

Dry eyes occur when there is insufficient lubrication of the eye and the bulbar and palpebral conjunctiva become less moist. It often results in a subjective sensation of irritation, a gritty sensation or discomfort, especially during blinking. Dryness occurs from trauma to the eye

surface or facial trigeminal nerve paralysis, in certain systemic diseases, or after taking certain medications.

Eye Drainage

Drainage from the eyes is abnormal and is commonly associated with eye infections or allergies.

Eye Appearance Changes

Changes in the appearance of the external eye, such as in the iris, anterior chamber, and sclera, can signal a variety of problems, including trauma, infection, and systemic disorders.

Blurred Vision

Blurred vision refers to an object's shape and detail being indistinct and fuzzy. It can occur for near objects as well as for distant ones.

Past Health History

This section of the health history focuses on gathering relevant information about the patient's past eye health and any injuries, diseases, or medications that could affect the eyes. The following are specific areas related to the eye that should be explored.

Past Health History

RISK FACTORS/QUESTIONS TO ASK

Childhood Illnesses

- Were you ever diagnosed with strabismus?

Surgeries/Serious Injuries/Hospitalizations

- Have you had surgery for strabismus, cataracts, or vision correction?
- Have you had corneal abrasions or chemical burns?

Serious/Chronic Illnesses

- Do you have a history of HTN, diabetes, hyperthyroidism, or MS?

Immunizations

- Are your immunizations up to date?

Allergies

- Do you have any allergies? If so, what are you allergic to and how do you react?

Medications

- Are you taking any medications, including prescription drugs and OTC or herbal supplements? If so, what is the dosage? How often do you take them? How long have you been using these medications? When was your last dose?

Recent Travel/Military Service

- Have you been exposed to sun or infectious/contagious diseases through recent travel?
- Have you been exposed to irritants, chemicals, or projectiles in military service?

RATIONALE/SIGNIFICANCE

- Uncorrected strabismus results in disuse of the affected eye and eventual blindness in that eye.

- May explain physical findings such as scarring.

- May explain physical findings.

- Vascular changes associated with HTN and diabetes can be seen on funduscopic examination. Exophthalmos is seen with hyperthyroidism (Graves' disease). Visual changes associated with MS include temporary vision loss, patchy blindness (scotomas), diplopia, blurred vision, nystagmus, and optic disc neuritis.

- Communicable diseases such as varicella can affect eyelids and conjunctiva.

- May explain physical findings.

- Determines medications used to treat eye disorders and those that have secondary effects on the eyes. Mydriatic and cycloplegic medications, used for eye examinations and treatment of uveitis and keratitis, dilate the pupil and prevent accommodation, respectively. See *Drugs That Adversely Affect the Eyes*.

- Miotics to constrict the pupil are used to treat glaucoma.
- Drug interactions with secondary effects on the eye include pupillary constriction from narcotics and pupil dilation from marijuana and atropine.

- Sun (ultraviolet) exposure is associated with cataracts. May explain examination findings or identify health promotion or teaching needs.

Drugs That Adversely Affect the Eyes

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Antiarrhythmics	Quinidine	Blurred vision, color perception disturbances, night blindness, mydriasis, photophobia, diplopia, reduced visual fields, scotomas, optic neuritis
	Flecainide	Blurred vision, difficulty focusing, spots before eyes, diplopia, photophobia, nystagmus
Anticholinergic agents	All types	Blurred vision, cycloplegia, mydriasis, photophobia
Anti-infectives	Chloramphenicol	Optic neuritis, decreased visual acuity
Antineoplastics	Cisplatin	Optic neuritis, papilledema, cerebral blindness
	Methotrexate	Conjunctivitis
	Tamoxifen	Retinopathy, corneal opacities, decreased visual acuity
Antitubercular agents	Isoniazid, pyrazinamide, ethambutol	Optic neuritis, decreased visual acuity, loss of red-green color perception, central and peripheral scotomas (ethambutol only)
Cardiotonic glycosides	Digoxin	Altered color vision, photophobia, diplopia, halos or borders on objects (all are signs of potential toxicity)
Genitourinary smooth muscle relaxants	Flavoxate	Blurred vision, disturbed accommodation
	Oxybutynin	Transient blurred vision, cycloplegia, mydriasis
Glucocorticoids	Prednisone, methylprednisolone, dexamethasone	Exophthalmos, increased intraocular pressure, cataracts, increased susceptibility to secondary fungal and viral eye infections
Antipsychotics	All phenothiazines	Abnormal corneal lens pigmentation, blurred vision
	Chlorpromazine	Cataracts, retinopathy, visual impairment
	Quetiapine	Lens changes
Miscellaneous agents	Carbamazepine	Blurred vision, conjunctivitis, transient diplopia, visual hallucinations, lens opacities
	Oral contraceptives (estrogen with progesterone)	Worsening of myopia or astigmatism, intolerance to contact lenses, neuro-ocular lesions
	Isotretinoin	Conjunctivitis, dry eyes, corneal opacities, eye irritation, altered color vision, optic neuritis, photophobia
	Loxapine	Blurred vision
	Pentazocine	Blurred vision, focusing difficulty, nystagmus, diplopia, miosis



CRITICAL THINKING ACTIVITY #2

What part of the assessment would reveal the effects of HTN and diabetes mellitus on Mrs. Scammel's vision?

Family History

When gathering information on your patient's family history, consider familial conditions that may affect the eyes.

Family History

RISK FACTORS/QUESTIONS TO ASK

Familial/Genetic Eye Disorders

- Does anyone in your family have vision problems such as blindness, cataracts, glaucoma, or colorblindness?

Familial/Genetic Disorders That Affect Eyes

- Does anyone in your family have a history of hypertension, diabetes, or multiple sclerosis?

RATIONALE/SIGNIFICANCE

- Person with family history of these problems has greater risk of developing them.
- Color blindness is an X-linked recessive disorder.

- Person with family history of these problems has a greater risk of developing them.

Review of Systems

Changes in the structure and function of the eye may relate to every other system of the body. The review of systems (ROS) will help you identify problems in other

systems that directly affect the eye and allow you to pick up on symptoms that you might have missed in your health history by triggering your patient's memory.

Review of Systems

AREA/SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>General Health Survey</p> <ul style="list-style-type: none"> How have you been feeling? How would you describe your general health? 	<ul style="list-style-type: none"> Changes in usual state of health may reflect underlying problems that can affect the eyes.
<p>Integumentary</p> <ul style="list-style-type: none"> Do you have any rashes or lesions? 	<ul style="list-style-type: none"> Rashes or lesions may be associated with irritation, infection, or allergies that cause eye irritation. May explain physical findings or source of symptoms.
<p>HEENT</p> <ul style="list-style-type: none"> Do you have a history of headaches or migraines? Do you have a history of head trauma? Do you have watery eyes or runny nose? 	<ul style="list-style-type: none"> May be associated with disorders of visual acuity. Visual aura often precedes migraines. Visual changes are early sign of increased intracranial pressure. Associated with environmental allergies or upper respiratory infections.
<p>Respiratory</p> <ul style="list-style-type: none"> Do you have a history of chronic obstructive pulmonary disease or other respiratory problems? 	<ul style="list-style-type: none"> Cyanosis can be seen in the conjunctiva.
<p>Cardiovascular</p> <ul style="list-style-type: none"> Do you have a history of cardiac problems, such as HTN or stroke? 	<ul style="list-style-type: none"> May be responsible for changes in the blood vessels of the eye and the retinal background. Linked to deficits in visual fields or damage to the optic tract.
<p>Gastrointestinal</p> <ul style="list-style-type: none"> Do you have a history of liver disease? 	<ul style="list-style-type: none"> Jaundice may be seen as icteric sclera.
<p>Genitourinary</p> <ul style="list-style-type: none"> Do you have a history of kidney disease? 	<ul style="list-style-type: none"> Periorbital edema may be seen with renal disease.
<p>Reproductive</p> <ul style="list-style-type: none"> Do you have a history of sexually transmitted diseases (STDs), discharge, or lesions? 	<ul style="list-style-type: none"> Cross-contamination may transmit infections to eyes. Newborns may be exposed to STDs during delivery through the birth canal.
<p>Musculoskeletal</p> <ul style="list-style-type: none"> Do you have joint pain, joint deformity, or rheumatoid arthritis? 	<ul style="list-style-type: none"> Episcleritis and keratoconjunctivitis are associated with rheumatoid arthritis.
<p>Neurological</p> <ul style="list-style-type: none"> Do you have MS, myasthenia gravis, trigeminal neuralgia, or Bell's palsy? 	<ul style="list-style-type: none"> MS may produce visual disturbances such as partial blindness, diplopia, or nystagmus as a result of demyelination of nerve fibers. Myasthenia gravis may cause ptosis. Trigeminal neuralgia may cause orbital pain. Bell's palsy may cause orbital dryness and risk of injury caused by inability to blink.
<p>Endocrine</p> <ul style="list-style-type: none"> Do you have hyperthyroidism? 	<ul style="list-style-type: none"> Results in exophthalmos and may be associated with visual acuity changes and diplopia.

AREA/SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Do you have diabetes mellitus? <p>Lymphatic</p> <ul style="list-style-type: none"> ■ Do you have a history of HIV/acquired immunodeficiency syndrome (AIDS), recent immunotherapy, or chemotherapy? 	<ul style="list-style-type: none"> ■ Damages the eye vessels, resulting in retinal hemorrhages, exudate, and changes in visual acuity. ■ Toxoplasma and cytomegalovirus infection of the retina occur frequently in immunosuppressed patients.

Psychosocial Profile

The psychosocial profile may reveal patterns in the patient’s lifestyle that affect health, increase the risk of health problems, or influence adaptation to health prob-

lems. Obtaining a psychosocial profile includes asking the patient about activities of daily living, personal habits, relationships, roles, coping, and home and work environment.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Health Practices and Beliefs/Self-Care Activities</p> <ul style="list-style-type: none"> ■ How often do you have your eyes examined? When was your last exam? ■ If you wear contact lenses, what special eye care practices do you follow? ■ Do you wear eye makeup? <p>Typical Day</p> <ul style="list-style-type: none"> ■ What activities do you engage in regularly? ■ Has your ability to engage in these activities changed recently because of your eye problem? <p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ What did you eat in the past 24 hr? ■ Does your diet include sources of vitamin A (leafy green vegetables, yellow fruits and vegetables, liver, milk, cheeses, and egg yolks)? <p>Activity and Exercise Patterns</p> <ul style="list-style-type: none"> ■ Does your eye problem limit your ability to exercise or perform the usual activities? <p>Recreation/Pets/Hobbies</p> <ul style="list-style-type: none"> ■ Does your eye condition affect your ability to read, watch TV, use equipment such as computers, or engage in other hobbies? <p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ Has your eye condition affected the amount of uninterrupted sleep you get? <p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you use street drugs? 	<ul style="list-style-type: none"> ■ Identifies risk factors and teaching needs of patient. ■ Adults with normal vision should have eye exam every 2 years and begin screening for glaucoma at age 40. ■ Eye makeup can be a source of eye irritation or infection. It should be changed every 6 mo. ■ Helps identify activities that pose a danger to the patient’s health or safety. Or may clarify progression of the condition. ■ Vitamin A deficiency can result in night blindness and keratomalacia. ■ Vitamin A excess can result in exophthalmos. ■ People with diminished visual acuity may be unable to exercise as usual and may be at increased risk for injury from falls or other trauma. ■ Difficulty engaging in hobbies may affect the patient’s sense of well-being and satisfaction. ■ Pain may cause interrupted sleep and increased fatigue. ■ Marijuana causes pupil dilation. ■ Crack cocaine, heroin, and opioids cause pupil constriction.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ Does your occupation expose your eyes to irritants, chemicals, or the sun? ■ Do you work at a computer? 	<ul style="list-style-type: none"> ■ Awareness of occupational hazards helps identify the patient's health risk factors and teaching needs such as wearing protective goggles. ■ Long hours at computer monitors can result in eyestrain.
<p>Environmental Health Patterns</p> <ul style="list-style-type: none"> ■ Are you exposed to chemicals, light, heat, or projectile objects? ■ Are you around environmental irritants, pollutants, tobacco smoke, or allergens? 	
<p>Family Roles and Relationships</p> <ul style="list-style-type: none"> ■ Has your eye condition affected your ability to interact with others or to assume your usual roles (parent, employee, student, etc.)? 	<ul style="list-style-type: none"> ■ Greater risk of injury to the eye. ■ May indicate health teaching needs related to ultraviolet protection and protective goggles. ■ May cause eye irritation, producing tearing and redness of the conjunctiva.
<p>Cultural Influences</p> <ul style="list-style-type: none"> ■ Do you have any cultural influences that affect your healthcare practices? 	
<p>Religious Influences</p> <ul style="list-style-type: none"> ■ Do you have any religious practices/beliefs that influence your healthcare practices? 	<ul style="list-style-type: none"> ■ Eye problems may result in a loss of self-esteem if the patient is no longer achieving satisfaction in roles.
<p>Social Supports</p> <ul style="list-style-type: none"> ■ If you are visually impaired, does it curtail your social activities? 	
<p>Stress and Coping</p> <ul style="list-style-type: none"> ■ If you are visually impaired, how are you dealing with it? 	<ul style="list-style-type: none"> ■ It is important to identify cultural influences that may affect your patient's healthcare practices. ■ It is important to identify religious practices/beliefs that may affect your patient's healthcare practices. ■ Visual changes or losses may cause patient to withdraw from social activities and become isolated. ■ Need to identify patient's ability to deal with vision changes.



Case Study Findings

Reviewing Mrs. Scammel's

Biographical data:

- 55-year-old Caucasian widow and mother of one child, age 30.
- Works full-time as a secretary at a large law firm.
- Has medical insurance coverage through her employer; does not have coverage for dental or vision examinations.
- Catholic religion, of Irish/Italian descent.

Current health status:

- Complains of blurred vision that has increased over the past month. Blurred vision is constant and worsened by fatigue. States that nothing seems to cause the problem or has helped relieve it. Wonders if she needs new glasses.

- Typically works at a computer for at least 4 hours each day.
- Blurred vision has not prevented her from caring for herself at home; however, it is difficult to read the material she is typing on the computer screen, even with reading glasses.
- Denies tearing, eye pain, drainage, double vision, sensitivity to light, or changes in the appearance of the eyes. Currently takes Lasix 20 mg bid, Captopril 25 mg bid, Glucalrol 5 mg bid, and 400 to 600 mg of ibuprofen as needed for joint pain nearly every day.

Past health history:

- History of HTN for past 10 years and poorly controlled adult-onset diabetes diagnosed 3 years ago.

- Denies trauma or surgery affecting the eyes or hospitalizations except for childbirth. One child, no abortions.
- Denies allergies to foods, drugs, or environment.
- Last physical examination 10 months ago; last eye examination 5 years ago.
- Mammogram and Pap smear negative.

Family history:

- One child, age 30, alive and well.
- One brother, deceased 2 years ago at age 50 from a myocardial infarction.
- Father deceased, history of colon cancer, and died of heart attack at age 56.
- Mother, age 78, lives in nursing home and has history of NIDDM, multiple sclerosis (MS), and stroke 4 years ago.
- No family history of glaucoma or vision loss.

Review of systems:

- *General Health Survey*: States that she is usually in good health.
- *Integumentary*: No skin lesions or allergies.
- *Head, Eyes, Ears, Nose, and Throat (HEENT)*: No eye tearing or drainage. No history of headache or head injury. No symptoms of upper respiratory infection.
- *Respiratory*: No history of respiratory disease.
- *Cardiovascular*: No history of cardiac symptoms; positive history of HTN.
- *Gastrointestinal*: No history of liver or renal disease.
- *Musculoskeletal*: No joint pain or deformity or rheumatoid arthritis.
- *Neurological*: No muscle weakness, paresthesia, MS, or myasthenia gravis.
- *Endocrine*: Positive history for type 2 diabetes; no history of thyroid disease.
- *Lymphatic*: No history of human immunodeficiency virus (HIV), chemotherapy, or immunotherapy.

Psychosocial profile:

- Last physical 10 months ago; last eye exam 5 years ago. Gets yearly mammograms and Pap tests.
- Typical day starts at 6:30 A.M., when she rises, showers, and dresses, and has breakfast (bagel and coffee). Leaves for work at 8 A.M. Works 8:30 A.M. to 5 P.M. Monday through Friday. Eats lunch (sandwich) at diner most days. Usually returns home by 6 P.M. unless she runs errands or shops for groceries. Rarely cooks a

full dinner unless her son visits. Watches TV or reads after dinner. Goes to bed at 11 P.M. Spends weekends gardening in warm weather, visiting mother in nursing home, visiting son, running errands, doing housework. Goes to movies with friends occasionally.

- Admits difficulty following prescribed diabetic and low-sodium diets. Tries to eat regularly but has a limited appetite and finds it difficult to cook balanced meals for one person.
- Main source of exercise is gardening and yard work.
- Since loss of her husband, has tried to become more involved in her favorite hobbies: reading, watching TV, and gardening.
- Generally sleeps 8 hours a day without napping.
- Drinks socially three to four times a year. Smoked 1 pack of cigarettes a day since age 18 until 10 years ago when diagnosed with high blood pressure (BP).
- Was a full-time homemaker during her 22-year marriage, but returned to workforce after husband died of a heart attack 2 years ago. Now works full-time as a secretary, with occasional overtime. Reports using computers for an average of 4 hours a day.
- Lives in a rural community just outside of town, in her own home. Commutes 30 minutes to and from work.
- Sees son once a week (lives in a neighboring town). Visits mother in nursing home once a week.
- Is Catholic, but is not currently active in church. Denies any specific ethnic or cultural practices.
- Main social supports are friends from neighborhood and son and daughter-in-law.
- Stress and coping mechanisms include gardening and talking over problems with son and longtime friends from neighborhood.



Case Study Evaluation

Before you proceed with the physical examination of the eye and the ear, document the key information you have learned from Mrs. Scammel's health history.

**CRITICAL THINKING ACTIVITY #3**

Considering Mrs. Scammel's health history, how might her visual problem affect her?

**CRITICAL THINKING ACTIVITY #4**

What special needs might Mrs. Scammel have because of the problems with her eyes?

**CRITICAL THINKING ACTIVITY #5**

What strengths does Mrs. Scammel possess that will help her cope with and manage her health problems?

**CRITICAL THINKING ACTIVITY #6**

What factors from her health history might affect her eyes?

Anatomical Landmarks

Before beginning your physical assessment of the eye, review the anatomical landmarks of the external eye (Fig. 12.10).




FIGURE 12.10. Front view of eyes with fields.

Physical Assessment

During the physical examination, you will assess the functions and structures of the body, including a focused examination of the eye. You will use the information obtained in the health history to guide you and help determine what body structures and functions should be focused upon. After you have explored the patient's key health history information and determined what aspects should be explored, begin the physical examination. The purpose of the physical examination is to identify normal, age-appropriate structures and functions of the eye as well as potential and actual health problems.

Approach

To assess the eye, use the techniques of inspection and palpation. Begin by testing visual acuity and performing other assessments that can be completed while you stand at a distance from the patient. For visual acuity, test and record the findings for each eye separately and then together. Standard abbreviations for recording findings are OD for the right eye, OS for the left eye, and OU for both eyes.

**HELPFUL HINTS**

- Perform the ophthalmic examination last to avoid eye fatigue and tearing, which would interfere with assessment of visual acuity, eye movements, and inspection of the external eye.
- To guard against having the patient memorize the eye chart, keep it out of visual range until ready to use. Test each eye separately, having the patient cover one eye and then the other. Then test both eyes together. If the patient wears glasses or contact lenses, test visual acuity first without corrective lenses, then with corrective lenses.

**TOOLBOX**

To perform the physical assessment, you'll need the following tools: visual acuity charts (Snellen and Snellen E chart, Allen cards, Jaeger chart), color vision chart (Ishihara's cards), ophthalmoscope, penlight, cotton swab, and cotton ball.

Performing a General Survey

Before performing the eye assessment, perform a general survey, noting the patient's overall appearance. Observe

nutritional status, emotional status, and body habitus, noting changes that would relate to the eyes. Then inspect for use of corrective lenses, noticeable visual deficits, and gross eye abnormalities such as ptosis, exophthalmos, edema, and redness.

Also take vital signs. A temperature elevation may indicate an infection and give the eyes a glazed appearance. High BP should alert you to look for vascular

changes when performing the fundoscopic examination.

Performing a Head-to-Toe Physical Assessment

Now examine the patient for more specific signs of diseases affecting other organ systems that might have an impact on the eyes and vision.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

General Health Survey

INSPECT

- Awake, alert, and oriented to time, place, and person.

Integumentary

INSPECT

- Eyelids and eyelashes intact

HEENT

Head

INSPECT/PALPATE

- Head symmetrical, nontender

Ear/Nose/Throat

- No drainage from ears or nose, mucous membranes pink and intact.

Neck

- No neck masses.

Respiratory

- Lungs clear.

Cardiovascular

- Regular heart rate/rhythm.
- +2 pulses.

Gastrointestinal

- Abdomen soft, no hepatomegaly.

Reproductive

- External genitalia pink, moist, and intact.

Musculoskeletal

- Equal muscle strength, full range of motion, no joint deformity.

Neurological

- Movements smooth, sensory function intact.

ABNORMAL FINDINGS/RATIONALE

- *Changes in mental status:* Head injury or conditions that cause an increase in intracranial pressure.

- *Changes in eyes (e.g., in pupils):* Early sign of increased intracranial pressure.

- *Xanthelasma, chalazion, and other skin lesions:* Eyelids are covered by skin, so they are vulnerable to various skin lesions.

- *Changes in pupil reaction and papilledema:* Head trauma, which can result in increased intracranial pressure and is readily detected in the eye.

- *Watery eyes:* Upper respiratory infection or ear infection.

- *Exophthalmos and eyelid lag:* Hyperthyroid disease.

- *Pale, cyanotic conjunctiva:* Hypoxia associated with respiratory disease.

- *Vascular changes (e.g., cotton wool, AV nicking, retinal hemorrhages) on fundoscopic examination:* Cardiovascular disease.

- *Yellow sclera:* Jaundice associated with liver disease.

- Eye infections can be caused by cross-contamination.

- *Ptosis, episcleritis, and keratoconjunctivitis:* Muscle weakness associated with rheumatoid arthritis.

- *Ptosis:* Myasthenia gravis and Bell's palsy.

- *Visual problems and fundoscopic changes:* MS.

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> General state of health Recent infections/illnesses Vision problems	General	<i>Inspect:</i> General appearance Level of consciousness
Integumentary	<i>Ask about:</i> Allergies Rashes	Integumentary	<i>Inspect:</i> Skin lesions on lids Periorbital edema
HEENT	<i>Ask about:</i> Headaches/migraines Head trauma Ear infections Runny nose Thyroid disease	HEENT	<i>Inspect:</i> Masses Abnormalities in oral, nasal and ear structures <i>Palpate:</i> Head Thyroid
Respiratory	<i>Ask about:</i> Breathing problems Lung disease	Respiratory	<i>Inspect:</i> Cyanotic color changes <i>Auscultate:</i> Breath sounds
Cardiovascular	<i>Ask about:</i> History of HTN Vascular disease	Cardiovascular	<i>Measure:</i> B/P <i>Palpate:</i> Pulses <i>Auscultate:</i> Heart sounds Bruits
Abdomen	<i>Ask about:</i> History of liver disease History of renal disease	Abdomen	<i>Palpate:</i> Abdomen Liver
Musculoskeletal	<i>Ask about:</i> Muscle weakness Joint swelling, pain, rheumatoid arthritis	Musculoskeletal	<i>Inspect:</i> Joints for deformities ROM <i>Palpate:</i> Joints for nodules Muscle strength
Neurological	<i>Ask about:</i> History of MS, MG or neurological problems	Neurological	<i>Test:</i> Sensory function CN II, III, IV, VI

Assessment of the Eyes' Relationship to Other Systems

Performing Physical Assessment of the Eye

A comprehensive physical examination of the eye involves assessing the functions, such as vision (distant, near, color, and peripheral), eye muscle functioning, and pupil reflexes, as well as inspecting the external and internal eye structures. The sequence for testing visual acuity progresses from testing done at a distance, such as the visual acuity examination, to observations made at close range, such as the ophthalmic examination. Proceeding in this sequence allows the patient to become comfortable with the nurse before examination at close range is performed. It also allows the nurse to establish the patient's degree of visual functioning, which establishes a baseline for conducting the remainder of the examination. The ophthalmic examination often requires administration of mydriatic or pupil-dilating eye drops.



HELPFUL HINTS

- Mydriatic drops should be administered toward the end of the examination to prevent interference with visual acuity and pupillary reflex testing.
- If the patient wears corrective lenses, first test her or his vision without the lenses, and then with them. This reduces the likelihood that the measurement without correction will be influenced by reading the chart with corrective lenses.



ALERT

If mydriatics are administered for the examination, instruct the patient to wear sunglasses to protect the eyes until the medication wears off.

Visual Acuity Testing

Visual acuity testing involves determining distant, near, peripheral, and color vision. The Snellen eye chart is used to test distant vision in adults and children of school age. The patient stands 20 feet from the chart, covers one eye, and reads the smallest line of print. He or she continues reading successively smaller lines until he or she reads them incorrectly (no more than two mistakes allowed per line) or says that the print is too blurry to distinguish letters. Record the fraction next to the smallest line of letters that the patient read. The top number, or numerator, indicates the distance in feet

from the chart, and the bottom number, or denominator, indicates the distance in feet that a person with normal vision would be able to read the chart. The higher the denominator, the worse the person's distant vision.

Have the person cover the opposite eye and repeat the procedure. After testing each eye individually, test both eyes simultaneously and record the fraction next to the smallest line read. A pocket vision screener may also be used. The letters are scaled down and simulate the Snellen chart, but the card is held only 14 inches from the patient.

Near vision is tested using Jaeger cards, in which lines of text are repeated in progressively smaller fonts. Test each eye separately by having the patient cover one eye and read the smallest line of text while holding the card at a distance of 14 inches. If Jaeger cards are not available, an alternative method is to have the patient read a newspaper and then record how far away he or she holds it.

Color vision is tested using Ishihara's embedded colors test, which consists of a series of cards displaying colored dots that contain an embedded colored figure or number. The patient is asked to identify the figure in each card. An alternative measure is to point to one of the red or green colored bars on the Snellen eye chart and ask the patient what color she or he sees (Fig. 12.11).

Peripheral vision in each eye is measured on two planes—horizontal and vertical—and in four directions—superior, inferior, medial (nasal), and lateral (temporal), using the confrontation test. The patient and nurse stand face to face, about 1 to 1½ feet apart. Ask the patient to fix his or her gaze straight ahead and cover one eye at a time, using his or her hand or an opaque




FIGURE 12.11. Card from Ishihara's embedded colors test. (From Scanlon, V.C., and Sanders, T. [1999]. *Essentials of Anatomy and Physiology*, ed. 3. Philadelphia: FA. Davis, with permission.)




FIGURE 12.12. Positions for peripheral vision testing.

cover. Then wiggle your fingers or bring a pen or other small object from the periphery to the center of the visual field. Tell the patient to say “now” as soon as your hand or the object enters his or her peripheral vision.

Repeat this procedure in each of the four visual fields, moving in a clockwise direction. Be sure to start testing from positions that are outside the normal peripheral vision range; then slowly move your hand or the

object into each of the four peripheral fields. Measure the degree of peripheral vision using the patient’s fixed gaze as a base (Fig. 12.12).

You can also test peripheral vision by comparing the patient’s peripheral vision with yours. This is helpful in detecting gross peripheral deficits, but the method is somewhat subjective because you have to assume that you have normal peripheral vision.

TESTING VISUAL ACUITY

1. Distance

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Have patient stand 20 ft from chart.
- Test each eye separately, having patient cover opposite eye being tested, then together with and without corrective lenses.
- Note smallest line of print patient is able to read with no more than two mistakes.
- Alternate method using pocket vision screener: Have patient hold pocket vision screener about 14 inches from eye and proceed testing as above.



Using Snellen chart



Using pocket vision screener

(continued)

TESTING VISUAL ACUITY

1. Distance (continued)

- Reads letters without correction on Snellen chart or Snellen E chart corresponding with 20/20 in OS, OD, and OU.



HELPFUL HINT

Use Snellen E chart for children who do not know the alphabet, people who cannot read, and those who do not read English.

ABNORMAL FINDINGS/RATIONALE

- *A smaller fraction (e.g., 20/40):* Person standing 20 ft from chart can see lines of print that a normal eye can read at 40 ft. Indicates diminished distant vision or myopia (nearsightedness).

2. Near Vision

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Have patient hold newsprint about 14 inches away and read.
- *Adult reads newsprint or Jaeger cards easily at a distance of 14 inches (recorded as 14/14 OS, OD, and OU).*



Reading newsprint

ABNORMAL FINDINGS/RATIONALE

- *A smaller fraction (e.g., 14/18):* Person must hold print farther away to see clearly because of decreased ability of lens to accommodate to near objects. Indicates diminished near vision, called hyperopia (farsightedness) or presbyopia if it occurs with aging.

3. Color Vision

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Have patient identify color bars on Snellen eye chart.
- Have patient identify figure embedded in the Ishihara chart.
- *Correctly identifies embedded figures in the Ishihara cards or identifies colored bars on the Snellen eye chart.*

ABNORMAL FINDINGS/RATIONALE

- *Inability to detect the embedded number or letter in the Ishihara chart:* Defect in color perception (color blindness). Often inherited in an X-linked recessive pattern, predominantly affecting males. Also results from macular degeneration or diseases that affect the cones that mediate color vision.

TESTING VISUAL ACUITY

4. Visual Fields

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Stand in front of patient, face to face about 1–1½ ft apart.
- Ask patient to fix gaze straight ahead and cover one eye.
- Bring a pen or wiggle your finger in from the periphery from four different fields (superior, inferior, temporal, and nasal).
- Have patient say “now” once fingers or object are seen.
- Measure degree of peripheral vision using patient’s fixed gaze as a base.
- *Alternate method:* Compare your patient’s peripheral vision with yours.
- *Peripheral vision intact in both eyes and all fields.*



Examiner coming in from periphery



Comparing patient with examiner



HELPFUL HINT

Go slowly. If you move your hands too rapidly into the patient’s visual field, it will be difficult to obtain an accurate measure of peripheral vision.

ABNORMAL FINDINGS/RATIONALE

- *Diminished visual fields:* Chronic glaucoma or stroke.
- Peripheral vision intact in both eyes and all fields.

Assessment of the Extraocular Muscles

To perform the corneal light reflex test, instruct the patient to fix her or his gaze straight ahead. Shine a penlight at the bridge of the nose and note where the light reflects on the cornea of each eye. Using the face of a clock as a guide, determine if the light reflex appears at the same clock position in each eye. The corneal light reflex test determines if the eyes are being maintained in a conjugate position.

The cover/uncover test helps determine if there is a weakness in the eye muscles of one or both eyes that can result in disconjugate eye movement. To perform the test, have the patient fix his or her gaze straight ahead. Stand in front of the patient, cover one of his or her eyes with a piece of paper, and observe the uncovered eye for movement indicating re-fixation of the gaze. Remove the

cover and observe the previously covered eye for movement indicating re-fixation of the gaze. Repeat the procedure for the other eye. The gaze should remain steady in each eye throughout the test.

Further testing of the extraocular muscles is done by testing for symmetrical (conjugate) rotation of the eyes, symmetrical movement of the upper eyelid, and nystagmus in the six cardinal fields of gaze test. The six cardinal fields of gaze tests the cranial nerves III, IV, and VI and the extraocular muscle. To perform this test, stand in front of the patient and instruct her or him to look straight ahead and follow your finger as you move slowly through the six cardinal fields. The patient should hold her or his head still and move only her or his eyes. Observe for smooth, symmetrical movement of the eyes and eyelids.

ASSESSING THE EXTRAOCULAR MUSCLES

1. Corneal Light Reflex Test

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Shine light directly in patient's eyes; note position of the light reflection off the cornea in each eye.



HELPFUL HINT

As you shine the light in your patient's eyes, look for the sparkle. That is, the light reflecting off the cornea.



Corneal light reflex test

- Light should be seen symmetrically on each cornea.

ABNORMAL FINDINGS/RATIONALE

- Asymmetrical corneal light reflex:** Weak extraocular muscles or strabismus.



Exotropia (divergent strabismus)



Congenital exotropia

2. Cover/Uncover Test

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Cover patient's one eye and have patient focus on object afar.
- Uncover eye and note any drifting.
- Gaze should be steady when eye is covered and uncovered. No drifting.



A. Cover



B. Uncover

ABNORMAL FINDINGS/RATIONALE

- Shift in gaze:** Weak eye muscle or muscles. If uncovered eye shifts in response to covering opposite eye, covered eye is dominant. If covered eye shifts after being uncovered, that eye is weak.
- Movement of eyes to refocus gaze:** Weakness of extraocular muscles or CN III, IV, and VI, which innervate extraocular muscles.

ASSESSING THE EXTRAOCULAR MUSCLES

3. Cardinal Fields of Gaze Test

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Stand in front of patient and instruct him or her to fix gaze straight ahead.
- Instruct patient to follow your finger or an object such as a pen through the six cardinal fields.
- Note patient's ability to do so, and note any nystagmus.
- *Smooth, conjugate (parallel) movement of eyes in all directions, equal palpebral fissures without eyelid lag. Nystagmus or horizontal jerking eye movements noted only in extreme lateral gaze. Extraocular muscle intact.*



A. Up



B. Side



C. Down



D. Down



E. Side



F. Up

ABNORMAL FINDINGS/RATIONALE

- *Limited or disconjugate movement in one or more fields of gaze, nystagmus in fields other than extreme lateral, ptosis (drooping of upper eyelid), and eyelid lag: Damage, irritation, or pressure on corresponding extraocular muscle or cranial nerve that innervates the muscle.*

Assessment of the External Structures

The next phase of the eye examination involves inspection and palpation of external eye structures. Careful inspection and palpation can reveal a variety of eye disorders as well as systemic disorders that affect the eye.

Testing the pupils to determine reaction to light is important for evaluating neurological function. Increased intracranial pressure from a head injury, tumor, or stroke may manifest in specific pupillary changes. Other conditions such as hypoxia or brain death or the use of certain medications can also affect the pupillary light reflex.

To test pupillary reflex, observe for direct (same side) and consensual (opposite side) response to a focused beam of light. Darken the room if possible. Then, using a penlight, flashlight, or ophthalmoscope light,

shine the light onto one eye as you observe whether the pupil constricts (referred to as a **direct response**). Repeat the procedure and note whether the other eye exhibits a consensual response or constriction to light. Repeat the test for the opposite eye.

Inspection of the anterior chamber of the eye can reveal a variety of conditions including infection, trauma, and risk for glaucoma. The shadow test is useful in identifying a shallow anterior chamber, which is commonly seen in glaucoma. To inspect the anterior chamber using this test, have the patient look straight ahead. Hold your penlight at the temporal side of one eye at a 90-degree angle across the anterior chamber. Without moving the penlight, shine the light across the limbus of the eye, toward the nose. A crescent-shaped shadow on the nasal side of the iris indicates a shallow anterior chamber.

(Text continued on page 355)

INSPECTING THE EXTERNAL STRUCTURES

1. General Appearance

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Note clarity and parallel alignment.
- Eyes clear and bright, in parallel alignment. Note presence of contact lenses.



Checking parallel alignment



A. Soft contact lenses



B. Hard contact lenses

ABNORMAL FINDINGS/RATIONALE

- *Glazed eyes:* Febrile state.

2. Eyelashes

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Note distribution, inversion or eversion.
- *Present and curving outward.*
- *No crusting or infestation.*



Inspecting eyelashes

ABNORMAL FINDINGS/RATIONALE

- *Absence of eyelashes:* Alopecia universalis.
- *Lice or ticks at base of eyelashes:* Infestation.
- *Inflammation:* Blepharitis—inflammation of edge of eyelids involving hair follicles and meibomian glands of eyelids.
- *Inverted eyelashes:* Entropion; can scratch cornea.



Lice



Blepharoconjunctivitis



Entropion

(continued)

INSPECTING THE EXTERNAL STRUCTURES

2. Eyelashes (continued)

■ *Everted eyelashes:* Ectropion; can lead to excessive drying of eyes.



Ectropion

■ Visible *sclera between iris and upper lid (exophthalmos):* Seen in hyperthyroidism and hydrocephalus (“setting-sun eyes”).



Exophthalmos

3. Eyelids

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Note edema, lesions.
- *Upper eyelid normally covers one-half of upper iris.*
- *Palpebral fissures symmetrical. Eyelids in contact with eyeball.*
- *No lesions.*

ABNORMAL FINDINGS/RATIONALE

- *Asymmetry of lids:* CN III damage, stroke.
- *Ptosis of both eyelids:* Myasthenia gravis.



Ptosis

- *Lesions on eyelids:* Basal cell carcinoma, squamous cell carcinoma, xanthelasma, chalazion, hordeolum.



Basal cell carcinoma



Squamous cell carcinoma



Xanthelasma



Chalazion



Hordeolum



Dacryocystitis

INSPECTING THE EXTERNAL STRUCTURES

4. Eyeball

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Note protrusion.
- *Normally doesn't protrude beyond frontal bone. Mild protrusion seen in some African Americans.*

ABNORMAL FINDINGS/RATIONALE

- *Protrusion:* Hyperthyroidism (exophthalmus) or inherited disorders of mucopolysaccharide metabolism.

5. Lacrimal Gland and Nasolacrimal Duct

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Lacrimal glands located below eyebrow, nasolacrimal ducts located on inner canthus of eyes.
- Note swelling, redness or drainage.
- *No swelling, redness, or drainage.*

ABNORMAL FINDINGS/RATIONALE

- *Swelling, redness, drainage, tenderness:* Obstruction or inflammation.

6. Conjunctiva

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- To examine bulbar conjunctiva, gently pull lower lids down.



Pulling eyelid down

- To examine the palpebral conjunctiva, use a Q-tip and gently roll eyelid up.



Rolling eyelid up

- Note color, foreign objects.
- *Palpebral conjunctiva is smooth, glistening, pinkish-peach color, with minimal blood vessels visible. Bulbar conjunctiva over globes are clear, with few underlying blood vessels and white sclera visible.*

ABNORMAL FINDINGS/RATIONALE

- *Red palpebral and bulbar conjunctiva:* Conjunctivitis.
- *Pale pink conjunctiva:* Anemia.
- *Growth or thickening of conjunctiva from inner canthal area toward iris:* Pterygium or pinguecula.



Acute allergic conjunctivitis



Pterygium



Pinguecula

(continued)

INSPECTING THE EXTERNAL STRUCTURES

6. Conjunctiva (continued)

- *Subconjunctival hemorrhage:* Eye injury.



Subconjunctival hemorrhage

- *Benign pigmented congenital discoloration:* Nevus



Nevus

- *Benign growth:* Papilloma.



Papilloma

7. Sclera

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Note color of sclera.
- *Should be smooth, white, glistening. Dark-skinned patients may have a yellowish cast to the peripheral sclera with whiter sclera at the limbus or small brown spots called **muddy sclera**.*

ABNORMAL FINDINGS/RATIONALE

- *Reddish sclera:* Diffuse episcleritis.
- *Bluish sclera:* Osteogenesis imperfecta.
- *Icteric (yellow) sclera at the limbus:* Elevated bilirubin (jaundice).



Diffuse episcleritis

8. Cornea and Lens

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Shine a light on the cornea from an oblique angle.
- Note clarity and abrasions.
- *Corneal reflex:* To test the corneal reflex take a wisp of rolled cotton and gently touch the cornea, or take a needleless syringe filled with air and shoot a puff of air over the cornea, note blinking and tearing.
- *Blink reflex:* Brush your index finger across patient's eyelashes and note blinking.



Corneal reflex with cotton ball



Corneal reflex with puff of air



Blink reflex

INSPECTING THE EXTERNAL STRUCTURES

8. Cornea and Lens (continued)

- *Corneal reflex positive.*
- *Cornea and lens clear, smooth, and glistening. White ring encircling outer rim (arcus senilis) is a normal variant in older adults.*



Normal lens



Arcus senilis

ABNORMAL FINDINGS/RATIONALE

- *Cloudy cornea:* Vitamin A deficiency; infection, which may be accompanied by hypopyon (pus in anterior chamber).
- *Roughness or irregularity of cornea:* Corneal abrasions and ulcers.
- *Yellow ring in outer margin (Kayser-Fleischer ring):* Wilson's disease, characterized by increased copper absorption.
- *Lens opacities:* Cataracts.
- *Negative corneal reflex:* May indicate neurological problem, CN V and VII, but also may be absent or diminished in people who wear contact lenses.

**ALERT**

Corneal scratches or abrasions may not always be visible with the naked eye.



Corneal abrasion



Healing corneal ulcer



Mature cataract

9. Iris

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Note color and shape.
- *Color should be normal variations of blue, green, or brown.*
- *Different iris color or different in one section of iris can be a normal variant.*
- *Shape should be circular.*

ABNORMAL FINDINGS/RATIONALE

- *White specks (Brushfield spots):* Common in Down syndrome.
- *Bloodshot appearance of vessels:* Iritis.
- *New blood vessel on anterior surface:* Diabetes.
- *Different-colored irides (heterochromia iridis):* Previous damage in lighter-colored eye or (rarely) Waardenburg syndrome.
- *Absence of part or all of iris (aniridia):* Congenital problem.
- *Keyhole wedges in iris:* Previous eye surgery.

(continued)

INSPECTING THE EXTERNAL STRUCTURES

10. Pupils

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note pupil size and equality.
- *Test pupillary reaction to light:* Have patient look straight ahead while you bring light in from the side over the eyes
- Note reaction and speed in both eyes.



Checking pupil size



Testing pupillary reaction to light

- *Direct reaction eye receiving stimulus constricts briskly.*
- *Consensual reaction, opposite eye not receiving stimulus directly also constricts.*
- *Should be round and equal bilaterally. Size is larger in children, smaller in older adults. Normal range is 3–5 mm in adults (usually 3 mm).*
- *In about 20 percent of the population, unequal pupils (anisocoria) can be a normal variation. If normal, the pupils react appropriately and the difference is slight, <0.5 mm.*
- *Shape is round.*
- *Normal direct and consensual pupillary response to light is brisk constriction of the pupil to about 1 mm or less.*



Anisocoria



HELPFUL HINT

Pupils constrict in response to looking at a near object, so shine light from temporal side, not directly in front of patient's line of vision. This ensures that pupil constriction you observe is a result of pupillary light reflex and not focusing on a near object.

ABNORMAL FINDINGS/RATIONALE

- *Small, pinpoint pupils (miosis):* Brain injury to the pons; use of narcotics, atropine, and other drugs.
- *Larger, dilated pupils (mydriasis):* Brain herniation, anoxia, or use of marijuana and mydriatic eye drops.
- *Unequal pupils (anisocoria):* CN III (oculomotor) damage, dilated pupil, ptosis, and later deviation of affected eye; unilateral brain herniation or increased intracranial pressure.
- *Unilateral large pupil (tonic pupil) that reacts to light slowly (benign):* Adie's pupil.
- *Unequal pupils; affected pupil small but reacts to light and has ptosis on affected eye related to sympathetic nerve lesion:* Horner's syndrome.



Adie's pupil



Horner's syndrome

INSPECTING THE EXTERNAL STRUCTURES

10. Pupils (continued)

- *Argyll Robertson pupil*: Pupils small and irregular with no reaction to light or accommodation, associated with neurosyphilis.
- *Oval pupils may occur early in head injury*. Irregularly shaped pupils may be caused by certain eye surgeries.
- *Sluggish or fixed pupil reaction to light*: Lack of oxygen to optic nerve or brain or topical or systemic drug effects.
- *Absence of consensual response*: Seen in conditions that compress or deprive those areas of oxygen.
- *Absent light reflex but no change in power of contraction during accommodation (Argyll Robertson pupil)*: Paralysis and locomotor ataxia caused by syphilis.

11. Accommodation (Adjustment of Eye for Various Distances)

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Hold your finger or an object in front of patient from a distance of about 14 inches from patient.
- Instruct patient to focus on finger or object while you move finger or object closer to patient.
- Note convergence of eyes and constriction of pupils as object gets closer.
- *Convergence of eyes and constriction of pupil to focus on a near object and dilation of pupil when looking at a far object. Accommodation may be sluggish in advanced age.*



Checking accommodation

ABNORMAL FINDINGS/RATIONALE

- *One or both pupils fail to dilate or constrict to near or distant objects*: May reflect a problem with CN III or extraocular eye muscles. Patients with hyperthyroidism (exophthalmos) also have poor convergence.
- *Sluggish accommodation not caused by advanced age*: Drug effect, as from anticholinergics.


12. Anterior Chamber

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Have patient look straight ahead as you shine a light from the side across the eye.
- Note clarity and shadowing from iris.
- *Absence of pus or blood.*
- *Absence of a shadow when light is shined across the eye at the limbus area.*
- *Crescent-shaped shadow may be a normal variant.*



Checking anterior chamber



Normal



Shallow

INSPECTING THE EXTERNAL STRUCTURES

12. Anterior Chamber (continued)

ABNORMAL FINDINGS/RATIONALE

<ul style="list-style-type: none"> ■ <i>Pus (hypopyon)</i>: Corneal ulcer or other infection. 	<ul style="list-style-type: none"> ■ <i>Blood in anterior chamber (hyphema)</i>: Trauma or intraocular hemorrhage. 	<ul style="list-style-type: none"> ■ <i>Crescent-shaped shadow on nasal side of iris</i>: Protrusion of iris into anterior chamber from increased intraocular pressure, seen in narrow-angle glaucoma.
		
<i>Hypopyon</i>	<i>Hyphema</i>	<i>Acute angle closure</i>

PALPATING THE EXTERNAL STRUCTURES

1. Eyeball

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

<ul style="list-style-type: none"> ■ Gently palpate below eyebrow and note firmness of eyeball. ■ <i>Globe is firm and nontender.</i> 	
<div style="border: 1px solid red; padding: 5px; display: inline-block;"> <p>ALERT</p> <p>Do not palpate eyeball in patients with eye trauma or known glaucoma.</p> </div>	<p><i>Palpating eyeballs</i></p>

ABNORMAL FINDINGS/RATIONALE

- *Excessively firm or tender globe*: Glaucoma.

2. Lacrimal Glands and Nasolacrimal Ducts

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

<ul style="list-style-type: none"> ■ To palpate glands, gently palpate below eyebrows on brows. 	<ul style="list-style-type: none"> ■ To palpate ducts, glands, gently palpate inner canthus of eyes. 	<ul style="list-style-type: none"> ■ Nontender; gland nonpalpable, no tearing
		
<i>Palpating glands</i>	<i>Palpating ducts</i>	

ABNORMAL FINDINGS/RATIONALE

- *Swelling and tenderness*: Inflammation.



CRITICAL THINKING ACTIVITY #7

Conjunctivitis, or “pink eye,” is an inflammation of the conjunctiva. If your patient has this condition, how would you explain that, even though it is called *conjunctivitis*, the sclera is actually what appears pink?

Assessment of the Internal Structures

The internal structures of the eyes are examined last. Using the ophthalmoscope permits the visualization of the red reflex, optic disc, blood vessels, general background, and macula (Fig. 12.13). Changes in the appearance of these structures can indicate localized eye disorders and eye trauma as well systemic health problems.

The first thing you should see with the ophthalmoscope is a red reflex over the pupil area, which is the reflection of light off the retina. Note the color and clarity of the red reflex. Now gradually move closer to the patient while holding the red reflex in sight. When you are approximately 2 inches from the eye, you should see either blood vessels or the general background of the retina. Follow the blood vessels nasally back to their origin at the optic disc. Now examine the optic disc for color, shape, discreteness of the disc margins, and a paler area near the center called the **physiological cup**. The optic disc is located nasally and has a yellow to yellow-orange color, with sharp distinct edges except for the nasal side, which may be slightly blurred.

Note the size of the physiological cup compared with the size of the optic disc (cup-to-disc ratio). The cup should be half the size of the disc or less. The cup is normally lighter or white in color. Follow the vessels as they leave the optic disc. Note the size of the arteries compared




FIGURE 12.13. Normal fundusoscopic structures of the left eye with fields.

with the veins (AV ratio). Arteries will appear smaller and brighter red than veins and will have a light streak reflecting off them. Also note constriction of the vessels and whether or not any nicking is noted when vessels cross. Inspect the general background for color and lesions. Color should be a uniform yellow-orange-red color, depending on individual skin tone. The lighter the skin tone, the lighter the background. After inspecting the optic disc, vessels, and general background, examine the macula. The macula is a circular area of slightly darker pigmentation located about two disc diameters (DD) temporally from the optic disc. It is the site of central vision and the source of one of the most common causes of blindness, macular degeneration. The macula is difficult to see, especially through undilated pupils. To see it, have the patient look directly into the ophthalmoscope light. This places the macula into direct view.

(Text continued on page 359)

How to Use the Ophthalmoscope

Because the ophthalmic portion of the eye examination is conducted in a darkened room, you need to be familiar with the parts of the instrument and feel comfortable manipulating the lens and aperture discs while holding the scope in one hand. Hold the ophthalmoscope in the palm of your hand with the brow rest facing toward you. Keep your index finger free so that you can use it to adjust the focusing disc and the light aperture disc. When examining undilated eyes, use the smallest white light aperture. This will cause less constriction of the pupils.

Start with the small round light and the lens focus set at “0.” Stand about 1 foot from the patient at a 15-degree lateral angle from his or her line of vision. With the patient’s vision fixed straight ahead, shine the light at the pupil to obtain a red reflex. Hold the scope up to your brow, and move closer to the patient while maintaining the red reflex, until you are only a few inches from him or her. Use your right eye to examine the patient’s right eye and your left eye to examine the patient’s left eye. As you move toward the patient, place your free hand on his or her head near the eye you are examining to stabilize yourself and provide a sense of direction as you approach the eye.

If the retinal structures are not focused, use your index finger to adjust the lens focus. The lens is measured in diopters and indicated as black and red numbers. The black diopters, or positive numbers, are convex lenses that help you to focus closer; the red diopters, or negative numbers, are concave lenses that help you to focus farther away. Turn the lens focus disc clockwise to select a positive diopter for farsighted people (indicated with black numbers in the illuminated lens indicator), or rotate the lens focus counterclockwise to use the negative diopters for nearsighted people (usually indicated in red numbers). The amount and degree of focusing needed depend on the combined refractive state of both you and the patient.

Ophthalmoscope Apertures

APERTURE TYPE	PURPOSE	APERTURE TYPE	PURPOSE
 <p>Small white aperture</p>	Best for examining the undilated eye. Start the eye examination with this setting.	 <p>Blue light</p>	When fluorescein dye is injected into patient intravenously, blue filter enables examiner to see movement of dye into eye vessels. Useful for detecting hemorrhages, leaking vessels, or vessel abnormalities.
 <p>Large white aperture</p>	For general examination of the eye and when pupils are dilated.	 <p>Grid</p>	Used to measure the size or location of lesions.
 <p>Red-free filter</p>	A green light differentiates hemorrhages, which appear black, from melanin, which looks gray. Also differentiates arteries, which appear black, from veins, which appear blue.	 <p>Slit or streak</p>	Used to determine levels or depth of lesions.

PERFORMING AN OPHTHALMIC EXAMINATION

1. Red Reflex

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Stand about 1 ft from patient at a 15-degree angle from patient's line of vision.
- Place index finger on lens wheel and turn wheel as needed to focus.
- Keep free hand on patient's forehead to determine distance/closeness to patient.
- Begin with lens wheel at 0, shining light on patient's pupil to identify red light reflex.



Checking red reflex

(continued)

PERFORMING AN OPHTHALMIC EXAMINATION

1. Red Reflex (continued)



HELPFUL HINTS

- Perform exam in darkened room. Ask patient to focus on object straight ahead.
- Use your right eye to examine patient's right eye, your left eye to patient's left eye.
- **Lens wheel:** Black numbers or positive numbers, convex lenses, help you to focus near; red numbers or negative numbers, concave lenses, help you to focus farther away.
- **Light wheel:** Use small white light for undilated pupils.

■ Positive red reflex over pupil area



Red reflex over pupil area

ABNORMAL FINDINGS/RATIONALE

- **Blackened/opaque areas:** Cataracts
- **Dark spots or shadows that interrupt red reflex:** Opacities in lens or vitreous humor.

2. Optic Disc

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Move closer within inches from patient, turning the lens wheel to focus as needed.
- Identify structures:
 - Disc located nasally, follow blood vessels to disc and note color, borders, and shape.
 - Physiological cup, center of disc, note size and color.
- **Yellow-peach color in fair-skinned people, orange in dark-skinned people. Round or oval shape.**
- **Disc margins sharply demarcated; nasal edge may be slightly blurred.**
- **Cup-disc ratio:** Cup is one-half the DD or less. When visible, cup is located centrally in disc and is a lighter color than rest of disc.



Examining optic disc



Normal fundus

(continued)

PERFORMING AN OPHTHALMIC EXAMINATION

2. Optic Disc (continued)

ABNORMAL FINDINGS/RATIONALE




Acute papilledema



Chronic papilledema




Glaucomatous optic nerve




Optic neuritis



Optic nerve pallor



Hypertensive changes



Diabetic retinopathy



Malignant hypertension



Age-related macular degeneration ("dry")—vision 20/20



Advanced macular degeneration ("wet")—vision 20/400

- *White spots with indistinct edges (cotton-wool spots):* Microinfarctions that occur with diabetes, HTN, lupus, and papilledema.
- *Deep intraretinal hemorrhages (dot hemorrhages):* Diabetes.
- *Flame-shaped, superficial hemorrhages:* HTN.
- *Excess or clumped pigment:* Trauma or retinal detachment.
- *Whitish/gray color:* Partial or complete death of optic nerve (optic atrophy).
- *Blurred margins other than nasally:* Hypertension, glaucoma, or papilledema.
- *Cup-disc ratio greater than 1:2:* Open-angle glaucoma.

3. Blood Vessels

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Arteries and veins originate from disc in pairs, so make sure to note size, color, and crossings.
- *Arteries and veins are paired.*
- *Arteries are bright red with a light reflex; AV ratio is 2:3 or 4:5.*
- *Arteries and veins may cross paths within 2 DD of optic disc if no nicking or interruption of blood flow occurs.*

PERFORMING AN OPHTHALMIC EXAMINATION

3. Blood Vessels (continued)

ABNORMAL FINDINGS/RATIONALE

- *Large arteriovenous (AV) ratio*: Hypertension.
- *Narrowed arteries*: Severe HTN, retinitis pigmentosa, and central retinal artery occlusion.
- *AV crossings more than 2 DD from optic disc or nicking or pinching of underlying vessel*: Hypertension.

4. General Background

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Note color, texture, lesions, and vessels.
- Color ranges from yellow to orange red. Color is even throughout and corresponds with patient's skin pigmentation. Light-skinned patients have lighter retinas; dark-skinned patients have darker retinas.

5. Macula

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Located temporally opposite optic disc; note color, lesions, and vessels.
- *Fovea centralis*: White light reflex in center of macula.
- Indistinct but darker circular area located 2 DD temporal to the optic disc. Size is about 1 DD.
- May see the foveal light reflex, a tiny white dot indicating the ophthalmoscope light shining on the macula.



HELPFUL HINT

The macula is the area of most acute vision and is very sensitive to light, so always examine it last.

ABNORMAL FINDINGS/RATIONALE

- *Hemorrhage or exudate in macula*: Macular degeneration.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Scammel have concluded. The following are your findings:

■ General Health Survey

- Moderately obese 55-year-old female.
- Alert and oriented x4.
- Affect pleasant but anxious.
- Head-to-toe scan revealing no signs of problems in other systems/areas that would affect the eyes except a +S₄.

■ Vital Signs

- Height, 5'2".
- Weight, 170 pounds.
- Temperature, 98.2°F.
- BP, 170/108.
- Pulse, 96.
- Respirations, 24.

■ Eye

- Near vision blurred, Snellen deferred, can see fingers at 10 feet, peripheral

vision 70 degrees temporally, other fields within normal limits.

- Cardinal fields of gaze without nystagmus or eyelid lag.
- Eyelids without lesions, conjunctiva pink without excessive tearing or drainage.
- No periorbital edema; lacrimal gland non-palpable.
- Cornea hazy in OD, clear in OS, smooth bilaterally, arcus senilis bilaterally.
- Iris and pupil within normal limits; pupils equal, round, react to light and accommodation (PERRLA).
- *Ophthalmic exam*—*OD*: Red reflex obscured, disc margins blurred, AV ratio 2:3 with nicking noted 1½ DD inferiorly at 5 o'clock. *OS*: Red reflex present, disc margins blurred, nicking noted superiorly 1½ DD at 2 o'clock. *OU*: Scattered dot hemorrhages throughout, deep pigmentation of macula, no fovea light reflex.



Case Study Analysis and Plan

Now that your examination of Mrs. Scammel is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses. Cluster the supporting data and list possible additional nursing diagnoses.



CRITICAL THINKING ACTIVITY # 8

Identify some nursing diagnoses for Mrs. Scammel, select one, and develop a brief nursing care plan and a teaching plan for her, including learning outcomes and teaching strategies.

Nursing Diagnosis

1. Disturbed Visual Sensory Perception
2. Risk for Injury
3. Impaired Social Interaction

RESEARCH TELLS US

Age-related macular degeneration (AMD) is the leading cause of blindness in adults older than 60 years. AMD affects central vision, and there is no known cure. Environmental factors have been associated with AMD, and recently, studies have confirmed a strong genetic component. Four independent studies identified a commonly inherited variant of the complement factor H (CFH) gene. CFH is a protein that influences the immune system's inflammatory response to diseased and damaged cells.

Further research on AMD is focusing on identifying genetic differences between wet and dry AMD, using chromosome mapping to identify CFH, and investigating the connection of AMD and an immune mechanism. The significance of such research may result in earlier detection of AMD, before the patient becomes symptomatic. Treatment may then begin at an earlier stage, possibly slowing the progression of the disease.

As nurses, it's important to ask about a patient's family history of AMD when reviewing possible risk factors for the disease.

Source: National Eye Institute, *NEI Statement*, "Gene found to increase risk of the most common cause of blindness," March 10, 2005, http://w.nei.nih.gov/news/statements/genes_amd.asp The Foundation Fighting Blindness (available online at <http://www.blindness.org>).

HEALTH CONCERNS

The American Academy of Ophthalmology (2005) recommends a comprehensive vision examination, including visual acuity and glaucoma, by an ophthalmologist:

- Infants at high risk for retinopathy associated with prematurity or those who have a positive family history of eye disorders, such as retinoblastoma, childhood cataracts, and glaucoma, should have an eye exam as soon as possible.
- All children should have an eye exam within first few months of life, and then again at age 6 months and at age 3 years.
- School-age children should be evaluated every 2 years.
- African Americans ages 20 to 39 should be evaluated every 3 to 5 years; every 2 to 4 years for ages 30 to 65.
- Regardless of race, adults ages 40 to 64 should be evaluated every 2 to 4 years, and then every 1 to 2 years starting at age 65.
- Persons with type 1 diabetes with onset before age 30 should have an eye examination 5 years after onset of disease and then annually; with onset after age 30, exam at time of diagnosis and annually. Persons with type 2 diabetes should have eye exam at time of diagnosis and annually.

Common Abnormalities


ABNORMALITY

Astigmatism

- Refraction of light rays diffused rather than sharply focused on retina.

ASSESSMENT FINDINGS

- History of blurred vision.
- Corneal irregularity.



Astigmatism

ABNORMALITY

ASSESSMENT FINDINGS

Cataract

- Abnormal, progressive opacity of the lens.

- Pupil may appear cloudy.
- Red reflex absent or darkened.

**Conjunctivitis**

- Irritation and inflammation of bulbar and palpebral conjunctiva.
- One of most common sources of eye discomfort.
- Typically caused by allergies, viruses, or bacterial infections.

- Manifests as redness of the palpebral conjunctiva and bloodshot sclera.
- Purulent drainage usually present if caused by infection.

**Cranial Nerve Palsy**

- Limited range of motion in one or more cardinal fields of gaze.
- May have eyelid lag, asymmetrical palpebral fissures, or unequal pupil size.

Glaucoma

- Second leading cause of blindness worldwide.
- Two main types: primary open angle glaucoma (POAG) and primary angle closure glaucoma (PACG)

- Changes in optic disc and visual fields.
- Signs and symptoms include protrusion of posterior chamber into anterior chamber, crescent shadow, firm eyeball, blurred optic disc margins, loss of peripheral vision and depth perception, contrast sensitivity.
- May be asymptomatic.
- POAG associated with loss of central vision and elevated eye pressure, but excessive eye pressure is not required for diagnosis.
- Acute PACG may present with acute symptoms including red eye, dilated pupil, nausea, vomiting, eye pain, and halos around lights. Assessment of the anterior chamber with oblique flashlight test often reveals forward bowing of iris.

**Iritis**

- Inflammation of iris that manifests as cloudy or reddened iris with constricted pupil.
- Blood vessels may be visible in iris.

- Symptoms include severe eye pain, tearing, sensitivity to light (photophobia), and in severe cases, diminished visual acuity.
- If untreated, scarring and permanently diminished vision occur.

**Macular Degeneration**

- Diminished visual acuity
- Loss of central vision
- Increased pigmentation of macula

(continued)

Common Abnormalities (continued)


ABNORMALITY

ASSESSMENT FINDINGS

Pinguecula

- Painless yellow nodule caused by thickening of bulbar conjunctiva.

- Usually located nasally.
- Often caused by exposure to sunlight or wind.



Presbyopia

- Occurs with aging.

- Decreased elasticity of the lens.
- Results in decreased ability to focus on near objects.

Pterygium

- Triangular growth of the bulbar conjunctiva from the nasal side of the eye toward the pupil.

- Can obstruct vision if growth occludes the pupil.



Retinal Detachment

- Separation of retinal layer and choroid layer in back of eye

- Signs usually develop gradually.
- Initial symptoms include seeing large numbers of floaters (resolves after several days), flashing lights when the eyes move, and a slowly expanding shadow in the lateral fields of gaze.
- Untreated retinal detachment results in irreversible blindness.

Retinitis Pigmentosa

- Degeneration of retina.
- Begins in childhood and may progress to blindness by middle adulthood.
- Photoreceptors in retina, known as rods, are initially affected, followed by the cones.

- Earlier signs may include night blindness, reduced visual fields, pigmentation of the retina, and macular degeneration.

Sjögren's Syndrome

- Immunologic disorder in which lacrimal, salivary, and other glands do not produce enough moisture

- Causes dryness of the mouth, eyes, and other mucous membranes.
- Damage to external eye tissues, such as the cornea and conjunctiva, may result from excessive and prolonged dryness.

Strabismus

- Axis of eye deviates and does not fixate on an object.
- Also called crossed or wall eye.
- Caused by weak intraocular muscles or a lesion on the oculomotor nerve.
- *Types:* Convergent (esotropia) and divergent (exotropia) strabismus.

- Causes disconjugate vision (one eye deviates from fixated image).
- Initially, diplopia results as each eye transmits the images received.
- Eventually, brain suppresses images received from deviating or weak eye.
- After a period of disuse, visual acuity in weak eye deteriorates and loss of vision results.
- Treatment before age 6 is necessary to prevent permanent damage.



ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the eyes.

Ask the patient the following:

- Do you have...
 - Vision loss?
 - Eye pain?
 - Double vision?
 - Eye tearing?
 - Dry eyes?
 - Eye drainage?
 - Eye appearance changes?
 - Blurred vision?
- Have you noticed any changes in your vision?
- Do you wear glasses or corrective lenses?
- Have you ever had eye surgery? Injury?
- Have you ever seen spots or floaters, flashes of light, or halos around lights?
- Do you have a history of recurrent eye infections, styes?
- When was your last eye exam?
- Do you have a history of diabetes or HTN?
- What medications are you currently taking?
- Do you take any prescribed or OTC eye drops?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam:

- Visual acuity chart (Snellen)
- Color vision chart
- Ophthalmoscope
- Penlight
- Cotton swab/ball
- Gloves if indicated

Position

Sitting

Technique

First, scan your patient head to toe checking for specific signs of diseases affecting other organ systems that might alter the eyes.

Inspection, Palpation, Ophthalmoscopy

Visual Acuity

- Test far (distant) vision.
- Test near vision.
- Test peripheral vision.
- Test color vision.

Extraocular Muscles

- Check parallel alignment.
- Check for lid lag.
- Test six cardinal fields.
- Perform cover-uncover test.

External Structures

Inspect:

- General appearance
- Eyelids and lashes*: Position and distribution
- Eyeball*: Protrusion
- Lacrimal glands and ducts*: Swelling, redness, or drainage
- Conjunctiva (bulbar and palpebral)*: Color, foreign objects
- Sclera*: Color
- Cornea*: Clarity, abrasions, corneal reflex
- Iris*: Color, size, equality, anterior chamber
- Pupils*: Size, equality, reaction to light, and accommodation

Palpate:

- Eyeball*: Consistency
- Lacrimal glands and ducts*: Tenderness

Ophthalmoscopy

Inspect:

- Red light reflex
- Optic disc*: Color, shape, borders
- Physiological cup*: Color, size
- Vessels (arteries and veins)*: Color, size, crossings
- Retina (fundus)*: Color, lesions
- Macula*: color, fovea centralis, lesions

DOCUMENT

Document your findings and develop your plan of care.

S U M M A R Y

- The eyes are complex sensory organs that provide specialized functions crucial to neurosensory development in infancy; to the development of psychosocial, motor, and cognitive skills in childhood; and to the maintenance of those skills in adulthood.
- A thorough health history provides direction for the physical examination, including exploration of factors that may be related to eye health.
- A comprehensive history and physical examination enable early detection and treatment of sight problems.
- Information from both the history and the physical examination is then analyzed to determine appropriate nursing diagnoses.

R E V I E W Q U E S T I O N S

1. Tom Conroy, age 70, is scheduled for a vision exam. Considering his age, which vision change might you expect to find?
 - a. Presbyopia
 - b. Myopia
 - c. Strabismus
 - d. Astigmatism
2. Mr. Conroy has a history of cardiovascular disease and is taking digoxin. What visual symptoms would be associated to digoxin toxicity?
 - a. Nystagmus and scotoma
 - b. Conjunctivitis and dry eye
 - c. Myopia and miosis
 - d. Diplopia and altered color vision
3. Mr. Conroy's far vision results are 20/10 OU (both eyes). How would you explain these findings to Mr. Conroy?
 - a. You can see at 20 feet what a person with normal vision can see at 10 feet.
 - b. You can see at 10 feet what a person with normal vision can see at 20 feet.
 - c. You can see at 20 feet in the right eye and at 10 feet in the left eye.
4. You are testing your patient's extraocular muscles. Which test is best for assessing these muscles?
 - a. Corneal light reflex
 - b. Corneal reflex
 - c. Pupillary reaction
 - d. Red light reflex
5. You are testing far vision on a 3-year-old preschooler. Considering the age of the child, which finding would be considered normal?
 - a. 20/200
 - b. 20/40
 - c. 20/20
 - d. 20/10
6. When testing the corneal reflex, which response would be considered normal?
 - a. Patient's pupils dilate.
 - b. Patient blinks and eyes tear.
 - c. Patient develops a nystagmus.
 - d. Patient's cornea reflects light equally.
7. When testing accommodation, which response would be considered normal?
 - a. Constriction and convergence
 - b. Dilation and convergence
 - c. Constriction and divergence
 - d. Dilation and divergence
8. While testing pupillary response to light, the eye receiving the light stimulus constricts, what is the normal reaction of the other eye?
 - a. Dilating
 - b. Constricting
 - c. No reaction
9. You are assessing your patient's tear ducts. To assess the lacrimal ducts gently palpate:
 - a. Below the eyebrows.
 - b. Below the eyes.
 - c. At the inner canthus.
 - d. At the outer canthus.
10. Which light aperture should be used when performing a fundoscopic exam on an undilated pupil?
 - a. Small white light
 - b. Large white light
 - c. Green light
 - d. Blue light

Anatomy and Physiology Review: The Ear

The main functions of the ears are hearing and equilibrium. Hearing requires an intact and unobstructed external canal, middle and inner ear, vestibulocochlear nerve (CN VIII), and temporal lobe (Figs. 12.14 and 12.15). Sound waves move through the external, middle, and inner ear, where they stimulate the vestibulocochlear nerve and transmit the impulses to the temporal lobe for interpretation. Maintaining normal equilibrium requires proper functioning of the structures in the inner ear.

Structures and Functions of the Ear

The three parts of the ear—external, middle, and inner—contain anatomical structures that work together to allow us to hear.

Understanding Sounds and Sound Waves

Hearing occurs by air conduction and bone conduction of sound waves. Sound waves are characterized by differences in pitch and loudness. **Frequency**, the number of sound waves per second, determines the pitch of the sound. Intensity or loudness is determined by the size of the sound waves. Sound waves can be classified on a

Structures and Functions of the Ear

STRUCTURE	FUNCTION
External Ear	
■ Auricle	■ Cartilaginous external ear whose shape is useful in collecting and conveying sound waves to the middle ear.
■ External ear canal (meatus)	■ One-inch (2.5-cm) canal that connects the auricle to the TM. ■ Outer third is cartilage; inner two-thirds covers the temporal bone. ■ Contains fine hair and glands that secrete cerumen (ear wax). ■ Fine hair and cerumen protect the canal from insects and foreign matter. Cerumen also keeps canal lubricated. ■ Canal protects TM from external environmental factors, such as temperature and humidity changes, that might affect the elasticity of the drum.
Middle Ear	
■ TM (eardrum)	■ Thin, fibrous, pearl-gray membrane that separates the external ear from the middle ear. ■ Slightly concave and pulled in near the center by the malleus. ■ Conducts sound waves to ossicles.
■ Tympanic cavity	■ Small, air-filled space in temporal bone that contains the ossicles. ■ Contains the oval and round windows. ■ The eustachian tube also opens into the middle ear.
■ Eustachian tube	■ The auditory tube from the middle ear to the nasopharynx that maintains equal pressure on both sides of the TM.
■ Auditory ossicles (malleus, incus, and stapes)	■ The smallest bones of the body, they transmit sound waves to the inner ear.
Inner Ear	
■ Bony labyrinth	■ Contains the vestibule, the central chamber of the labyrinth, which in turn contains the utricle and saccule, sacs filled with endolymph and sensory receptors. ■ Three semicircular canals that contain receptor structures called <i>crista ampullaris</i> . ■ The utricle, saccule, and semicircular canals affect equilibrium.
■ Cochlea	■ Shell-like structure containing perilymph-filled scala vestibuli and scala tympani and endolymph-filled scala media. ■ Transmits sound vibrations.
■ Spiral organ of Corti	■ Organ of hearing composed of supporting cells and hair cells that transmit to CN VIII (vestibulocochlear nerve).




FIGURE 12.14. External ear.

continuum from high pitched to low pitched—for example, the high-pitched sounds of a whistle blowing or the screech of chalk on a chalkboard or the low-pitched sounds of a deep drum, thunder, or an explosion. The highness or lowness of pitch is a function of frequency or how fast or slow a sound wave vibrates. The frequency of a sound wave is measured in waves per second or Hertz (Hz). Low-frequency sound waves are interpreted by the brain as low-pitched sounds, and higher frequency waves are interpreted as high-pitched sounds.

The loudness of a sound wave is measured in a scale of units called **decibels (dB)**. The structures of the inner ear are especially sensitive to loudness. Although we tolerate conversation, which is typically around 60 dB, exposure to excessive or repeated noise above 80 dB such as

traffic or machinery noise or exposure to rock music concerts at 120 dB or more can cause damage to the hearing structures and result in permanent hearing loss.

How We Hear

Air conduction is the primary mechanism of hearing; it involves carrying sound waves through the external auditory canal to the tympanic membrane (TM). There, the sound vibrations cause the TM and the **malleus (hammer)**, **incus (anvil)**, and **stapes (stirrup)** bones to move, thus transmitting the vibrations to the inner ear structures.

Bone conduction provides an additional pathway whereby sound waves vibrate the skull bones and transmit the vibrations to the inner ear structures. Both air and bone conduction use a common final pathway involving transmission of the vibrations to the inner ear structures, then on to the cranial nerve and the temporal lobe (Fig. 12.16).

Interaction With Other Body Systems

The Musculoskeletal System

Normal hearing involves proper functioning of certain skeletal structures. Hearing by air conduction involves the transmission of sound waves that vibrate the TM, which in turn transfers the vibrations to the three auditory ossicles. First, the malleus, which is in direct contact with the TM, receives the vibrations. The malleus then transfers the vibrations to the incus and then to the stapes, which transmits them directly to the oval window in the inner ear.

Hearing by bone conduction is much less precise and is dominant only when hearing by air conduction is not possible because of obstruction or perforation of the




FIGURE 12.15. Cross-section of ear.




FIGURE 12.16. Transmission of sound. **A.** Air conduction. **B.** Bone conduction.

TM, fluid behind the membrane, or otosclerosis of the auditory ossicles. Nevertheless, hearing by bone conduction also relies on an intact skeletal structure, the temporal bone, to transmit vibrations to the inner ear. Cooperation between the neurological and the skeletal systems is necessary to provide precise and accurate hearing.


The Neurological System

The primary functions of the ears, hearing and equilibrium, rely strongly on an intact and functioning neurological system. Whether hearing occurs solely by air conduction or bone conduction, the sound waves must eventually travel to the inner ear, where they are picked up by the **organ of Corti**, the sensory organ for hearing. Hair cells in the organ of Corti bend and mediate the vibrations into

electrical impulses that are conducted to the vestibulo-cochlear nerve (CN VIII). The process of picking up sound waves, transmitting them, converting them to electrical impulses, and transmitting them to the brain represents only the first of three levels of auditory functioning.

The second level involves the interaction of both ears and the brainstem in determining the location of the sound in space. When a sound is heard, the vestibulo-cochlear nerve from each ear sends electrical impulses to each side of the brainstem. The brainstem pinpoints the origin of the sound by evaluating factors such as head position, intensity, and timing of the information received from each ear.


The cortex of the brain, specifically the temporal lobe, is involved in the third level of auditory function, interpretation of the sound. Once sound waves have been

**RESPIRATORY**

Because eustachian tube opens into middle ear, upper respiratory infections often precede ear infections, especially in children.

LYMPHATIC

Ear infections may cause lymph node enlargement.


**INTEGUMENTARY**

Skin protects cartilaginous external ear.

Small hairs in external ear protect ear from foreign objects.


**CARDIOVASCULAR**

Tinnitus or vertigo may be associated with cerebrovascular disease.


**NEUROLOGICAL**

VIII, the acoustic nerve, is responsible for hearing.

Abnormal ear position and low-set ears can reflect an underlying problem, such as Down syndrome.

**URINARY**

Ears and kidneys develop at same time in utero, so if ear deformity is present, renal problems may also be present

**MUSCULOSKELETAL**

Ossicles in middle ear (malleus, incus, and stapes) are smallest bones in body.

Gouty tophi deposits can be found on ear lobes.

converted to electrical impulses, they are received in the temporal lobe, where they are identified on the basis of past experience. Impulses are then sent to motor areas of the brain so that an appropriate response to the sound can be made. This level of response allows us to differentiate a telephone ring from a doorbell and take the action needed to either pick up the phone or answer the door. As for maintaining equilibrium, the vestibule of the inner ear plays a crucial role. The vestibule is involved in sensing and perceiving how the body is moving in space. The vestibule also orients the body to maintain a vertical position, stabilizes the position of the head, and helps to maintain the center of gravity.

Disorders of equilibrium can result from two primary causes. The first is chronic loss of vestibular information caused by CN VIII and vestibular hair cell degeneration related to aging or neurotoxicity. The second is distortion of information produced by disruption of the fluid dynamics in the labyrinth of the inner ear. With chronic degeneration, problems with balance are more constant and the person can adapt to them to some degree if the terrain remains constant. However, persons with disorders involving disruption or distortion tend to have symptoms more intermittently and unpredictably, thus preventing the person from adapting as readily. In addition, movement of the head may trigger the vestibular receptors to signal the brain that the head and body are moving when in fact they are not.

Developmental, Cultural, and Ethnic Variations

Infants and Children

The structures needed for hearing are predominantly developed in utero during the first trimester. During this time, damage to the structures from genetic, congenital, and infectious processes can cause hearing problems. Assessment of the ear is especially important in infancy. Observing the appearance of the external ear and noting

behaviors that indicate intact hearing are important ways to screen for potential problems that can cause developmental delays or indicate genetic conditions. Abnormalities in the structure and positioning of the ears are more common in infants who have a hearing deficit. Low-set ears, ears that are positioned at greater than a 15-degree angle, and malformed ears are often associated with genetic disorders and developmental delays.

Infants and children are more prone to inner ear infections than are adults. One reason is that the eustachian tube, which opens from the nasopharynx to the middle ear, is shorter, wider, and more horizontal than in adults, making migration of bacteria from the nasopharynx common. Infants also have greater amounts of lymphoidal tissue surrounding the lumen of the eustachian tube, which can occlude and trap bacteria. By school age, the external auditory canal has assumed a straighter, more adult configuration, which is less prone to infection (Fig. 12.17).

Young and Middle-Aged Adults

Hearing loss caused by excessive or chronic exposure to noise damages the cochlear structures of the inner ear that are involved in conduction of vibrations to CN VIII. Noise-induced hearing loss from exposure to loud music or machinery is the most common cause of hearing loss for the 20- to 40-year-old age group.

Older Adults

Hearing loss in older adults is extremely common and can be associated with sensorineural loss or conductive loss. Hearing loss associated with aging, referred to as **presbycusis**, occurs around the fifth decade and gradually progresses. Typically, presbycusis involves hearing loss for high-pitched sounds, such as consonants, and affects men more often than women.

Older adults are prone to stiffening of the cilia in the external canal. This impedes the transmission of




FIGURE 12.17. A. Infant ear canal. B. Adult ear canal.

sound waves to the TM and causes cerumen to accumulate more readily and obstruct the membrane. Excess accumulation of cerumen impairs hearing by air conduction and is one of the most common correctable causes of conductive hearing loss in older adults.

People of Different Cultures/Ethnic Groups

Several variations in ear structure and disorders of the ear are known to occur in certain ethnic groups. **Otitis media**, or middle ear infection, is a relatively common affliction in infants as a result of bottle feeding and exposure to second-hand smoke. However, the incidence and severity of otitis media for Native American, Hispanic, and Alaskan infants is higher than for infants in the general population.

The characteristics of **cerumen**, or ear wax, also vary with different ethnic groups. Cerumen can be of two types—dry, white, and flaky, as seen in the majority of Asians and Native Americans; or brown, wet, and sticky, as seen in the majority of blacks and whites. People living in highly industrialized communities are routinely exposed to sounds above 80 dB, such as traffic and occupational machinery, and are more prone to hearing loss. In the United States, hearing loss caused by noise affects more than 1 million people and is one of the most prevalent occupational disorders.



Introducing the Case Study

Pablo Arnez is a 50-year-old construction supervisor. He is the sole support of his two teen-aged children (ages 14 and 16) and his physically disabled wife. He has worked in construction for 15 years. He has a clinic appointment to discuss “problems with my ears” that make it increasingly difficult for him to function in the work environment.

Performing the Ear Assessment

Assessment of the ear involves obtaining a complete health history and performing a physical examination. As you perform the assessment, be alert for signs and symptoms of actual or potential problems in the various structures of the ear.

Health History

The health history identifies any related symptoms or risk factors and the presence of diseases involving the ear. It must also detect any other disorders that may affect the ear. Your history will include obtaining biographical data and asking questions about the patient’s current health, past health, and family and psychosocial history. It also includes a review of systems and an ear history. If

your patient has severe hearing problems, you may have to communicate by writing or have a family member answer questions. If you don’t have time to perform a complete health history, make sure to perform at least a focused health history of the ears.

Focused Ear History

If a detailed history is not feasible or appropriate, focus only on the most essential questions. Remember to do a symptom analysis (PQRST) for any positive symptom identified. Ask these questions:

- Do you have problems with your ears such as ringing? Do you have hearing problems?
- Do you have balance problems?
- Do you have drainage from your ears? If yes, how much and what color?
- Have you had recent head trauma?
- Do you have any health problems?
- Are you exposed to noise pollution at work or in your home environment?
- Are you on any prescribed or OTC medications?
- Do you have allergies?

Biographical Data

Review the patient’s biographical data to identify age, occupation, gender, and ethnic background. Use these patient characteristics to guide the types of health history questions that you ask and facilitate your interpretation of subsequent history and examination findings. For example, the techniques used to examine an infant’s ear differ from those used to inspect an adult’s ear. The types of assessments you perform for hearing acuity also vary greatly with the patient’s age. Knowledge of the patient’s cultural and ethnic heritage can provide information on potential language barriers and can facilitate the evaluation of physical examination findings that vary with different groups. Knowing the patient’s occupation can provide information about environmental risk for ear trauma and can help identify areas for health promotion education.

Current Health Status

First, ask the patient to describe the chief complaint in his or her own words. Next, use the PQRST format to investigate further about any symptoms reported. If your patient reports an ear problem, focus your questions on the ear symptoms presented in the following section.

Symptom Analysis

Symptom analysis tables for the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.



HELPFUL HINTS

- If a hearing deficit is known to exist, determine whether the patient can be interviewed orally or if written questioning would be more effective. If you use oral questioning, be sure to face the patient at all times while asking questions and have your face adequately illuminated.
- Hearing impairment in older adults tends to involve high-tone frequency loss. To facilitate communication, use a deeper tone of voice.

Hearing Loss

Hearing loss is a diminished ability to perceive sounds. It may be a result of problems that affect transmission of sound waves to the middle ear (conductive loss) or problems involving interpreting the sound and converting it to neurological impulses (sensorineural loss). Hearing loss may be sudden or gradual, unilateral or bilateral, and may be limited to sounds of certain frequencies or pitches.

Vertigo

Vertigo is a subjective feeling of the body moving or swaying in space or of stationary objects moving or

spinning in space. Vertigo most often results from a disturbance of structures involved with equilibrium and balance, which includes the inner ear and central nervous system.

Tinnitus

Tinnitus is a subjective perception of a high-pitched ringing or buzzing sound in one or both ears. It occurs in certain disorders of the external, middle, and inner ear.

Ear Drainage (Otorrhea)

Drainage from the ear is abnormal. Purulent drainage indicates infection; drainage that is clear or contains cerebral spinal fluid or blood indicates trauma.

Earache (Otalgia)

Ear pain, known as earache or **otalgia**, is a common symptom of ear disorders, particularly ear infections.

Past Health History

During this section of the health history interview, focus on gathering relevant information about the patient's childhood illnesses and injuries, adult illness and injuries, and medication use as they relate to the ears.

Past Health History

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Childhood Illnesses

- Do you have a history of frequent ear infections?
- Were you exposed to second-hand smoke as a child?

- May result in damaged TM (ruptured or scarred) and hearing loss.
- Contributes to more frequent ear infections in infants and children.

Hospitalizations/Surgeries/Serious Injuries

- Have you ever had a severe head injury? A stroke?
- Have you ever received intravenous antibiotics?

- May explain reports of tinnitus and patient's difficulty interpreting speech.
- Use of aminoglycosides, antineoplastics, and loop diuretics may cause hearing loss and tinnitus.

Serious and Chronic Illnesses

- Do you have a history of tinnitus? Vertigo?

- Increased risk of injury from falls.

Immunizations

- Are your immunizations up to date?

- Mumps can cause sensorineural deafness.

Allergies

- Do you have any allergies? If so, what are they and how do you react?

- May explain physical findings.

Medications

- What medications are you taking? What dosage? How often do you take them? For how long?

- Ototoxic drugs in high doses can cause tinnitus, vertigo, and hearing loss. See *Drugs That Adversely Affect the Ear*.

Military Service/Exposure to Irritants

- Were you ever in the military?
- Have you ever been exposed to loud noises?

- Noise pollution from guns can cause hearing loss.

Drugs That Adversely Affect the Ear

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Aminoglycosides	All aminoglycosides	Tinnitus, vertigo, hearing loss
Anti-inflammatory agents	All nonsteroidal anti-inflammatory agents (such as diflunisal, ibuprofen, and indomethacin)	Tinnitus, vertigo, hearing loss
Antimalarials	Quinine	Tinnitus, vertigo, hearing loss
Diuretics	Furosemide and bumetanide	Tinnitus, vertigo, hearing loss (with too-rapid intravenous administration)
Nonnarcotic analgesics and antipyretics	All salicylates and all combination products containing salicylates	Tinnitus, dizziness, hearing loss (with high-dose or long-term therapy)
Miscellaneous agents	Capreomycin, cisplatin, erythromycin, ethacrynic acid, quinidine sulfate, and vancomycin	Tinnitus, vertigo, hearing loss

Family History

When gathering information on your patient's family his-

tory, consider the following familial conditions that may affect the ears.

Family History

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Familial/Genetic Ear Disorders

- Is there a history of hearing loss in your family?
- Does anyone in your family have malformed or malpositioned ears?

- Many genetic disorders manifest with deafness as a feature.
- Unusually shaped ears, low-set ears, or ears set at greater than a 10-degree angle are associated with genetic disorders and possible hearing loss.

Familial/Genetic Disorders That Affect the Ears

- Is there a history of family members with delayed development?
- Do any neurological disorders run in your family?
- Do you have any skeletal problems?

- Tay-Sachs disease in Ashkenazi Jews causes developmental decline in infancy. Hypersensitivity to sound is an early sign.
- Williams' syndrome, a disorder of calcium metabolism, causes failure to thrive and hyperactivity. Affected people are often sensitive to loud noises.
- Charcot-Marie-Tooth disease, a slow, progressive syndrome causing lower extremity muscle weakness around age 10, later progresses to upper extremities. Renal problems often occur in early adulthood, and progressive sensorineural hearing loss begins in the teens.
- People with osteogenesis imperfecta (brittle bone disease) experience conductive hearing loss caused by damage to the ossicles.
- In Hunter's syndrome, growth rate in boys slows after toddler age. Joints stiffen, contractures develop, and facial features become coarse. Deafness is common.
- People with Treacher-Collins syndrome have facial deformities including a small mandible and maxillae. Half of them have small, malformed external ears and middle ears, causing conductive hearing loss.
- Waardenburg's syndrome is characterized by abnormalities of the upper limbs, eyes, and ears. Sensorineural deafness occurs in one-fourth of cases.

Review of Systems

Changes in the structure and function of the ear may relate to every other system of the body. The review of systems (ROS) will help you identify problems in other systems

that directly affect the ear. Remember, the ROS allows you to pick up anything that you might have missed, and the findings will give meaning to the symptoms by relating them to the affected system.

Review of Systems

AREA/SYSTEM/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- How would you describe your general state of health?
- How have you been feeling?

- Change in general health may indicate underlying problem that may affect the ears.

Integumentary

- Do you have any patches of skin that are darker colored?

- Six or more brown patches of skin (café-au-lait spots) in the axillary region may signify neurofibromatosis, commonly associated with acoustic neuroma and hearing loss.

HEENT

- Do you have a history of environmental allergies? Head injury?

- Can explain presence of ear examination findings.

Respiratory

- Have you had any recent respiratory infections or colds?

- Ear infections often follow upper respiratory infections, especially in children.

Gastrointestinal

- Has your appetite decreased?

- Children with ear infections often have a decrease in appetite.

Genitourinary

- Do you have a history of kidney disease?

- The ears and kidneys are formed in utero at similar time periods.

Reproductive

- *For female patients:* Are you pregnant?

- Increased vascularity may cause fullness and diminished hearing.

Musculoskeletal/Neurological

- Do you have any problems with balance or vertigo?

- Indicates problem with inner ear or cerebellum.
- Tinnitus and balance problems may be associated with MS.

Lymphatic

- Have you had a recent ear infection?

- Ear infections may cause lymph node enlargement.

Psychosocial Profile

The psychosocial profile includes asking questions about the patient's lifestyle, including a typical day, personal

habits, medications, sleep patterns, and home and work environment. These factors affect the patient's overall health.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Health Practices and Beliefs/Self-Care Activities

- When was your last physical examination?
- When was your last hearing test and what were the results?
- How do you clean your ears?
- Do you have problems removing ear wax?

- May indicate need for referral.
- Use of Q-tips to clean ears may traumatize auditory canal or drum.
- May need instructions on cerumen removal.

Typical Day

- How does your hearing/ear problem affect your ability to engage in usual activities such as work or school?

- Ability to hear has major influence on how effectively and efficiently a person interacts and communicates with others.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ What did you eat in the past 24 hours? ■ What is your usual diet? 	<ul style="list-style-type: none"> ■ It is important to identify nutritional patterns that may affect your patient's overall health.
<p>Activity/Exercise Patterns</p> <ul style="list-style-type: none"> ■ Do problems with hearing loss, tinnitus, vertigo, and balance adversely affect your ability to do your usual activities? To exercise? 	<ul style="list-style-type: none"> ■ Hearing loss and tinnitus may have a negative impact on ability to engage in sports or activities involving auditory communication. ■ Vertigo and balance problems can make it difficult to navigate in a smooth manner.
<p>Recreation/Pets/Hobbies</p> <ul style="list-style-type: none"> ■ Do your hobbies rely on a high degree of communication with others? ■ Do your hobbies include listening to music or playing musical instruments? 	<ul style="list-style-type: none"> ■ May be unable to enjoy hobbies, resulting in altered coping. Sense of social isolation may increase because of decrease in effective communication. ■ Exposure to loud music can result in permanent hearing loss.
<p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ Does the ear problem cause pain or discomfort that interferes with sleep patterns? 	<ul style="list-style-type: none"> ■ May result in fatigue and sleep deprivation. ■ Tinnitus may interfere with ability to rest.
<p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you smoke or are you exposed to second-hand smoke? 	<ul style="list-style-type: none"> ■ Children have an increased incidence of ear infection when exposed to second-hand smoke.
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ Are you exposed to loud noises or to continuous noise at lower decibels? 	<ul style="list-style-type: none"> ■ Exposure to noise levels higher than 80 dB results in varying degrees of permanent hearing loss depending on dB level and duration of exposure.
<p>Environmental Health Patterns</p> <ul style="list-style-type: none"> ■ Are you exposed to loud noises in your home? 	<ul style="list-style-type: none"> ■ Living in areas of high traffic or where machinery is being used can result in permanent hearing loss.
<p>Family Roles and Relationships</p> <ul style="list-style-type: none"> ■ Has your ear disorder affected your ability to interact with others or to assume your usual roles (parent, employee, student)? 	<ul style="list-style-type: none"> ■ May affect self-esteem if patient is not able to engage in usual roles and relationships.
<p>Cultural Influences</p> <ul style="list-style-type: none"> ■ Do you have any cultural influences that affect your healthcare practices? 	<ul style="list-style-type: none"> ■ It is important to identify cultural influences that may affect your patient's healthcare practices.
<p>Religious Influences</p> <ul style="list-style-type: none"> ■ Do you have any religious practices/beliefs that influence your healthcare practices? 	<ul style="list-style-type: none"> ■ It is important to identify religious practices/beliefs that may affect your patient's healthcare practices.
<p>Stress and Coping</p> <ul style="list-style-type: none"> ■ How do you usually deal with stress and resolve problems? ■ What support systems do you have? 	<ul style="list-style-type: none"> ■ Loss of hearing may cause stress. Identifying coping mechanisms and supports can be used to help the patient deal with hearing impairment.



Case Study Findings

Reviewing Mr. Arnez's

Biographical data:

- Married, 50-year-old father of two (ages 14 and 16).
- Full-time construction supervisor with bachelor's degree.

- Family's main source of financial support; wife is disabled.
- Carries HMO insurance through his employer that covers entire family.
- Native American and Spanish descent; practicing Catholic.

- Source of biographical data is patient, who appears reliable.

Current health status:

- Chief complaint is intermittent fullness and pressure in both ears for past 2 weeks.
- Denies precipitating factors such as recent upper respiratory infection or recent altitude changes.
- Related symptoms include intermittent, bilateral hearing loss described as “like hearing in a tunnel” and intermittent dizziness.
- Rates severity of his ear fullness and pressure as 7 on a 10-point scale.
- Denies ear pain, drainage, or recent trauma to the ears.

Past health history:

- History of frequent ear infections as an infant and young child. Treated with antibiotics.
- Hospitalized once at age 3 with pneumonia and concurrent bilateral otitis media.
- Denies asthma, diabetes, or HTN.
- Received polio, measles/mumps/rubella, tetanus, and diphtheria vaccinations.
- Allergic to sulfa, penicillin, dust, and pollen.
- Current medications include an over-the-counter (OTC) allergy medication.

Family history:

- Father, age 78, has controlled HTN.
- Mother, age 70, diagnosed with MS at age 45, currently in remission.
- One sister, age 40, diagnosed with neurofibromatosis including acoustic neuroma.
- One brother, age 36, recently diagnosed with HTN.
- Patient has two teen-aged girls, ages 14 and 16, both alive and well.

Review of systems:

- *General Health Survey:* Denies fatigue or recent weight loss. Health is usually good.
- *Integumentary:* Denies rashes, lesions, or slow wound healing.
- *HEENT:* Denies ear pain, drainage, tinnitus, or ear trauma. Reports recent upper respiratory infection 4 weeks ago with earache, resolved with antibiotics. History of frequent childhood ear infections and occasional ear infections as an adult, occurring yearly or less. Environmental allergy to dust and pollen.
- *Gastrointestinal:* Denies nausea, vomiting, diarrhea, or recent changes in bowel habits.
- *Genitourinary:* Denies changes in bladder habits.
- *Neurological/Musculoskeletal:* Intermittent problem with balance since onset of ear

pressure. Denies vertigo, muscle weakness, or paresthesia.

Psychosocial profile:

- Last physical examination was 6 months ago. Last ear exam was 4 weeks ago during recent ear infection. Denies ever having a hearing test.
- Typical day starts at 5:30 A.M. on workdays, when he rises, showers, and has breakfast (cereal and coffee). Leaves for work by 6:15 A.M. Works 7 A.M. to 3 P.M. Monday through Friday. Eats quick lunch from sidewalk vendor. Returns home by 4 P.M. most days unless he has errands to run. Cooks dinner for wife and two daughters, then often drives daughters to various school or social activities. Spends evenings watching TV. Goes to bed by 10 P.M. Spends weekends attending sporting events, running errands, chauffeuring daughters. (Wife cannot drive because of disability.)
- Generally eats a well-balanced diet, although lunch at work is rushed. Rarely exercises because of lack of time, fatigue, and intermittent balance problems.
- Ear pressure and fullness have caused intermittent problems in managing his household and socializing. States that he can usually compensate for the hearing loss at home (e.g., by turning up sound on TV and phone handset).
- Ear problems have not caused sleep difficulties. Usually gets 7 hours of sleep a night.
- States that hearing loss causes safety issues at work.
- Denies chronic or recent exposure to loud noise at home or at work.
- Main social supports are family, especially brother and father, and a few close friends at work.
- Has a hectic life with major family responsibilities and many stressors. Talking with friends and family is main coping mechanism.



Case Study Evaluation

Before you proceed with the physical examination, document the key information you have learned from Mr. Arnez’s health history and his related ear history.



CRITICAL THINKING ACTIVITY #9

What factors in Mr. Arnez’s history may explain his hearing problem?




FIGURE 12.18. A. Normal ear attachment. B. Deviated ear attachment. C. Low-set ear attachment.

CRITICAL THINKING ACTIVITY #10
 Considering Mr. Arnez's history, how might this hearing problem affect him?

CRITICAL THINKING ACTIVITY #11
 What strengths does Mr. Arnez possess that will help him cope with and manage his health problems?

tems, so assess for signs in every system. Examination of the ear involves inspection and palpation of the external ear and external auditory canal, followed by tests for hearing and equilibrium. Examination of the auditory canal and TM are done early in the examination to ensure that structures for hearing are intact and unobstructed before testing hearing acuity.

The examination is usually performed with the person in a sitting position. However, when performing the otoscopic examination on a young child, it is better if the child is supine with the head turned to the side being examined. A quiet environment is essential when performing the hearing screening tests.

Anatomical Landmarks

Before beginning your physical examination of the ear, review the internal structures and normal positioning of the external ear (Figs. 12.18 and 12.19).

Physical Assessment

The health history has provided you with clues that will help direct your physical examination. When examining patients who have severe hearing deficits, you may need to explain what you are doing by some other means, such as writing.

Approach

Physical assessment of the ear entails screening for hearing deficits and examining the external ear and TM. Keep in mind that ear problems may relate to other body sys-

TOOLBOX
 The tools needed for examination of the ears include a tuning fork (512 Hz or greater), an otoscope with pneumatic attachment, a thermometer, and a watch. Because examination of the ear involves inspection and palpation, you will need a keen eye and a gentle touch.




FIGURE 12.19. TM, left ear.

Performing a General Survey

Before assessing the ear, perform a general survey. Note the person's overall appearance, including nutritional status, body habitus, and emotional status. Also take vital signs—temperature elevations may indicate an infection. Be especially alert for signs that suggest problems with the ear. Ask yourself these questions:

- Is the person guarding her or his ear? If the patient is a child, is she or he tugging or rubbing the ear? These are signs of an ear infection.
- Is the patient attentive and responding appropriately? Inappropriate responses or inattentiveness may result in hearing deficits.
- Is the patient speaking loudly? People with hearing deficits tend to speak louder.

Performing a Head-to-Toe Physical Assessment

AREA/SYSTEM/NORMAL VARIATIONS

ABNORMAL FINDINGS/RATIONALE

General Health Survey

INSPECT

- Alert, responds accurately when asked to identify person, place, and time.

Integumentary

External Ear

INSPECT

- External ear is pale, peach pink to dark brown color, depending on racial and genetic heritage.

HEENT

Eyes and Ears

INSPECT/PALPATE

- Eyes clear without tearing.

Neck

- Lymph nodes nonpalpable, nontender.

Nose

- Pink or coral-colored nasal mucosa.

- No swelling or drainage.

Mouth/Throat

- Pharynx uniform pink coral without whitish exudate, tonsils +1 adults, +2 or less in children, without exudate.

Respiratory

INSPECT/AUSCULTATE

- Equal bilaterally. No adventitious sounds or cough. Respiratory rate 12–20/min for adults.

Cardiovascular

AUSCULTATE

- Heart: Regular, no extra sounds, no carotid bruits, +2 pulses.

Musculoskeletal

INSPECT

- Muscle strength strong and equal bilaterally. No tremors.

Neurological

INSPECT

- Smooth movement of extremities, steady gait, sense of touch intact.

- Behaviors similar to confusion or disorientation during conversation: Hearing loss.

- Axillary freckling, café-an-lait spots, acoustic neuroma, hearing loss: Neurofibromatosis.

- Tearing eyes, ear fullness: Environmental allergy.

- Palpable, tender, warm lymph nodes: Infection.

- Pale, boggy mucosa with watery drainage: Environmental allergy.

- Purulent or mucoid discharge, often with concurrent otitis media: Upper respiratory infection.

- Reddened pharynx, whitish exudate on pharynx or tonsils, or enlarged tonsils: Upper respiratory infection, often with concurrent otitis.

- Cough with sputum, crackles/wheezes: Upper respiratory infection. (Ear infections often exist concurrently or as a precursor.)

- Tinnitus/vertigo: Cardiovascular/cerebrovascular disease.

- Altered muscle strength, tremors, auditory problems, balance problems: MS.

- Tinnitus/balance and gait problems: MS; balance problems also may indicate problem with inner ear.

- Do you notice any problems with the patient's ability to maintain balance? Balance problems are associated with inner ear problems.

Performing a Head-to-Toe Physical Assessment

Now scan your patient from head to toe. Check for more specific signs of diseases that involve other organ systems and that might affect the ears.

Performing Physical Assessment of the Ear

Now that you have completed your general survey and head-to-toe scan, you can focus on the ear. First, inspect and palpate the external structures of the ear. Then examine the internal ear with an otoscope. Finally, perform hearing acuity tests.

FOCUSED ASSESSMENT OF THE EXTERNAL EAR

1. Inspection

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- **Angle of attachment:** Draw an imaginary line from the top of the helix to the external canthus of the eye, then draw a line perpendicular to the ear and note angle of attachment.
- Note color, size, shape, drainage.
- Size greater than 4 cm and smaller than 10 cm.
- **Normal shape and presence of landmarks:** *Helix, antihelix, antitragus, tragus, and lobule. Absence of pits, creases, or lesions.*
- *Darwinian tubercle is a benign protrusion on upper part of helix.*
- **Position:** *Helix of ear is level with imaginary line drawn through inner and outer canthus to occiput. Ears should be rotated from 0–15 degrees posteriorly.*
- **Condition of skin:** *Intact, no lesions. Note if skin piercings are present.*
- **Color:** *Consistent with skin color.*
- **Drainage:** *Cerumen is the only normal drainage.*



Normal angle of attachment



Darwinian tubercle

ABNORMAL FINDINGS/RATIONALE

- **Ears less than 4 cm vertical height in adults:** Microtia, seen in some genetic disorders.
- **Ears greater than 10 cm vertical height in adults:** Macrotia.
- **Missing or malformed landmarks:** Associated with hearing deficit.
- **Creased earlobe:** Associated with heart conditions.
- **Ear pits or sinuses usually located anterior to the tragus:** Associated with internal ear anomalies.
- **Low-set ears or ears rotated posteriorly more than 15 degrees:** Associated with mental retardation.
- **Drainage:** Bloody drainage can result from trauma and purulent drainage from an infection. Clear drainage may be spinal fluid from a head injury.
- Impacted cerumen.
- **Redness:** Inflammation may indicate infection, fever.
- **Lesions:** E.g., skin cancer from sun exposure.
- Cysts.
- Topic associated with gout.



Congenital ear anomaly



Low-set ears



Impacted cerumen

FOCUSED ASSESSMENT OF THE EXTERNAL EAR

2. Palpation

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- *Ears:* Palpate for consistency and tenderness.
- Always palpate tragus, helix, and mastoid process for tenderness prior to otoscopic exam.
- *Soft and pliable, nontender.*
- *Structures should be nontender, with no nodules or swelling.*



Palpating the ear



Palpating the tragus



Palpating the mastoid



Pulling helix forward

ABNORMAL FINDINGS/RATIONALE

- *Tenderness of mastoid, helix, tragus or pinna:* Ear infections.

Otoscopic Examination

The otoscope illuminates and magnifies the auditory canal and the TM. The auditory canal is assessed for color, lesions, and foreign objects. The TM is assessed for color, intactness, appropriateness of landmarks, and mo-

bility of the drum. Select the largest and shortest speculum that will fit into the patient's ear canal comfortably, and attach it to the head of the otoscope. Usually, a 4-, 5-, or 6-mm, ½-inch speculum is appropriate for an adult ear canal.

PERFORMING AN OTOSCOPIC EXAMINATION

1. Otoscopic Exam

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use shortest and largest speculum that will accommodate ear canal (usually 3, 4, 5 mm, ½ inch).
- Hold otoscope like a pen with the handle up, bracing hand on patient's head. Look through magnifying lens. (*Alternate method:* Hold otoscope like a pen with the handle down and look through magnifying lens.)
- Have patient tilt head opposite direction of ear being examined.
- For an adult, pull helix up and back to straighten ear canal.
- For a child under 2 years, pull ear lobe down.
- Insert ½ inch for adult; ¼ inch for child.



Otoscope insertion with handle up



Otoscope insertion with handle down

PERFORMING AN OTOSCOPIC EXAMINATION

2. External Auditory Canal

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Look before inserting speculum into the external ear canal.
- Insert only in the outer third of the canal.
- Note color, drainage, lesions.
- *No foreign objects. Canal patent. Free of redness and drainage.*
- *Ear wax may be visible. Native Americans and Asians have higher rates of ear infections than Caucasians. Cerumen is white, dry, and flaky in patients of Asian and Native American descent and honey-colored and sticky in whites and African Americans.*
- **Exostosis (“surfer’s ear”):** Abnormal benign growth of bone in external auditory canal—often results from swimming in cold water.



Foreign object



Exostosis

ABNORMAL FINDINGS/RATIONALE

- *Reddened canal:* Otitis externa.
- *Exudate and edema:* “Swimmer’s ear” or external otitis.
- Excessive impacted cerumen in older adults can contribute to conductive hearing loss.



External otitis

3. Tympanic Membrane

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note color, position of landmarks, intactness of drum.
- If unable to visualize, apply more traction on the ear; do not move otoscope.
- If unable to visualize owing to cerumen, a cerumen scoop can be used to clear canal.
- Never irrigate the canal, unless you are sure the drum is intact.
- *Normally shiny, pearl gray, intact, and mobile. TM lighter colored in older adults. Bony landmarks of the umbo (the central depressed portion of the concavity on the lateral surface of the TM) and malleus should be visible. Capillaries may normally be seen on the malleus.*



HELPFUL HINTS

- The ears are mirror images, so the cone of light, a triangular reflex, is at 5 o’clock in the right ear and 7 o’clock in the left ear.
- Assess drum mobility in pediatric patients with a pneumatic tube attachment to the otoscope. Gently use the pneumatic tube to puff air into the canal, and note movement of the cone of light.



Normal TM (left ear)

PERFORMING AN OTOSCOPIC EXAMINATION


3. Tympanic Membrane (continued)

ABNORMAL FINDINGS/RATIONALE

- *Yellowish membrane with fluid and air bubbles visible behind TM:* Serous otitis.
- *Reddish TM with absent or distorted light reflex:* Otitis media.
- *Round/oval dark area:* Perforated TM.
- *White, irregularly shaped area on TM:* Scar tissue.



Serous otitis



Otitis media



Perforated TM



Hemotympanum



Adhesive otitis media



Cholesteatoma


- *Change in position or shape of cone of light reflex and absence or exaggeration of bony landmarks:* Imbalance in middle ear pressure. Bony landmarks more prominent if negative pressure in inner ear, less prominent in infections or conditions in which fluid or pus collects behind membrane.
- *Limited mobility of a drum:* Associated with a bulging drum.
- *Blue to black TM from bleeding (hemotympanum):* Usually results from trauma.
- *Golden-brown TM with caramel-colored drainage that is thick and elastic like rubber cement (secretory, adhesive otitis media, also known as “glue ear”):* Associated with viral infection.
- *Cystic mass of epithelial cells that develops in middle ear (cholesteatoma):* Complication of chronic otitis media.
- Tubes are placed in ear to promote drainage and equalize pressure.



Pressure equalization tube



Tympanotomy



Intact patient pressure equalization tube

Turn the otoscope light on. Have the patient tilt his or her head away from the side you are examining. Always look into the external canal before inserting the otoscope. Two insertion techniques may be used:

- **Technique #1:** Hold the otoscope upside down like a pencil, with the magnifying lens facing the examiner. Pull the pinna of the ear up and back for adults and down for children under age 2. Brace your insertion hand on the patient's head for stabilization. This technique is ideal for children, because if they move the head, your hand moves with it.
- **Technique #2:** An alternative technique is to hold the otoscope handle upright and slowly and gently insert the scope along the axis of the external auditory canal (about $\frac{1}{2}$ inch in an adult and $\frac{1}{4}$ inch in a child). Be careful to enter only the outer third of the ear canal. With the scope inserted, put your eye up to the viewing lens. If you cannot visualize the TM, do not move the otoscope. Instead, apply more traction, pull on the ear, or carefully adjust the angle of the otoscope more toward the patient's nose. Do not release the traction on the ear until the speculum of the otoscope has been removed from the ear. Remove the speculum in the same angle as it was inserted, and then release the traction to the pinna.



ALERT

- Before inserting the otoscope, palpate the tragus, mastoid, and helix. If any of these are tender, insert the scope very carefully.
- Always inspect the external canal for foreign objects before inserting the otoscope. Otherwise, you may inadvertently push an object farther up the canal.
- Insert the otoscope only in the outer third of the canal. The inner two-thirds of the ear canal are over the temporal bone and are very sensitive.

Hearing Tests

Examination of the ear includes conducting hearing tests for high-tone, low-tone, sensorineural, and conductive hearing loss. Typical tests include the following:

- **Watch tick test:** This test determines the patient's ability to hear high-pitched sounds and screen for high-tone hearing loss. To do this test, have the patient obstruct one ear at a time by placing her or his index finger in the external canal. Using a ticking watch, hold it close to the unobstructed ear and slowly move it away until the patient says that she or he can hear the

watch tick (usually about 5 inches [13 cm]). Repeat the test on the other ear.

- **Whisper test:** This test assesses low-pitch or low-tone hearing loss. Again, have the patient obstruct one ear with the index finger. Then stand 1 to 2 feet behind and to the side of the patient's unobstructed ear and whisper three or four unrelated words. Have the patient repeat the words heard. Test the other ear in a similar fashion.
- **Tuning-fork tests:** These tests check for intact sensorineural and conductive hearing. The Weber test assesses lateralization of sound. Sound transmission with the Weber test is both through bone conduction (BC) and air conduction (AC). To perform the test, place a vibrating tuning fork firmly on top of your patient's head or forehead.



HELPFUL HINT

Hold the tuning fork by the stem and tap the prongs gently on your palm or thigh. Do not touch the prongs once they are vibrating; this will dampen or stop the vibrations.

Ask the patient if the vibration sounds the same in both ears or different. Normally, the vibration is heard equally in both ears. If the vibration is louder or more distinct in one ear than in the other, it has lateralized to that ear (positive lateralization). If there is conductive hearing loss, the sound lateralizes to the impaired ear; if there is sensorineural hearing loss, the sound lateralizes to the unaffected or good ear. Conductive hearing loss is usually caused by a problem with the external or middle ear, such as acute otitis media, perforated eardrum, or a blocked canal with cerumen. Sensorineural hearing loss is usually caused by a problem in the inner ear, such as acoustic nerve damage from ototoxic drugs.

Having a conductive loss in the lateralizing ear (e.g., as a result of accumulation of cerumen or a perforated eardrum) means that sound waves are not conducted effectively by AC to the inner ear. Thus, limited or no sound waves are received by AC. The only mechanism remaining for sound wave conduction in the lateralizing ear is by BC. But without competing AC, sound waves will be interpreted by the patient as sounding louder. To test this yourself, create a temporary conductive hearing loss by occluding the external auditory canal and placing the vibrating tuning fork as described. In a person with normal hearing, sound will lateralize to the obstructed ear because you have blocked any competing AC sounds from being transmitted.

With a sensorineural hearing loss, the sound lateralizes to the good ear. The ear with sensorineural deafness is unable to transmit sound waves by AC or BC to the acoustic nerve to the brain. Consequently, the sounds transmitted to the opposite ear will be interpreted as louder, or lateralizing.

The Weber test can detect lateralization, but it cannot differentiate the cause. So if lateralization occurs, you will need to perform the Rinne test to establish if the problem is caused by a conductive hearing loss.

The Rinne test is a timed tuning-fork test used to compare AC and BC. To perform this test, place a vibrating tuning fork on your patient's mastoid process. When it is on the mastoid process, sound transmission is through BC to the inner ear. Ask the patient to tell you when he or she no longer hears the vibrations.

Record the amount of time the patient heard the sound in seconds. When the sound is no longer heard, immediately place the vibrating fork about 1 inch in front of the external auditory canal and continue to time until your patient can no longer hear the sound. When in front of the ear, sound transmission is through AC through the auditory canal, to the TM, middle ear, and inner ear. Avoid touching hairs in front of the ear with the vibrating tuning fork because this may provide a clue that the tuning fork is vibrating. Record the amount of time the patient hears the vibrating tuning fork in seconds. Normally, sound transmission through air is twice as long as sound transmission through bone. If AC time is less than twice as long as BC time, it generally indicates hearing loss by AC. Inability to hear the tuning fork during BC indicates sensorineural hearing loss.

HEARING TESTS

1. Whisper Test (for Low-Pitch Deficits)

TEST TECHNIQUES/NORMAL VARIATIONS

- Have patient cover opposite ear being tested.
- Stand about 1–2 ft behind patient, and whisper.
- Note patient's ability to hear sound.
- *Patient repeats most words whispered in each ear at a distance of 1–2 ft.*



Performing whisper test

ABNORMAL FINDINGS/RATIONALE

- *Inability to repeat words:* Low-tone frequency loss.

2. Watch Tick Test (for High-Pitch Deficits)

TEST TECHNIQUES/NORMAL VARIATIONS

- Have patient cover opposite ear being tested.
- Hold ticking watch within 5 inches from ear.
- Note patient's ability to hear sound.
- *Patient hears tick of a watch in each ear at a distance of 5 inches.*



Performing watch tick test

(continued)

HEARING TESTS

ABNORMAL FINDINGS/RATIONALE

- *Inability to hear watch ticking:* High-tone frequency loss.

3. Weber Test

TEST TECHNIQUES/NORMAL VARIATIONS

- Hold tuning fork by stem.
- Tap tuning fork prongs on palm of your hand.
- Place vibrating tuning fork in the middle of the patient's forehead or on the top of patient's head.
- Note patient's ability to hear sound; note lateralization of sound.
- *Vibrations should be felt or heard equally in both ears.*
- Negative lateralization.



Performing Weber test on head



Performing Weber test on forehead

ABNORMAL FINDINGS/RATIONALE

- *Lateralization of vibrations:* Conductive or sensorineural loss.

4. Rinne Test

TEST TECHNIQUES/NORMAL VARIATIONS

- Place vibrating tuning fork on the mastoid process.
- Count the time until patient can no longer hear tuning fork.
- Immediately bring vibrating tuning fork in front of ear.
- Continue to note the length of time until patient no longer hears sound.
- *The length of time the patient hears the vibrations by AC is normally twice as long as for BC.*
- *The ratio of AC to BC is similar in both ears.*



Performing Rinne test on mastoid



Performing Rinne test in front of ear

HEARING TESTS

4. Rinne Test (continued)

ABNORMAL FINDINGS/RATIONALE

- *AC to BC ratios that differ markedly in each ear:* Unilateral hearing deficit.
- *AC less than twice BC:* Hearing loss by AC, possibly caused by ear wax, otitis media, serous otitis, or damage to the ossicles of the middle ear.

5. Romberg's Test (Tests Inner Ear Vestibular Function)

TEST TECHNIQUES/NORMAL VARIATIONS

- Have patient stand with feet together, eyes opened, and then eyes closed.
- Stand close by in case patient loses balance.
- Note patient's ability to maintain balance.
- Patient stands with feet together and eyes closed. Maintains balance with minimal sway. Negative Romberg. (For more information, see Chapter 21.)

ABNORMAL FINDINGS/RATIONALE

- *Loss of balance:* Inner ear disorder, cerebellar damage, or ingestion of intoxicants.



Case Study Findings

Your general health survey and head-to-toe assessment of Mr. Arnez have concluded. The following are your findings:

■ General Health Survey

- Well-developed, 50-year-old male in no acute distress.
- Alert and oriented x4.
- Pleasant and cooperative.
- Good historian.
- Head-to-toe scan reveals no signs of problems in other systems/areas that would affect the ears except three café-au-lait spots on trunk.

■ Vital Signs

- BP, 142/88.
- Pulse, 78, strong and regular.
- Respirations, 16/minute.
- Ear
 - External ear normal shape, size, and position.
 - Mastoid, tragus, and pinna nontender.
 - External auditory meatus without drainage or lesions.
 - TM pearl gray with scattered white scar tissue and distorted light reflex.
 - No fluid or pus visible behind TM.
 - Hearing tests: Weber, no lateralizing; Rinne, AC 1.5 x BC bilaterally.
 - Dizziness with head rotation.
 - Positive sway with Romberg's test.



Case Study Analysis and Plan

Now that your examination of Mr. Arnez is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mr. Arnez. Use this information to identify a list of nursing diagnoses. Some possible nursing diagnoses are provided below. Cluster the supporting data.


1. Disturbed Auditory Sensory Perception, related to ear pressure and fullness
2. Chronic Pain, related to ear pressure and fullness
3. Risk for Infection

Identify any additional nursing diagnoses.



CRITICAL THINKING ACTIVITY # 12

Once you have identified appropriate nursing diagnoses for Mr. Arnez, select one and develop a nursing care plan and a brief teaching plan, including learning outcomes and teaching strategies.



AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> General state of health Recent infections/illnesses Hearing difficulty	General	<i>Inspect:</i> Ear guarding, inattentiveness, inappropriate responses, balance problems <i>Measure:</i> Vital signs
Integumentary	<i>Ask about:</i> Lesions on ears (sun-exposed area, common site for skin cancer)	Integumentary	<i>Inspect:</i> Lesions, areas of inflammation <i>Palpate:</i> Tenderness
HEENT <i>Head and Neck</i>	<i>Ask about:</i> Pain Swollen nodes	HEENT <i>Head and Neck</i>	<i>Palpate:</i> Lymph node enlargement, sinus tenderness
<i>Nose and Throat</i>	<i>Ask about:</i> Sore throat Nasal congestion	<i>Nose and Throat</i>	<i>Inspect:</i> Oral and nasal structures
Respiratory	<i>Ask about:</i> Cough, congestion Mucus production	Respiratory	<i>Measure:</i> Respiratory rate <i>Auscultate:</i> Lung sounds
Cardiovascular	<i>Ask about:</i> History of vascular disease	Cardiovascular	<i>Measure:</i> Heart rate <i>Palpate:</i> Pulses, carotid thrills <i>Auscultate:</i> Heart sounds, bruits
Musculoskeletal	<i>Ask about:</i> History of gout Hearing problems	Musculoskeletal	<i>Inspect:</i> Ear lobes for tophi
Neurological	<i>Ask about:</i> History of TIAs Balance problems Neurological problems	Neurological	<i>Test:</i> Romberg's

Assessment of the Ears' Relationship to Other Systems

RESEARCH TELLS US

Hearing loss is the third most common chronic condition in older adults, with 25 to 40 percent of adults aged 65 years or older suffering from hearing impairment—and the incidence increases with age. By age 85, 80 percent of adults will have hearing problems. Despite the incidence of hearing problems with the older adult, hearing impairment is underdiagnosed and undertreated, with only 9 percent of internists performing hearing tests and only 25 percent of patients with hearing loss receiving treatment with hearing aids.

A review of the evidence on screening and management of hearing loss of older adults in primary care revealed that use of an audioscope (an otoscope with an audiometer) and a self-administered questionnaire (the Hearing Handicap Inventory for the Elderly—Screening Version) are reliable tests that screen for hearing loss. Even though most hearing loss in older adults is sensorineural due to presbycusis, conductive hearing loss accounts for up to 30 percent of all cases of older adults with hearing loss. Cerumen impaction and chronic otitis media are frequent causes of conductive hearing loss that are treatable.

The effects of hearing loss have a direct effect on ability to communicate, but they also have been associated with depression and functional decline.

The Hearing Handicap Inventory for the Elderly—Screening (HHIE-S) is a 10-item, 5-minute, self-administered questionnaire that assess functional hearing loss and the degree of social and emotional handicap from hearing loss.

Items:

1. Does a hearing problem cause you to feel embarrassed when meeting new people?
2. Does a hearing problem cause you to feel frustrated when talking to members of your family?
3. Do you have difficulty hearing when someone speaks in a whisper?
4. Do you feel handicapped by a hearing problem?
5. Does a hearing problem cause you difficulty when visiting friends, relatives, or neighbors?
6. Does a hearing problem cause you to attend religious services less often than you would like?

7. Does a hearing problem cause you to have arguments with family members?
8. Does a hearing problem cause you difficulty when listening to TV or radio?
9. Do you feel that any difficulty with your hearing limits or hampers your personal or social life?
10. Does a hearing problem cause you difficulty when in a restaurant with relatives or friends?

Responses include:

- Yes (4 points)
- Sometimes (2 points)
- No (0 points)

Interpretation of score:

- 0 (no handicap) to 40 (maximum handicap)
- 0–8 = 13% probability of hearing impairment
- 10–24 = 50% probability of hearing impairment
- 26–40 = 84% probability of hearing impairment

The HHIE-S has reported sensitivity and specificity ranging from 0.63 to 0.80 for score 10 and above.

The audioscope tests for hearing deficits at 25 to 40dB pure tone at 500 Hz, 1000 Hz, 2000 Hz, and 4000 Hz. The frequencies are in the range of human speech. The audioscope also allows the examiner to assess the external ear canal and TM for cerumen, foreign bodies, or otitis. The audioscope has reported sensitivity of 0.94 and specificity of 0.69 to 0.80 for hearing loss (**sensitivity** is the ability to detect true positives; **specificity** is the ability to detect true negatives).

Nursing Implications

The physical implications of hearing loss are apparent, but the effects go much further, also affecting the patient's psychosocial well-being. Nurses, being aware of the frequency of hearing impairment in older adults, can be alert for hearing loss in their older patients by assessing for communication impairment and social withdrawal and referring them as needed for further evaluation and treatment.

Source: Yueh, B., Shapiro, N., MacLean, C., and Shelelle, P. (2003). Screening and management of adult hearing loss in primary care: Scientific review. *JAMA* 289(15), April 16, pp. 1976–1985.

HEALTH CONCERNS

- Screening for hearing loss should begin in infancy. Because development of speech and psychosocial skills depend on adequate hearing, any child with a suspected developmental delay should be screened for hearing loss.
- Evaluate the environment and feeding habits of infants diagnosed with frequent ear infections. Chronic exposure to a smoker in the home and putting the child to bed with a bottle increase the incidence of ear infections.
- People exposed to loud noise should have regular hearing examinations and be instructed in ways to reduce their exposure, including the use of protective ear wear.
- People receiving ototoxic medications should be evaluated for problems with balance, gait, vertigo, tinnitus, and diminished hearing on a regular basis throughout therapy and be instructed to stay alert for signs of ototoxicity several weeks after therapy ends.

Common Abnormalities: Ears

ABNORMALITY

ASSESSMENT FINDINGS

Acoustic Neuroma

- Benign tumor that grows into auditory canal. Involves CN VIII

- *Symptoms:* tinnitus, progressive hearing loss, papilledema, headache, facial numbness, and balance and gain problems (depending on the tumor's location and size).

Acute Otitis Media

- Inflammation or infection of middle ear

- Ear pain with reddened TM.
- Distorted cone of light reflex and signs of conductive hearing loss.



Cholesteatoma

- Cystlike sac of keratin debris caused by a congenital defect or chronic otitis media

- Can enlarge and obstruct middle ear.
- Can release enzymes that erode the ossicles.



Hemotympanum

- Dark blue TM caused by bleeding in middle ear behind it.

- Sign of conductive hearing loss.
- Evidence of recent head injury.



ABNORMALITY

ASSESSMENT FINDINGS

Ménière's Disease

- Chronic, progressive disease of inner ear that leads to permanent hearing loss.

- Affects proprioception, the ability to accurately sense one's body position.
- Characterized by a sensation of fullness or pressure in the ears and recurrent episodes of vertigo, tinnitus, and hearing loss, lasting from a few minutes to several hours.
- Person may also experience nausea, vomiting, and profuse sweating.
- Subjective sensation of ear fullness, vertigo, tinnitus.
- Disturbance of balance and gait.
- Positive sway with Romberg test.

Otitis Externa

- Inflammation or infection of external ear, often caused by excessive swimming, chronic irritation, or removal of cerumen

- Ear pain, especially with movement of the tragus or pulling of ear lobe.
- Redness of external auditory canal or auricle of external ear.



Presbycusis

- Diminished hearing acuity in older adults specifically for high-pitched sounds.

- Hearing acuity decreased.
- Diagnosed by watch tick test.

Serous Otitis Media

- Yellowish TM with serous fluid level or air bubble visible through membrane.

- Cone of light reflex may be distorted.
- Transient conductive hearing loss present.



ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the ears.

Ask the patient the following:

- Do you have...
 - Hearing loss?
 - Vertigo?
 - Tinnitus?
 - Ear drainage?
 - Earache?
- Do you have problems with your ears such as ringing? Do you have hearing problems?
- Do you have balance problems?
- Do you have drainage from your ears? If yes, how much and what color?
- Have you had recent head trauma?
- Do you have any health problems?
- Are you exposed to noise pollution at work or in your home environment?
- Are you on any prescribed or OTC medications?
- Do you have allergies?

ASSESSMENT CHECKLIST (continued)

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam:

- Tuning fork
- Otoscope with pneumatic attachment
- Thermometer
- Watch
- Gloves if indicated

Position

Sitting for adult
Supine for infant or toddler to stabilize head

Technique

First, scan your patient head to toe checking for specific signs of diseases affecting other organ systems that might alter the ear.

Inspection, Palpation, Otoscopic Exam, Hearing Tests

External Ear

Inspect:

- Angle of attachment
- Size, shape, symmetry
- Condition of skin, drainage

Palpate:

- Consistency, tenderness, nodules
- Helix, tragus, mastoid process

Otoscopic Exam

External Ear Canal

Inspect:

- Color, drainage, foreign objects, lesions

Tympanic Membrane

Inspect:

- Color, position of landmarks, integrity of TM, mobility of TM

Hearing tests (Test Each Ear Separately)

- Whisper Voice Test
- Ticking Watch Test
- Weber Test
- Rinne Test

DOCUMENT

Document your findings and develop your plan of care.

S U M M A R Y

- A thorough health history provides direction for the physical examination to include exploration of factors that may be related to ear health. Information from both the history and the physical examination is then analyzed to determine appropriate nursing diagnoses.
- The ears are complex sensory organs that provide specialized functions crucial to neurosensory develop-


ment in infancy and to the development of psychosocial, motor, and cognitive skills in childhood and the maintenance of those skills in adulthood.

- A comprehensive history and physical examination enables early detection and treatment of hearing problems.

R E V I E W Q U E S T I O N S

1. John Spivey, age 75, is scheduled for a hearing exam. Considering his age, which hearing change might you expect to find?
 - a. Presbycusis
 - b. Otorrhea
 - c. Cholesteatoma
 - d. Tinnitus
2. Your patient is complaining of tinnitus. Which question would best assess this symptom?
 - a. Do you have earwax?
 - b. Are you taking aspirin?
 - c. Do you have any ear drainage?
 - d. Did you ever have an ear infection?
3. You are assessing a newborn infant's position of his ears. Which finding would be cause for concern and referral?
 - a. Low-set ears
 - b. Angle of attachment <10 degrees
 - c. Ears aligned with external canthus
4. Prior to insertion of the otoscope, which areas should be palpated?
 - a. Helix, tragus, mastoid
 - b. Antihelix, mastoid, antitragus
 - c. Antitragus, lobule, helix
 - d. Tragus, antitragus, mastoid

5. You have done the Weber test on your patient, and the patient reports the sound is heard equally in both ears. How would you explain these findings to your patient?
 - a. This is a normal finding.
 - b. You have a conductive hearing loss.
 - c. You have a perceptive hearing loss.
 - d. You have a hearing loss, but I'm unable to differentiate what kind of hearing loss with this test.
6. You have tested your patient's hearing using the Rinne test. The results reveal bone conduction is longer than air conduction. What do these findings suggest?
 - a. Nothing, they are normal findings,
 - b. Conductive hearing loss,
 - c. Perceptive hearing loss,
 - d. Sensorineural hearing loss,
7. You are performing an otoscopic exam on a 2-year-old-child. To straighten the ear canal, you should:
 - a. Pull helix up.
 - b. Pull helix down.
 - c. Pull helix forward.
 - d. Pull helix back.
8. You are performing an otoscopic exam on an adult patient. How far should you insert the speculum?
 - a. $\frac{1}{4}$ "
 - b. $\frac{1}{2}$ "
 - c. 1"
 - d. $1\frac{1}{2}$ "
9. As you examine the TM, you note the color of the drum. Which TM color would be considered normal?
 - a. Pink
 - b. Red
 - c. Pearly gray
10. When performing an otoscopic exam, what type of drainage is considered normal in the external ear canal?
 - a. No drainage
 - b. Cerumen
 - c. Clear
 - d. Bloody



Assessing the Respiratory System

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent respiratory history questions
- Obtain a respiratory history
- Perform a respiratory physical assessment
- Document respiratory findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

Breathing is a natural function that most people never think about—unless it is impaired. As you will see, an ineffectively functioning respiratory system causes disruption throughout the body, just as changes in other body systems have a significant impact on the respiratory system.

Nurses can play a major role in preventing dysfunction or halting the progression of respiratory disease by identifying subtle changes. They can also identify people who are at risk because of unhealthy behaviors or genetic predisposition. Thus, the goal of a thorough respiratory assessment is early detection of risk factors, potential problems, or dysfunction of the system. With accurate and complete information, you will then be able to plan appropriate interventions—teaching health promotion/disease prevention and/or implementing treatment measures.

Anatomy and Physiology Review

Before you begin your assessment, you need to recognize normal respiratory function in order to distinguish abnormal findings. The anatomy will direct your assessment process. Visualizing the anatomical landmarks and underlying structures will enable you to perform the assessment accurately, and understanding the physiology will allow you to interpret your findings.

The primary function of the respiratory system is the exchange of oxygen and carbon dioxide through respiration. This system also plays a role in maintaining acid-base balance. The physiology of respiration consists of both mechanical and physiological processes.

The mechanical process of respiration is accomplished by pulmonary ventilation, the exchange of air between the lungs and the atmosphere. It consists of two phases—inspiration and expiration.

The physiological process occurs on three levels: external, internal, and cellular:

- **External respiration** is the exchange of gases (oxygen and carbon dioxide) between the alveoli and the blood through the alveolar-capillary membrane.
- **Internal respiration** is the exchange of gases between the systemic capillaries and the tissue at the cellular level.
- The **cellular physiological process** is the exchange of gases within the cell.

Structures of the Respiratory System

The respiratory system (Fig. 13.1) begins at the nose and continues as a series of airways or passages extending to the alveoli where gas exchange takes place. The **nasal**,




FIGURE 13.1. Structures of the respiratory system.

oropharynx, and **conducting airways** are considered dead space because no exchange of gases occurs there. The nasal and oropharynx airways include the nasal cavity, pharynx, and larynx. The conducting airways include the trachea and bronchi.

The **bronchioles** are transitional airways where some gas exchange occurs. The alveolar ducts, sacs, and alveolus are the functional units of the lung in which exchange of gases occurs with the pulmonary capillary bed. The primary muscle of respiration is the **diaphragm**; the secondary muscles are called the **accessory muscles**. The negative lung pressure that is needed for breathing is maintained by the **pleura**.

Structures and Functions of the Respiratory System

STRUCTURE	DESCRIPTION/PRIMARY FUNCTION
Nasal Cavity	<ul style="list-style-type: none"> ■ Warms, filters, and moistens air.
Pharynx	<ul style="list-style-type: none"> ■ Musculomembranous tube that acts as a passage between nose and larynx for air and between mouth and esophagus for food. ■ Consists of three parts: nasopharynx, oropharynx, and laryngopharynx.
Nasopharynx	<ul style="list-style-type: none"> ■ Passage for air behind nasal cavities. ■ Adenoids or pharyngeal tonsils located on posterior wall are lymphoid tissue and have a protective function. ■ Eustachian tubes extend from middle ears and open into nasopharynx to equalize atmospheric pressure.
Oropharynx	<ul style="list-style-type: none"> ■ Passage for both air and food, located behind mouth. ■ Palatine tonsils are on lateral wall of oropharynx.
Laryngopharynx	<ul style="list-style-type: none"> ■ Most inferior portion of pharynx. ■ Opens anteriorly into larynx and posteriorly into esophagus.
Larynx	<ul style="list-style-type: none"> ■ Passage between pharynx and trachea. ■ Contains vocal cords to produce sound.
Single Cartilages	<ul style="list-style-type: none"> ■ Cartilages that support larynx.
<ul style="list-style-type: none"> ■ Thyroid ■ Epiglottis ■ Cricoid 	<ul style="list-style-type: none"> ■ Prevents food from entering trachea.
Paired Cartilages	<ul style="list-style-type: none"> ■ Support and affect vocal cords.
<ul style="list-style-type: none"> ■ Arytenoid ■ Cuneiform ■ Corniculate 	
Trachea	<ul style="list-style-type: none"> ■ Conducting airway between the larynx and the bronchi, about 10 to 12 cm long. ■ Divides into two primary bronchi midthorax at the carina.
Bronchi	<ul style="list-style-type: none"> ■ Conducting airways to hilum of each lung. ■ Each primary bronchus branches into secondary (lobar) and tertiary (segmental) bronchi.
Bronchioles	<ul style="list-style-type: none"> ■ Smaller transitional airways to alveoli. ■ Consist of bronchioles (>1 mm in diameter), terminal bronchioles (>0.5 mm), and even smaller respiratory bronchioles, in which some gas exchange occurs.
Alveolar Ducts, Sacs	<ul style="list-style-type: none"> ■ Functional units of lung. ■ Each lung has more than 350 million alveoli, in which exchange of gases between alveoli and pulmonary capillary bed occurs.
Pleura Serous	<ul style="list-style-type: none"> ■ Protective linings of lungs that maintain negative pressure and aid in mechanics of breathing.
Parietal Pleura	<ul style="list-style-type: none"> ■ Consists of parietal pleura and visceral pleura.
Visceral Pleura	<ul style="list-style-type: none"> ■ Line thoracic wall and superior portion of diaphragm. ■ Extension of parietal pleura that covers lung. ■ Serous fluid fills and lubricates pleural cavity, preventing friction.

Acid-Base Balance

The respiratory system plays a key role in maintaining acid-base balance by responding first to changes in blood acidity. When carbon dioxide is converted to bicarbonate or carbonic acid, the blood's pH becomes either more alkaline or more acidic. Depending on the type of acid-

base problem, the respiratory system responds by changing the rate and depth of respiration.

For example, in **metabolic acidosis**, a metabolic imbalance characterized by excess acid retention, the rate and depth of respiration *increase* to blow off excess carbon dioxide, which reduces the amount available to make carbonic acid. This results in a decrease of hydrogen

ions and subsequent movement of the pH toward its normal range. On the other hand, in **metabolic alkalosis**, an imbalance characterized by an excess of base, the respiratory system *decreases* the rate and depth of respiration to conserve carbon dioxide. Breathing pattern changes such as hyperventilation or hypoventilation are either compensation for an out-of-balance pH or a primary cause of acid-base imbalances. Hyperventilation can result in respiratory alkalosis; hypoventilation can result in respiratory acidosis.

Interaction With Other Body Systems

The neurological, musculoskeletal, and circulatory systems have an essential role in maintaining respiratory function. A problem in any of these systems may affect the functioning of the respiratory system.

The Neurological System

In the brain, the respiratory control centers are located in the reticular substance of the medulla oblongata and in the pons. The respiratory center within the medulla can be divided further into two opposing centers: the **inspiratory center** and the **expiratory center**. These two centers control the motor neurons to the phrenic nerve, which arises in the third, fourth, and fifth cervical nerve segments of the spinal cord and innervate the diaphragm, and the intercostal muscles, which arise in the upper thoracic segments. The inspiratory center stimulates the contractions of the respiratory muscles, at the same time depressing the expiratory center. This results in **inhalation**. The process then reverses; as the inspiratory center is depressed, the expiratory center is stimulated, resulting in relaxation of the respiratory muscles and **exhalation**.

The pons also has two respiratory centers—the apneustic and the pneumotaxic—that are responsible for the rhythm of breathing. The **apneustic center** prolongs inspiration, and the **pneumotaxic center** contributes to exhalation. These brainstem centers are connected to the hypothalamus and cerebral cortex, the parts of the brain that control emotions such as fear, pain, and anger. These types of emotions can affect breathing to the same extent as activities like talking or singing, which require more conscious breath control. A child who holds his or her breath during a tantrum is exhibiting conscious breath control. However, if the child loses consciousness as a result, the lower brain centers resume control again, so the child breathes automatically.

The respiratory center is sensitive to changes in carbon dioxide (PaCO_2) levels and responds accordingly by affecting the rate and depth of respiration until homeostasis is achieved. Central chemoreceptors are found bilaterally in the medulla; peripheral chemoreceptors are located in the aortic arch and at the bifurcation of the

carotid arteries. The most potent chemical affecting respiration is PaCO_2 . Even slight elevations reduce the pH of the cerebrospinal fluid, sending strong signals to the central chemoreceptors. This, in turn, stimulates the respiratory centers to increase the respiratory rate.

Changes in oxygen (PaO_2) have a weak influence on respiratory rate. Only when the peripheral chemoreceptors sense a PaO_2 less than 60 mm Hg will PaO_2 become a major stimulus to increase respiration. In general, blood oxygen levels act by increasing or decreasing the chemoreceptor's sensitivity to PaCO_2 . Homeostasis of the acid-base balance of the body is maintained by the very complex interactions between carbon dioxide, oxygen, and blood pH unless disease intervenes.


The Musculoskeletal System

The musculoskeletal system provides the bony structure (thorax) and the muscles needed for breathing. The phrenic nerve innervates the **diaphragm**, the primary muscle used for breathing. A dome-shaped muscle at the floor of the thorax, the diaphragm separates the thorax from the abdomen. During inspiration, the diaphragm flattens as it contracts, enlarging the chest capacity. During exhalation, the diaphragm relaxes and rises to assume a dome shape.

Accessory muscles include the sternocleidomastoid, anterior serrati, scalene, trapezius, intercostal, and rhomboid muscles. They come into play during strenuous physical activity (such as jogging) or when the body has intrapulmonary resistance to air movement. The accessory muscles enhance ventilation by increasing chest expansion and lung size during inspiration. When contracted, the external intercostal muscles pull the ribs up and out; the internal intercostal muscles pull the ribs down and inward. Contraction of these muscles decreases intrathoracic pressure, causing an inward flow of air. As the muscles relax, the lungs recoil, and exhalation occurs. The abdominal muscles also assist with rapid breathing, deep breathing, exercise, coughing, and sneezing. An intact thoracic cage is essential for normal respiratory function (Fig. 13.2).

The Circulatory System

The circulatory system of the lungs consists of the bronchial and pulmonary circulation. **Bronchial circulation** attends to the metabolic demands of the lung tissue. **Pulmonary circulation** fosters the exchange of gases between the alveoli and the blood in the pulmonary capillaries. Oxygen, which diffuses from the alveoli into the pulmonary circulation, binds with hemoglobin and is transported to the cells. Similarly, carbon dioxide moves from the cells, into the pulmonary circulation, and finally into the alveoli to be exhaled from the body. Any disruption of the circulation within the lungs affects the body's ability to provide




NEUROLOGICAL

Medulla is respiratory center.
Chemoreceptors respond to changes in carbon dioxide, oxygen and hydrogen ions by affecting rate and depth of respirations.



INTEGUMENTARY

Protects structures of upper respiratory tract.
Nasal hair filters air.
Skin color reflects oxygenation.




ENDOCRINE

Converting enzymes in lung convert angiotensin I to angiotensin II.
Epinephrine and norepinephrine increase respiratory rate and dilate airways.



DIGESTIVE

Abdominal muscles aid with respirations.
GI tract provides nutrients to respiratory system.



CARDIOVASCULAR

Cardiovascular system transports oxygen and carbon dioxide between lungs and peripheral tissue.

Activation of angiotensin by lungs important in B/P regulation.

Peripheral chemoreceptors in aortic arch and carotid arteries respond to oxygen, carbon dioxide and hydrogen ion concentrations.

SKELETAL


Ribs provide protective framework for lungs.



MUSCULAR

Diaphragm, chest and abdominal muscles needed for breathing.

Accessory muscles used when oxygen demands are increased, as with strenuous exercise.



URINARY


GU system works with respiratory system to eliminate wastes and maintain acid-base balance.

LYMPHATIC

Alveolar macrophages trap microorganisms and other foreign substances.

Foreign material removed by cilia or lymphatic system.


Tonsils and adenoids protect upper airways from infection, but also may become enlarged and obstruct breathing.



REPRODUCTIVE

Sexual activity increases oxygen demands.

Pregnancy affects oxygen demands and breathing.



Relationship of Respiratory System to Other Systems




FIGURE 13.2. Structures of the respiratory system.

adequate oxygen to tissues or to remove enough carbon dioxide for healthy functioning.

Developmental, Cultural, and Ethnic Variations

Infants

Infants are obligatory nose breathers. Their respirations are primarily abdominal, and breath sounds are louder, harsher, and more bronchial. Their respiratory rhythm is often irregular, with brief periods of apnea. The shape of the chest is more round than oval, with the ribs in a horizontal position. After about age 2, the ribs become more oblique, and the breathing pattern becomes primarily intercostal.



ALERT

In infants, periods of apnea that exceed 20 seconds may indicate a cardiopulmonary or central nervous system (CNS) disorder, sepsis, gastroesophageal reflux, hypoglycemia, or other metabolic problems and need further evaluation. The infant may be at high risk for sudden infant death syndrome (SIDS).

Pregnant Women

During pregnancy, increases in tidal volume and respiratory rate allow for a 20 percent increase in oxygen consumption. Later in the pregnancy, the diaphragm rises

and the costal angle widens to accommodate the enlarging uterus.

Older Adults

Alveoli usually become fibrotic with age, resulting in a decreased surface area for exchange of gases. Breathing and lung capacity decrease as a result of muscle weakness and decreased elasticity. This results in increased dead space, trapped air, and decreased vital capacity. Musculoskeletal changes that occur with aging can cause an increase in the anterior-posterior diameter from weakening of the intercostal muscles and softening of the rib cartilage. The thoracic curve may also increase, resulting in mild kyphosis, an accentuated thoracic curve. Together, these changes give the appearance of a barrel chest, but without the dysfunction.

People of Different Cultures/Ethnic Groups

Chest size can vary among cultures in proportion to the person's size. For example, Chinese Americans generally have smaller chests than Caucasians or African Americans. The higher incidence of respiratory problems among certain ethnic groups relates more specifically to occupational and environmental factors than genetic ones. For example, African Americans living in urban areas have a higher risk of respiratory disease. Appalachians have a higher risk for black lung, emphysema, and tuberculosis (TB). The Irish have a higher risk for respiratory problems related to

coal mining. Navajo Indians have an increased risk for respiratory diseases because of close living quarters.

to perform a complete health history, perform a focused respiratory history.

Introducing the Case Study



Mrs. Katie Kane is scheduled for a cholecystectomy and has come to the hospital for a preoperative workup. She is a 45-year-old married mother of three and part-time accountant. As you begin the interview, she says, “Even though I’m a little scared about surgery, I’ll be glad when my gallbladder is out. It’s been causing me a lot of pain.”

Mrs. Kane is scheduled to have general anesthesia, so a thorough respiratory assessment is necessary.

Biographical Data

A quick review of the biographical data may identify actual or potential problems. How old is the patient? Respiratory structure and function change with age. For example, young children are more susceptible to respiratory infections because their airways are smaller and an unpracticed immune system, but as people age, their forced expiratory volume decreases. Residence and occupation also have an impact on the respiratory system because a person’s job or environment may harbor risk factors. Chronic diseases may place financial strains on patients, so you may need to make referrals to agencies that can assist them.

Performing the Respiratory Assessment

Assessment of the respiratory system involves a comprehensive health history and physical examination. As you perform the assessment, be alert for signs or symptoms of actual or potential respiratory problems.

Health History

The health history identifies respiratory symptoms and risk factors. It provides you with subjective data to guide your examination. As you proceed, be alert for actual or potential problems. If time is an issue and you are unable

Current Health Status

Begin with the patient’s chief complaint. Three major respiratory symptoms to watch for are dyspnea, cough, and chest pain. Perform a symptom analysis for any presenting symptom, using the PQRST format. If you notice signs of respiratory distress, such as shortness of breath (SOB), confusion, or anxiety, postpone the detailed history and focus on the acute problem. Obtain a detailed history later, when the patient’s condition improves, or get it from a secondary source such as a family member.

A patient with chronic respiratory disease often adapts to a compromised respiratory status. Therefore, compare the patient’s present respiratory status with her or his status at previous examinations to evaluate the impact of the disease’s progression.

Focused Respiratory History

Your patient’s condition or time restraints may prohibit taking a detailed respiratory history. If so, ask questions that focus on a history of respiratory disease, the presence of respiratory symptoms, identification of risk factors, and health promotion activities. The patient’s response to these questions will help to direct your assessment. Key questions include:

- Do you have a history of respiratory disease? If so, are you taking any medications? What are they and why are you taking them?
- Do you have any other medical problems (especially cardiac problems)?
- Do you have allergies? If so, describe your reaction.
- Do you have a cough, shortness of breath, or chest pain?
- Do you use tobacco? If so, what kind, how much, and for how long?
- What is your occupation?
- Where do you live?
- When was your last PPD, and what were the results?
- Have you ever had a chest x-ray? If so, what were the results?
- Have you been immunized for influenza or pneumonia?

**HELPFUL HINT**

Respiratory disease makes people tire easily because most of their energy is expended on breathing. Therefore, divide the history into several short interviews and ask questions that can be easily answered with few words.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Cough

Coughing is one of the most common respiratory complaints. The causes range from insignificant to life-threatening conditions. Coughing is a protective, reflexive mechanism that helps maintain a patent airway. It occurs in three phases: Deep inspiration that increases lung volume, closure of the glottis, and then muscular contraction forcing the sudden opening of the glottis, resulting in a cough. Cough receptors, located in the larynx, respiratory tree, pleura, acoustic duct, nose, sinuses, pharynx, stomach, and diaphragm respond to mechanical, inflammatory, or irritating stimuli.

Dyspnea

Dyspnea is a subjective sensation of breathing difficulty often described as SOB. It may be normal with overexertion or anxiety, but it may also signal underlying cardiopulmonary or neuromuscular problems or allergic reactions.

COPD and CHF**COPD**

Chronic obstructive pulmonary disease (COPD) is a chronic, progressive disease characterized by airway obstruction and diminished lung function. It includes emphysema and chronic bronchitis. Increasing dyspnea may correlate with the progression of the disease.

CHF

Congestive heart failure (CHF) causes the heart to pump ineffectively, resulting in volume overload. The right side, left side, or both sides of the heart can fail. Right heart failure (cor pulmonale) is frequently associated with lung disease.

Chest Pain

Chest pain may have cardiac, pulmonary, gastrointestinal, or musculoskeletal origins. However, chest pain of respiratory origin is associated with the parietal pleura, chest wall, and mediastinal structures because the lungs and visceral pleura do not have pain fibers.

Related Symptoms

Other symptoms associated with respiratory disease include **edema** and fatigue. Edema results from right-side congestive heart failure, a common complication of chronic obstructive lung disease. Usually, edema is located in the lower extremities or abdomen. Ask your patient, “Do you have swelling in your abdomen, legs, ankles, or feet?”

Hypoxia, increased energy expended for breathing, and associated cardiac involvement accompany long-standing lung disease and contribute to the development of fatigue. People adapt to fatigue by lowering their activity level, so they have more difficulty performing activities of daily living (ADLs). Changes in rest and sleep patterns may also be seen. Ask your patient, “Do you have enough energy to do your usual daily activities? Do you need to sleep or rest more than usual?”

Cor Pulmonale

Cor pulmonale is right heart hypertrophy or failure resulting from disorders of the lungs or pulmonary vessels. Common causes include COPDs and living at high altitudes for extended periods.

Past Health History

The purpose of the past health history is to compare it with the patient’s present respiratory status or uncover risk factors that might predispose him or her to respiratory disorders. Be sure to follow up on any unclear or vague answers. Rewording the question may help the patient find a relevant response.

**ALERT**

Adverse effects of oral or inhaled bronchodilators include nausea/vomiting, tremors, nervousness, insomnia, tachycardia, hypertension, and oral fungus (inhaled).

Past Health History

RISK FACTORS/ QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Childhood Illnesses

- Did you have frequent respiratory infections, asthma, TB, or pneumonia as a child?

- May indicate underlying chronic disease.

Hospitalizations/Surgeries

- Have you ever been hospitalized? What for?
- Have you ever had a TB test, chest x-ray, or other respiratory diagnostic testing? When?

- Can be used as baseline data.
- Also indicates disease prevention activities.

Serious Injuries/Chronic Illnesses

- Have you had fractured ribs, steering wheel injuries, or knife wounds to the chest?
- Have you ever had a previous respiratory illness?
- Do you have any other chronic medical problems, such as heart disease, renal disease, cancer, or HIV?

- May explain physical findings.
- May have respiratory manifestations.

Immunizations

- Were you immunized as a child?
- Did you receive adult boosters against respiratory diseases (e.g., pneumovax for pneumonia and bacille Calmette-Guerin [BCG] for TB)?

- Identify health promotion or teaching needs.
- BCG explains a positive TB test (PPD).

Allergies

- Do you have any allergies (e.g., to pollens, food, drugs, environmental factors)? What type of reaction do you have?

- Can cause chronic sinus or nasal congestion, trigger an asthma attack, or progress to COPD.

Medications

- What prescribed and OTC drugs are you taking? For how long?

- Determines compliance with treatment and therapeutic vs. toxic effects.
- Excessive use of medications can lead to tolerance and rebound effects.
- Patient may be on cardiac drugs or oxygen.
- Corticosteroids (cortisone), which are often prescribed to relieve signs and symptoms of COPD, have serious adverse effects that need to be monitored.
- Allows evaluation of possible adverse drug interactions (e.g., beta adrenergics and antihypertensives). See *Drugs That Adversely Affect the Respiratory System*.

Recent Travel/Military Service/Exposure to Infectious or Contagious Diseases

- Have you traveled by plane recently?
- Have you stayed in a hotel room recently?
- Have you been exposed to people with colds, flu, or cough?

- Respiratory infections (e.g., URIs or Legionnaire's disease) are more easily transmitted by the closed airflow in airplanes or faulty ventilation systems in hotels.
- Can identify contagious disorders.
- TB is most commonly acquired from living with infected family members.

Family History

The purpose of the family history is to identify any predisposing or causative factors of respiratory origin. If possible,

help patients draw a family tree. This helps them remember more relevant information about family members.

Family History	
RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Genetically Linked Respiratory Disorders</p> <ul style="list-style-type: none"> ■ Does anyone in your family have: <ul style="list-style-type: none"> ■ Alpha antitrypsin deficiency? ■ Allergies or asthma? ■ Cystic fibrosis? <p>Smoking</p> <ul style="list-style-type: none"> ■ Are you or were you exposed to second-hand smoke in your home? If so, for how long? What type of tobacco product was used? <p>TB</p> <ul style="list-style-type: none"> ■ Does anyone in your family have TB? If so, what type and how long were you exposed? 	<ul style="list-style-type: none"> ■ Increased risk for emphysema ■ Avoid allergens, smoking ■ Autosomal dominant disorder <ul style="list-style-type: none"> ■ Second-hand smoke can cause cancer, COPD, asthma. <ul style="list-style-type: none"> ■ Will need follow-up PPD or chest x-ray.

Drugs That Adversely Affect the Respiratory System

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Adrenergic agents (sympathomimetics)	Epinephrine	Dyspnea, paradoxical bronchospasms
Antineoplastic agents	Bleomycin	Fine crackles, dyspnea (early signs of pulmonary toxicity); interstitial pneumonitis, pulmonary fibrosis
	Busulfan	Irreversible pulmonary fibrosis (busulfan lung)
	Carmustine	Pulmonary fibrosis
	Cyclophosphamide	Pulmonary fibrosis (with high doses)
	Melphalan	Interstitial pneumonitis, pulmonary fibrosis
	Methotrexate	Interstitial pneumonitis
	Mitomycin	Dyspnea, cough, pulmonary infiltrates, interstitial pneumonitis
Antiarrhythmics	Amiodarone	Interstitial pneumonitis, pulmonary fibrosis
Antihypertensives	Enalapril	Cough
	Guanabenz	Nasal congestion, dyspnea
	Reserpine	Nasal congestion
Anti-infectives	Polymyxin B sulfate	Respiratory paralysis
Beta-blockers	All beta-blockers	Bronchospasm, particularly in patients with a history of asthma; dyspnea, wheezing
		Dyspnea, bronchoconstriction (with subcutaneous administration)
Cholinergic agents	Bethanecol	Increased bronchial secretions, bronchospasm, dyspnea
	Neostigmine	Pulmonary infiltrates, interstitial pneumonitis, interstitial fibrosis, "gold" bronchitis
Gold salts	Aurothioglucose, gold sodium thiomalate	Respiratory depression
Narcotic analgesics	All types	Bronchospasm
Nonsteroidal anti-inflammatory agents	Aspirin	

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Penicillins	Ibuprofen	Bronchospasm, dyspnea
Sedatives and hypnotics	Indomethacin	Bronchospasm, dyspnea
Urinary tract antiseptics	All types	Anaphylaxis
	All types	Respiratory depression, apnea
	Nitrofurantoin	Pulmonary sensitivity reactions, such as cough, chest pain, dyspnea, pulmonary infiltrates; interstitial pneumonitis (with prolonged use)
Miscellaneous agents	Cromolyn sodium	Dyspnea, pharyngitis
	Levodopa	Excessive nasal discharge, hoarseness, episodic hyperventilation, bizarre breathing patterns

TB in Immigrants

Although not a common practice in the United States, the BCG vaccine for TB is administered in many other countries, such as Southeast Asia. Immigrants are often PPD positive without infection. They should still have a chest x-ray at least every 5 to 10 years. They may need to take antituberculosis drugs if active disease is present in a family member.

Review of Systems

Changes in the respiratory system have an impact on every other body system. In addition to helping you detect problems that directly affect the respiratory system, the review of systems (ROS) identifies changes in other systems that result from changes in the respiratory system. The ROS allows you to catch anything that you might have missed so far, and it gives meaning to the symptom by relating it to the affected system.

Review of Systems

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>General Health Survey</p> <ul style="list-style-type: none"> Do you have fatigue or activity intolerance? Have you had unexplained fever, night sweats, or weight loss? Have you gained weight recently? <p>Integumentary</p> <ul style="list-style-type: none"> Have you noticed any skin color changes? Do you have swollen ankles or tight shoes? <p>Head, Eyes, Ears, Nose, and Throat (HEENT)</p> <p>Head/Neck</p> <ul style="list-style-type: none"> Do you have any swelling, masses, or difficulty swallowing? 	<ul style="list-style-type: none"> Chronic lung disease often causes fatigue and activity intolerance because so much energy is expended on breathing. May be associated with a more serious underlying disease such as TB. May correlate with fluid retention associated with right side CHF, a common complication of COPD. Cyanosis: Cardiopulmonary disease. Fluid retention: Right-side cardiac involvement of long-standing respiratory disease. Enlarged lymph nodes: Infection or malignancy. Enlarged thyroid may compromise respiratory function.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Eyes ■ Do you have excessive tearing?	■ Associated with allergies.
Nose/Throat ■ Do you have postnasal drip, sinus pain, or sore throat?	■ May indicate allergies or acute/chronic URI.
Cardiovascular ■ Do you have chest pain?	■ May indicate cardiac involvement.
Gastrointestinal ■ Do you have right upper quadrant (RUQ) pain, nausea, or gastrointestinal upset? ■ Have you experienced loss of appetite and weight loss?	■ Associated with right CHF, a common complication of COPD. ■ Associated with chronic lung disease, TB, or lung malignancy.
Urinary ■ Do you awaken during the night to urinate?	■ <i>Nocturia</i> : Associated with heart failure or diuretic use to treat right-side heart failure.
Reproductive ■ Have you had any changes in sexual performance? ■ Do you practice safe sex?	■ Chronic respiratory disease requires most of the patient's energy for breathing. ■ Unprotected sexual activity increases risk for human immunodeficiency virus (HIV).
Musculoskeletal ■ Have you had any weakness or muscle wasting?	■ Chronic lung disease.
Neurological ■ Have you experienced any irritability, confusion, forgetfulness, or changes in your mental abilities? ■ Have you had tremors?	■ <i>Change in mental status</i> : Early sign of hypoxia. ■ Associated with theophylline toxicity or CO ₂ narcosis.
Immune/Hematologic ■ Have you had any anemia or allergies?	■ Directly affect respiratory system.

Psychosocial Profile

The psychosocial profile reveals lifestyle patterns that may affect the respiratory system and place the patient at

risk for respiratory disorders. The patient's lifestyle may be affected by respiratory disease, especially when it is chronic.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Health Practices and Beliefs/Self-Care Activities ■ Do you have regular healthcare checkups; knowledge of medications, protection from environmental pollutants and allergens?	■ Ascertains compliance with treatment or screening for risk factors. ■ Identifies teaching needs.
Typical Day ■ What constitutes a usual day's activities? Has this changed over the last year?	■ Identifies whether a patient is doing too much for energy level or limiting activities to compensate for loss of energy. ■ Looks at possible disease progression. ■ Shows ways patient may be able to adapt lifestyle if needed.

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ Have you lost weight and appetite? ■ Have you gained weight? <p>Activity and Exercise Patterns</p> <ul style="list-style-type: none"> ■ How would you rate your level of activity? ■ Do you exercise regularly? <p>Pets/Hobbies</p> <ul style="list-style-type: none"> ■ Do you have cats, dogs, or birds in your home? 	<ul style="list-style-type: none"> ■ Associated with lung cancer, but usually by the time weight loss is evident, metastasis has occurred. ■ Dyspnea, fatigue, and decreased lung capacity from overinflation result in appetite loss and smaller amounts of food being consumed. Also evaluate ability to purchase and prepare food and effects of social isolation on eating habits. ■ Weight loss is a common symptom in catabolic infections such as TB and acquired immunodeficiency syndrome (AIDS). ■ <i>Sudden gain</i>: Fluid retention from cardiac failure or secondary to medication (e.g., cortisone). ■ Gradual decrease or change in activity/exercise patterns seen in chronic respiratory disorders. ■ Identifies need for physical therapy referral to improve respiratory function or for health promotion. ■ Pets and hobbies can promote health if they do not irritate respiratory function.
<ul style="list-style-type: none"> ■ Do you have hobbies that require use of chemicals? <p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ How much uninterrupted sleep do you get per night? ■ Do you have insomnia? ■ Do you feel rested in the morning? ■ Do you take daytime naps? <p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you or did you ever smoke? In what form (e.g., cigarettes, cigars, marijuana)? Determine the pack-years (number of packs a day times number of years). If you quit smoking, how long ago? ■ Do you drink alcohol? What kind, how much, and how often? ■ Do you use street drugs? 	<div data-bbox="837 1018 1460 1228" style="border: 1px solid red; padding: 5px;">  <p>ALERT</p> <p>Allergies to pet fur or dander can trigger asthma attacks; exposure to bird droppings or feathers may cause hypersensitivity pneumonitis with eventual pulmonary fibrosis.</p> </div> <ul style="list-style-type: none"> ■ Exposure to glue, paint, sprays, silica, sawdust, or other irritants may necessitate changing hobbies. ■ Dyspnea, cough, pain, and medications can cause sleep pattern disturbances. ■ Sleep apnea results in interrupted sleep and daytime fatigue. ■ Quitting smoking can improve respiratory function even if chronic respiratory problems already exist. ■ Cigars may not be inhaled, but heat and tar can cause oral/pharyngeal/laryngeal damage. ■ Marijuana is much more destructive to lungs than regular cigarettes, especially if smoked regularly. ■ Alcohol affects ciliary action, reducing mucus clearance and decreasing immune competence. ■ Cough reflex may be depressed with heavy alcohol consumption, increasing risk of aspiration pneumonia. ■ Intravenous drug abuse increases chance of HIV infection, which is associated with high incidence of

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ What is your occupation? Do you use protective equipment? ■ Do you work in a smoke-free environment? ■ Does your employer promote a healthy environment, clean air? ■ Is your work environment safe? Will you be able to continue in your present occupation? ■ Do you have disability insurance? 	<p>opportunistic respiratory diseases such as <i>Pneumocystis carinii</i> pneumonia.</p> <ul style="list-style-type: none"> ■ Use of marijuana increases risk for lung cancer. ■ Snorting drugs, such as cocaine and heroin, and huffing, such as inhaling aerosols or glue, can result in lung damage, even respiratory failure. ■ Looks for exposure to respiratory irritants such as chemicals and coal dust. ■ Evaluates past employment for long-term exposure to asbestos and coal dust, which can cause cancer or black lung disease many years later. ■ Healthcare workers may be exposed to TB and other diseases. ■ Exposure to second-hand smoke increases the risk for lung cancer. ■ A safe work environment provides masks, good ventilation, and freedom from second-hand smoke. ■ Chronic respiratory disease affects job performance. Many employers allow respiratory-disabled people to work at home via computers. Explore this possibility with patients.
<p>Environmental Health Patterns</p> <ul style="list-style-type: none"> ■ Where do you live? ■ Is your home suitable for posthospital care or for a person with progressive respiratory dysfunction? ■ Is your home environment urban or rural? ■ Are you exposed to radon? ■ What kind of heating and air conditioning do you have? ■ How many people live with you? ■ Do you travel on freeways? 	<ul style="list-style-type: none"> ■ Some respiratory diseases are more prevalent in certain areas (e.g., histoplasmosis in Mississippi River Valley, coccidioidomycosis in Southwest). ■ Identifies discharge or home care needs, including durable medical equipment, supplies, caregivers. ■ Can identify air pollutants. ■ Radon increases risk for cancer. ■ Molds and dust in heating and air conditioning systems can trigger reactive airway disease. ■ Multiple families in crowded conditions contribute to risk of transmitting infectious respiratory diseases. ■ Freeway travel increases exposure to air pollutants.
<p>Roles/Relationships/Self-Concept</p> <ul style="list-style-type: none"> ■ What is your role in the family? In the community? ■ Have respiratory problems affected your family life, social life, or self-concept? 	<ul style="list-style-type: none"> ■ Chronic respiratory disease affects roles, relationships, and self-concept. ■ Economic strain can occur as a result of loss of work. ■ Limited mobility from chronic respiratory disease may lead to isolation and loss of community involvement.
<p>Cultural Influences</p> <ul style="list-style-type: none"> ■ Do you have any cultural influences that may affect your healthcare practices? 	<ul style="list-style-type: none"> ■ It is important to identify any cultural influences that may affect your patient's healthcare practices.



ALERT

Be alert for signs of depression.

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Religious/Spiritual Influences</p> <ul style="list-style-type: none"> Do you have any religious practices/beliefs that may influence your healthcare practices? <p>Sexuality Patterns</p> <ul style="list-style-type: none"> Has your respiratory problem affected your ability to perform sexually? <p>Social Supports</p> <ul style="list-style-type: none"> Are you involved in the community? <p>Stress and Coping</p> <ul style="list-style-type: none"> How do you manage anger, frustration, loss, and change? 	<ul style="list-style-type: none"> It is important to identify any religious practices/ beliefs that may affect your patient's healthcare practices. Loss of energy/SOB can affect sexual performance. Identifies positive support systems in family and community. Stress and coping patterns may be affected by respiratory status. Identifies and supports positive coping behaviors to decrease asthma attacks/dyspnea. For example, an action-reaction cycle occurs with stressed asthmatics: Stress triggers dyspnea; then as dyspnea worsens, person becomes more anxious. Learning appropriate coping skills to interrupt this cycle is important.

Fungal and Bacterial Respiratory Diseases

Histoplasmosis is a systemic, fungal respiratory disease whose organism is found in high-organic-content soil contaminated with bird droppings. Coccidioidomycosis (desert fever) is a self-limiting fungal respiratory disease that also has a more severe, progressive systemic form. Legionnaire's disease is a severe, often fatal, gram-negative bacterial disease resulting in pneumonia and multisystem failure. The disease was isolated after an outbreak at an American Legion convention.



Case Study Findings

Reviewing Mrs. Kane's

Biographical data:

- 45-year-old white female.
- Married, mother of three children, ages 15, 10, and 8.
- Part-time accountant; BA degree in accounting.
- Born and raised in United States.
- Catholic religion.
- Healthcare insurance (HMO) through husband's work.
- Referral:* Preadmission workup for surgery.
- Source:* Self, seems reliable.

Current health status:

- Brief morning cough for past 5 years; productive of small amount of thin, whitish-gray mucus, no odor or bad taste. Cough does not occur with activity or exercise. Positive history of smoking.
- No dyspnea or chest pain.
- No edema, loss of usual energy, change in sleeping patterns (except when gallbladder discomfort awakens her).

Past health history:

- Frequent upper respiratory infections (URIs) as a child.
- No history of asthma, pneumonia, or TB.
- Never had surgery. Hospitalized only for childbirth—uncomplicated vaginal delivery of three children.
- No known exposure to people with respiratory infections.
- Denies allergies to food, drugs, or environmental factors.
- No medical problems except for gallbladder disease.
- Immunizations up to date. Last purified protein derivative (PPD) test a year ago was negative.
- Denies taking prescribed drugs. Takes acetaminophen about twice a month for headache relief.

(continued on page 408)



Case Study Findings (continued)

Family history:

- Grandfather smoked and died of lung cancer at age 63.
- Father, age 65, recently quit smoking after smoking a pack a day for 40 years (40 pack-years). Diagnosed with lung cancer a year ago.
- All other known relatives alive and well.

Review of systems:

- *General Health Survey*: States that health is usually good. Has gained 15 lb in past 2 years.
- *Integumentary*: No changes in skin, hair, or nails.
- *HEENT*
 - *Head/Neck*: No swelling or masses.
 - *Eyes/Ears*: Vision 20/20; hearing good.
 - *Nose/Throat*: Productive morning cough with whitish-gray mucus. Has occasional cold.
- *Cardiovascular*: No chest pain.
- *Gastrointestinal*: Appetite good, no nausea or vomiting. Daily bowel movement.
- *Urinary*: No changes reported.
- *Reproductive*: Is in a monogamous relationship; satisfied with sexual performance.
- *Musculoskeletal*: No weakness or muscle wasting reported.
- *Neurological*: Memory good. No changes noted. No tremors.
- *Immune/Hematologic*: No allergies or anemia.

Psychosocial profile:

- Has a physical exam/Pap smear every 2 years. Made a special appointment when noticed RUQ (right upper quadrant) colicky abdominal pain after eating high-fat meals. Scheduled surgery as soon as recommended by physician.
- Typical day consists of arising at 6 A.M., having breakfast with family, and getting children off to school. Walks five blocks to and from work at a certified public accounting office 4 days a week. Works 8 A.M. to 4 P.M. at desk with computer. Spends evenings playing with children, helping them with homework, shopping, fixing dinner, doing housework, visiting with husband, and watching TV or reading. Showers, then goes to bed at 11 P.M.
- Usually eats a healthy diet, but has gained about 15 pounds over the last 2 years. Attributes this to sedentary job, sweet tooth, and lack of exercise.
- Walks to and from work and likes to ride a bike, but doesn't have much time because of family, home, and work responsibilities.

- No pets; hobbies include reading and biking.
- Sleeps about 6 to 7 hours a night, uninterrupted, with one pillow. Denies problems falling asleep, but awakens with morning cough.
- Has smoked one pack of filtered cigarettes a day for past 10 years (10 pack-years). Lived with smokers all her life. Has tried to quit by cutting back without success. Drinks alcohol about three times a month at social events. Denies recreational drug use.
- Denies respiratory irritants at work.
- Lives in two-story, single-family home in suburbs with ample living space for five people; has hot water, heating, and air conditioning; home negative for radon gas; denies recent travel outside local community. Will stay in downstairs bedroom for the first week after surgery.
- Supportive, caring husband; good relationship with children. Concerned about father's prognosis and mother's ability to care for him. Aware of health risks of second-hand smoke for children, so tries not to smoke at home. Husband also smokes, both would like to quit. No community involvement at this time, but has several friends who will help out after surgery.
- Copes with stress by talking with husband/friends or by ignoring it; also by smoking.



Case Study Evaluation

Before you proceed with the physical examination, document the key information you have learned from Mrs. Kane's health history.



CRITICAL THINKING ACTIVITY #1

Based on the subjective information from Mrs. Kane's history, what respiratory problems do you think she might have after surgery?



CRITICAL THINKING ACTIVITY #2

What strengths have you identified that might help Mrs. Kane avoid postoperative respiratory problems?



CRITICAL THINKING ACTIVITY #3

What suggestions would you make for postoperative care?

Anatomical Landmarks

Before you begin the physical examination, you need to visualize the underlying structures and identify your landmarks to ensure an accurate and thorough assessment. Approach the respiratory assessment from anterior, lateral, and posterior views. The right lung has three lobes, and the left has two. These lobes are divided by fissures (Fig. 13.3).

Imaginary lines are used to correctly identify landmarks and document findings. When approached anteriorly, the apices of the lungs extend about 2 cm ($\frac{3}{4}$ to 1 inch) above the inner aspect of the clavicles. They then continue downward to the sixth intercostal space at the midclavicular line (MCL). Keep in mind that access to the lungs from this approach may be limited by other structures such as breast tissue or the heart (Fig. 13.4).

Laterally, the lower border of the lung is at the eighth rib at the midaxillary line (Fig. 13.5).

The posterior approach affords the best access to the lungs because there are no underlying organs or structures to get in the way. Posteriorly, the apices of the lungs start at T1 and extend to T10 and T12 on deep inspiration. The right lung may be slightly higher because of the liver (Fig. 13.6).



HELPFUL HINTS

- Because of the anatomical position of the right middle lobe, the anterior and lateral approaches are best for assessment. The right middle lobe is not readily accessible using the posterior approach.
- An easy way to find T1 is to have the patient bend her or his neck. The most prominent spinous process is C7 and the one below is T1.




FIGURE 13.3. Lungs. A. Anterior view. B. Right lateral view. C. Left lateral view. D. Posterior view.




FIGURE 13.4. A. and B. Anterior views with landmarks.



CRITICAL THINKING ACTIVITY #4

The gallbladder is located in the right upper quadrant of the abdomen. How might this location affect the respiratory system?

Physical Assessment

Now that you have completed subjective data collection, begin collecting objective data by means of the physical examination. Keep key history findings in mind. These findings, along with those from the physical examination, complete your assessment picture. Now you can analyze data, formulate nursing diagnoses, and develop a plan of care.

Approach

You will use all four techniques of physical assessment to examine the respiratory system. Begin with inspection, proceed to palpation and percussion, and then perform auscultation from the anterior, lateral, and posterior approaches. The best position for the examination is with the patient sitting. For the posterior approach, ask the patient to lean forward and cross his or her arms over the chest. This spreads the scapula, providing the greatest access to the lung surface. Perform all assessment techniques at each approach before changing the patient's position.

If the patient is unable to sit up and has to be in a supine, semi-Fowler's, or sidelying position, your findings will be distorted. Because the physical findings are more

evident on the dependent side, help the patient turn, and then reassess her or him from the other side.

During auscultation, have the patient breathe slowly and somewhat deeply through the mouth. Be alert for dizziness and other signs of hyperventilation. To guard against this, caution the patient to take slow, deep breaths and to tell you if he or she becomes dizzy.

Changes in the respiratory system can affect every other system, so you may see physical signs in many systems. Normal findings may vary from patient to patient, so a systematic approach is essential. Proceed from apex to base, comparing one side with the other, using the patient as her or his own control. Establish a baseline that you can use to compare the patient's respiratory status in subsequent assessments.



TOOLBOX

The tools for respiratory assessment include a stethoscope, felt-tipped marker, and metric ruler. Also, sharpen your senses. You will use all of them during this examination.



HELPFUL HINT

Minimize position changes when examining older people or those with chronic respiratory disease. It helps conserve their energy.




FIGURE 13.5. A. Right lateral view with landmarks. B. and C. Left lateral views with landmarks.

Performing a General Survey

Before assessing specific areas, scan the patient from head to toe, looking at general appearance and for signs of respiratory dysfunction. For example:

- Check for consistency of appearance with chronological age. Chronic illness or long-term smoking can make people appear older than they are.
- Examine weight distribution and muscle composition. Cortisone therapy can alter weight distribution and cause muscle atrophy of extremities.
- Evaluate nutritional status. People with lung cancer and chronic lung disease are frequently cachectic. Those on cortisone or with edema may appear overweight.
- Note anxiety, fear, distress, or pain. People with dyspnea or chest pain exhibit these signs.
- Consider the person's posture and assumed position. Is he or she erect and relaxed, slouched, hypererect, or leaning forward and/or using the chair arms for support (tripod position)? People with chronic obstructive pulmonary disease (COPD) or dyspnea often prefer an erect sitting or tripod position.




FIGURE 13.6. A. and B. Posterior view with landmarks.

- Take vital signs (blood pressure [BP], temperature, pulse, and respirations) and weigh the patient. Include these results in your database. Changes in pulse and BP may have a cardiopulmonary basis. Weight changes are often seen in patients with chronic respiratory disease.

Temperature

An elevated temperature may indicate a bacterial, viral, or fungal infection. Fever can also occur in noninfectious conditions such as pulmonary embolism. Remember that people who are immunosuppressed, elderly, or on corticosteroid therapy may not have a fever even with an infection.

Respirations

Eupnea refers to normal rate, depth, and rhythm of respirations. Rate varies with age. Normal adult rate is 14 to 20 breaths per minute. Depth/tidal volume for an adult is 300 to 500 mL/min, considered moderate. Rhythm should be regular, with signs every 15 minutes at rest. Respiration should be quiet and relaxed unless the patient is involved in vigorous activity.

Respiration rate, depth, and rhythm are affected by respiratory, cardiac, neurological, metabolic, and emotional disorders, as well as by medications.

Rate

- Tachypnea (increased rate):** Can be caused by hypoxia, metabolic acidosis, anxiety, fear, pain, compromised neurological control of breathing, sepsis, fever, or increased metabolism.

- Bradypnea (decreased rate):** Results from excessive sedation, hypercapnea, compromised neurological control of breathing, or metabolic alkalosis.

Depth and Rhythm

- Shallow respiration:** Decreased depth may be a result of habit; fatigue; metabolic alkalosis; ascites; restrictive lung disease; chest, abdominal, or pleuritic pain; or neurological disorders.
- Increased depth:** May be related to neurological disorders, hyperventilation with anxiety, or metabolic acidosis (especially with tachypnea).

Abnormal Patterns

- Kussmaul:** Rapid, deep respiration associated with metabolic acidosis (body's attempt to blow off CO₂), seen in diabetic ketoacidosis or lactic acidosis.
- Cheyne-Stokes:** Progressively increasing rapid, deep respiration that peaks and then gradually ceases, followed by a period of apnea, after which the pattern recurs. Can be drug-induced, related to heart or renal failure, a sign of brain damage or impending death, or normal in frail elderly people during sleep.
- Biot's:** Ataxic breathing pattern that is irregular in rate and depth and alternates with irregular periods of apnea. Seen in respiratory depression, damage to medullary respiratory centers, or head injury.

Performing a Head-to-Toe Physical Assessment

The respiratory system affects every other system, so look for changes from head to toe in each system that might signal a respiratory problem.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

ABNORMAL/RATIONALE

General Health Survey

INSPECT

- Awake, alert, oriented to time, place, person.
- Immediate, recent, remote memory intact.
- Judgment intact; compliant with treatment plan.

- Early signs of hypoxia include confusion, restlessness, irritability, short attention span.
- Short-term memory may also be affected by hypoxia.
- People with chronic respiratory disease may deny illness, severity of symptoms, or associated risk factors.
- Anxiety can trigger a respiratory problem, such as hyperventilation, or can result from a respiratory problem, such as an acute asthmatic attack or pulmonary emboli.
- Fatigue is associated with chronic lung disease, lung cancer, and pneumonia because more energy is expended on breathing.

Integumentary Skin and Nails

INSPECT



HELPFUL HINT

When assessing for color changes in dark-skinned people, look at mucous membranes or conjunctiva.

Skin color varies with racial and genetic heritage. Buccal mucosa, tongue, palates, conjunctiva and nail beds are usually pink, but may be darker pink, dark brown, or even blue in people of Mediterranean or African descent.

- Peripheral cyanosis may be normal response to exposure to cold temperatures.
- No edema.
- Positive capillary refill, nailbeds pink, negative clubbing.

- *Pale diaphoretic*: Sympathetic response to respiratory distress and hypoxia.
- *Central cyanosis*: Dusky or blue buccal mucosa and tongue when PO_2 is less than 50. Caused by respiratory failure, shock, pulmonary edema, airway obstruction, ventilation-perfusion problems. May see cyanosis in vessels of palpebral conjunctiva in end-stage COPD.
- *Peripheral cyanosis*: Blue dusky-red, or purple color of lips, nail beds, tips of nose and ears, sometimes face, and especially cheeks. Caused by slow or congested blood flow in peripheral vessels or disorders with compensatory polycythemia, such as COPD, CHF.
- *Ruddy, reddish color*: Associated with polycythemia.
- Peripheral edema frequently seen in people with chronic lung disease secondary to right-side CHF.
- *Clubbing*: Long-standing lung disease.
- *Cyanotic or dusky nails*: Reflect peripheral cyanosis seen in vasoconstriction and slowing of peripheral blood flow. Also associated with central cyanosis.
- *Yellow-brown stains on nails and fingers*: Nicotine stains from long history of smoking.
- *Purple/dusky lower extremities*: Venous stasis, especially if PO_2 is low.

HEENT

Head and Neck

INSPECT/PALPATE

- Trachea should arise out of sternal notch and be midline.
- No pursed-lip breathing, nasal flaring, neck vein distension, or use of accessory muscles unless associated with strenuous activity. Visible neck vein in older people or those with overdeveloped neck muscles. No hypertrophy of neck muscles unless occupationally related (e.g., weightlifters, singers, professional speakers). No visible/palpable lymph nodes.
- *Tracheal shifts*: Collapsed lung, tumors, pneumothorax, or hemothorax.
- Pursed-lip breathing, or physiological positive end-expiratory pressure, is a compensatory mechanism used by people with COPD to prolong expiration help expel trapped air, and keep alveoli open longer for maximum oxygenation of pulmonary blood. Nasal flaring occurs in infants and small children and indicates acute respiratory distress or cyanotic heart disease.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL/RATIONALE
<p>Eyes</p> <ul style="list-style-type: none"> ■ Conjunctiva pink, no excessive tearing. ■ No papilledema. <p>Ears/Nose/Throat</p> <ul style="list-style-type: none"> ■ Oral and nasal mucous membranes pink, TM membrane pearl, gray, nares patent, no drainage. <p>Cardiovascular</p> <p>PALPATE/AUSCULTATE</p> <ul style="list-style-type: none"> ■ Heart, regular rate and rhythm (HRRR), no extra sounds, +2 pulses, negative Homans' sign. <p>Gastrointestinal</p> <p>PALPATE/PERCUSS</p> <ul style="list-style-type: none"> ■ No hepatomegaly or ascites. <p>Musculoskeletal</p> <p>INSPECT/PALPATE</p> <ul style="list-style-type: none"> ■ +5 muscle strength, no weakness or muscle wasting <p>Neurological</p> <p>INSPECT</p> <ul style="list-style-type: none"> ■ No tremors, +2 deep tendon reflex (DTR) 	<ul style="list-style-type: none"> ■ <i>Neck vein distension</i> is general sign of respiratory distress or air hunger. When client sits, elevating head and chest 45 degrees, neck vessels are firm and tortuous. This sign of increased venous pressure is seen in right heart failure and cor pulmonale. People with chronic lung disease may also develop hypertrophied neck muscles from increased efforts to breathe. ■ <i>Use of scalene, sternocleidomastoideus (SCM), and trapezius accessory muscles</i>: Sign of respiratory distress or COPD. ■ <i>Visible/palpable lymph nodes</i>: Infection or malignancy ■ Central cyanosis can be seen in the conjunctiva ■ <i>Excessive tearing</i>: Allergies. ■ <i>Papilledema</i>: Hypercapnea or CO₂ narcosis. ■ <i>Red mucous membranes, red TM, drainage</i>: Upper respiratory infection. ■ <i>Right-sided S₃ and S₄</i> Right CHF and pulmonary HTN. ■ <i>Accentuated P₂</i>: Pulmonary HTN. ■ <i>Absent/diminished pedal pulses and ankle/pedal edema</i>: Heart failure and cor pulmonale. ■ Clients with polycythemia are at risk for thrombophlebitis and possibly pulmonary embolism caused by sluggish blood flow. A positive Homans' sign may indicate thrombophlebitis, but it is not conclusive. ■ <i>Enlarged liver and ascites</i>: Right-side CHF, a common complication of chronic lung disease. ■ <i>Weakness/muscle wasting</i>: Long-standing respiratory disease. ■ <i>Tremors, seizures, decreased DTRs</i>: Respiratory failure. ■ <i>Asterixis (flapping hand tremors)</i>: CO₂ narcosis.

Clubbing

In people with chronic hypoxia associated with chronic lung disease, the kidneys compensate by increasing secretion of erythropoietin, a hormone that stimulates red blood cell (RBC) production. The increase in RBCs (polycythemia) results in thick, sluggish blood flow that “plugs” the capillaries of the nail folds, causing the tissue to become swollen and spongy. The result is “clubbing”—a nail plate that becomes more convex, with a rounded distal phalanx and an increased angle of attachment.

Perfusion/Ventilation Problems

Ventilation-perfusion problems occur when one or more areas of the lung receive adequate ventilation but inadequate blood flow (perfusion problem) or inadequate ventilation and adequate blood flow (ventilation problem). This results in increased dead space, shunting, and hypoxemia. Emphysema causes increased dead space and creates a ventilation problem. A pulmonary embolus causes inadequate blood flow and shunting and creates a perfusion problem.

Performing a Respiratory Physical Assessment

Now that you have completed your general survey and head-to-toe scan, you can focus on the respiratory assess-

ment. Perform the assessment anterior/posterior/lateral, Apex to base, comparing side with side. Begin with inspection, then palpate, percuss, and finally auscultate the lungs.

Respiratory Patterns

The following are normal and abnormal respiratory patterns.

Eupnea

Normal respiratory rate and rhythm



Apnea

Absence of breathing (may be periodic)



Apneustic

Prolonged, gasping inspiration followed by extremely short, inefficient expiration



Biot's

Faster and deeper respirations than normal, with abrupt irregular pauses between them



Bradypnea

Slow but regular respirations



Cheyne-Stokes

Respirations that gradually become faster and deeper than normal, then slower; alternates with periods of apnea



Hyperventilation

Deeper respirations; normal rate



Kussmaul's

Faster and deeper respirations without pauses



Tachypnea

Increased respiratory rate






FIGURE 13.7. Chest shapes.

Inspection

Begin with inspection. Inspect the chest for size, shape (Fig. 13.7), symmetry, and excursion. Ask yourself: Does the chest rise equally? Are respirations quiet? What

muscles are being used to breathe—chest or abdominal? Are there signs of respiratory distress, such as retractions? Inspect the skin, noting scars or color changes.

INSPECTING THE CHEST

1. Anterior Chest

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect respiratory rate, rhythm, depth, and symmetry of chest movements.
- Inspect anteroposterior (AP) to lateral ratio, costal angle, spinal deformities, and condition of skin.
- **Shape and symmetry:** Normal adult chest has AP-to-lateral ratio of approximately 1:2 and costal angle less than 90 degrees. Ribs slope obliquely, chest symmetrical in appearance with symmetrical rise and fall when breathing. Skin intact.
- **Movement with breathing:** Women have more thoracic respiratory movements; men and infants have more abdominal respiratory movements.
- No sternal or intercostal retraction or bulging, unless associated with strenuous activity.
- Musculoskeletal changes associated with aging can increase the AP diameter, giving chest a barrel appearance.
- AP diameter greater in infants (apple-shaped chest).
- **Condition of chest skin:** Skin color and hair distribution should be consistent with patient's gender, ethnicity, and exposure to sun. Skin intact, no scars.



Inspecting anterior chest

INSPECTING THE CHEST

1. Anterior Chest (continued)

ABNORMAL FINDINGS/RATIONALE

- **Altered chest shape:** Seen in COPD (trapped air with overinflated lungs). AP diameter is increased, resulting in “barrel chest.” Rib slope is nearly parallel, with costal angle greater than 90 degrees (see Fig. 13.7).
- **Altered chest symmetry:** Seen in musculoskeletal disorders of spine such as kyphosis, scoliosis, and kyphoscoliosis. Can affect respiratory function by restricting movement.
- **Altered breathing symmetry:** May be caused by rib fractures, especially at sternal border (flail chest), pneumonectomy, pneumothorax/hemothorax, and atelectasis. Affected area of chest may not move with respiration.
- **Sternal and intercostal retractions:** Seen in severe hypoxia or respiratory distress, especially with airway obstruction.
- **Altered skin color/condition:** Extreme hypoxia or cold temperature may cause blue flush on chest wall (cyanosis). Steroids may produce excessive hair in women.
- **Intercostal bulging:** Seen during expiratory effort of person with COPD.
- Scars may indicate trauma or surgery. Look for signs of lung surgery (often on lateral thorax) or 1-cm stab wounds from chest tubes.

2. Lateral Chest

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect respiratory rate, rhythm, depth, and symmetry of chest movements.
- Inspect AP to lateral ratio and condition of skin.
- *As with anterior and posterior chest, skin is intact and chest expansion equal.*



Inspecting right lateral chest



Inspecting left lateral chest

ABNORMAL FINDINGS/RATIONALE

- Same as with anterior and posterior chest. Look for scars from pneumonectomy or lobectomy, located laterally and curving under scapula.

3. Posterior Chest

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS



Inspecting posterior chest

- Inspect respiratory rate, rhythm, depth, and symmetry of chest movements.
- Inspect AP to lateral ratio, spinal deformities, and condition of skin.
- *Skin should be intact, chest expansion equal, and spine straight without lateral curves or deformities.*

INSPECTING THE CHEST

3. Posterior Chest (continued)

ABNORMAL FINDINGS/RATIONALE

- Same as with anterior chest, although abnormal spinal curves are more obvious. Look for scars from pneumonectomy or lobectomy, located laterally and curving under scapula.

Palpation

Palpation is useful in assessing for tracheal position, tenderness and crepitus, chest excursion, and tactile fremitus. Use light palpation to assess surface characteristics of the chest, such as temperature, turgor, and moisture, as well as to identify tenderness, masses, or crepitus. Palpate from the anterior, posterior, and lateral approaches.

To assess tracheal position, place a finger on either side of the trachea. The distance between the trachea and the sternocleidomastoid muscle on both sides should be equal.

Chest excursion refers to the chest's expandability. Use this technique to assess symmetrical chest expansion with both anterior and posterior approaches. Place your hands vertically on the patient's chest, with fingers spread and thumbs at the costal margin anteriorly and at the eighth to the tenth rib posteriorly. Your palms should be firm against the skin, with thumbs touching and fingers lightly in contact with the chest wall. Gently gather a small fold of skin between your contacting thumbs. Have the patient take a deep breath. You should feel equal pressure on your hands,

and your thumbs should move apart evenly. If you note any abnormality, move your hands up and assess the apices of the lung.

Tactile or vocal fremitus is the palpable vibration you feel when the patient speaks. Assessment of tactile fremitus is used to evaluate airflow and density of underlying tissue. Because bony prominences are best for detecting vibratory sensations, use the balls or the ulnar surfaces of your hands. Have the patient say "99, 99" while you move your hands in a systematic fashion from apex to base, comparing side with side and anteriorly and posteriorly. Note the level where fremitus is palpable and where it diminishes and goes away. In the average adult, fremitus is not palpable below the third to fourth intercostal space anteriorly and at about T6 to T8 posteriorly.

Several factors besides airflow influence the quality of tactile fremitus. These include chest wall thickness, voice pitch, and airway size. Normally, the thicker the chest wall, the more diminished the fremitus; and the lower the voice pitch, the greater the fremitus. Also, fremitus is usually increased over the larger airways (Fig. 13.8). (See *Palpating the Chest*.)




FIGURE 13.8 . Palpation sequences. A. Anterior. B. Posterior.

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Changes in energy level Activity intolerance Fatigue Weight changes 	General	<p><i>Inspect:</i></p> <ul style="list-style-type: none"> Signs of acute distress, such as SOB, DOE Posture: Tripod; Position: Orthopnea <p><i>Measure:</i></p> <ul style="list-style-type: none"> Height, weight (check for changes) Temperature, pulse, respirations, B/P (check for increases)
Integumentary	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Skin color changes Fevers, night sweats 	Integumentary	<p><i>Inspect:</i></p> <ul style="list-style-type: none"> Skin color changes (e.g., cyanosis, pallor, ruddiness) Central (mucous membranes) vs. peripheral cyanosis Nail clubbing <p><i>Palpate:</i></p> <ul style="list-style-type: none"> Skin temperature, turgor, edema Capillary refill
HEENT <i>Head, Face, Neck, Eyes, Ears, Nose and Throat</i>	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Lumps or swelling in neck Excessive eye tearing Ear infections, sore throats, URIs, sinus problems Difficulty swallowing or breathing 	HEENT <i>Head, Face, Neck, Eyes, Ears, Nose and Throat</i>	<p><i>Inspect:</i></p> <ul style="list-style-type: none"> Facial expression (e.g., anxious) Neck vein distention, hypertrophy and use of accessory neck muscles Color of conjunctiva Ear or nose drainage, nasal flaring Color of mucous membranes, color of tonsils and enlargement <p><i>Palpate:</i></p> <ul style="list-style-type: none"> Sinus tenderness, lymph nodes, tracheal position, thyroid gland, patent nares, tonsillar glands <p><i>Examine:</i></p> <ul style="list-style-type: none"> Fundus, optic disk External ear and TM Internal nasal mucosa and structures
Cardiovascular	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Chest pain Palpitations Swelling, tight shoes 	Cardiovascular	<p><i>Inspect:</i></p> <ul style="list-style-type: none"> Edema <p><i>Palpate:</i></p> <ul style="list-style-type: none"> Edema, Homans' sign <p><i>Auscultate:</i></p> <ul style="list-style-type: none"> Right side S4 and S3
Gastrointestinal	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Changes in appetite GI complaints RUQ pain 	Gastrointestinal	<p><i>Palpate:</i></p> <ul style="list-style-type: none"> Enlarged liver, ascites
Genitourinary/ Reproductive	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Nocturia Changes in sexual activity Safe-sex practices Pregnancy 	Musculoskeletal	<p><i>Inspect:</i></p> <ul style="list-style-type: none"> Hypertrophy and use of accessory muscles, muscle atrophy, spinal deformities <p><i>Measure:</i></p> <ul style="list-style-type: none"> Muscle strength
Musculoskeletal	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Weakness 	Neurological	<p><i>Inspect:</i></p> <ul style="list-style-type: none"> Impaired mental status, AAO x 3 Asterixis (flapping tremors)
Neurological	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Memory changes Morning headaches Tremors 		
Endocrine	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Thyroid disease 		
Lymphatic/ Hematologic	<p><i>Ask about:</i></p> <ul style="list-style-type: none"> Bleeding, anemia Allergies 		

Assessment of Respiratory System's Relationship to Other Systems

PALPATING THE CHEST

1. Tracheal Position

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Place your thumb and index finger on either side of the trachea, and note position and distance between trachea and sternocleidomastoid muscle.
- *Trachea should be midline.*



Palpating tracheal position

ABNORMAL FINDINGS/RATIONALE

- *Tracheal deviation:* Tumor or thyroid enlargement may cause tracheal deviation. Tension pneumothorax deviates trachea *away from* affected lung. In severe atelectasis, trachea may deviate *toward* affected lung.

2. Chest Tenderness and Crepitus

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use light palpation to assess for tenderness and crepitus.
- *Nontender. No deformities or crepitus.*



HELPFUL HINTS

- Chest pain that is respiratory or cardiac in origin is unaffected by palpation.
- Crepitus feels crackly, like crumpling cellophane. Although air leaking into the tissue does not hurt it, the leak must be found because that air is not going into the lungs.



Light palpation of anterior chest.



Light palpation of posterior chest.

ABNORMAL FINDINGS/RATIONALE

- *Pain/tenderness at costochondral junctions/vertebral connections of ribs:* Fracture. May also be present with palpation of inflamed cartilage at the rib/sternal junctions (costochondriasis).
- *Spontaneous pathological fractures:* Corticosteroid therapy or osteomalacia.
- *Crepitus (subcutaneous emphysema):* Results from air leaking into subcutaneous tissue. Check around wound sites, chest tubes, central lines, or tracheostomy tubes.

PALPATING THE CHEST

3. Chest Excursion

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Anteriorly, place hands vertically on the chest with fingers spread on the costal margin and thumbs together at the costal angle (like a butterfly).
- Posteriorly, place hands vertically on the chest with fingers spread and the thumbs together at the spine at the eighth to tenth rib (like a butterfly).
- Feel for equal expansion of the chest; if absent, decreased, or unequal, move up the chest.
- *Symmetrical without lag. Normal causes of decreased excursion are habitual shallow breathing or wearing tight, restrictive undergarments.*



Palpating anterior chest excursion at apex



Palpating anterior chest excursion at base



Palpating posterior chest excursion at apex



Palpating posterior chest excursion at base

ABNORMAL FINDINGS/RATIONALE

- *Asymmetrical excursion:* Associated with thoracotomy (removal of lung or lobes), complete or partial airway obstruction, pleural effusion, and pneumothorax.
- *Decreased excursion:* May occur in patients with overinflated lungs and fixed diaphragm (COPD). May also be present with splinting because of pain of fractured ribs.

4. Tactile Fremitus

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Place the balls of your hands with your fingers hyperextended or the ulnar surface of your hand on the patient's chest.
- Have patient say "99" as you palpate vibrations.
- Note level where fremitus is palpable, increased, diminished, or absent.
- *Equal bilaterally and diminished midthorax.*
- *Normal causes of diminished fremitus are thick chest wall or very soft voice. Young children and very thin people may have increased fremitus.*



Palpating anterior tactile fremitus



Palpating posterior tactile fremitus

PALPATING THE CHEST

4. Tactile Fremitus (continued)

ABNORMAL FINDINGS/RATIONALE

- *Increased fremitus:* Occurs with conditions causing fluid or exudates in lungs, such as consolidating pneumonia, atelectasis, pulmonary fibrosis, pulmonary edema, or pulmonary infarction. May also occur with lung tumor, although tumor may stop vibrations, depending on its size and mobility.
- *Decreased or absent fremitus:* Occurs where there is air trapping, solid tissue, or decreased air movement—for example, emphysema, asthma, pleural effusion, pneumothorax, or distal to airway obstruction.




FIGURE 13.9. Percussion sequence. A, Anterior. B, Right lateral. C, Left lateral. D, Posterior.

Percussion

Percussion is used to assess the density of underlying lung tissue. It allows you to determine if the tissue is predominantly solid or filled mainly with air or fluid. Percussion also enables you to identify the extent of the lung fields and is useful in assessing **diaphragmatic excursion**, the distance the diaphragm moves from expiration to inspiration. Percuss the entire anterior and posterior thorax in a systematic manner, moving from apex to base and comparing side with side. The normal percussion note over adult lung fields is called **resonance**; small children will percuss more hyperresonant. It is helpful to percuss over the intercostal spaces, because percussion over bone produces a less resonant sound.

Percussion is a skill that is perfected through practice. You need to develop that skill, as well as the ability to differentiate the sounds produced by percussion. Percussion sites are the same as palpation sites anteriorly (Fig. 13.9). For more information on percussion sounds, see Chapter 3, Approach to the Physical Assessment.

Diaphragmatic Excursion

To assess diaphragmatic excursion, percuss the level of the diaphragm on full expiration and on full inspiration, noting the difference in distance in centimeters between the two. First, have the patient exhale fully and then hold her or his breath as you percuss from resonance over the lung tissue to dullness over the diaphragm. Mark this area. Next, have the patient take a deep breath and hold it. Continue percussing at the same location, moving from resonance to diaphragmatic dullness. Again, mark the area. Normally, the percussion that was previously dull at the diaphragm should change to resonance as the lung fills with air and the diaphragm moves downward. Measure the distance between the two marks in centimeters. Repeat the procedure on the opposite side. Normal diaphragmatic excursion is 3 to 6 cm. Remember: Work quickly to get the most accurate results and not tire the patient. (See *Percussing the Chest*.)

PERCUSSING THE CHEST

1. Anterior Thorax

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Using indirect or mediate percussion, percuss over intercostal spaces.
- Note areas of resonance, hyperresonance, or dullness.
- *Resonance to second intercostal space on left; slight dullness over third through fifth intercostal space over heart.*
- *Resonance to fourth intercostal space on right with dullness from approximately fifth to just above costal margin over liver. It will vary with person's size and size of liver.*



Percussing anterior chest

ABNORMAL FINDINGS/RATIONALE

- *Dullness:* Seen with exudate, fluid, tumors, pneumonia, pulmonary edema, pleural effusion.
- *Hyperresonance:* Noted with air trapping of emphysema.

2. Lateral Thorax

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Using indirect or mediate percussion, percuss over intercostal spaces.
- Note areas of resonance, hyperresonance, or dullness.
- *Resonance to eighth intercostal space.*

PERCUSSING THE CHEST

3. Posterior Thorax

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Using indirect or mediate percussion, percuss over intercostal spaces.
- Note areas of resonance, hyperresonance, or dullness.
- *Resonance to T10-T12 with deep inspiration.*



Percussing posterior chest

ABNORMAL FINDINGS/RATIONALE

- *Hyperresonance to T12 level:* Overinflated lungs.

4. Diaphragmatic Excursion

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Have patient take a deep breath and fully exhale, then percuss the level of the diaphragm and mark.
- Have the patient take a deep breath and hold it, then percuss the level of the diaphragm.
- Measure the distance between the two marks.
- *3–6 cm diaphragmatic excursion.*



Measuring diaphragmatic excursion

ABNORMAL FINDINGS/RATIONALE

- *Decreased diaphragmatic excursion unilaterally or bilaterally:* Paralyzed diaphragm, atelectasis, COPD with overinflated lungs.

Auscultation

Use auscultation to assess airflow through the upper airways and lungs. As you listen with the diaphragm of the stethoscope, note normal, abnormal, and adventitious breath sounds, as well as abnormal vocal sounds. Be sure the room is private, warm, and quiet, and warm the diaphragm of the stethoscope between your hands before you begin. Auscultation sites are the same as for percussion.



HELPFUL HINTS

- If the patient has a hairy chest, moisten the chest hair before auscultating to reduce interference.
- Be sure to use all of your assessment techniques, and do not rely on one finding. For example, you may hear transmitted sounds over an area with no airflow, when in fact, sounds are absent. If there is no airflow, there will be no chest movement.

AUSCULTATING THE LUNGS

1. Testing for Normal Breath Sounds in Appropriate Locations

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Using the diaphragm portion of the stethoscope, auscultate over auscultatory sites anterior, posterior, and lateral.
- Listen to one full respiratory cycle at each site.
- Note normal, abnormal, and adventitious sounds.
- If adventitious sound heard, have patient cough then listen to see if sound has cleared.
- *Bronchial breath sounds are loud, high-pitched, and hollow, with a short inspiratory phase and a long expiratory phase. They are normally heard in anterior neck and nape of neck posteriorly.*
- *Bronchovesicular breath sounds are moderate sounding and medium pitched, with equal inspiratory and expiratory phases. They are normally heard over first to second interspaces anteriorly and between scapula posteriorly.*
- *Vesicular breath sounds are soft and low pitched, with a long inspiratory phase and a short expiratory phase. They are heard over most lung fields.*
- *With no adventitious sounds, lungs are clear to auscultation. No crackles, wheezes, or rubs.*



Auscultating bronchial breath sounds over trachea



Auscultating bronchovesicular breath sounds



Auscultating vesicular breath sounds.

ABNORMAL FINDINGS/RATIONALE

- *Bronchial sounds heard outside of their normal locations:* Fluid or consolidated tissue, such as in pneumonia.
- *Diminished breath sounds:* Thick, obese, or muscular chest wall, poor inspiratory effort, emphysema with air trapping, pleural effusion.
- *Absent breath sounds:* Missing lung/lobe, airway obstruction, pneumothorax.
- *Crackles/Rales:* Pulmonary edema, pneumonia, atelectasis, and upon arising in elderly people.

AUSCULTATING THE LUNGS

1. Testing for Normal Breath Sounds in Appropriate Locations (continued)

- *Velcro rales*: Pulmonary fibrosis.
- *High-pitched, sibilant wheezes*: Bronchospasm, asthma, COPD without infection.



ALERT

The disappearance of wheezes may be an ominous sign if associated with decreased breath sounds caused by decreased air movement. This may indicate severe obstruction and impending respiratory failure.

- *Low-pitched sonorous wheezes/rhonchi*: Uncleared secretions, bronchitis, pneumonia, tumors.
- *Stridor*: Laryngeal or tracheal obstruction, epiglottitis, viral croup.
- *Pleural friction rub*: Inflammation of pleura.

2. Testing for Abnormal Voice Sounds

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Egophony*: Have patient say “ee”; will sound like “aa” over affected area.
- *Bronchophony*: Have patient say “1, 2, 3”; clearer transmission of spoken voice sound over affected area.
- *Whispered pectoriloquy*: Have patient whisper “1, 2, 3”; clearer transmission of whispered voice sound over affected area.

ABNORMAL FINDINGS/RATIONALE

- *Abnormal voice sounds*: Consolidating pneumonia.

Normal Breath Sounds

With the patient in a sitting position, auscultate directly over the skin, exposing only the areas being assessed. Listen through at least one full respiratory cycle at each site. Be systematic in your approach, and follow the same sequence as percussion: Auscultate the anterior, lateral, and posterior thorax, moving from apex to base and comparing side with side. To help differentiate breath sounds, note the relationship of inspiration to expiration (I:E) and which phase is longer, identify the pitch and intensity of the sound, and locate the area where the sound is heard best (Fig. 13.10). (See *Normal Breath Sounds*.)

Abnormal Breath Sounds

Breath sounds that are diminished or misplaced are referred to as *abnormal*. Breath sounds may be decreased or diminished in a person with a thick chest wall or pleura, with shallow breathing, or with restricted/obstructed airflow. They may also be diminished in an obese or mus-

cular person or in an older person with poor inspiratory effort.

A bronchial breath sound heard anywhere except over the trachea is a misplaced sound. For example, bronchial breath sounds are heard over areas of consolidation.

Be sure to work from side to side, comparing the patient with himself or herself. Examples of pathology associated with diminished breath sounds include emphysema, pleural effusion, and airways blocked by mucus plugs. Remember: Do not rely on one assessment finding; evaluate the patient’s entire picture before coming to a decision.



HELPFUL HINT

If you hear any abnormal sounds, have the patient cough, and then listen again.




FIGURE 13.10. Normal breath sounds. A. Anterior. B. Right lateral. C. Left lateral. D. Posterior.

Adventitious Breath Sounds


Adventitious breath sounds are additional sounds superimposed over normal breath sounds and include crackles or rales, rhonchi, wheezing, stridor, grunting, and friction rubs. Unfortunately, there is no universal classification of adventitious sounds. You may hear the terms *rales* and *crackles* used interchangeably. Adventitious sounds are often classified as **continuous** and **discontinuous**. Understanding the underlying mechanism may help you to differentiate sounds. Quality, pitch, location, and timing are important characteristics that will also help you differentiate the sounds and determine their significance. (See *Adventitious Breath Sounds*.)

Rales/Crackles


Rales or **crackles** are discontinuous sounds resulting from air bubbling through moisture in the alveoli or from collapsed alveoli popping open. Crackles tend to occur at the end of inspiration, in the terminal bronchioles and alveoli. Depending on the cause, crackles may be affected by coughing; loose exudate rales may clear with cough, but the rales of pulmonary edema do not.




Rales associated with long-standing chronic lung disease or with interstitial pulmonary fibrosis are sometimes referred to as *Velcro rales* because they sound like Velcro being pulled apart. Velcro rales are louder than fluid rales and usually do not change over time. They

Normal Breath Sounds

INSPIRATION:EXPIRATION	PITCH	INTENSITY	LOCATION
 <p>Vesicular</p>	Low	Soft	Most of lung field
 <p>Bronchovesicular</p>	Medium	Medium	Mainstem bronchi
 <p>Bronchial</p>	High	Loud	Trachea

Adventitious Breath Sounds

SOUND/DESCRIPTION	INSPIRATION:EXPIRATION	CONDITIONS	LOCATION
 <p>Crackles</p> <p>Popping; discontinuous; may be affected by coughing.</p>	More prominent on inspiration	CHF, atelectasis, pneumonia, interstitial fibrosis	Periphery
 <p>Rhonchi</p> <p>Snore, gurgle,</p>	More prominent on expiration	Pneumonia	Larger airways
 <p>Mild Wheeze</p> <p>Musical, high pitched; continuous; may be affected by coughing.</p>	More prominent on expiration, but as severity increases, extends through inspiration and expiration	Asthma, cardiac asthma	Smaller airways

SOUND/DESCRIPTION	INSPIRATION:EXPIRATION	CONDITIONS	LOCATION
 <p>Moderate Wheeze</p>			
 <p>Severe Wheeze</p>			
 <p>Friction Rub</p> <p>High pitched, scratching, squeaking; continuous; not affected by coughing.</p>	Inspiratory and expiratory	Pleuritis Pleurisy	Affected area

result from air trying to force through fibrotic tissues or fibrotic tissue rubbing against the visceral pleural. Velcro rales are unaffected by coughing.

Rhonchi/Wheezes

Wheezes are continuous sounds caused by the narrowing of an airway by spasm, inflammation, mucus secretions, or a solid tumor. Pitch is determined by the relative tightness/narrowness of the airway. For example, in asthma, tightly constricted small airways produce very high-pitched, or sibilant, wheezes. However, the loose mucus secretions in a larger airway produce lower-pitched, sonorous wheezes, often called **rhonchi**. These wheezes may even have a snoring or rattle-like quality.

Although rhonchi may occur in either the inspiratory or the expiratory phases of breathing, they are more commonly heard during expiration when air is being forced through the obstruction or the narrowed airway. Wheezes may clear with cough if the airway narrowing is caused by secretions. So if you hear adventitious sounds, ask the patient to cough, and then reassess the lung sounds.



HELPFUL HINTS

The sudden absence of wheezes does not always indicate that the patient's condition has improved. It may actually be a sign of a worsening condition. If the patient is having difficulty speaking, she or he may be having trouble moving air.

Stridor

Stridor is a harsh, high-pitched, continuous honking sound resulting from an upper airway obstruction, a partial obstruction, or a spasm of the trachea or larynx. A person with stridor is usually in acute respiratory distress and requires immediate intervention.

Grunting

Grunting is a larger airway sound heard predominantly on expiration. It results from retention of air in the lungs, which prevents alveolar collapse.

Friction Rub

A **friction rub** is different from all other adventitious sounds because it occurs between the pleural layers, not in the lung. It results from the rubbing together of the parietal and visceral layers of an inflamed pleura, which produces a high-pitched grating or squeaking sound. The rub may occur during both inspiration and expiration, but because it is not in the lung, it will never be affected by coughing. A pleural friction rub is associated with pleuritis, so people often complain of pain in the area of the rub, especially with deep inspiration.

Abnormal Vocal Sounds

Abnormal vocal sounds include bronchophony, egophony, and whispered pectoriloquy. Transmission of voice sounds through healthy lung tissue is normally muffled. Consolidated lung tissue produces clearer transmission of voice sounds over the affected area. **Bronchophony** is the abnormal clarity of the spoken word as heard through the

stethoscope. **Egophony** is translated literally as “voice of the goat.” Ask the patient to say “eee” while you auscultate the lungs through the stethoscope at various places. Egophony is present if the sound changes from a muffled “eee” to an “ay.”

To test for **whispered pectoriloquy**, ask the patient to whisper “one, two, three” while you auscultate the lungs. Normally, you should hear “puff, puff, puff.” If you hear “one-two-three” clearly through the stethoscope, whispered pectoriloquy is present.

Vocal sounds should be muffled and indistinct through the stethoscope. *Bronchophony*, *egophony*, *whispered pectoriloquy*: Consolidated lung tissue, pulmonary edema, pulmonary hemorrhage.



HELPFUL HINT

Sound transmission over consolidated lung tissue is like sound transmission around an enclosed swimming pool—the sound comes across louder.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Kane have concluded. The following are your findings:

- **General Health Survey**
 - Well-developed, 45-year-old Caucasian female.
 - Appears stated age.
 - In no acute respiratory distress.
 - Sits upright and relaxed during interview; answers questions appropriately.
 - Affect pleasant and appropriate.
 - Alert and responsive without complaints.
 - Oriented to time, place, situation, and person.
 - Head-to-toe scan reveals no signs of problems in other systems/areas that would affect respiratory system except nicotine-stained fingers.
- **Vital Signs**
 - Temperature, 98.6° F
 - Pulse, 64 beats per minute (BPM), strong and regular.
 - Respirations, 16/min, unlabored.
 - BP, 120/70.
 - Height, 5'5".
 - Weight: 165 lb.
- **Respiratory**
 - Trachea midline.
 - *Chest*: AP: lateral 1:2, costal angle < 90, equal chest excursion, normal spinal

curvatures, skin intact, no use of accessory muscles, no retraction.

- Chest nontender, symmetrical chest excursion, no masses, crepitus.
- Tactile fremitus equal anterior and posterior to midthorax.
- Resonance noted throughout lung fields. Diaphragmatic excursion 4 cm bilaterally.
- Lungs clear to auscultation except for expiratory wheezing that clears with coughing; no crackles or rubs.



Case Study Analysis and Plan

Now that your examination of Mrs. Kane is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Your next step is to analyze the data from Mrs. Kane's assessment and develop nursing diagnoses. The following are possible nursing diagnoses for Mrs. Kane. Cluster the supporting data for each nursing diagnosis.

1. Anxiety, related to knowledge deficit of preoperative routine and postoperative course
2. Risk for Ineffective Airway Clearance related to ineffective cough secondary to anesthesia, postoperative pain, and respiratory changes from smoking
3. Ineffective Health Maintenance related to insufficient knowledge of effects of tobacco use and available self-help resources
4. Health-Seeking Behaviors
5. Risk for Ineffective Health Maintenance related to lack of understanding of the significance of family and personal risk factors
6. Imbalanced Nutrition: more than body requirements, related to possible lack of exercise program

Identify any additional nursing diagnoses and any collaborative nursing diagnoses.



CRITICAL THINKING ACTIVITY #5

Now that you have identified some nursing diagnoses for Mrs. Kane, select one and develop a nursing care plan and a brief teaching plan for her, including learning objectives and teaching strategies.

RESEARCH TELLS US

Considine (2005) performed a literature review of the relationship between signs and symptoms of respiratory dysfunction and adverse events, and the nurse's ability to detect and prevent adverse events related to respiratory dysfunction. The literature analysis identified two major issues: that respiratory dysfunction is a precursor of adverse events and that respiratory dysfunction has a significant negative effect on mortality. The findings also revealed the following:

- 31 percent to 57 percent of patients with physiological abnormalities experienced respiratory dysfunction.
- Respiratory dysfunction is the most common reason for intensive care unit (ICU) admission.

- Respiratory dysfunction accounts for 20 percent of readmission to ICU.
- 83 percent of patients with unrecognized physiological abnormalities experience respiratory dysfunction.

Considine also found that the clinical indicators of respiratory dysfunction—respiratory rate, dyspnea, hypoxemia and acidosis—were key factors in the development of adverse events.

Although the findings were limited to the studies reviewed, the implications for nursing are great. Assessment is key to identifying changes in respiratory rate, dyspnea, and signs and symptoms of hypoxia and acidosis. Early recognition and intervention of respiratory dysfunction may be crucial in preventing adverse events and improved patient outcomes.

Source: Considine, J. (2005). The role of nurses in preventing adverse events related to respiratory dysfunction: Literature review. *Journal of Advanced Nursing* 49(6), March, pp. 624-633.

HEALTH CONCERNS

Facts About Lung Cancer

Leading cause of cancer deaths in men and women:

- 163,510 deaths from lung cancer in 2005 (90,490 men; 73,020 women).
- 60 percent diagnosed with lung cancer die within first year.
- 75 percent die within 2 years.
- 5-year survival rate is 15 percent.

Risk Factors for Lung Cancer

- **Smoking:** Smoking or exposure to second-hand smoke accounts for 87% of lung cancer cases.
- **Asbestos:** Persons who work with asbestos are seven times more likely to die from lung cancer. Persons who work with asbestos as well as smoke have an even greater risk.
- **Radon:** Exposure may double or triple risk for lung cancer.
- **Occupational exposure to cancer-causing agents:** Workplace cancer-causing agents include uranium, arsenic, beryllium, vinyl chloride, chloromethyl ethers, nickel, chromates, coal products, mustard gas, gasoline, and diesel exhaust.
- **Marijuana:** Suspected risk because marijuana has more tar than cigarettes.

- **Radiation therapy:** Suggested increased risk, especially if the patient smokes. Radiation after a lumpectomy does not have associated increase risk.
- **Recurring lung inflammation:** May result in scarring, which increases the risk for adenocarcinoma.
- **Mineral exposure:** Silicosis and berylliosis increase risk.
- **Family history:** Positive sibling history of lung cancer increases risk.
- **Genetic abnormality:** Chromosome 6 abnormality more readily develops lung cancer.
- **Diet:** Diets low in fruits and vegetables suggest increased risk for lung cancer.
- **Air pollution:** Slight increase in risk.

Source: American Cancer Society, 2005. <http://www.cancer.org>

Warning Signs of Lung Cancer

- Persistent cough
- Changes in respiratory pattern
- Unexplained dyspnea
- Blood-streaked sputum
- Hemoptysis
- Rust-colored or purulent sputum
- Chest, shoulder, or arm pain
- Recurring pleural effusion, pneumonia, or bronchitis

Common Abnormalities

ABNORMALITY	ASSESSMENT FINDINGS
<p>Asthma Reactive airway disease causing inflammation and airway obstruction because of increased reactivity to a variety of stimuli.</p>	<ul style="list-style-type: none"> ■ Pallor or cyanosis caused by hypoxia ■ Diaphoresis ■ Sits upright and leans forward ■ Use of accessory muscles ■ Intercostal and supraclavicular retraction ■ Chest tightness ■ Dyspnea, respiratory rate > 30/min ■ Increased pulse with premature ventricular contractions, increased or decreased BP ■ Pulsus paradoxus >12 mm Hg ■ Crackles, rhonchi, wheezes, decreased or absent breath sounds ■ Early in disease: expiratory wheezes ■ Late in disease: inspiratory and expiratory wheezes ■ Hyperresonance ■ Decreased tactile fremitus ■ Decreased chest excursion
<p>Bronchitis Excessive mucus production with recurrent, persistent cough during 3 mo of the year for 2 consecutive years. Results in obstructive lung disease and hypoxic and right-side CHF.</p>	<ul style="list-style-type: none"> ■ Color changes related to hypoxia ■ Average or obese ■ Changes in mental status related to hypoxia ■ Productive cough with copious amounts of mucus ■ Cyanosis ("blue bloater") ■ Increased AP diameter ■ Increased costal angle ■ Increased use of accessory muscles ■ Cardiac enlargement ■ Decreased excursion ■ Decreased tactile fremitus ■ Hyperresonance at bases, dullness over exudate areas ■ Crackles, rhonchi, wheezes ■ Breath sounds decreased at bases
<p>Emphysema Permanent enlargement of alveoli distal to terminal bronchioles with destruction of alveolar wall.</p>	<ul style="list-style-type: none"> ■ Thin with muscle wasting ■ Anxiety ■ Cough (rare) ■ Neck vein distension ■ Difficulty speaking because of respiratory distress ■ Pursed-lip breathing ■ Pallor, ruddiness (pink puffer) ■ Decreased turgor ■ Rapid, shallow respiration ■ Use of accessory muscles ■ Distant heart sounds, right-sided S3 ■ Tachycardia with arrhythmias ■ Hyperresonance at bases or in all lung fields ■ Decreased excursion ■ Decreased tactile fremitus ■ Increased AP diameter ■ Increased costal angle

ABNORMALITY	ASSESSMENT FINDINGS
<p>Lung Cancer Malignant tumor of lung tissue.</p>	<ul style="list-style-type: none"> ■ Decreased breath sounds ■ Thin as a result of weight loss ■ Unexplained dyspnea ■ Persistent dry or productive cough ■ Blood-streaked, rust-colored, or purulent sputum ■ Hemoptysis ■ Change in respiratory pattern ■ Wheezes
<p>Pleural Effusion Collection of fluid in pleural space.</p>	<ul style="list-style-type: none"> ■ Decreased breath sounds over affected lung ■ Chest, shoulder, or arm pain ■ Dyspnea ■ Complaints of chest pain ■ Decreased excursion on affected side ■ Decreased tactile fremitus on affected side ■ Dullness on percussion ■ Decreased or absent breath sounds ■ Friction rub with initial inflammation; rub disappears as fluid develops
<p>Pneumonia Infectious process of lung tissue. Clinical manifestations vary depending on causative agent.</p>	<ul style="list-style-type: none"> ■ Changes in mental status related to hypoxia or fever ■ Pharyngitis ■ Productive cough; pink, rusty, purulent, green, yellow, or white sputum ■ Diaphoretic or dry with poor turgor ■ Pallor or flushing ■ Circumoral or nailbed cyanosis ■ Tachypnea ■ Use of accessory muscles ■ Asymmetrical chest movement ■ Decreased excursion ■ Increased tactile fremitus ■ Crackles, rhonchi ■ Bronchial breath sounds ■ Positive bronchophony ■ Dullness on percussion
<p>Pneumothorax Complete or partial collapse of lung.</p>	<ul style="list-style-type: none"> ■ Shallow, rapid respiration ■ Dyspnea ■ Complaint of chest pain with or without hemoptysis ■ Asymmetrical excursion ■ Decreased fremitus ■ Hyperresonance ■ Absent breath sounds ■ If tension pneumothorax, severe respiratory distress, tachycardia, cyanosis, tracheal shift away from affected lung, and displaced point of maximal impulse occur.
<p>Pulmonary Fibrosis (“Stiff Lung”) Restrictive lung disease; chronic, nonmalignant, noninfectious inflammatory process of interstitial lung tissue</p>	<ul style="list-style-type: none"> ■ Weakness ■ Fatigue ■ Dyspnea ■ Weight loss ■ Anorexia

(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Tuberculosis Mycobacterial infection of lung</p>	<ul style="list-style-type: none"> ■ Vesicular breath sounds, occasionally bronchovesicular ■ “Velcro rales” ■ Decreased excursion ■ Pulmonary hypertension occurring late; if present, prominent P wave, neck vein distension, right ventricular heave, split S2, and accentuated pulmonic component ■ Clubbing (late sign) ■ Fatigue ■ Persistent, long-term, low-grade fever ■ Chills and night sweats ■ Anorexia and weight loss ■ Dyspnea. ■ Productive cough with nonpurulent, blood-streaked sputum. ■ Hemoptysis. ■ Pleuritic or dull chest pain. ■ Chest tightness. ■ Apical posterior segment of upper lobes or superior segments of lower lobes usually affected. Alters normal breath sounds.

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the respiratory system.

Ask the patient the following:

- Do you have...
 - Cough?
 - Dyspnea?
 - Chest pain?
 - Related symptoms, such as edema and fatigue?
- Do you have any history of respiratory disease? If so, are you taking any medication? What are they and why are you taking them?
- Do you have any other medical problems?
- Do you have allergies? If so, what are you allergic to? Can you describe your reaction?
- Do you use tobacco? If so, what kind? How much tobacco do you use? How long have you been using it? What is your occupation?
- Where do you live?
- When was your last PPD? And what was the result?
- Have you ever had a chest x-ray? If so, what were the results?
- Have you ever been immunized for influenza or pneumonia?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam:

- Stethoscope
- Marker
- Ruler

Position

Sitting

Technique

First, scan your patient head to toe checking for specific signs of diseases affecting other organ systems that might alter the respiratory system.

Inspection, Palpation, Percussion, Auscultation

Apply techniques:

- Anterior, posterior, lateral
- Side to side
- Apex to base

Chest

Inspect:

- Shape and symmetry
- Muscles for breathing
- Anteroposterior:lateral ratio

ASSESSMENT CHECKLIST (continued)

- Respiratory rate, rhythm
- Costal angle
- Spinal deformities
- Condition of the skin

Palpate:

- Tenderness, masses, crepitus
- Excursion
- Fremitus

Percuss:

- Identify percussion sound.
- Diaphragmatic excursion

Auscultate:

- Note breath sounds.
- Note abnormal sounds.
- Note adventitious sounds.
- Note abnormal voice sounds.

Trachea

Palpate:

- Position

DOCUMENT

Document your findings and develop your plan of care.


SUMMARY

- A thorough respiratory assessment provides invaluable data about your patient's overall health status.
- Before you begin the assessment, visualize the underlying structures and landmarks and review expected normal findings.
- As you work through the assessment, differentiate normal from abnormal, and work in a systematic manner from apex to base and from side to side.
- Remember, if a person has a respiratory problem, it may affect every other body system.
- Establish a baseline, using the patient as his or her own control. If he or she has an acute respiratory problem, postpone the comprehensive respiratory assessment and perform a focused assessment.
- After completing the assessment, document your findings.
- The plan of care should flow from your assessment. Share your findings with the patient. This provides an excellent opportunity for teaching, especially if you include the patient in developing a plan of care that promotes a healthy respiratory system.

REVIEW QUESTIONS

1. You are assessing your patient's lung bases and you hear a low-pitched, soft sound. The inspiratory phase is longer than the expiratory phase. This sound most likely is a:
 - a. Vesicular sound.
 - b. Bronchovesicular sound.
 - c. Bronchial sound.
 - d. Tracheal sound.
2. You are assessing your patient's lungs over the manubrium and you hear a medium-pitched sound with equal inspiratory and expiratory phases. This sound most likely is a:
 - a. Vesicular sound.
 - b. Bronchovesicular sound.
 - c. Bronchial sound.
 - d. Tracheal sound.
3. You are assessing your patient's lungs. While listening over the trachea, you hear a high-pitched, loud sound with the inspiratory phase shorter than the expiratory phase. This sound most likely is a:
 - a. Vesicular sound.
 - b. Bronchovesicular sound.
 - c. Bronchial sound.
 - d. Tracheal sound.
4. Don Andrews, age 76, has been hospitalized for pneumonia and developed acute respiratory distress syndrome. He is now on a ventilator. While auscultating his lungs, you hear an extra expiratory sound, anterior in the upper airways, near the second intercostal space. This sound is most likely:
 - a. Crackles.
 - b. Rhonchi.
 - c. Wheezes.
 - d. A friction rub.

5. James Lynch, age 59, has COPD. You are assessing his nailbeds for clubbing. Which finding would be positive for clubbing of the nails?
 - a. 45-degree angle of attachment
 - b. 90-degree angle of attachment
 - c. 160-degree angle of attachment
 - d. 180-degree angle of attachment
6. Mr. Lynch has classic barrel chest secondary to his COPD. What would be his AP:lateral ratio?
 - a. 1:1
 - b. 1:2
 - c. 1:3
 - d. 2:1
7. When percussing Mr. Lynch's lungs, what sound would you expect to hear over the bases?
 - a. Resonance
 - b. Hyperresonance
 - c. Tympany
 - d. Dullness
8. When assessing Mr. Lynch for tactile fremitus, what would you expect to find?
 - a. Increased fremitus at the bases.
 - b. Decreased fremitus at the bases.
 - c. Increased fremitus at the apex.
 - d. Decreased fremitus at the apex.
9. When assessing Mr. Lynch's breath sounds, what type of breath sounds would you expect to hear at the bases?
 - a. Increased breath sounds.
 - b. Decreased breath sounds.
 - c. Bronchial breath sounds.
 - d. Bronchovesicular breath sounds.
10. What area is best for assessing color changes in dark-skinned individuals?
 - a. Nailbeds
 - b. Palms of hands
 - c. Lips
 - d. Conjunctiva



Assessing the Cardiovascular System

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent cardiovascular history questions
- Obtain a cardiovascular history
- Perform a cardiovascular physical assessment
- Document cardiovascular findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

The cardiovascular system is the lifeline of the body. Its primary function is to act as a transport system, delivering oxygen by way of the red blood cells and delivering nutrients, metabolites, and hormones to every cell in the body. At the same time, it transports metabolic wastes for detoxification and excretion. The cardiovascular system also contains white blood cells, whose main function is to fight infection.

Because cardiovascular disease is the leading health problem in the United States, accurate assessment of the cardiovascular system is essential to identify and evaluate changes in cardiovascular function and potential risk factors for cardiovascular disease. A thorough, accurate cardiovascular assessment will allow you to develop a plan of care that addresses not only treatment measures but also health promotion and disease prevention measures.

Anatomy and Physiology Review

A thorough, accurate assessment depends on a solid understanding of the cardiovascular system. You especially need to understand normal cardiovascular function before you can identify abnormal findings. Having a mental image of the underlying structures will help you to perform the assessment accurately, and an understanding of the physiology—especially of the mechanisms of heart sounds—will guide your assessment and help you interpret your findings.

Structures and Functions of the Cardiovascular System

The cardiovascular system is a closed system consisting of the heart and blood vessels.




FIGURE 14.1. Structures of the heart.

The Heart

The heart is a muscle that pumps blood through a vast network of blood vessels extending over 60,000 miles

through arteries, arterioles, capillaries, venules, and veins (Fig. 14.1).

Anatomy and Physiology of the Heart

STRUCTURE	DESCRIPTION/PRIMARY FUNCTION
Heart	<ul style="list-style-type: none"> ■ A cone-shaped muscle with four chambers; a double pump about the size of a clenched fist (12 cm long and 9 cm wide). Weighs 250–390 g (8.8–13.8 oz) in adult males and 200–275 g (7.0–9.7 oz) in adult females. ■ Pumps blood throughout circulatory system.
Right Side	
■ Right atrium	<ul style="list-style-type: none"> ■ Upper chamber of right heart. ■ Receives unoxygenated blood from superior and inferior vena cava.
■ Tricuspid valve	<ul style="list-style-type: none"> ■ Right AV valve with three cusps (tricuspid). Attached by chordae tendineae to papillary muscles, which are attached to inner heart muscle. ■ Valve between right atrium and right ventricle.
■ Right ventricle	<ul style="list-style-type: none"> ■ Lower chamber of right heart. ■ Receives blood from right atrium and pumps it into pulmonary circuit.
■ Pulmonary semilunar valve	<ul style="list-style-type: none"> ■ Composed of three cusps. ■ Valve between right ventricle and main pulmonary artery.
■ Main pulmonary artery	<ul style="list-style-type: none"> ■ Artery leading from right ventricle to lungs. ■ Divides into right and left branches supplying respective lungs. ■ Carries unoxygenated blood from right ventricle to lungs.
■ Pulmonary veins	<ul style="list-style-type: none"> ■ Veins leading from lungs to left atrium. ■ Carry oxygenated blood to left atrium.
Left Side	
■ Left atrium	<ul style="list-style-type: none"> ■ Upper chamber of left heart. ■ Receives oxygenated blood from lungs through pulmonary veins.
■ Mitral valve	<ul style="list-style-type: none"> ■ AV valve with two cusps (bicuspid) attached by chordae tendineae to papillary muscles, which are attached to inner heart muscle. ■ Valve between left atrium and left ventricle.
■ Left ventricle	<ul style="list-style-type: none"> ■ Lower chamber of left half of heart. ■ Receives blood from left atrium and pumps oxygenated blood through systemic circulation.
■ Aortic valve	<ul style="list-style-type: none"> ■ Composed of three cusps. ■ Valve between left ventricle and aorta.
■ Interventricular septum	<ul style="list-style-type: none"> ■ Wall between left and right ventricles. ■ Vertically separates left and right sides of heart.

Layers of the Heart

LAYER	DESCRIPTION/PRIMARY FUNCTION
Endocardium	<ul style="list-style-type: none"> ■ Inner layer of heart; a smooth, thin layer of endothelium and connective tissue. Smooth inner lining of the heart.
Myocardium	<ul style="list-style-type: none"> ■ Middle and thickest layer of heart; heart muscle. ■ Responsible for cardiac contraction.
Epicardium	<ul style="list-style-type: none"> ■ The layer of serous pericardium on heart's surface. ■ Contains main coronary blood vessels.

(continued)

Layers of the Heart (continued)

LAYER	DESCRIPTION/PRIMARY FUNCTION
Pericardium	<ul style="list-style-type: none"> ■ Sac that surrounds the heart and roots of the great vessels. Composed of two layers: fibrous pericardium (outer layer of fibrous connective tissue) and serous pericardium. ■ Serous pericardium also has two layers: outer or parietal layer that lines the fibrous layer and visceral or inner layer that lines the heart and is also called the epicardium. ■ Serous pericardium contains pericardial fluid (10–20 mL of serous fluid). Pericardial fluid moistens the pericardial sac between the two layers and prevents friction during systole and diastole.

The Circulatory System

The circulatory system has two main networks, the pulmonary circulation and the systemic circulation (Fig. 14.2). The coronary circulation is part of the systemic circulation and supplies the heart itself.




FIGURE 14.2. Systemic and pulmonary circulation.

The pulmonary circulation involves blood vessels that circulate blood through the pulmonary arteries, the lungs, and the pulmonary veins. Unoxygenated blood enters the pulmonary circulation from the right and left pulmonary arteries. The unoxygenated blood then flows through the pulmonary arterioles to the lung capillaries, where the exchange of carbon dioxide and oxygen occurs. The oxygenated blood then enters the pulmonary venules that lead to the pulmonary veins. Oxygenated blood is then carried back to the left atrium through the right and left pulmonary veins.

Systemic Circulation

The systemic circulation is responsible for supplying oxygen to every cell in the body through the arterial system and then returning unoxygenated blood to the heart through the venous system. Oxygenated blood flows into the left atrium from the pulmonary circulation. The left atrium then pumps the oxygenated blood into the left ventricle, which in turn expels the oxygenated blood through the aorta into the arterial systemic circulation. From the aorta, blood then flows through smaller arterioles to the systemic capillaries. The systemic capillaries link the arterial and venous systems. At this point, exchange of oxygen, nutrients, and wastes occurs. From the capillaries, unoxygenated blood then flows through the venules, into the larger veins, and then to the superior and inferior vena cavae. Unoxygenated blood then enters the right atrium and is pumped to the right ventricle into the pulmonary circulation to continue the cycle. Layers of arteries and veins are shown in Figure 14.3. (See *Layers of Arteries and Veins*.)

Coronary Circulation

The **heart** is a muscle with the primary responsibility of circulating blood throughout the entire body. Consequently, the oxygen requirements of the heart are great, second only to those of the brain. The **coronary circulation** is responsible for supplying the heart with the oxygen needed to perform this task effectively. The coronary circulation consists of the right and left coronary arteries and the coronary sinus

Layers of Arteries and Veins

STRUCTURE	DESCRIPTION/PRIMARY FUNCTION
Arteries	<ul style="list-style-type: none"> ■ Blood vessels with three coats: tunica intima, tunica media, and tunica adventitia. ■ Carry oxygenated blood away from left heart and unoxygenated blood to lungs via pulmonary arteries.
Tunica intima	<ul style="list-style-type: none"> ■ Inner layer of endothelial cells containing basement membrane of fine, collagenous fibers and external elastic layer called internal elastic lamina.
Tunica media	<ul style="list-style-type: none"> ■ Middle, thickest layer containing collagenous fibers, connective tissue, smooth muscle cells, and elastic fibers.
Tunica adventitia	<ul style="list-style-type: none"> ■ Outer layer containing collagen and elastic fibers, nerves, lymphatic vessels.
Arterioles	<ul style="list-style-type: none"> ■ Smallest arteries; contain large amount of smooth muscle cells that can dilate and constrict. ■ Carry blood to capillaries and control blood flow to capillaries through dilation/constriction.
Capillaries	<ul style="list-style-type: none"> ■ Single layer of microscopic endothelial cells. ■ Connect arterial and venous system for exchange of gases, fluids, nutrients, and wastes.
Veins	<ul style="list-style-type: none"> ■ Contain same three layers as arteries, but are thinner with less elastic and collagenous tissue and smooth muscle. ■ BP in venous system is low; veins have valves to prevent backflow. ■ Carry unoxygenated blood back to right heart, except for pulmonary veins, which carry oxygenated blood from lungs to left heart.
Venules	<ul style="list-style-type: none"> ■ Smallest veins, consisting of endothelium and thin tunica adventitia. Blood from capillaries drains into venules.




FIGURE 14.3. Layers of arteries and veins.

and cardiac veins. The coronary arteries are the first branches off the aorta. The cardiac veins drain into the coronary sinus, which in turn drains directly into the right atrium.

Mechanisms of Heart Sounds

To perform an accurate cardiac assessment, you need an understanding of the normal anatomy and physiology of the heart as well as knowledge of the mechanisms of heart sounds in order to interpret your findings. Heart sounds are the result of events within the heart. The movement and pressure of the blood (**hemodynamics**), the activity of the electrical conduction system, and the movement of the valves affect the sounds that you hear (Fig. 14.4).

The Cardiac Cycle

The cardiac cycle comprises all the physiological events needed for the heart to beat. The valves, the hemodynamic events within the heart and the conduction system work together in the cardiac cycle. The cardiac cycle comprises systolic and diastolic phases. The systolic phase is the contraction or emptying phase, and the diastolic phase is the resting or filling phase. The atria and ventricles alternate through the systolic and diastolic phases; while the atria are contracting, the ventricles are relaxing, and vice versa (see Fig. 14.4).

Ventricular Diastole

Ventricular diastole marks the beginning of the filling phase. Very shortly after the onset of ventricular diastole, the pressure in the ventricles is less than that of the atria, and the mitral and tricuspid valves open to allow filling of the ventricles (the rapid-filling phase of diastole). Toward the end of ventricular diastole, the atria contract to propel additional blood into the ventricles, thus completing the filling phase of the ventricles. This is referred to as the **atrial kick** and is responsible for about 25 percent of the total blood volume in the ventricles. The ventricles are now filled, marking the end of ventricular diastole. This is referred to as the **end-diastolic volume**.

Ventricular Systole

With ventricular contraction, the pressure in the ventricles will exceed that in the atria. This results in closure of the mitral and tricuspid valves and marks the beginning of ventricular systole. While the ventricles are contracting and blood is being propelled from them, the atria are relaxed (atrial diastole) and are filling. When pressure in the ventricles exceeds that of the aorta and pulmonary arteries, the aortic and pulmonic valves open to allow for

Branches of the Right and Left Coronary Arteries

Branch	Areas Supplied
RIGHT CORONARY ARTERY	
Nodal branch	SA (pacemaker) node
Right marginal branch	Ventral and dorsal surface of right ventricle
Nodal branch	AV node
Small branches	Right AV bundle
Posterior branches	Join to anterior descending branch and circumflex artery of left coronary artery
LEFT CORONARY ARTERY	
Anterior interventricular branch	Left ventricle
Small branches	Left AV bundle
Circumflex	Lateral and posterior side of heart

Branches of the Cardiac Veins

Branch	Area Drained
Great cardiac vein	Anterior portion of heart
Middle cardiac vein and oblique vein	Posterior aspect of heart
Small cardiac vein	Drains directly into right atrium
Thebesian veins	Drain into each of heart's four chambers

forward flow into the systemic and the pulmonic systems, respectively. While the ventricles are contracting and propelling blood into the aorta and pulmonary arteries, the pressures in the ventricles are gradually decreasing to eventually attain a level below that of the pulmonary arteries and aorta. At this point, the pulmonic and aortic valves close, marking the end of ventricular systole.

The amount of blood in the ventricles at the end of diastole is referred to as **end-diastolic volume** and reflects the largest volume of blood in the ventricles. The amount of blood remaining in the ventricles at the end of ventricular systole is referred to as the **end-systolic**




FIGURE 14.4. Mechanisms of heart sounds.

volume and is indicative of the least amount of residual blood in the ventricles. The pressure of the end-diastolic volume is referred to as **end-diastolic pressure**; the pressure of the end-systolic volume is **end-systolic pressure**. The amount of blood ejected from the heart with each contraction is referred to as the **stroke volume**. This is the difference between the end-diastolic volume and the end-systolic volume—basically the difference in volume between the beginning and the end of ventricular contraction.

The Cardiac Output

The **cardiac output** is the amount of blood ejected per minute. Cardiac output equals stroke volume multiplied by heart rate. Because so many factors can affect cardiac output, this is an important concept to understand. For

example, an increase in heart rate (tachycardia rates over 100 beats per minute [BPM]) while exercising increases cardiac output to meet the body's demands. On the other hand, an athlete often has a slow heart rate (bradycardia rates less than 60 BPM) with a greater stroke volume increasing cardiac output to meet the body's demand.

Although both heart rate and stroke volume affect cardiac output, the change in heart rate is the major regulator of cardiac output. The internal pacemaker of the heart controls the rate, but the autonomic nervous system also plays a major role. The heart rate also responds to various external factors, such as temperature, pain, exercise, pH, hormones, and emotions. Changes in heart rate can readily be seen in response to these factors.

The Stroke Volume

Stroke volume is less variable and responds to changes in preload, afterload, and contractility of the heart. **Preload** refers to the volume of blood in the ventricles at the end of diastole. The greater the preload, the greater the “stretch” or contractility of the ventricles, and the greater the stroke volume. An increase in venous return to the heart will, in turn, increase preload. Factors that affect venous return include:

- **Venous blood reservoirs:** Distensibility of veins allows them to act as reservoirs and can either decrease or increase venous return to the heart.
- **Skeletal muscle pump:** Contraction of skeletal muscles pumps blood back to the heart.
- **Venous tone:** Sympathetic stimulation of smooth muscles in the walls of veins increases venous return to the heart.
- **Respiratory pump:** During inspiration, the increase in negative intrathoracic pressure increases venous return to the heart.

Afterload reflects the end-systolic volume. It is affected by the amount of resistance the ventricles have to contract against. An increase in afterload results in a decrease in stroke volume. Afterload may be affected by:

- Arterial elasticity.
- Peripheral vascular resistance.
- Aortic valve resistance.
- Viscosity and volume of blood.
- **Contractility**, which reflects the force of contraction.

- **Positive inotropes**, which increase the force of contraction.
- **Negative inotropes**, which decrease the force of contraction.



HELPFUL HINTS

- **Remember:** If contractility is increased, so are the oxygen requirements of the heart.
- In the normal heart, left-side cardiac activity occurs slightly before right-side cardiac activity.

The Heart's Electrical Conduction System

The cardiac electrical conduction system is essential for each contraction of the heart. The cardiac muscle has its own electrical system that is responsible for initiating and maintaining the rhythmic beat of the heart. The specialized cells of the conduction system have automaticity, excitability, conductivity and refractoriness. **Automaticity** is the cell's ability to initiate an impulse. **Excitability** is the cell's ability to respond to an impulse and create an action potential. **Conductivity** is the cell's ability to transmit an impulse. **Refractoriness** is the cell's ability to respond to the transmitted impulse.

The **sinoatrial (SA) node** is the normal pacemaker of the heart located in the right atrium near the superior vena cava entrance point. The SA node paces the normal adult heart at 60 to 100 BPM. The activation of the SA node passes through the atria and results in atrial




FIGURE 14.5. Electrical conduction system.

contraction. The electrical activity is then conducted to the **atrioventricular (AV) node**. This node is located at the base of the right atrium between the atria and the ventricles. It has the ability to pace the heart at a rate of 40 to 60 BPM. The electrical impulse is then transmitted from the AV node to the **bundle of His**, which divides into two branches, the right and the left, which traverse the interventricular septum. Finally, the impulse is transmitted to small branches that eventuate into the **Purkinje fibers**, which stimulate the ventricles to contract. The pacer ability of the bundle of His is 20 to 40 BPM (Fig. 14.5).

The Valves

The most accepted theory concerning the origin of heart sounds relates primarily to the closure of the valves. The

“lub-dub” sounds that are usually readily heard are referred to as S_1 (lub) and S_2 (dub), the first and the second heart sounds, respectively. They result from valve closure.

The First Heart Sound (S_1)

S_1 (Fig. 14.6) marks the beginning of systole, and S_2 marks the end of it. S_1 , the first heart sound, results from the closure of the mitral (M_1) and tricuspid (T_1) valves. M_1 and T_1 normally close within approximately 0.02 second or less. These valve sounds are often heard as a single sound. S_1 is best heard at the apex or left lateral sternal border (LLSB) with the diaphragm of the stethoscope.

The Accentuated S_1 . The intensity of S_1 may be increased normally in anemia, hyperthyroidism, and exercise. An apparent increased intensity of S_1 also may be noted with mitral or tricuspid stenosis. It is important to




FIGURE 14.6. Graphic depiction of S_1 heart sounds.

note that the intensity does not really change in these entities; rather, the S_1 becomes sharper (shorter and higher pitched), and consequently, is often described clinically as increased in loudness.

The Diminished S_1 . The intensity of S_1 may be decreased normally in patients with thick chest walls or emphysema because there is greater distance for the heart sound to traverse from the myocardium to the chest wall. S_1 may also be diminished in patients with first-degree heart block. The delay in conduction between atrial contraction and ventricular contraction allows more time for the valves to close, thus diminishing the sound's intensity.

The Varying S_1 . The intensity of S_1 may vary with irregular rhythms, such as complete heart block or atrial fibrillation, a grossly irregular rhythm.

The Split S_1 . In normally functioning hearts, the left side contracts before the right side; therefore, the mitral valve will close before the tricuspid, and the aortic valve will close before the pulmonic. Often the closures of the mitral and tricuspid valves occur so close together that the first sound is heard as a single sound. There are times when each component (mitral and tricuspid) of the sound may be heard clearly, defining this as a normal split S_1 .

If the time interval between mitral and tricuspid valve closure is greater than 0.20 second, each component of valve closure may be heard. If you can hear the mitral valve close and then the tricuspid valve close, this is referred to as a **split S_1** . The mitral component of S_1 is louder than the tricuspid component. So when listening for a split S_1 , listen over the site of second valve closure, the tricuspid area or LLSB. Conduction problems such as right bundle branch block (RBBB), premature ventricular contractions (PVCs), and ventricular tachycardia may result in asynchronous closure of the mitral and tricuspid valves, producing a widely split S_1 .

The Second Heart Sound (S_2)

When the systolic pressure in the ventricles decreases below that of the aorta and the pulmonary artery (toward the end of systole), the aortic (A_2) and pulmonic (P_2) valves close, producing the second heart sound (Fig. 14.7). Clinically, this sound marks the end of systole and the beginning of diastole. A_2 and P_2 normally close about 0.02 second from each other; consequently, they may occasionally be heard as a single sound. (See following text for an important distinction between A_2 and P_2 with respiration.)

The Accentuated S_2 . The intensity of S_2 may also be normally increased with high-output states such as exercise, resulting also in a concomitant increased heart rate. An

accentuated S_2 is also associated with pathology. A_2 may be accentuated with systemic hypertension (HTN); P_2 is likewise accentuated with pulmonary HTN.

The Diminished S_2 . A thick chest wall and chronic lung disease also decreases the intensity of S_2 . A_2 is also decreased characteristically in calcific aortic stenosis, and P_2 is diminished in pulmonic stenosis.

The Split S_2 . When the time interval between A_2 and P_2 is greater than 0.02 second, you will hear each valve component. This is referred to as a **split S_2** . This split differs from a split S_1 because it has a respiratory variation. During inspiration, S_2 will split, so you will hear the aortic valve close, then hear the pulmonic valve close ("blub"). Characteristically, S_2 will be heard as "blub" (A_2 followed by P_2). During expiration, the valves close as one to create a single sound ("dub").

This respiratory variation in the S_2 results from the normal physiological changes that occur during inspiration, when an increase in negative intrathoracic pressure results in increased blood return to the right side of the heart. With more blood in the right side of the heart, the right ventricle contraction time increases, resulting in a greater time interval between A_2 and P_2 , compared with the converse occurring during expiration. Thus, during inspiration you hear "lub-blub" (A_2 and P_2 as separate sounds) in comparison with "lub-dub" during expiration, when A_2 and P_2 are closer to one another and often heard as a single sound. The increased venous return to the right side of the heart that occurs during inspiration may also account for the frequent sinus variation that is noted, especially in children. During inspiration, the cardiac rate increases to accommodate the increase in venous return.

The Widely Abnormal Split of S_2 . This occurs with RBBB. The delay between aortic and pulmonic valve closure will average greater than 0.03 second during expiration, with further widening of the split during inspiration.

The Paradoxical (Reversed) Split of S_2 . When P_2 comes before A_2 , this is defined as a paradoxical or reversed split of S_2 . The mechanism relates to delayed contraction of the left ventricle, resulting from pathological blockage or conduction through the left heart's electrical nerve conduction system. Consequently, the left heart is depolarized later by way of the electrical impulse going from the right heart and through the interventricular septum to stimulate the left ventricular muscle to contract.

Atrial Septal Defect. The fixed, widely split S_2 occurs in atrial septal defect. It is unaffected by respiration. In this entity and in some other congenital cardiac abnormalities, the split averages 0.04 to 0.07 second.




FIGURE 14.7. Graphic depiction of S_2 heart sounds.

Extra Heart Sounds

Additional sounds that may be heard during auscultation (Fig. 14.8) include early ejection click, midsystolic ejection click, opening snap, S_3 , and S_4 . These sounds do not always indicate pathology.

The Early Ejection Click. An early ejection click is a high-pitched, short systolic sound occurring shortly after S_1 . The ejection click associated with the aortic and pulmonic valves is an early sound occurring about 0.03 to 0.08 second after S_1 . The ejection click of aortic valve origin is heard at the base and better heard at the apex. The ejection click of pulmonic valve origin is best heard at the left sternal border at the second and third inter-spaces and, rarely, at the apex.

The Midsystolic Ejection Click. This high-pitched short sound is a cardinal finding in mitral valve prolapse (MVP). The ejection click of MVP occurs at least 0.14 second after the first heart sound, but its exact placement in midsystole varies with the position of the patient (e.g., sitting, standing, lying down, squatting). It is heard best at the apex or just medial to it, but it may be widely transmitted over the precordium.

The Opening Snap. This is a short, high-pitched, mid-diastolic sound occurring 0.05 to 0.14 second after S_2 . It is a classic finding in mitral stenosis. It is also found in tricuspid valve stenosis, a much rarer condition. It is heard best at or near the apex, with the patient in the left lateral position. It is accompanied by the other classic findings of




FIGURE 14.8. Graphic depiction of extra heart sounds.

mitral stenosis—namely, the sharp first heart sound and the mid-late diastolic murmur of mitral stenosis.

The Physiological S_3 . The ventricles receive blood from the atria. As the blood flows torrentially from the atria into the ventricles in early diastole and forcefully impacts on the ventricular walls, it sets up vibrations that result in an audible sound—the normal S_3 . A normal finding in children and young adults, S_3 is a low-pitched, short, soft sound, occurring about 0.14 to 0.16 second after S_2 . It is heard best at the apex with the patient in the left lateral position using the bell portion of the stethoscope.

The S_3 Gallop Rhythm. This rhythm was formerly referred to as a *ventricular gallop* or *protodiastolic gallop*, but these terms are now obsolete. The sound of the gallop is, as its name implies, a lilt or canter similar to the sound of a galloping horse. Often, a rapid heart rate accompanies heart failure, adding to the gallop quality of the sound. The mechanism of the S_3 gallop is essentially the same as that of the normal S_3 . However, it relates to hemodynamic filling of a failing, noncompliant ventricle.

It is therefore a sign of congestive heart failure (CHF). The cadence of the S_3 gallop sounds like “lup-dah-dah, lup-dah-dah.” It is designated as an S_3 **gallop rhythm** based on a constellation of other symptoms in keeping with the diagnosis of CHF associated with the characteristic lilt or canter quality present. The S_3 gallop of right-side origin is best heard at the LLSB and the S_3 of left-side origin is best heard at the apex, with the patient in the left lateral position using the bell portion of the stethoscope.

The S_4 Gallop Rhythm (S_4). At the end of diastole the atria contract and expel some of the residual blood within them to complete ventricular filling. This atrial contraction at the end of diastole is referred to as the *atrial kick* and is responsible for about 25 percent of the blood entering the ventricles. The sound is not normally heard. However, if the atria have to contract against ventricular resistance or decreased ventricular compliance, significant vibrations are generated, resulting in the audible S_4 sound known as an S_4 **gallop rhythm**. The S_4 gallop sound is a low-pitched sound occurring 0.08 to 0.20 second before S_1 .

S_4 may be detected as an inaudible event on a phonocardiogram. When it is heard, it is abnormal and termed an S_4 gallop rhythm. (The terms *atrial* and *presystolic gallop* are obsolete.) The S_4 gallop may be associated with HTN, coronary artery disease (CAD), myocardial infarction (MI), and aortic or pulmonic stenosis. The cadence of S_4 sounds like “lah-dah-dup, lah-dah-dup.” The S_4 of right-side origin is best heard at the left sternal border and the S_4 of left-side origin is best heard at the apex with the patient in the left lateral position using the bell portion of the stethoscope.

Murmurs

Murmurs are defined as a series of audible, prolonged sounds resulting from turbulence created within the vascular system. Causes of turbulent flow include:

- Increased flow through normal blood vessels, creating frictional, audible sounds.
- Flow through constricted blood vessels (e.g., aortic stenosis).
- Flow into a dilated blood vessel from one of normal size.
- Combination of the preceding causes.

Figure 14.9 is a graphic depiction of the causes of turbulent flow.

Although some murmurs are “innocent,” identification and thorough description of the murmur are essential in differentiating an innocent murmur from a pathological one. Murmurs are characterized in terms of:

- **Quality:** Blowing, rough or harsh, or musical.
- **Frequency:** High, medium, or low.
- **Intensity or loudness:** Grades 1 through 6.
- **Duration:** Early, mid, late, holo, or pan (heard throughout systole or diastole).
- **Configuration:** Crescendo, decrescendo, or plateau.
- **Placement in the cardiac cycle:** Systole or diastole.
- **Location of sound.**
- **Radiation of sound.**

Murmurs are also designated as systolic, diastolic, or continuous. They may originate from either the right or the left side of the heart, and tend to radiate downstream in the direction of the jet created at their point of origin. Any changes in the quality of the murmur may indicate an acute alteration in the valve structure, such as acute valve rupture. Figure 14.10 is a graphic depiction of murmurs.

Continuous Sounds

Sounds that are heard during both systole and diastole are called **continuous** (Fig. 14.11). A continuous murmur, venous hum, and friction rub are continuous sounds with three different mechanisms.

Continuous Murmur. The continuous murmur is generated whenever there is uninterrupted forward flow of blood from a high pressure system into one of low pressure, as in patent ductus arteriosus. A continuous murmur crescendos in late systole, envelops the second sound,




FIGURE 14.9. Graphic depiction of the causes of turbulent flow.

and decrescendos throughout diastole. This murmur is medium pitched and heard best in the infraclavicular area and in the second left interspace.

Venous Hum. Turbulent flow within the venous system in the neck and superior vena cava may result in a venous hum, a continuous, low-pitched sound. Venous hum may be a normal finding, especially in children. The quality of the venous hum is either blowing, rough, or musical. It is a low-pitched sound, heard best over the right internal jugular vein in the supraclavicular area but readily disappearing when light pressure is applied to the internal jugular vein, establishing its benign nature.

Friction Rub. A pericardial friction rub is a high-pitched, scratchy, leathery sound that is generated when the inflamed parietal and visceral pericardial layers rub together. A pericardial friction rub may be associated with pericarditis, pericardial effusion and acute MI. Sharp, knife-like chest pain may accompany the rub. The pain may




FIGURE 14.10. A. Graphic depiction of systolic murmurs. B. Graphic depiction of diastolic murmurs.

radiate to the left shoulder and neck and is aggravated with deep inspiration, coughing, swallowing, and position changes. The pericardial friction rub may be heard in one or all phases of the cardiac cycle. A triple component rub results from atrial contraction, ventricular contraction, and ventricular relaxation and is usually best heard at the third interspace at the left sternal border.

HELPFUL HINT
 Anything that affects the valves, the hemodynamic events, or the conduction system may affect what you hear.

The Arterial System

Pressure in the arteries is much higher than in the venous system. The pressure created from the contraction of the ventricles can be converted into a waveform. The pressure creates one positive wave with a dichrotic notch. The positive wave represents systole. The wave has a smooth and rapid upstroke with a less acute descent. The dichrotic notch represents the closure of the aortic valve. Figure 14.12 illustrates a normal arterial pressure wave.

The Venous System

The venous system is a low-pressure system compared with the arterial system. The venous system reflects the





FIGURE 14.11. Graphic depiction of continuous sounds.



Normal

NORMAL PULSE

As shown in this illustration, a normal pulse has two components: systole and diastole. Indicated by the initial upstroke, systole signifies the arterial pressure during ventricular contraction. Diastole, the downstroke, indicates the arterial pressure during ventricular relaxation when the heart fills.

FIGURE 14.12. Normal arterial pressure wave.

flow of blood back to the heart. The flow of blood through the venous system is regulated by the pressure gradient for venous return, resistance to blood flow, and the venous pumps. The pressure gradient is affected by right arterial pressure and venous pressure. Resistance to blood flow also affects venous return. The venous pumps include the skeletal muscle pump and the respiratory pump.

Assessment of venous pressure in the jugular veins is an indirect assessment of right atrial pressure. The pressure in the jugular veins can be converted into a waveform. The jugular waveform has three positive waves and two negative waves that correspond with the cardiac cycle.

Keep in mind that the atria work opposite the ventricles, so when one is contracting, the other is relaxing. So, at the end of diastole (ventricular filling), the atria contract to fill the ventricle. The force of the right atrial contraction results in the “a” wave. The ventricles are filled, diastole is completed, and the mitral and tricuspid valves close, marking the beginning of systole. The “c” wave represents the force of tricuspid valve closure. As the ventricles are contracting, the atria are relaxing. The pressure in the atria at this point is low, represented by the “x” wave. The atria are now filling as the ventricles complete systole. As the atria fill, the pressure rises. The “v” wave represents right atrial

filling. Once the atria are filled and the ventricles have contracted, diastole begins and the tricuspid valve opens. As the tricuspid valve opens, there is a sudden drop in right atrial pressure as the atrium empties into the ventricle, represented by the “y” wave. Figure 14.13 illustrates correlation of the jugular waveform with the cardiac cycle.





FIGURE 14.13. Correlation of jugular wave form with cardiac cycle.



Relationship of Cardiovascular System to Other Systems

Interaction With Other Body Systems

The Respiratory System

The respiratory system works closely with the cardiovascular system to supply oxygen to and remove carbon dioxide from the body. The exchange of gases that occurs within the respiratory system could not occur without the circulatory system, which transports the gases. The respiratory system also works with the cardiovascular system to ensure that oxygen requirements are met. If the body needs more oxygen (e.g., during exercise), the respiratory and cardiac rates will increase to meet this need. Because the two systems work so closely together, disease that affects one system eventually affects the other. For example, patients with chronic obstructive pulmonary disease (COPD) often develop right-sided CHF.

The Nervous System

The nervous system (cerebrum, hypothalamus, medulla oblongata, and autonomic nerves) has a regulatory effect on the heart. The **medulla oblongata** is the major control, the cardiorespiratory center. In response to external factors, the **cerebrum** and **hypothalamus** send afferent information to the medulla oblongata, which then responds by changing the heart rate. The medulla oblongata also responds to afferent information from the chemoreceptors and baroreceptors in the aortic arch and carotid artery sinus. The chemoreceptors respond to changes in oxygen, carbon dioxide, and hydrogen ions. The baroreceptors respond to changes in blood pressure (BP). The medulla oblongata responds by changing the heart rate as needed to maintain homeostasis.

The **autonomic nervous system**, sympathetic and parasympathetic, can affect the heart rate. The **sympathetic nervous system**, with the release of norepinephrine, accelerates the heart rate; the **parasympathetic nervous system**, with the release of acetylcholine, decreases it.

The Endocrine System

When the sympathetic nervous system is activated, the adrenal medulla responds by secreting epinephrine and norepinephrine, both of which have a positive inotropic (contractility) and chronotropic (rate) effect on the heart.

Developmental, Cultural, and Ethnic Variations

Infants

Several developmental changes in the cardiovascular system occur shortly after birth. The foramen ovale that connected the right and left atria in fetal circulation closes after birth, allowing the heart to function as a double pump. The ductus arteriosus, a shunt between the pulmonary trunk and the aorta, also closes shortly after birth. This may result in systolic murmurs until closure occurs, usually within 48 hours after birth. The normal heart rate for the newborn is 120 to 160 BPM.

Children

Because the heart lies more horizontally in children, the apical impulse is at the fourth intercostal space until age 7, to the left of the midclavicular line till age 4, at the midclavicular line until age 6, and to the right of the midclavicular line by age 7. Sinus arrhythmia with respiratory variations is common in children. Because of the thin chest wall, heart sounds are louder, and a physiological S_3 may be heard. Innocent murmurs are also common findings in children. These murmurs are systolic and usually of low intensity (no greater than grade 3). A venous hum is also a common benign finding in children. The heart rate gradually slows to the adult rate of 60 to 100 BPM as the child gets older.



ALERT

Although murmurs are very common in children, they need to be reported to differentiate an innocent from an abnormal murmur.

Pregnant Women

Several changes in the cardiovascular system can be seen during pregnancy. The increase in cardiac workload often results in the development of a systolic murmur. A mammary souffle is a murmur that develops from increased blood flow through the mammary artery occurring late in pregnancy or during lactation. As the pregnancy progresses and the diaphragm rises, the heart rises and rotates, displacing the apical impulse. The BP is usually lower during the first and second trimesters, and the heart rate is slightly higher than in the nonpregnant state.

Older Adults

Changes in BP are frequently seen in older adults. Postural (orthostatic) hypotension results from a decrease in

Korotkoff's Sounds

When you take your patient's BP, you may hear five distinct phases called **Korotkoff's sounds**. These phases occur because the BP cuff partially obstructs blood flow and disturbs the laminar flow pattern, causing turbulence. Korotkoff phases include the following:

- **Phase I:** A faint, clear, rhythmic tapping noise that gradually increases in intensity. Intraluminal pressure and cuff pressure are equal.
- **Phase II:** A swishing sound that is heard as the vessel distends with blood.
- **Phase III:** Sounds become more intense. Vessel is open in systole but not in diastole.
- **Phase IV:** Sounds begin to muffle, and pressure is closest to diastolic arterial pressure.
- **Phase V:** Sounds disappear because vessel remains open.

baroreceptor response, decreased elasticity of the arteries, and the increased incidence of cardiovascular disease and medications. Auscultatory gaps also frequently occur. These gaps are periods of silence after the first Korotkoff's sound (true systolic) and subsequent Korotkoff's sounds, and may result in inaccurate BP readings.

People of Different Cultures and Ethnic Groups

Several cultural or ethnic variations are seen in relation to the development of cardiovascular disease. These variations result from both biological and ethnic differences and lifestyles associated with the specific culture. For example:

- Japanese have a lower incidence of HTN and high cholesterol.
- Hispanics have a lower mortality rate from heart disease than, non-Hispanics.
- Middle-aged Caucasians have the highest incidence of CAD.
- African Americans have an earlier onset and greater severity of CAD than other groups.
- African American women have a greater incidence of CAD than white women.
- Native Americans under age 35 have twice the mortality rate from heart disease as other groups.

Introducing the Case Study



Henry Brusca is a 68-year-old, married father of 7 who was in relatively good health until 3 weeks ago. At that time, he visited the emergency room with the complaint of “just not feeling right.” His BP on admission was 170/118, so he was admitted to the coronary care unit with the diagnosis of uncontrolled HTN. His BP was controlled with medication, and he was discharged several days later. He is now being seen for follow-up care and management of HTN. Because Mr. Brusca is newly diagnosed with HTN, you will need to complete a history and thorough cardiovascular examination.

Performing the Cardiovascular Assessment

Assessment of the cardiovascular system includes a comprehensive health history and physical examination. The history will enable you to identify any cardiovascular symptoms and risk factors, and the physical examination will help you evaluate the normal functioning of the heart and detect any abnormalities. As you proceed with the assessment, identify not only actual problems but also potential problems. Remember, if there is a problem with the cardiovascular system, every other system will be affected. Be alert for changes in other systems that may reflect cardiovascular problems.

Health History

During the health history, collect biographical data and ask your patient about his or her current health, past health, review of systems, and family and psychosocial history as it relates to the cardiovascular system. If time is an issue and you are unable to perform a complete health history, perform a focused cardiovascular history.

Biographical Data

A quick review of the patient's biographical data helps identify actual or potential problems. Begin by considering your patient's age. The risk for cardiovascular disease increases with age. Gender is also a risk factor, with men having a greater incidence of cardiovascular disease than women. On the other hand, women have a higher mortality rate within 1 year after MI than men. CAD usually strikes women 10 years later than men. In addition, women who are having a heart attack often have different presenting symptoms than men. Besides chest pain, fatigue, and shortness of breath, they may also experience back pain, anorexia, and light-headedness.

Focused Cardiovascular History

If your patient's condition or time prohibits a detailed cardiovascular history, take a history that focuses specifically on the cardiac status of your patient. The patient's response to the following questions will help direct your assessment:

- Are you having any chest discomfort? If yes, when did it start?
- What were you doing before the pain started? Did anything make it better or worse?
- Have you ever had this pain before? What does it feel like? Where does it hurt? On a scale from 1 to 10, with 10 being the worst, how severe is the pain?
- Do you have a history of cardiovascular disease? If yes, are you taking any medications? If yes, what are you taking and why?
- Do you have any other medical problems?
- Are you having any breathing difficulties?
- Do you have any allergies? If yes, describe reaction.

Ethnic background is another risk factor. Certain cardiovascular diseases are more prevalent in specific races or ethnic backgrounds. For example, African American men in the United States have a high incidence of HTN. Occupation may also be a contributing factor to cardiovascular problems if the patient has a high-stress job or if progressive cardiovascular disease forces her or him to make career changes. Identify your patient's supports and financial status. Chronic cardiovascular disease may take a toll on the patient's resources, and you may need to make referrals to ensure that she or he is able to maintain the medical treatment plan.

Current Health Status

If the patient has a cardiovascular problem, begin with his or her chief complaint. Keep in mind that the person with a cardiac problem may present with many symptoms or none at all. Identifying key symptoms may be crucial to detecting cardiac problems early and preventing complications, but interpreting these findings may be difficult. For example, symptoms that seem to be of obvious cardiac origin may actually originate in another system, whereas symptoms that seem to be from another system may actually be cardiac in origin.

If your patient presents with acute cardiac symptoms such as chest pain, defer the detailed history and focus on the acute problem. A detailed history can be obtained at a later time, when the patient's condition improves, or from a secondary source, such as a family member.

Frequent cardiac symptoms include chest pain, dyspnea, cough, edema, syncope, palpitations, fatigue, and changes in the extremities such as numbness, tingling, and intermittent claudication. These symptoms usually occur not in isolation but rather in combination. One symptom may lead to another; for example, palpitations may result in syncopal attacks, or chest pain may be associated with dyspnea.

Symptom Analysis

Symptom analysis tables for all of the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Chest Pain

Chest pain is the most common presenting cardiac symptom. It may be cardiac, respiratory, gastrointestinal (GI), musculoskeletal, or psychogenic in nature. In adults, chest pain is always treated as cardiac in origin until otherwise determined. Chest pain of cardiac nature is transmitted through the thoracic region by the five thoracic spinal nerves and may be referred to areas served by the cervical or lower thoracic segments. Also, these spinal nerves innervate the skin and the skeletal muscles. This explains the difficulty in determining the origin of chest pain and also explains the occurrence of cardiac chest pain in areas aside from the heart.



HELPFUL HINT

Chest pain that increases with position change or palpation or is described as pleuritic or sharp is probably not cardiac in origin.



ALERT

If your patient presents with chest pain, be sure to determine the time of onset because this may influence the course of treatment. Thrombolytic agents must be given within 6 hours of onset of pain to be effective.

Risk Factors for Coronary Artery Disease

Uncontrollable Factors

Age: Risk increases with age.

Sex: Incidence is greater in men than in women until women reach menopause.

Heredity: Positive family history increases the chance of heart disease.

Race: Incidence is higher for African American men and women under age 45 than in white men and women in the same age group.

Controllable Factors

Smoking: Smokers have a two to six times greater risk of developing CAD than nonsmokers.

Cholesterol: Serum cholesterol levels > 200 mg/dL increase risk for CAD.

Hypertension: Uncontrolled hypertension (BP > 120/80) increases risk for CAD.

Physical Inactivity: People who have a sedentary lifestyle are at greater risk for CAD.

Obesity: People who are 30 percent over desired weight for height and body build increases risk for CAD and other problems, such as type 2 diabetes and HTN.

Controllable Contributing Factors

Diabetes: Diabetes is a contributing factor for CAD.

Stress: Stress and being a type A personality (perfectionist, hard-working, driven).

Homocystine levels: High levels of homocystine, a sulfur-containing amino acid produced by the breakdown of the essential amino acid methionine found in dietary proteins, may contribute to atherosclerosis.

Palpitations

A patient with **palpitations** has the sense that her or his heart is racing or skipping beats.

Syncope

Syncopal attacks (dizziness) are another symptom that may signal cardiovascular problems. These episodes can result from decreased cerebral blood flow. Keep in mind that syncopal attacks may also result from noncardiac conditions or be side effects of medications.

Edema

Edema may be seen with right-sided CHF and vascular disease. Edema may result from an increased capillary hydrostatic pressure that shifts fluid from the capillaries to the tissue.

Fatigue

Fatigue is a feeling of being tired and having no energy. Often, a very subtle symptom, fatigue is associated with cardiovascular disease. However, it may not be apparent to you because patients tend to alter their activity levels to adapt to decreased energy levels.



HELPFUL HINTS

- To accurately assess fatigue, investigate changes in activities of daily living (ADLs), rest and sleep patterns.
- Visible vascular changes in the extremities may be a clue that more extensive systemic vascular changes are occurring.

Extremity Changes

Changes in the extremities may provide clues about underlying cardiovascular disease. Symptoms such as paresthesia (numbness, tingling), coolness, and intermittent claudication (pain in calves during ambulation) may be associated with vascular disease, coronary heart disease, or cerebral vascular disease.

Dyspnea and Cough

Although often associated with respiratory disorders, **dyspnea** may also occur with cardiac disease such as left-sided CHF. In this case, dyspnea results from pulmonary congestion or increased pulmonary venous and capillary pressure. See Chapter 13, Assessing the Respiratory System, for complete symptom analysis.

Past Health History

The purpose of the past health history is to help you identify factors in the patient's past that may have a direct link to his or her current cardiovascular status. Or you may uncover risk factors that place your patient at risk for cardiovascular disease. This part of the assessment focuses on areas as they relate to the cardiovascular system.

Past Health History

RISK FACTORS/QUESTIONS TO ASK

Childhood Illnesses

- Do you have a history of rheumatic fever or frequent streptococcal infections?
- Do you have any congenital heart defects?
- Do you have a heart murmur?

Hospitalizations/Surgeries

- Have you ever been hospitalized for cardiovascular problems?
- Have you ever had an ECG?
- Have you ever had surgery?

Serious Injuries

- Have you been in an accident recently?

Serious/Chronic Illnesses

- Do you have a history of diabetes, HTN, or hyperlipidemia?
- Do you have a history of chronic respiratory disease?
- Do you have a history of renal disease or bleeding disorders?

RATIONALE/SIGNIFICANCE

- May cause rheumatic heart disease.
- May have direct correlation to present status.
- Murmurs are very common during childhood. They are often innocent but may indicate pathology.
- An ECG can be used to establish baseline and comparative data and to evaluate the status of chronic cardiovascular disease.
- Blunt chest trauma or acceleration/deceleration may have caused myocardial contusion or pericardial tamponade.
- If controlled, these cardiac risk factors are alterable.
- Long-standing respiratory disease often leads to cardiac involvement.
- Can affect cardiovascular system.

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Immunizations</p> <ul style="list-style-type: none"> ■ Did you get a flu shot or Pneumovax this year? <p>Allergies</p> <ul style="list-style-type: none"> ■ Do you have any allergies? (Specifically ask about allergies to iodine or shellfish.) <div style="border: 1px solid red; padding: 5px; margin: 10px 0;">  <p>ALERT</p> <p>If your patient is allergic to shellfish, she or he may be allergic to the dye used for cardiac catheterization.</p> </div> <p>Medications</p> <ul style="list-style-type: none"> ■ Are you taking any prescribed or over-the-counter (OTC) medications, including herbal supplements? 	<ul style="list-style-type: none"> ■ For patients with chronic cardiovascular disease, flu or pneumonia can place added stress on an already compromised cardiac status, resulting in decompensation. ■ Allergies to foods, drugs, or environmental factors may influence treatment. ■ Assess patient's compliance with treatment plan. Medications or interaction between prescribed and OTC medications, such as a side effect or toxic effect, may be a direct or contributing factor to presenting symptoms. ■ Drugs such as digoxin, diuretics, nitrates, antihypertensives, and anticoagulants may cause a variety of symptoms, including cardiac arrhythmias, GI upset, visual disturbances, headaches, muscle cramps, dizziness, and bleeding. See <i>Drugs That Adversely Affect the Cardiovascular System</i>.

Drugs That Adversely Affect the Cardiovascular System

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Antidepressants	Trazodone hydrochloride	Hypotension, HTN, syncope, tachycardia, palpitations, ECG changes
	Tricyclic antidepressants	Postural hypotension, hypertension, ECG changes, dysrhythmias, syncope
Antineoplastics	Daunorubicin, doxorubicin	Dose-dependent CMP manifested by CHF, ECG changes, dysrhythmias
Antipsychotics	Phenothiazines, ziprasidone, haloperidol, risperidone, quetiapine, olanzapine, aripiprazole	Hypotension, postural hypotension, tachycardia, syncope, ECG changes, dysrhythmias
Anxiolytics	Diazepam	Hypotension, bradycardia, cardiac arrest, dysrhythmias (with intravenous route)
	Lorazepam	Hypotension
Antifungal	Midazolam	Hypotension, cardiorespiratory arrest
Bronchodilators, antiasthmatic agents	Aminophylline, theophylline	Palpitations, sinus tachycardia, extrasystoles, ventricular dysrhythmias, hypotension
Cerebral stimulants	Amphetamine	Tachycardia, palpitations, dysrhythmias, HTN
Hormones	Caffeine	Tachycardia, HTN
	Oral contraceptives	HTN, fluid retention, increased risk of stroke, MI, thromboembolism
	Conjugated estrogens, estradiol, oral contraceptives	HTN, thromboembolism, thrombophlebitis

(continued)

Drugs That Adversely Affect the Cardiovascular System (continued)

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
	Vasopressin	Angina in patients with vascular disease; in large doses, HTN, MI
Narcotic agents	Morphine	Hypotension, bradycardia
Miscellaneous agents	Bethanechol	Hypotension, tachycardia
	Hydralazine	Tachycardia, angina pectoris
	Levodopa-carbidopa	Orthostatic hypotension
	Levothyroxine	With excessive doses, angina pectoris, dysrhythmias, tachycardia, HTN
	Phenytoin	Hypotension, bradycardia, ventricular fibrillation (with intravenous route)

Family History

The family history will help you identify familial or genetically linked risk factors that predispose your patient to

cardiovascular disease. It is important to note not only the type of problem but also the age at which the problem occurred.

Family History

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Familial/Genetically Linked Cardiovascular Disorders</p> <ul style="list-style-type: none"> ■ Does your family have a history of: <ul style="list-style-type: none"> ■ Familial hyperlipoproteinemia? ■ HTN, CAD, or MI? ■ Diabetes, HTN, or renal disease? ■ Genetically linked cardiac disorders such as Marfan's syndrome? ■ Mitral valve prolapse? 	<ul style="list-style-type: none"> ■ Positive family history is an uncontrollable risk factor for CAD. ■ An inherited autosomal dominant trait that may increase risk for CAD at an early age. ■ Increases risk for CAD. ■ An inherited autosomal-dominant trait characterized by elongation of bones that often has associated cardiovascular abnormalities. ■ Causes one or both leaflets to billow into the atrium during ventricular systole. Runs in families.

Review of Systems

Changes in the cardiovascular system may affect every other system of the body. The review of systems (ROS) will help you identify problems in other systems that will directly effect the cardiovascular system and also

help you determine the effect and extent of cardiovascular disease on other body systems. The ROS allows you to pick up on symptoms you might have missed during your health history by jogging your patient's memory.

Review of Systems

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>General Health Survey</p> <ul style="list-style-type: none"> ■ Have you experienced fatigue or activity intolerance? ■ Have you had influenza, other recent illness, or weight gain? 	<ul style="list-style-type: none"> ■ Chronic cardiovascular disease (chronic CHF) causes decreased cardiac output, impaired circulation, and decreased oxygen and often leads to early fatigue and difficulty performing ADLs. ■ Increasing fatigue and activity intolerance may correlate with disease's progression. ■ Influenza or other recent illness can cause CMP. ■ Sudden weight gain may indicate fluid retention.

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Integumentary	
Skin	
<ul style="list-style-type: none"> ■ Have you had any changes in skin texture, color, or temperature? ■ Do you have any sores or ulcers that won't heal? ■ Have your ankles swollen or your shoes become tight? 	<ul style="list-style-type: none"> ■ Skin changes may indicate vascular insufficiency. ■ Poor wound healing may signal diabetes, which is a risk factor for CAD. ■ Edema is associated with vascular disease and CHF.
Nails	
<ul style="list-style-type: none"> ■ Have your nails changed in shape or in color? 	<ul style="list-style-type: none"> ■ Clubbing and cyanosis may reflect chronic cardiopulmonary problem.
HEENT	
Head	
<ul style="list-style-type: none"> ■ Do you have headaches? ■ Do you have dizzy spells? 	<ul style="list-style-type: none"> ■ May indicate HTN, a risk factor for CAD. ■ Syncopal attacks may occur with vascular disease or cardiac arrhythmias or may be a medication side effect.
Eyes	
<ul style="list-style-type: none"> ■ Have you experienced visual problems such as blurred vision, double vision, or colored spots? 	<ul style="list-style-type: none"> ■ Double vision and temporary loss of vision are associated with HTN, transient ischemic attack, cerebrovascular insufficiency, and digitalis toxicity. ■ Yellow spots may signal digitalis toxicity.
Ears	
<ul style="list-style-type: none"> ■ Have you experienced ringing in your ears? 	<ul style="list-style-type: none"> ■ Tinnitus is associated with cerebrovascular insufficiency.
Nose	
<ul style="list-style-type: none"> ■ Have you experienced nosebleeds? 	<ul style="list-style-type: none"> ■ Epistaxis is associated with HTN.
Throat	
<ul style="list-style-type: none"> ■ Have you experienced frequent strep throats? 	<ul style="list-style-type: none"> ■ Beta-hemolytic streptococcal infection is associated with rheumatic heart disease.
Respiratory	
<ul style="list-style-type: none"> ■ Do you experience breathing difficulties? ■ Do you have a history of COPD? 	<ul style="list-style-type: none"> ■ Dry cough, shortness of breath (SOB), dyspnea on exertion (DOE), paroxysmal nocturnal dyspnea (PND), orthopnea, and cough are symptoms of left-sided CHF. ■ Chronic COPD can result in cardiac involvement, such as pulmonary HTN and right-sided CHF.
Gastrointestinal	
<ul style="list-style-type: none"> ■ Have you experienced right upper quadrant (RUQ) pain, nausea, GI upset? 	<ul style="list-style-type: none"> ■ GI upset and RUQ pain may accompany right-sided CHF. ■ GI complaints are associated with medications such as digitalis.
Genitourinary	
<ul style="list-style-type: none"> ■ Have you experienced changes in urination, such as waking up at night to go to the bathroom? 	<ul style="list-style-type: none"> ■ CHF leads to decreased renal perfusion during the day. But at night, when patient is in a recumbent position, fluid moves from interstitial spaces back into circulatory system, increasing renal blood flow and causing diuresis (nocturia).
Reproductive	
Female	
<ul style="list-style-type: none"> ■ Are you postmenopausal? ■ Do you use oral contraceptives, or are you on hormone replacement therapy (HRT)? ■ During pregnancy, did you have gestational diabetes or pregnancy-induced hypertension (PIH)? 	<ul style="list-style-type: none"> ■ Increases CAD risk. ■ Oral contraceptives/estrogen supplements are associated with thrombus formation. ■ HRT is associated with increased risk of cardiovascular disease. ■ Increased risk for developing diabetes later in life, which increases risk for CAD.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Male</p> <ul style="list-style-type: none"> ■ Do you have problems with impotence or sexual performance? ■ Do you have chest pain during sexual activity? <p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Do you have muscle weakness? ■ Do you experience leg-muscle cramps when walking? <p>Neurological</p> <ul style="list-style-type: none"> ■ Do you experience fainting episodes, loss of consciousness, or headaches? ■ Do you experience behavioral changes such as confusion, decreased attention span or loss of memory? <p>Endocrine</p> <ul style="list-style-type: none"> ■ Do you have diabetes or thyroid disease? <p>Lymphatic/Hematologic</p> <ul style="list-style-type: none"> ■ Have you had bleeding problems? ■ Have you had a recent infection? 	<ul style="list-style-type: none"> ■ Impotence/erectile dysfunction may be caused by vascular disease, diabetes, or medication. ■ Sexual activity increases the heart's workload and can precipitate an angina attack. ■ Chronic cardiovascular disease may result in weakness secondary to decreased use. ■ Intermittent claudication is associated with arterial insufficiency. ■ Syncopal attacks may signal vascular problems or cardiac arrhythmias. ■ HTN or chronic CHF may cause hypoxia and impair cerebral circulation. ■ Diabetes is a known risk factor for CAD. Hyperthyroid disease can lead to hypertrophic CMP. ■ Anemia increases the heart's workload. Polycythemia increases risk for thrombus, HTN, and cardiopulmonary disease. ■ CMP.

Psychosocial Profile

The psychosocial profile enables you to examine your patient's lifestyle and identify risk factors for cardiovascular disease. Identifying risk factors is essential in developing an

effective plan of care. Chronic cardiovascular disease can also affect your patient's lifestyle, forcing her or him to make changes to adapt to the illness.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Health Practices and Beliefs/Self-Care Activities</p> <ul style="list-style-type: none"> ■ Do you get yearly physicals? ■ Do you see a doctor or nurse regularly? ■ What medications are you taking and why? <p>Typical Day</p> <ul style="list-style-type: none"> ■ What is your typical day? Has it changed over the last year? For example, are there activities you find difficult to do or are unable to do? <p>Nutrition/Weight Patterns</p> <ul style="list-style-type: none"> ■ Do you have weight problems—for example, obesity or a sudden weight increase? 	<ul style="list-style-type: none"> ■ Determines preventive practices. ■ Ascertains compliance with treatment program. ■ Identifies teaching needs. ■ Activity level can correlate with energy level. ■ Chronic heart disease decreases energy levels. ■ Changes in ADLs may reflect patient's attempt to adapt to progression of illness. ■ Obesity is a risk factor for CAD. ■ Sudden weight increases are usually associated with fluid retention.

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ What did you eat in the last 24 hr? ■ Do you get chest discomfort after eating? ■ Do you have anorexia, loss of appetite, or nausea? 	<ul style="list-style-type: none"> ■ 24-hr recall helps identify diets high in cholesterol and sodium, which may contribute to cardiovascular disease. ■ Large, heavy meals can precipitate an angina attack in patients with CAD. ■ May indicate right-sided CHF or be a side effect or toxic effect of cardiac medications.
<p>Activity/Exercise Patterns</p> <ul style="list-style-type: none"> ■ Do you exercise routinely? ■ Do you have chest discomfort after certain types of activity or exercise? 	<ul style="list-style-type: none"> ■ A gradual decrease or change in activity or exercise patterns is seen in patients with chronic cardiovascular disease. ■ Exercise can precipitate an angina attack in patients with CAD. ■ May identify need for referral for cardiac rehabilitation.
<p>Recreation/Pets/Hobbies</p> <ul style="list-style-type: none"> ■ Do you have any pets or hobbies? What kinds? <p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ Do you awaken during the night to go to bathroom? ■ Do you awaken with SOB? ■ Do you awaken with chest pain? 	<ul style="list-style-type: none"> ■ Pets and hobbies can be a good way to reduce stress. ■ Nocturia is associated with CHF. ■ PND is associated with CHF. ■ Angina may occur during rest (nocturnal angina and Prinzmetal's angina). During the rapid eye movement (REM) cycle of sleep, myocardial oxygen demands increase. This may explain the high incidence of MIs in the early morning hours. ■ The incidence of sudden cardiac death is higher in patients with sleep apnea.
<ul style="list-style-type: none"> ■ Do you snore? 	
<p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you smoke cigarettes? How many packs per day and for how many years? ■ Do you use street drugs such as cocaine? ■ Do you drink alcohol? How many glasses a day? 	<ul style="list-style-type: none"> ■ Smoking cigarettes is a known risk factor for CAD. Nicotine increases catecholamine release, which leads to vasoconstriction and increased heart rate and BP. This increases the heart's workload, and increased carbon monoxide levels result in decreased oxygen supply. ■ <i>"Crack" heart</i>: Cocaine increases catecholamine release, which increases heart rate and the heart's workload. May result in MI, CHF, or CMP. ■ Alcohol abuse can lead to increased pulse and BP, CMP, CAD, HTN, and stroke.
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ What is your job? Are you currently working? ■ How many hours a day do you usually work? Does your job make physical demands? 	<ul style="list-style-type: none"> ■ Cardiac limitations may prevent patient from working. ■ Helps establish cardiovascular workload and helps you devise an activity plan.



HELPFUL HINT

Prinzmetal's (variant) angina usually has no precipitating factors and occurs at rest as a result of spasms of the coronary artery. Patients with Prinzmetal's angina frequently have a history of migraine headaches and Raynaud's disease.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> Are you exposed to smoke, noise, extreme temperatures, or dust? <p>Environmental Health Patterns</p> <ul style="list-style-type: none"> Where do you live? Are there stairs? <p>Roles/Relationships/Social Supports</p> <ul style="list-style-type: none"> Do you belong to any church or community groups? What is your role in family, church, and community? What are your support systems? <p>Sexuality Patterns</p> <ul style="list-style-type: none"> How has your cardiovascular problem affected your sex life? <p>Stress and Coping</p> <ul style="list-style-type: none"> What do you do when you feel upset, angry, frustrated, or stressed out? 	<ul style="list-style-type: none"> May cause cardiovascular symptoms. As CAD progresses, patient may have difficulty within own environment. Referrals may be indicated. Identify discharge or home-care needs. Chronic cardiovascular disease may affect patient's ability to perform role tasks. Roles may change as disease progresses, and referrals may be needed. Patient may isolate self as disease progresses, becoming homebound as a "cardiac cripple." Be alert for signs of depression. Fear of heart damage may cause patients and spouses to avoid sex or become impotent. Antihypertensive or antianginal medications may also cause impotence. Stress increases CAD risk by increased stimulation of sympathetic nervous system. Volatile emotions such as anger also increase sympathetic nervous system response, which in turn increases the heart's workload.



Case Study Findings

Reviewing Mr. Brusca's

Biographical data:

- 68-year-old white male.
- Married, father of seven grown children.
- Self-employed entrepreneur; BS degree in engineering.
- Born and raised in the United States, Italian descent, Catholic religion.
- Blue Cross/Blue Shield medical insurance plan.
- Referral:* Follow-up by primary care physician.
- Source:* Self, reliable.

Current health status:

- No chest pain, dyspnea, palpitations, or edema.
- Complains of fatigue, loss of energy, and occasional dizzy spells.

Past health history:

- No rheumatic fever or heart murmurs.
- No history of injuries.
- Inguinal hernia repair.
- Left ventricular hypertrophy revealed by electrocardiogram (ECG).

- Hospitalized 3 weeks ago for HTN.
- No known food, drug, or environmental allergies.
- No other previous medical problems.
- Immunizations up to date.
- No prescribed medications except Vasotec 5 mg bid and weekly use of antacid for indigestion.

Family history:

- Positive family history of HTN and stroke.
 - Mother had HTN and died at age 78 of a stroke.
 - Paternal uncle died at age 79 of MI.

Review of systems:

- General Health Survey:* Fatigue, weight gain of 60 lb over past 3 years.
- Integumentary:* Feet cold, thick nails, tight shoes.
- Head, Eyes, Ears, Nose, and Throat (HEENT):* Two dizzy spells over past 6 months.
 - Eyes:* Wears glasses, no visual complaints, yearly eye examination.
 - Respiratory:* "Short winded" with activity.
 - Gastrointestinal:* Indigestion on weekly basis.

- **Genitourinary:** Awakens at least once a night to go to bathroom.
- **Reproductive:** Little sexual activity because he is “too tired”; becoming more and more difficult to have an erection.
- **Musculoskeletal/Neurological:** General weakness, cramps in legs with walking.
- **Lymphatic:** No reported problems.
- **Endocrine:** No reported problems.

Psychosocial profile:

- States that he does not have time for routine checkups. “I only go to the doctor’s when I’m sick.” Typical day consists of arising at 7 A.M., showering, having breakfast, and then going to work. Returns home by 6 P.M., eats dinner, watches TV till 11:30 P.M., but usually falls asleep before news is over. Usually in bed by 12 midnight.
- 24-hour recall reveals a diet high in carbohydrates and fats and lacking in fruits and vegetables. Heavy-handed with salt shaker; salts everything. Admits that he has gained weight over the years and is 60 lb overweight.
- No regular exercise program. States: “I’m too busy running my business.”
- Hobbies include reading, crossword puzzles, and antique collecting.
- Sleeps about 7 hours a night, but usually feels he is not getting enough sleep. Lately is more and more tired. Wife states that he snores.
- Never smoked. Has a bottle of wine every night with dinner.
- Works at sedentary job, usually 7 days a week. No environmental hazards in workplace.
- Lives with wife of 45 years in a two-story, single home in the suburbs with ample living space.
- Has a large, close, caring family.
- Admits that running his own business is very stressful, but feels he can handle it alone and doesn’t need anyone to help him.



CRITICAL THINKING ACTIVITY #1

From the subjective information you have obtained from Mr. Brusca’s history, what are his identifiable risk factors for heart disease? Which risk factors are modifiable and which are unmodifiable?



Case Study Evaluation

Before you begin the physical assessment, summarize Mr. Brusca’s subjective findings obtained from his history. What key points in his history will help you plan his care? Document his cardiovascular history.




FIGURE 14.14. Anatomical site of heart.

Anatomical Landmarks

Before you begin the physical assessment, visualize the underlying structures and identify cardiac landmarks to ensure an accurate and thorough assessment. The heart is situated in the chest in the third, fourth, and fifth intercostal spaces to the left of the sternum. In a normal-sized heart, the right side is located more anteriorly, with only the tip of the left ventricle situated anteriorly (Fig. 14.14).

Even though heart sounds may be heard anywhere on the precordium, specific sites have been identified as best for performing the cardiac assessment: right base/aortic area, left base/pulmonic area, Erb’s point/third intercostal space, LLSB (tricuspid), and apex (mitral) (Fig. 14.15).

Cardiac Auscultation Sites

Traditional Sites	Alternative Sites
Apex/mitral area	Apex
LLSB/tricuspid area	Lower left sternal border
Erb’s point	Left base
Base left/pulmonic area	Right base
Base right/aortic area	Xiphoid

Central Artery and Jugular Veins

Before you begin assessing the central arteries and jugular veins, identify your anatomical landmarks to ensure accurate assessment. The carotid and the internal and external jugular need to be located. The sternocleidomastoid and trapezius muscles are helpful in locating these vessels. The carotid artery and the internal jugular run parallel to each other along the sternocleidomastoid muscle toward the sternal notch. The external jugular crosses the internal jugular and lies posterior to the




FIGURE 14.15. Sites for cardiac assessment.

sternocleidomastoid muscle. These neck structures are shown in Figure 14.16.

Physical Assessment

The significant subjective data that you obtained from the health history will help to direct your physical examination. The history provides you with clues about what physical findings you can expect to find in your patient. These physical findings provide the objective database.

Together, the health history and physical examination complete your assessment picture. You can then analyze the data, formulate nursing diagnoses, and develop a plan of care for your patient.

The physical examination begins with a general survey, proceeds to a head-to-toe assessment of each body system, and then ends with a focused cardiovascular assessment.

Approach

You will use all four techniques of physical assessment to assess the cardiovascular system. Because changes in this system may affect every other body system, assess for physical signs in every system. Perform the assessment in three positions: sitting, supine, and left lateral recumbent. Placing the patient in several positions facilitates the auscultation of specific heart sounds. However, keep in mind that patients with chronic cardiovascular problems may have little cardiac reserve. You may need to minimize position changes to conserve your patient's energy.

Work from the right side of your patient. This is especially helpful during auscultation because it allows you to stretch the tubing of the stethoscope and minimize interference. Allay your patient's anxiety by explaining that a cardiovascular examination always takes time. Be systematic in your approach. Your first assessment will serve as the baseline that will be used for subsequent assessments. This baseline will enable you to detect subtle changes in the patient's cardiovascular status.

Before you begin, wash your hands, assemble your equipment, and prepare a warm, quiet, private environment. Let your patient know what you will be doing. This




FIGURE 14.16. Neck structures.

will help to minimize his or her anxiety and maximize cooperation during the examination. Sharing your findings with your patient can help you validate them and provide an ideal opportunity for patient teaching.



TOOLBOX

The tools for cardiovascular assessment include a stethoscope with a diaphragm for high-pitched sounds, and a bell for low-pitched sounds, a sphygmomanometer, scale, metric ruler, thermometer, and marking pen. Sharpen your senses; you will need all of them to perform a cardiovascular assessment.

Performing a General Survey

Before assessing specific areas, perform a general survey that includes evaluating the patient's general appearance, vital signs, height, and weight. Be alert for signs of cardiovascular disease.

General Appearance

General appearance can give you important clues about the patient's cardiovascular status. Note the following:

- **Chronological age:** Is the patient's age consistent with her or his appearance? Chronic illness can "age" a person.
- **Weight distribution and muscle composition:** Look for edema that may indicate cardiovascular disease. Muscle wasting may occur from lack of use.
- **Facial expression:** Does it match the patient's behavior? If he or she presents with chest pain, his or her expression and behavior should be consistent with the complaint.
- **Posture or assumed position of comfort:** The patient's posture should be erect and midline. Variations from this may reflect fatigue associated with chronic cardiac disease. A sitting position, to ease breathing, may be the preferred position for the patient with CHF.
- **Dress and grooming:** The patient with chronic cardiovascular disease may have increasing difficulty performing ADLs, and this may be reflected in her or his dress and grooming.
- **Body stature:** Several genetic disorders have characteristic stature profiles and associated cardiac problems. Tall stature with long limbs in proportion to the trunk is seen with **Marfan's syndrome**, a genetic disorder associated with aortic dissection, aortic valve incompetence, and cardiomyopathy (CMP). Tall stature and long extremities are also seen with **Klinefelter's syndrome**, a genetic disorder in males characterized by an extra X chromosome with possible atrial or ventricular septal defects, patent ductus arteriosus, and tetralogy of Fallot. The short stature of patients with

Turner's syndrome is associated with coarctation of the aorta and pulmonic stenosis.

- **Muscle wasting:** Cardiac cachexia and muscle wasting on the upper arms, thighs, and chest wall may be seen in patients with chronic cardiac disease. Anthropometric measurements may be made to accurately determine the degree of cachexia. The normal percentage of body fat for men is 14 to 18 percent; for women it is 20 to 25 percent. Loss of body fat and muscle mass not only reduces the body's energy stores but also affects the immune system and healing process.
- **Abnormal movements:** Rhythmic bobbing of the head simultaneous with the heartbeat (Mussel's sign) may result from high back pressure associated with aortic insufficiency or aneurysm. Squatting position is seen in patients with tetralogy of Fallot.

Vital Signs

BP, pulse, respirations, and temperature are a good reflection of how well the cardiovascular system is functioning.

Blood Pressure

BP reflects stroke volume, distensibility of arteries, peripheral vascular resistance, and volume. BP measurements greater than 120/80 may indicate HTN; BP with a systolic less than 100 may indicate hypovolemia or CHF.



HELPFUL HINT

If the patient's BP is high, serial readings of elevated BP will help confirm a diagnosis of HTN.

Pulse

The quickest way to increase cardiac output is to increase heart rate. Therefore, changes in heart rate are often adaptive mechanisms by the body as it attempts to maintain homeostasis. Aside from heart rate, take note of rhythm. An irregular rhythm indicates an underlying arrhythmia. Also check for a pulse deficit—a difference between the radial pulse and the apical pulse—which can occur with atrial fibrillation. The strength of the pulse may reflect the vascular system; bounding pulses may be seen with HTN, and weak pulses or pulsus alternans (alternating weak and strong pulse) may reflect a failing heart.



ALERT

If you get an irregular radial pulse, take an apical rate for 1 full minute. If the apical pulse is also irregular, report these findings to the physician.

Respiratory Rate

Patients with CHF or pulmonary edema have an increased respiratory rate, dyspnea, and orthopnea. The respiratory rate usually increases in response to hypoxia. If the patient has chest pain, you will also see changes in the respiratory rate and pattern.

Temperature

Elevated temperature may indicate an infectious disease process. Slight elevation in body temperature is often seen within the first 24 hours after an acute MI because of the inflammatory process that occurs during an MI. The elevation is usually low grade at 100.4°F, but it may rise to 102.2°F and last for 1 week post MI.

Acute pericarditis may also result in temperature elevations to 102.2°F, and even higher elevations are seen with bacterial endocarditis. No matter what the cause, fever increases metabolism, which in turn increases cardiac workload. The cardiac response to this increase is tachycardia.

Height and Weight

Accurate height and weight measurements are helpful in planning treatment, calculating dosages, identifying nutritional deficits, and detecting and monitoring fluid retention.

Height

As mentioned previously, tall stature is linked to several genetic disorders that have associated cardiac problems, such as Marfan's syndrome, Klinefelter's syndrome, and

acromegaly. Short stature may also have genetic links and be associated with cardiac problems. Examples are Turner's syndrome and dwarfism.

Weight

Weighing your patient is the most accurate means of monitoring fluid status. Note any weight changes, especially sudden weight gains, which suggest fluid retention as is seen with CHF. Changes of 2 to 3 pounds in 48 hours usually reflect a change in fluid status, not body mass. A patient can gain 6 to 8 pounds before edema may be visible, so weighing your patient every day at the same time and with the same scale is the most accurate means of assessing fluid status.

Performing a Head-to-Toe Physical Assessment

The cardiovascular system can affect every other body system. Therefore, once you have done a general survey, look at the patient from head to toe, checking for changes in each body system that might signal a cardiovascular problem.

Performing a Cardiovascular Physical Assessment

Now that you have completed your general survey and head-to-toe assessment, focus your attention on assessing the cardiac system. All four techniques of physical assessment are used. Figure 14.17 shows the positions for cardiac examination.

(Text continued on page 470)



FIGURE 14.17. Positions for cardiac assessment. **A.** Sitting position. **B.** Supine position. **C.** Left lateral position.

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	Ask about: Fatigue, DOE, activity intolerance, recent illness or flu, weight changes	General	Inspect: Signs of chest pain, SOB, posture, orthopnea changes in VS Measure: Height and weight
Integumentary	Ask about: Skin changes Poor wound healing Skin temperature changes Nail color changes Ankle swelling Tight shoes	Integumentary	Inspect: Color changes (e.g., cyanosis or pallor) Color changes in legs (e.g., red or brown) Lesions (e.g., petechiae or leg wounds) Taut, shiny, hairless skin Nails for clubbing Palpate: Skin temperature, turgor, edema Capillary refill
HEENT <i>Head and Neck</i>	Ask about: Headaches, dizzy spells	HEENT <i>Head and Neck</i>	Inspect: Facial expression Palpate: Thyroid, neck vein distention
Eyes	Ask about: Blurred or double vision, yellow spots	Eyes	Test: Visual acuity Inspect: Periorbital edema, xanthelasma, arcus senilis, sclera icterus, exophthalmos Examine: Fundus for AV nicking, exudates, cotton wool, hemorrhages
Ears, Nose and Throat	Ask about: Ringing in ears, frequent "strep" throat, nosebleeds	Ears, Nose and Throat	Note: Gross hearing Inspect: Oropharynx, pink tonsils
Respiratory	Ask about: Breathing difficulties (e.g., SOB, DOE, PND), dry cough, history of COPD	Respiratory	Inspect: Chest deformities, respiratory rate, retraction or use of accessory muscles Auscultate: Adventitious sounds such as crackles
Gastrointestinal	Ask about: RUQ pain, nausea, GI upset	Gastrointestinal	Inspect: Ascites Palpate: Hepatomegaly
Neurological	Ask about: Fainting episodes Behavioral changes, confusion, memory loss	Neurological	Check: Mental status (AAO x 3) Immediate, recent, remote memory Judgment Decreased/diminished sensations
Musculoskeletal	Ask about: Weakness Muscle cramps	Musculoskeletal	Test: Muscle strength Inspect: Atrophy, wasting
Lymphatic/Hematologic	Ask about: Bleeding Recent infections	Lymphatic/Hematologic	Inspect: Skin for ecchymosis, petechiae Palpate: Enlarged lymph nodes
Endocrine	Ask about: History of diabetes or thyroid disease	Endocrine	Palpate: Thyroid Auscultate: Bruits over thyroid
Genitourinary	Ask about: Awakening to urinate (nocturia) Men: Sexual performance problems Women: Pre- or postmenopausal, birth control pills, hormone replacement therapy		

Assessment of Cardiovascular System's Relationship to Other Systems

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

ABNORMAL FINDINGS/RATIONALE

General Health Survey

INSPECT

- Awake, alert, and oriented to time, place, and person.
- Immediate, recent, and remote memory intact. Judgment intact, compliant with treatment plan.

- *Confusion, restlessness, irritability, short attention span:* Early signs of hypoxia and impaired cerebral circulation caused by decreased cardiac output.
- *Memory loss:* Hypoxia.
- Patients with cardiovascular disease may deny their illness, severity of symptoms, or associated risk factors.
- *Fatigue:* Often accompanies chronic cardiovascular disease.

Integumentary

Skin

INSPECT/PALPATE

- Normal skin color varies from patient to patient because of racial and genetic differences. Buccal mucosa pink, but may be darker (including dark brown or blue) in patients of Mediterranean or African descent.
- Peripheral cyanosis can be a normal response to cold.
- Changes from the patient's normal color should be noted.
- Skin intact, no lesions.
- Skin pink, warm and dry, +2 pedal pulses, no edema.
- Positive skin turgor. Poor skin turgor may be normal in older adults as a result of decreased elasticity.
- No edema.

- If cyanosis is present, differentiate between central and peripheral.
- *Central cyanosis:* Suggests a cardiopulmonary problem, such as CHF, MI, or pulmonary edema. Look for central cyanosis in the oral mucosa.
- *Peripheral cyanosis in absence of cold:* Hypovolemia, shock, or peripheral vascular disease caused by vasoconstriction, vascular occlusion, or decreased cardiac output.
- *Ashen color:* MI.
- *Dependent rubor and pallor with elevation of extremities:* Arterial insufficiency.
- *Pallor:* Anemia and high-output failure.
- *Jaundice:* Hepatic congestion secondary to right CHF.
- *Cyanosis when dependent and brown pigmentation:* Venous insufficiency.
- *Petechiae:* Bacterial endocarditis.
- *Thin, shiny, hairless, cool skin with decreased or absent pulses:* Arterial insufficiency.
- *Peripheral edema, leathery skin with cyanosis and brownish discoloration:* Venous insufficiency.
- *Peripheral edema:* Right-sided CHF.
- *Cool, clammy skin:* Shock, MI, CHF, arterial insufficiency, or occlusion as a result of vasoconstriction in response to decreased cardiac output. This is a compensatory mechanism to shunt blood to vital organs.
- *Taunt, shiny skin:* Edema caused by cardiovascular disease such as right CHF or vascular insufficiency.
- *Anasarca (generalized body edema):* Right CHF.



HELPFUL HINT

When assessing for edema, check the patient's most dependent areas such as the legs, sacrum, and scrotum.

Hair

INSPECT/PALPATE

- Normal hair distribution for age and sex.

- *Thinning or loss of hair on extremities:* Arterial insufficiency.

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p>Nails INSPECT/PALPATE</p> <ul style="list-style-type: none"> ■ Nailbeds pink to dark brown or blue in patients of Mediterranean or African descent; positive capillary refill; angle of attachment 160 degrees; no clubbing, 	<ul style="list-style-type: none"> ■ <i>Cyanotic nailbeds with poor capillary refill and clubbing:</i> Cardiopulmonary disease. ■ <i>Splinter hemorrhages:</i> Bacterial endocarditis.
<p>HEENT Head and Neck INSPECT/PALPATE</p> <ul style="list-style-type: none"> ■ Facial expression appropriate. ■ Thyroid not palpable. 	<ul style="list-style-type: none"> ■ <i>Struggling, frightened look:</i> Pulmonary edema. ■ <i>Apprehension:</i> CAD, MI. ■ <i>Enlarged thyroid:</i> Hyperthyroid disease, which can affect the cardiac system, resulting in CHF
<p>Eyes INSPECT</p> <ul style="list-style-type: none"> ■ Eyes clear and bright, no lesions or edema. <ul style="list-style-type: none"> ■ Arcus senilis (thin, whitish-gray ring around cornea) may be seen in older patients. ■ Sclera white; palpebral conjunctiva pink. ■ No lid lag or exophthalmos. ■ Funduscopic examination reveals yellow to orange retina without hemorrhages or exudates. Arteriovenous (AV) crossings smooth. 	<ul style="list-style-type: none"> ■ <i>Periorbital edema, xanthelasma, arcus senilis, sclera icterus:</i> Suggest cardiovascular disorders. Edema is also associated with CHF. ■ <i>Xanthelasma (small, yellow, raised plaques on eyelids):</i> Lipid deposits that may indicate hyperlipidemia. ■ <i>Arcus senilis in patients under age 65:</i> Hyperlipidemia. ■ <i>Sclera icterus (jaundiced sclera):</i> Right-sided CHF (cardiac cirrhosis). ■ <i>Cyanosis of palpebral conjunctiva:</i> CHF, MI, pulmonary edema. ■ <i>Exophthalmos and lid lag:</i> Hyperthyroidism, which can result in supraventricular tachycardia (SVT), angina, or high-output failure. ■ HTN and diabetes are risk factors for CAD, so it is important to note funduscopic changes. ■ <i>Funduscopic changes associated with HTN:</i> Papilledema, AV nicking, hard exudates, creamy yellow lesions, soft exudates, such as cotton wool (also seen with subacute bacterial endocarditis). ■ <i>Funduscopic changes associated with diabetes:</i> Diabetic retinopathy.
<p>Respiratory INSPECT/AUSCULTATE</p> <ul style="list-style-type: none"> ■ Respirations 16/min, lungs clear on auscultation. <ul style="list-style-type: none"> ■ Chest AP: Lateral 1:2, negative barrel chest or spinal deformities. ■ No retraction or use of accessory muscles 	<ul style="list-style-type: none"> ■ <i>Breathing difficulties, such as dyspnea, DOE, PND, and tachypnea:</i> Associated with CHF, pulmonary edema, MI. ■ <i>Cheyne-Stokes respiratory pattern:</i> Severe CHF. In patients with CHF, further assessment reveals crackles and wheezes. Cardiac asthma can develop with CHF. ■ <i>Barrel chest:</i> COPD. Chronic COPD often has right-sided cardiac involvement. ■ <i>Pectus excavatum (depressed sternum), scoliosis (lateral deviation of spine), and kyphosis (accentuated convex thoracic curve):</i> If severe, may affect cardiac functioning by impairing chest expansion and cardiac movement. ■ <i>Retraction of intercostals and use of accessory muscles:</i> Respiratory difficulty caused by a respiratory disorder, congenital heart defect, or CHF.
<p>Gastrointestinal INSPECT/PALPATE/AUSCULTATE</p> <ul style="list-style-type: none"> ■ Positive bowel sounds; abdomen soft, nontender; no bruits or venous hums; liver not palpable; positive pulsation in epigastric area; abdominal aorta < 2.5 cm. 	<ul style="list-style-type: none"> ■ <i>Ascites, hepatomegaly, RUQ tenderness, venous hum:</i> Right CHF. ■ <i>Positive bruit or wide, diffuse pulsation in epigastric area:</i> Abdominal aortic aneurysm.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
Musculoskeletal <i>INSPECT/PALPATE</i> <ul style="list-style-type: none"> Grade 4 to 5 muscle strength, no atrophy. 	<ul style="list-style-type: none"> <i>Muscle weakness:</i> Chronic cardiac disease resulting in muscle atrophy from lack of use.
Neurological <i>INSPECT/PALPATE</i> <ul style="list-style-type: none"> Sensory intact in extremities. +2 DTR. 	<ul style="list-style-type: none"> <i>Decreased sensation and diminished DTR:</i> Peripheral neuropathy associated with diabetes, which is a risk factor for CAD.

Inspection

You will need to inspect the great vessels in the neck and the precordium for pulsations. Examining these areas from different angles and with tangential lighting will help you visualize pulsations.

Neck

With the patient in a supine position, inspect the carotid and jugular venous systems in the neck for pulsations. Remember that the carotid artery and the internal jugular vein lie parallel to the sternocleidomastoid muscle. Although the internal jugular vein is larger than the external jugular vein, the external jugular is more superficial, making visualization easier. Because the internal jugular lies deep, you will be able to see only the jugular pressure wave. To visualize external venous pulsations, look for pulsations in the supraclavicular area. To visualize internal venous pulsations, look for pulsations at the suprasternal notch. Using a penlight to cast a shadow on the neck vessels may help you visualize the pulsations.

To measure jugular venous pressure (JVP), elevate the patient's head to 30 to 45 degrees. Identify the highest point of visible internal jugular pulsation, and then place a ruler vertically at the sternal angle (angle of Louis). This is the site at which the manubrium attaches to the sternum

and is used to measure JVP because it is located 5 cm above the heart. Remember, the jugular venous system reflects right-sided heart activity. Place another ruler at the highest point of the venous wave horizontal to the ruler placed vertically. Measure the distance up from the chest wall. The normal JVP is < 3 cm. A central venous pressure can be estimated by adding 5 cm to the JVP. If venous pulsations are not visible, measure the JVP by identifying the highest point of external jugular vein distension.

Is It a Jugular Wave or a Carotid Arterial Wave?

- Carotid pulsation is normally palpable; jugular pulsation is not. Because jugular pulsation is a low-pressure wave, applying pressure can easily obliterate it.
- Carotid pulsation is not affected by position; jugular venous pulsation is.
- Carotid pulsation is unaffected by respirations; jugular venous pulsation is.
- Carotid pulsation has one positive wave; jugular venous pulsation has three positive waves (undulated).

INSPECTING THE CAROTID ARTERY AND JUGULAR VENOUS SYSTEM

1. Neck

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Differentiate carotid arteries and jugular veins.
- Carotid arteries and internal jugular veins run parallel to sternocleidomastoid muscle, the external jugular crosses the sternocleidomastoid muscle.*
- Carotids have visible pulsation, jugulars have undulated wave.*
- Carotids have palpable pulsations, jugulars easily obliterated.*
- Carotids not affected by respirations, jugulars are.*
- Carotids not affected by position, jugulars normally only visible when client is supine.*

INSPECTING THE CAROTID ARTERY AND JUGULAR VEIN SYSTEM

1. Neck

ABNORMAL FINDINGS/RATIONALE

- *Large, bounding visible pulsation in neck of at suprasternal notch:* HTN, aortic stenosis, or aneurysm.
- *Abnormal venous waveforms:*
 - Giant A waves.
 - Tricuspid stenosis, right ventricular hypertrophy, cor pulmonale.
 - *Absent A wave:* Atrial fibrillation.

2. Measuring Jugular Venous Pressure

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Position patient with the head of bed at 30 to 45-degree angle.
- Place a ruler vertically, perpendicular to the chest at the angle of Louis (sternal angle).



HELPFUL HINT

The angle of Louis, sternal angle, is located at the junction of the manubrium and the sternum where the second rib is attached to the sternum.

- Identify the highest level of the jugular vein pulsation; if unable to see pulsations, use the highest level of jugular vein distension.
- Place another ruler horizontally at the point of the highest level of the venous pulsation.
- Measure the distance up from the chest wall.
- *Positive carotid pulsations. JVP 2 cm at 45-degree angle.*
- *Jugular venous wave undulated, easily obliterated, varies with position change and respirations.*



Inspecting the neck



Measuring JVP

ABNORMAL FINDINGS/RATIONALE

- *Elevated JVP:* Right-sided CHF, constrictive pericarditis, tricuspid stenosis, or superior vena cava obstruction.
- *Low JVP:* Hypovolemia.

Precordium

With the patient supine, inspect the precordium with tangential lighting. Check for pulsations or abnormal movements, such as lifts and heaves or thrusts. Thrills are palpable vibrations created by turbulent blood flow. Lifts

or heaves are diffuse, lifting impulses. A thrust is a rocking movement. To facilitate the assessment, have the patient sit up and lean forward or turn to the left side. This brings the apex of the heart closer to the chest wall. (See *Inspecting the Precordium.*)

INSPECTING THE PRECORDIUM

1. Precordium

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Look for pulsations on the precordium, paying particular attention to the apex area.
- *Positive pulsation at apex. May note slight pulsations over base in thin adults and children.*



Inspecting the precordium

INSPECTING THE PRECORDIUM

1. Precordium (continued)

ABNORMAL FINDINGS/RATIONALE

- *Pulsations to right of sternum or at epigastric or sternoclavicular areas: Aortic aneurysm.*
- *Apical pulsation displaced toward axillary line: Left ventricular hypertrophy.*



HELPFUL HINT

You are more likely to see pulsations at the apex of the precordium than at the base, because the apex is the more active area, whereas the base is the quiet area.

Figure 14–18 illustrates abnormal jugular venous waves.

Palpation

Both the neck and the precordium are palpated for pulsations. Palpation is also useful in detecting abnormal pulsations or movements, such as thrills, lifts, and thrusts. Light palpation with your fingerpads is best for detecting pulsations. The balls of your hand are best for detecting vibratory sensations such as thrills.


Neck

Assess the carotid artery for rate, rhythm, amplitude, contour, symmetry, elasticity, and thrills.

- **Rate:** Rate is normally age dependent. Keep in mind that rates > 100 BPM or < 60 BPM do not always indicate pathology. Sympathetic nervous stimulation, such as fear, anxiety, and pain, may increase the rate. Vomiting or suctioning may cause vagal stimulation, which in turn may decrease the rate. Although activity and exercise often cause an increase in rate, well-trained athletes often have a slower rate with a larger stroke volume.
- **Rhythm:** Note if the rhythm is regular, fluctuating, or consistently irregular. Although a slightly irregular rhythm may occur with respirations, especially in children, irregular rhythms may also indicate an arrhythmia.
- **Amplitude:** Amplitude reflects the strength of the pulse. Amplitude is described as absent, weak, strong, or bounding or is graded on a scale of 0 to +3, with +2 as normal.
- **Contour:** Contour is the pressure wave created by the force of ventricular contraction. Both amplitude and contour are affected by the strength of ventricular contractions.
- **Symmetry:** The pulses should be equal.
- **Elasticity:** Because arteries are elastic, the carotid should be soft and pliable. Palpate the carotid medial to the trachea below the jaw angle. Do not palpate the carotids


simultaneously. This can obstruct blood flow to the brain and cause a syncopal attack. Also avoid excessive carotid stimulation, which may stimulate the vagus nerve, lowering the heart rate and BP and impairing cerebral circulation. If a thrill is detected, listen for a bruit.

Constrictive Pericarditis



Exaggerated "x" wave

Right Ventricular Hypertrophy




Exaggerated "a" waves

Atrial Fibrillation



No "a" waves

Tricuspid Regurgitation



Exaggerated V waves

FIGURE 14.18. Abnormal jugular venous waves.

PALPATING THE NECK

1. Neck

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

Palpating the Carotid

- Lightly palpate each carotid separately.
- Note rate, rhythm, amplitude, contour, symmetry, elasticity, thrills.



Palpating carotid artery

Palpating the Jugulars

- Palpate jugular veins and check direction of fill.
- *Occluding under the jaw, the jugular should flatten, but the wave form will become more prominent.*
- *Occluding above the clavicle, the jugular normally distends while the jugular wave diminishes.*



Assessing jugular flow. Compress jugular below jaw. Jugular vein collapses and jugular wave is more prominent at supraclavicular area.



Checking jugular fill. Compress jugular above clavicle. Jugular distends and jugular wave disappears.

- **Rate:** Depends on age.
- **Rhythm:** Regular.
- **Amplitude:** Strong, +2 or +3 pulses may normally be seen in high-output states such as exercise.
- **Contour:** Smooth upstroke with smooth, less acute descent.
- **Symmetry:** Pulses equal.
- **Elasticity:** Carotid soft and pliable.

Testing Abdominojugular (Hepatojugular) Reflux

- Position patient at 45-degree angle, place hands over the midabdominal area and apply 20 to 30 mm Hg of pressure for about 15 to 30 sec. Estimate the pressure by placing a partially inflated BP cuff on the abdomen under your hand.
- Look at the jugular veins while applying pressure, note increase vein distension, and return to normal upon release of pressure.
- *Negative abdominojugular reflux.*
- *Jugulars are easily obliterated and fill appropriately.*



Testing hepatojugular reflex

ABNORMAL FINDINGS/RATIONALE

- **Cardiac rates > 100 BPM:** Sinus tachycardia, supraventricular tachycardia (SVT), paroxysmal tachycardia (PAT), uncontrolled atrial fibrillation, ventricular tachycardia. Causes include CHF; drugs, such as atropine, nitrates, epinephrine, isoproterenol, nicotine and caffeine; hypercalcemia.
- **Cardiac rates < 60 BPM:** Sinus bradycardia heart block. Causes include MI; drugs such as digoxin, quinidine, procainamide, and beta-adrenergic inhibitors; hyperkalemia.
- **Irregular rhythm:** Arrhythmia.
- **Abnormal pulses:** See Figure 14.19 on page 477.
- **Unequal pulses:** Obstruction or occlusion.
- **Stiff, cordlike arteries:** Atherosclerosis.
- **Positive abdominojugular reflux:** Right-sided CHF, tricuspid regurgitation, tricuspid stenosis, constrictive pericarditis, cardiac tamponade, inferior vena cava obstruction, hypervolemia.

Precordium

The thicker the chest wall, the more difficult palpating pulsations becomes. Obesity, breast tissue, and increased musculature may interfere with palpation. Also, conditions such as COPD with overinflation of the lungs increase the distance between the heart and the chest wall, making palpation more difficult. Having the patient sit brings the heart more anterior, facilitating the palpation of impulses. If the apical impulse is not palpable, turn the patient to the left lateral recumbent position and palpate

again. You may also ask the patient to hold his or her breath to assist in palpation of pulsations.

Palpate the chest for pulsations, lifts, thrusts, and thrills. If you detect a thrill, listen for a murmur. Palpate the base right (aortic), base left (pulmonic), LLSB (tricuspid), and apex (mitral) areas. Be systematic in your approach. The pulsation felt at the apex originates from ventricular contraction and is referred to as the **point of maximum impulse (PMI)**, **left ventricular impulse (LVI)**, or **apical impulse**. This is the easiest pulsation to

PALPATING THE PRECORDIUM

1. Precordium

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Identify and palpate each cardiac site for pulsations, thrusts, heaves, and thrills:
- Apex (left ventricular area), or mitral area fifth intercostal space, midclavicular line.
- LLSB (tricuspid area), fourth to fifth intercostal space at left sternal border.
- Base left (pulmonic area), second intercostal space left sternal border.



Palpating apex of heart



Palpating LLSB



Palpating base left

- Base right (aortic area), second intercostal space right sternal border.
- Epigastric area below the xiphoid process.



Palpating base right



Palpating epigastric area

- Apex (left ventricular area):** PMI or LVI is 1–2 cm, amplitude small, duration nonsustained, systolic. Negative thrills. Amplitude may normally be increased in high-output states, such as exercise. Apical pulsation may not always be palpable. Left lateral displacement of PMI may occur during the last trimester of pregnancy.



HELPFUL HINT

When palpating the epigastric area, pulsations originating from the abdominal aorta are felt with the fingerpads; pulsations coming from the right ventricle are felt with the fingertips.

- LLSB:** May not be palpable, although small, nonsustained, systolic impulse may be palpated, especially in thin patients. Negative thrills.
- Epigastric area:** Positive slight pulsation may be normal, no diffusion. Palpations not palpable at base left, the pulmonic area, and base right, the aortic area, except in thin patients.

PALPATING THE PRECORDIUM

1. Precordium (continued)

ABNORMAL FINDINGS/RATIONALE

- *Enlargement and displacement of PMI to left midaxillary line:* Ventricular hypertrophy with dilation.
- *Apical impulse located on right side of precordium:* Dextrocardia, a heart located on the right side, often associated with congenital heart disease.
- *Enlarged apical pulsation without displacement > 2–2.5 cm with patient supine or > 3 cm with patient in left lateral recumbent position:* Ventricular enlargement, HTN, aortic stenosis.
- *Sustained pulsation:* Hypertrophy, HTN, overload, CMP.
- *Presystolic impulse:* May correlate with S₄ and be seen with aortic stenosis.
- *Early diastolic impulse:* May correlate with S₃ and be seen with CHF.
- *Diffuse, sustained impulse displaced downward and laterally:* Congestive CMP.
- *Thrills:* Murmur.
- *Right ventricular impulse with increased amplitude and duration:* Pulmonary stenosis or pulmonary HTN.
- *Palpable lifts or heaves:* Right ventricular hypertrophy.
- *Pulsations felt on the fingertips:* May come from the right ventricle, indicating right ventricular hypertrophy.
- *Large diffuse epigastric pulsation:* Abdominal aortic aneurysm.
- *Accentuated pulsation in pulmonic area:* Pulmonary HTN.
- *Accentuated pulsation in aortic area:* HTN or aneurysm.

palpate. Pulsations are least likely to be palpated at the base, the quieter area of the precordium, except in thin patients and children. Record the location, size, amplitude, and duration of any pulses you palpate.

When you palpate a pulse, record the following:

- **Location:** Also note any displacement.
- **Size:** Estimate in centimeters.
- **Amplitude:** Describe as small or increased.
- **Duration:** In heart rates less than 100 BPM, duration of the apical impulse is about one-third of the cardiac cycle.

The apical impulse should be palpable during systole. Impulses occurring during diastole may indicate an S₃ or S₄. The location, size, amplitude, and duration of the pulsation may be affected by hyperkinetic states that result in increased preload. (See *Palpating the Precordium*.)



HELPFUL HINT

One finger's width is about 1 cm. To determine the size of the pulsation, determine the number of finger pads you use to palpate it.

Precordial Impulses

IMPULSE	NORMAL	INCREASED STROKE VOLUME	INCREASED AFTERLOAD	INCREASED PRELOAD
Size	<ul style="list-style-type: none"> ■ Left: 2 cm ■ Right: Slight or nonpalpable 	<ul style="list-style-type: none"> ■ Increased ■ Palpable 	<ul style="list-style-type: none"> ■ Increased ■ Palpable 	<ul style="list-style-type: none"> ■ Increased ■ Palpable
Location	<ul style="list-style-type: none"> ■ Left: Apex ■ Right: LLSB 	<ul style="list-style-type: none"> ■ Apex ■ LLSB 	<ul style="list-style-type: none"> ■ Apex ■ LLSB 	<ul style="list-style-type: none"> ■ Displaced to left toward axilla ■ Displaced to left toward heart
Duration	<ul style="list-style-type: none"> ■ Left: Short ■ Right: Not discernible 	<ul style="list-style-type: none"> ■ Short ■ Short 	<ul style="list-style-type: none"> ■ Sustained ■ Sustained 	<ul style="list-style-type: none"> ■ Sustained ■ Sustained

(continued)

Precordial Impulses (continued)

IMPULSE	NORMAL	INCREASED STROKE VOLUME	INCREASED AFTERLOAD	INCREASED PRELOAD
Amplitude	<ul style="list-style-type: none"> ■ <i>Left</i>: Small ■ <i>Right</i>: Not discernible or nonpalpable 	<ul style="list-style-type: none"> ■ Increased ■ Small 	<ul style="list-style-type: none"> ■ Increased ■ Increased 	<ul style="list-style-type: none"> ■ Increased ■ Increased
Effect			<ul style="list-style-type: none"> ■ Increased pressure leads to hypertrophy 	<ul style="list-style-type: none"> ■ Increased volume leads to dilatation
Examples		<ul style="list-style-type: none"> ■ Noncardiac conditions, such as exercise, anxiety, anemia, hyperthyroidism 	<ul style="list-style-type: none"> ■ <i>Left</i>: HTN, aortic stenosis ■ <i>Right</i>: Pulmonary HTN, pulmonic stenosis 	<ul style="list-style-type: none"> ■ <i>Left</i>: Aortic or mitral regurgitation ■ <i>Right</i>: Atrial septal defect

The jugulars are further assessed by evaluating for **abdominojugular reflux**. Position the patient at about a 45-degree angle or less, which allows you to easily visualize jugular venous pulsations. Place your right hand over the midabdominal area and apply 20 to 30 mm Hg of pressure for about 15 to 30 seconds. Estimate this pressure by placing a partially inflated BP cuff between the abdomen and the right hand and applying sustained pressure until the desired pressure is reached. As you apply pressure, assess the jugular veins. A sustained increase in the JVP above 4 cm is considered a positive abdominojugular reflux. Valsalva's maneuver

can result in false positives, so warn the patient not to hold her or his breath or bear down during this maneuver.

Abnormal pulses are shown in Figure 14.19.

Percussion

Percussion of the precordium may be performed to determine cardiac size. However, other methods are more accurate. For example, palpation of the apical impulse, noting size and location, indirectly reflects cardiac size, and chest x-ray accurately determines cardiac size. Consequently, percussion has limited value in the cardiac assessment.

PERCUSSING THE PRECORDIUM

1. Precordium

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use indirect or mediate percussion to determine cardiac borders.



Percussing precordium

- *Dullness at third, fourth, and fifth intercostal spaces to left of sternum at midclavicular line.*

ABNORMAL FINDINGS/RATIONALE

- Left sternal border extends to midaxillary lines in an enlarged, dilated heart.




FIGURE 14.19. Abnormal pulses.

If you percuss the precordium, use mediate or indirect percussion. Begin at the anterior axillary line and percuss to the sternum at the fifth intercostal space. A resonant sound will change to dullness once you reach the left border of the heart. This should normally be at the fifth intercostal space at the midclavicular

line and should correspond with the PMI at the apex. Because the right border of the heart lies under the sternum, a change cannot be detected through percussion. Anteriorly, dullness is normally percussed over the third, fourth, and fifth intercostal spaces to the left of the sternum.

Auscultation

Auscultation is very useful in assessing heart sounds and vascular sounds. It is used extensively in the cardiovascular assessment to evaluate cardiac function. Auscultate the vascular structures of the neck and the precordium.

Neck

Auscultate the carotid artery and jugular veins for “flow” sounds. Assess the carotid for **bruits**, audible low-pitched sounds created by turbulent flow. Assess the jugular veins for a **venous hum**, a continuous low-pitched sound that reflects turbulent jugular venous flow and is often heard normally in children. These vascular sounds are low-pitched, so use the bell portion of the stethoscope.

HELPFUL HINTS

- Breath sounds over the trachea are loud, bronchial sounds, so ask your patient to hold his or her breath while you listen over the neck for vascular sounds.
- Turbulent flow in the heart is referred to as a **murmur**; turbulent flow outside of the heart is a **bruit**.

ALERT

The absence of a bruit is no guarantee that the carotid is patent. In fact, it may signal total occlusion of the artery.

Precordium

After establishing the cardiac rate and rhythm, carefully auscultate the precordium and identify normal and abnormal sounds. Listen to one sound at a time in each cardiac cycle at each auscultatory site. Closing your

eyes to block out other sensory stimuli will help you focus on the sounds. When auscultating, listen for the following specific characteristics that will help you differentiate and identify sounds: location, frequency, intensity, quality, duration, and timing in the cardiac cycle. Note if the sound is affected by breathing patterns or position changes. As you listen, integrate your auscultatory findings with the hemodynamic events that they represent.

Location. Specific sites have been identified for cardiac auscultation. Ask yourself, “At what site do I hear the sound best?”

HELPFUL HINT

Use this mnemonic to help you remember auscultatory sites:

- All (aortic)
- Patients (pulmonary)
- Take (tricuspid)
- Meds (mitral)

Frequency. The frequency of a sound relates to the pitch. Heart sounds and vascular sounds may be high, medium, or low in pitch, so you need a stethoscope that can detect high- and low-pitched sounds. Ask yourself, “What part of the stethoscope allows me to hear the sound best?” The diaphragm of the stethoscope is best for detecting high-pitched sounds; the bell is best for low-pitched sounds.

HELPFUL HINT

When using the diaphragm, use firm pressure; when using the bell, use light pressure.

AUSCULTATING THE VASCULAR STRUCTURES OF THE NECK	
1. Neck	
ASSESSMENT TECHNIQUES/NORMAL VARIATIONS	
<ul style="list-style-type: none"> ■ Have client hold breath. ■ Auscultate the carotid with the bell portion of the stethoscope for bruits. ■ Auscultate the jugulars with the bell portion of the stethoscope for venous hums. ■ <i>Negative bruits.</i> ■ <i>Positive carotid bruit may be normal in children and is associated with high-output states.</i> ■ <i>Negative venous hum.</i> ■ <i>Positive venous hum may be normal in children.</i> 	 <p>Auscultating carotid</p>
ABNORMAL FINDINGS/RATIONALE	
<ul style="list-style-type: none"> ■ Bruit suggests carotid stenosis. Murmurs can also radiate up to the neck from the heart, as in aortic stenosis. 	

Heart Sounds

HEART SOUND	LOCATION	PITCH	TIMING IN CARDIAC CYCLE	INTERPRETATION
S ₁	■ Apex/mitral area	■ High	■ Systolic	■ Normal
S ₁ Split	■ LLSB/tricuspid area	■ High	■ Systolic	■ Normal ■ RBBB
S ₂	■ Base right/aortic area	■ High	■ Systolic	■ Normal
S ₂ Split	■ Base left/pulmonic area	■ High	■ Systolic	■ Normal ■ RBBB, ASD, VSD, PE, pulmonic stenosis
S ₃ (Left Ventricular Origin)	■ Apex	■ Low	■ Diastolic	■ May be normal at age under 30 years ■ Left-sided CHF
S ₃ (Right Ventricular Origin)	■ LLSB	■ Low	■ Diastolic	■ Right-sided CHF
S ₄ (Left Ventricular Origin)	■ Apex, LLSB	■ Low	■ Diastolic	■ May be normal in children and young adults ■ MI, HTN, CAD
S ₄ (Right Ventricular Origin)		■ Low	■ Diastolic	■ Pulmonary HTN
Opening Snap	■ Apex	■ High	■ Diastolic	■ Mitral stenosis
Ejection Click	■ Base	■ High	■ Early systolic	■ Aortic stenosis, pulmonic stenosis
	■ Apex	■ High	■ Midsystolic	■ MVP
Friction Rub	■ LLSB	■ High	■ Systolic/diastolic or both	■ Pericarditis or pericardial effusion

Intensity. Intensity relates to loudness of a sound. Ask yourself, “How loud is the sound?” Intensity can be described as soft, loud, very loud, or on a scale of 1 to 4. A +1 sound is slightly louder than normal, +2 louder than normal, +3 very loud, and +4 the loudest. A grading scale can also be used to describe intensity of murmurs. Murmurs are classified on a 1 to 6 scale:

- **Grade 1:** Not immediately heard. You need to “tune in” for several seconds before sound can be heard.
- **Grade 2:** Quiet sound, but heard immediately if you are paying attention.
- **Grade 3:** Loud, readily heard.
- **Grade 4:** Loud, with a palpable thrust or thrill.
- **Grade 5:** Loud with palpable thrust or thrill with stethoscope tilted slightly off chest wall.
- **Grade 6:** Loud with palpable thrust or thrill with stethoscope off chest wall.

Although any cardiac sound can be graded, in clinical situations, the grading scale is most frequently used to grade murmurs. If using the grading scale, document

your finding as a fraction—for example, 3/6 or III/VI would represent a grade 3 sound on the 6-step scale.

Quality. Quality distinguishes one sound from another. Ask yourself, “What did it sound like?” Descriptive words such as snapping, blowing, and rumbling can be used to describe the quality of a sound.

Duration. Duration is the length of time the sound lasts. Ask yourself, “How long is the sound? Does it extend through systole, diastole, or both?” Duration is often described as short, medium, or long. Normal heart sounds are relatively short in duration as compared with murmurs, which are long-duration sounds.


Timing in the Cardiac Cycle. Timing in the cardiac cycle is one of the most important characteristics in distinguishing sounds. If you cannot determine where the sound is occurring in the cardiac cycle, you won’t be able to distinguish one sound from another. Ask yourself, “Does the sound occur during systole or diastole or throughout the entire cardiac cycle?” Here are some

Auscultating Valvular Heart Disease

DISEASE


DESCRIPTION/ASSESSMENT FINDINGS

Mitral regurgitation




- Holosystolic; extends from first heart sound to second heart sound.
- Loudness constant throughout systole.
- Heard best at apical area.
- May radiate to left lateral chest or, less commonly, to base.
- Blowing, rough (harsh), or musical.
- If very loud, it may mask first heart sound.

Mitral stenosis




- Occurs in diastolic phase of cardiac cycle.
- With normal sinus rhythm, characteristically a low-pitched rough or harsh murmur beginning immediately after opening snap.
- Extends to sharp first heart sound.
- Demonstrates presystolic (late diastolic) accentuation with normal sinus rhythm and just before termination.
- Presystolic accentuation of mid-late diastolic murmur disappears with atrial fibrillation.
- Best heard in apical area with patient in left lateral position and with bell portion of stethoscope.
- Localized to a very small (quarter-sized) area, so may be hard to find.

Aortic stenosis



- Midsystolic; characterized by crescendo-decrescendo pattern.
- Begins shortly after first heart sound and ends before second sound.
- Rough (low-pitched).
- Second heart sound at base markedly diminished or absent as a result of poor movement of severely calcified valve leaflets.
- When transmitted to apex, may have musical overtone in addition to rough quality (Gallavardin effect).
- If congenital (represented by two valve leaflets instead of three), early ejection click is often heard shortly after first heart sound.
- Both acquired and congenital murmurs heard best at base in second right interspace. May also be transmitted to apex and neck.

Aortic regurgitation



- Onset early in diastole.
- Decrescendo, ending before first heart sound.
- Usually blowing.
- Heard best at base of heart in third left interspace in parasternal line.
- Often transmitted toward apex, if loud enough.
- Loudness varies from grade 1 to grade 3 or 4.
- Often is only grade 1 intensity. If so, to facilitate identification of murmur, have patient sit forward, exhale, and hold his or her breath while you listen in third left interspace in parasternal line.

AUSCULTATING THE PRECORDIUM

1. Precordium

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Auscultate at apex.
- Note rate, rhythm, extra sounds, or murmurs.
- Auscultate at each site (apex, LLSB, Erb's point, base left and base right).
- Note S₁, S₂, extra sounds, or murmurs.
- Listen at each site with both the bell and the diaphragm.



Auscultation with bell of stethoscope



Auscultation with diaphragm of stethoscope



HELPFUL HINT

The diaphragm of the stethoscope is best for detecting high-pitched sounds; the bell is best for detecting low-pitched sounds. Use firm pressure with the diaphragm and light pressure with the bell.

2. Apex (Mitral)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Rate*: Depends on age.
- *Rhythm*: Regular; S₁ > S₂; high-pitched systolic, short duration. No extra sounds.
- *Physiological S₃ and S₄ may be heard in children and young adults without heart disease.*



Auscultating at the apex while timing S₁ with the carotid pulsation

ABNORMAL FINDINGS/RATIONALE

- *Bradycardia rates < 60 BPM or tachycardia rates > 100 BPM*: See variations in pulse in Palpating the Neck and Precordium, page 474 and 475–476.
- *Irregular rhythm*: Arrhythmia.
- *Accentuated S₁*: High-output states, mitral or tricuspid stenosis.
- *Diminished S₁*: First-degree heart block, CHF, CAD.
- *Variable S₁*: Atrial fibrillation.
- *S₃, low-pitched, early diastolic sound*: CHF.
- *S₄, low-pitched late-diastolic sound*: CAD, HTN, MI.
- Quadruple rhythm, S₃ + S₄ with fast rate is called a *summation gallop*.

(continued)

AUSCULTATING THE PRECORDIUM

3. LLSB (Tricuspid)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- $S_1 \geq S_2$, + split S_1



Auscultating at LLSB

ABNORMAL FINDINGS/RATIONALE

- *Wide split*: RBBB.
- *Midsystolic ejection click, a high-pitched systolic sound*: MVP.
- *Opening snap, a high-pitched diastolic sound*: Mitral or tricuspid stenosis, ventricular septal defect, patent ductus arteriosus.
- *Pericardial friction rub, a high-pitched systolic and diastolic sound*: Pericarditis or postoperative cardiac surgery.

4. Erb's Point

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *No aortic murmurs*

ABNORMAL FINDINGS/RATIONALE

- Aortic murmurs

5. Base

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Base Right (Aortic)

Base Left (Pulmonic)

- $S_1 < S_2$
- + split S_2 on inspiration at pulmonic area.



Auscultating at base left



Auscultating at base right

ABNORMAL FINDINGS/RATIONALE

- *Diminished S_2* : Incompetent aortic or pulmonic valves and low output states.
- *Ejection click, a high-pitched systolic sound*: Aortic or pulmonic stenosis.
- *Accentuated S_2* : Associated with HTN or pulmonary HTN.
- *Wide split S_2* : Occurs with RBBB, pulmonic stenosis, atrial septal defect, ventricular septal defect.
- *Fixed split S_2 , a split with no respiratory variation*: atrial septal defect, ventricular septal defect, CHF.
- *Paradoxical split S_2 , occurs during expirations*: Left bundle branch block or aortic stenosis.

AUSCULTATING THE PRECORDIUM

6. Murmurs

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Innocent grade 2/6, systolic, blowing murmurs often heard in children.*
- *Innocent systolic murmurs may also be heard during pregnancy.*

ABNORMAL FINDINGS/RATIONALE

- *Systolic and diastolic murmurs:* See Characteristics of Murmurs, on page 482–483.

ways to establish the timing of the sound in the cardiac cycle:

- **Heart rate:** The best way to identify the first and second heart sounds is to note the time interval between them. In heart rates less than 100 BPM, systole is shorter than diastole. The shorter interval between the sounds represents systole (the timing between S_1 and S_2), and the longer interval represents diastole (the timing between S_2 and the next S_1). With practice and experience, you will find this method invaluable and practically infallible.
- **Location:** Different heart sounds are best heard at specific sites. For example, S_1 is best heard at the apex, and S_2 is best heard at the base. Therefore, if you are listening at the apex, the louder of the two sounds should be S_1 ; if you are listening at the base, it should be S_2 .
- **Carotid pulsation:** The first heart sound will immediately precede the carotid pulsation.
- **Precordial movement:** Precordial apical movement is simultaneous with S_1 .
- **ECG monitor:** If your patient is on a cardiac monitor, the QRS complex is simultaneous with S_1 .

Auscultating Murmurs

Murmurs involving the mitral and aortic valves are the most common. They may indicate the presence of mitral regurgitation, mitral stenosis, aortic stenosis, or aortic regurgitation.

Figure 14.20 shows how to differentiate heart sounds.



HELPFUL HINTS

- It can be very confusing when extra sounds are added to the normal heart sounds. To help differentiate one sound from the other, first establish the timing of the sound in the cardiac cycle.
- Diastolic murmurs are usually not considered innocent. Remember, the diastolic phase is the resting phase, and if there is turbulent flow during the resting phase, there is usually a problem.
- Murmurs above grade 3/6 are usually associated with pathology. Remember, murmurs greater than grade 3/6 are associated with a thrill. If the turbulent flow is so great that it can be felt, there is usually a problem.
- Innocent or functional murmurs are often related to increased flow through normal valves.

Characteristics of Murmurs

TYPE	LOCATION	PITCH	QUALITY	INTERPRETATION
Early Systolic	■ LLSB	■ High/Low	■ Rough	■ Ventricular septal defect (muscular site)
Midsystolic	■ Base right ■ Base left	■ High	■ Harsh	■ Aortic stenosis ■ Pulmonic stenosis
Late Systolic	■ Apex ■ LLSB	■ High	■ Blowing	■ MVP ■ Tricuspid regurgitation
Holosystolic	■ Apex	■ High		■ Mitral regurgitation, VSD
Early Diastolic	■ LLSB ■ Erb's point ■ Base left	■ High	■ Blowing	■ Tricuspid regurgitation ■ Aortic regurgitation ■ Pulmonic regurgitation

(continued)

Characteristics of Murmurs (continued)				
TYPE	LOCATION	PITCH	QUALITY	INTERPRETATION
Mid-Diastolic	<ul style="list-style-type: none"> Apex LLSB 	Low	Rumbling	<ul style="list-style-type: none"> Mitral stenosis Tricuspid stenosis
Late Diastolic	<ul style="list-style-type: none"> Apex LLSB 	Low	Rumbling	<ul style="list-style-type: none"> Mitral stenosis Tricuspid stenosis
Pandiastolic	Base left	Low	Continuous	Patent ductus arteriosus



FIGURE 14.20. How to differentiate sounds.

- A. Split S_1 :** Systolic, high pitch heard best at tricuspid area with diaphragm; S_4 : Diastolic, low pitch heard best at apex with the bell.
- B. Split S_1 :** Systolic, high pitch heard best at tricuspid area with diaphragm; *Ejection sound*: Systolic (mid-late), high pitch heard best at base if aortic or pulmonic stenosis, with the diaphragm, or at the apex if MVP.
- C. Split S_2 :** Systolic, high pitch heard best at pulmonic area during inspiration with the diaphragm; S_3 : Diastolic, low pitch heard best at apex with the bell.
- D. Split S_2 :** Systolic, high pitch heard best at pulmonic area during inspiration with the diaphragm; *OS*: Diastolic, high pitch heard best at apex with the diaphragm.
- E. S_3 :** Diastolic, low pitch heard best at apex with the bell; *OS*: diastolic, high pitch heard best at apex with the diaphragm.

Questions on Characteristics of Murmurs

Location: Where did you hear the murmur best? Note the area.

Pitch: Did you hear the murmur with the bell or the diaphragm? Murmurs can be high, medium, or low pitched.

Intensity: How loud was the murmur? Grade the intensity on the 1 to 6 scale.

Quality: What did the murmur sound like? Was it blowing, harsh, mechanical, or rumbling?

Timing: Where does the murmur occur in the cardiac cycle? Murmurs can be systolic, diastolic, or both.

Duration: How long is the murmur? Murmurs by their very nature are long-duration sounds. However, some murmurs are longer than others. Some occur in parts of the cardiac cycle, whereas others occur throughout the cycle.

Radiation: Do you hear the murmur in any other places? Murmurs can radiate across the chest, through the back, and to the neck.

Position: Is the murmur heard best in a particular position? Some murmurs change with position.

Respiratory variations: Does the murmur change with breathing? Some murmurs are affected by respirations.

Shape or configuration: Does the intensity of the murmur change? Does the murmur start soft, then become louder? Start loud, then become softer? Is it consistent throughout? Describe the pattern of the intensity as crescendo, decrescendo, or crescendo-decrescendo?

Cardiac Auscultation Tips

- A quiet, warm, well-lit environment will minimize extraneous sound, facilitate cardiac inspection, and maximize patient comfort.
- Remember, you cannot auscultate heart sounds through layers of clothes. Because the patient will have to disrobe, ensure her or his privacy during the examination.
- Position changes can facilitate auscultation. The three positions for cardiac examination are supine, left lateral recumbent, and sitting. The left lateral recumbent position and sitting positions bring the apex of the heart closer to the chest wall, enhancing auscultation.
- Always work from the right side of the patient. This allows you to stretch the stethoscope across the chest wall and minimize interference from the tubing.
- Remember, a stethoscope basically transmits sound, so the better the instrument, the better your ability to hear sound.
- Be systematic in your approach. Listen with both the diaphragm and the bell. The diaphragm is best for detecting high-pitched sounds and is used by applying firm pressure. The bell is best for detecting low-pitched sounds and should be held lightly on the chest wall.
- Become a selective listener. Closing your eyes during auscultation allows you to focus your attention on listening to sound.
- Explain to your patient what you will be doing and that auscultation takes time. Teaching can be incorporated into the assessment process by sharing your findings with your patient.
- The thickness of the chest wall makes heart sounds more difficult to hear in obese and muscular patients than in patients with thin chests.
- Overinflated lungs increase the distance between the heart and the chest wall, also making auscultation more difficult. Patients with COPD often have distant heart sounds.



Case Study Findings

Your general health survey and head-to-toe assessment of Mr. Brusca have concluded. The following are your findings:

■ General Health Survey

- Well-developed, well-groomed 68-year-old white male, appears younger than stated age.

- Sits upright and relaxed during interview, answers questions appropriately.
- Alert and responsive without complaint, oriented x 4 (time, place, situation, and person).

(continued on page 486)



Case Study Findings (continued)

- Affect pleasant and appropriate.
 - Head-to-toe scan reveals positive arcus senilis, positive AV nicking and cotton wool, extremity changes including thin, shiny skin, thick nails, and edema.
 - **Vital Signs**
 - Temperature, 98°F
 - Pulse, 86 BPM, strong and regular.
 - Respirations, 18/min, unlabored.
 - BP, 150/90.
 - Height, 6'.
 - Weight, 275 lb.
- Cardiovascular assessment findings include:
- **Neck Vessels**
 - Positive large carotid pulsation, +3, symmetrical with smooth, sharp upstroke and rapid descent, artery stiff, negative for thrills and bruits.
 - JVP at 30 degrees <3 cm, negative abdominojugular reflux.
 - **Precordium**
 - Positive sustained pulsations displaced lateral to apex, PMI 3 cm with increased amplitude.
 - Slight pulsations also appreciated at LLSB and base, but not as pronounced.
 - Negative thrills; cardiac borders percussed third, fourth, and fifth intercostal spaces to the left of the midclavicular line.
 - Heart sounds appreciated with regular rate and rhythm at apex $S_1 > S_2$ and $+S_4$, at LLSB $S_1 > S_2$.
 - S_2 negative split, at base left $S_1 < 2$ negative split, at base right $S_1 < 2$ with an accentuated S_2 , negative for murmurs and rubs.



CRITICAL THINKING ACTIVITY #2

What effect would HTN have on Mr. Brusca's heart?



CRITICAL THINKING ACTIVITY #3

Considering the relationship of the cardiovascular system to the respiratory system, what respiratory problems might Mr. Brusca have as a result of his cardiovascular disease?



CRITICAL THINKING ACTIVITY #4

How would you explain the fact that the carotid pulse is palpable, but the jugular pulse is not?



CRITICAL THINKING ACTIVITY #5

List all of the strengths and areas of concern that would affect Mr. Brusca's health status at this time.



Case Study Analysis and Plan

Now that your examination of Mr. Brusca is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

After you have completed Mr. Brusca's assessment, you need to analyze the data, cluster the data, and formulate the nursing diagnoses. Some possible nursing diagnoses for Mr. Brusca are listed below. Cluster the data that support each diagnosis.

1. Ineffective Tissue Perfusion related to cardiovascular disease
2. Imbalanced Nutrition: more than body requirements
3. Risk for decreased Cardiac Output
4. Ineffective Health Maintenance related to lack of knowledge regarding pathology, complications, and management of hypertension
5. Knowledge deficit regarding complications and management of hypertension

Identify any additional nursing diagnoses.

Collaborative Nursing Diagnoses

1. Potential Complication: Hypertensive crises
2. Potential Complication: MI, stroke
3. Potential Complication: Adverse effects from medication



CRITICAL THINKING ACTIVITY #6

Now that you have identified some nursing diagnoses for Mr. Brusca, select one and develop a nursing care plan and a brief teaching plan for him, including learning objectives and teaching strategies.

RESEARCH TELLS US

C-reactive protein (CRP), an inflammatory biomarker, has been identified as a predictor of coronary heart disease (CHD) and peripheral arterial disease (PAD). A study of 3435 middle-aged white, German men from 45 to 74 years of age examined CRP as a risk predictor for CHD. CRP was measured in conjunction with the Framingham CHD prediction score, a global risk assessment for those prone to CHD (Koenig et al., 2004). Nonfatal and fatal coronary events were monitored over an average follow-up of 6.6 years. During that period, 191 coronary events occurred.

CRP independent of the Framingham Research Study (FRS) was found to be a significant predictor of coronary events ($p = .0002$). CRP added significant prognostic data to FRS for subjects in two intermediate risk categories ($p = .03$ and $p = .02$). The findings suggest that CRP enhances coronary risk assessment by the FRS.

The Ridker, Stampfer, and Rifai (2001) study of risk factors for systemic atherosclerosis compared the predictive value of 11 lipid and nonlipid biomarkers as risk factors for PAD. A nested case-control sample consisted of 14,916, healthy U.S. male physicians between the ages of 40 to 84 years. Over a 9-year follow-up period, 140 subjects developed symptomatic PAD. High-density lipoprotein cholesterol (HDL-C) ratio and CRP were found to be the strongest predictors of the development of PAD. CRP in conjunction with lipid screening significantly improved risk prediction of PAD ($p < .001$).

Nurses working in primary care are focused on prevention and early detection of health problems. CRP may improve predictability and prognostic data for CHD and PAD.

Identifying those at risk for CAD and PAD may lead to early interventions that may prevent the progression of the disease.

Source: Koenig, W., Loel, H., Baumert, J. and Meisinger, C. (2004). C-reactive protein modulates risk prediction based on the Framingham Score: Implications for future risk assessment: Results from a large cohort study in southern Germany. *Circulation* 109(11):1349–1353; and Ridker, P., Stampfer, M., Rifai, N. (2001). Novel risk factors for systemic atherosclerosis: A comparison of C-reactive protein, fibrinogen, homocystine, lipoprotein(a), and standard cholesterol screening as predictors of peripheral arterial disease. *JAMA* 285(19):2481–2485.

HEALTH CONCERNS

- Over 70 million Americans have one or more types of cardiovascular disease, with hypertension accounting for 65 million cases and coronary heart disease for 13 million cases.
 - In 2002, cardiovascular disease was identified as the primary cause of death for over 1.4 million people.
 - CHD is the single largest killer of Americans. Every 26 seconds an American suffers a coronary event.
- One person dies about every minute from CHD.
 - Approximately 400,000 to 460,000 sudden deaths from CAD occur in the emergency room or before the victim reaches the hospital.
 - The average age for the first heart attack is 65.8 years for men and 70.4 years for women.

Source: American Heart Association, 2005. <http://www.americanheart.org>

Common Abnormalities

ABNORMALITY/AREA

Angina Pectoris

- Chest pain resulting from myocardial ischemia.

ASSESSMENT FINDINGS

- Anxiety, chest pain.
- Skin pale, diaphoretic, cool, clammy.
- Dyspnea, tachycardia, pulsus alternans, arrhythmias, S₄, S₃.
- Nausea, belching.
- Weakness, paresthesias.

(continued)

Common Abnormalities (continued)

ABNORMALITY/AREA	ASSESSMENT FINDINGS
<p>Congestive Heart Failure</p> <ul style="list-style-type: none"> ■ Failure of the heart to pump sufficiently to meet the demands of the body. CHF can be right, left, or both. <p>Right-Sided Failure</p> <p>Left-Sided Failure</p> <p>Coronary Artery Disease</p> <ul style="list-style-type: none"> ■ A progressive narrowing of the coronary arteries. Atherosclerosis is the major cause of CAD. CAD can present as angina pectoris, acute MI, or sudden cardiac death. MI is necrosis of myocardial tissue from ischemia. <p>Pericarditis</p> <ul style="list-style-type: none"> ■ An inflammation of the visceral or parietal pericardium, resulting in cardiac compression, decreased ventricular filling and emptying, and cardiac failure. It often occurs 2 to 3 days after MI. 	<ul style="list-style-type: none"> ■ Fatigue, weight gain, confusion. ■ Skin pale, cool. ■ Neck vein distension. ■ Tachycardia, right ventricular heaves, murmurs, S₃, right-sided pleural effusion. ■ Anorexia, bloating, RUQ tenderness, hepatomegaly, ascites. ■ Edema, diminished hair growth. <ul style="list-style-type: none"> ■ Fatigue, confusion. ■ Skin pale, dusky, cyanotic, cool. ■ Left ventricular heaves, pulsus alternans, increased heart rate, displaced PMI, S₃, S₄, dyspnea, crackles, orthopnea, dry, hacking cough, PND. ■ Nocturia. <ul style="list-style-type: none"> ■ Anxiety, dizziness, chest pain, fatigue. ■ Skin pale to ashen, cool, diaphoretic, feverish. ■ Neck vein distension. ■ Dyspnea, tachypnea, crackles, tachycardia or bradycardia, arrhythmias, elevated BP initially, S₃, S₄, murmur, rubs, and diminished heart sounds. ■ Nausea, vomiting, low urinary output. ■ Cool, pale, decreased pulses. <ul style="list-style-type: none"> ■ Chest pain aggravated by inspiration, coughing, or movement. ■ Fever. ■ Friction rub at LLSB.

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the cardiovascular system.

Ask the patient the following:

- Do you have . . .
 - Chest pain?
 - Palpitations?
 - Syncope?
 - Edema?
 - Fatigue?
 - Extremity changes?
 - Dyspnea and cough?
- Are you having any chest discomfort? If yes, when did it start?
- What were you doing before the pain started? Did anything make it better or worse?

- Have you ever had this pain before? What does it feel like? Where does it hurt? On a scale from 1 to 10, with 10 being the worst, how severe is the pain?
- Do you have a history of cardiovascular disease? If yes, are you taking any medications? If yes, what are you taking and why?
- Do you have any other medical problems?
- Are you having any breathing difficulties?
- Do you have any allergies? If, yes, describe reaction.

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam:

Stethoscope with the ability to detect high- and low-pitched sounds

ASSESSMENT CHECKLIST (continued)

Sphygmomanometer
Marker
Ruler

Position

Supine, sitting, left lateral recumbent

Technique

First, scan your patient head to toe looking for signs related to the cardiovascular system.

Inspection, Palpation, Percussion, Auscultation

Neck

Inspect:

- Locate carotid and jugular pulsations and differentiate.
- Measure JVP.

Palpate:

- Carotid arteries
- Jugulars
- Test abdominojugular reflex

Percuss:

- Cardiac dullness

Auscultate (With Diaphragm and Bell)

- Bruits or venous hums

Precordium

Inspect:

- Apex for pulsations.
- Precordium for pulsations or movement.

Palpate:

- Locate PMI (LVI).
- Pulsations, lifts, heaves, thrills, thrills

Auscultate (With Diaphragm and Bell):

- Each auscultatory site

Apex

- Rate, rhythm, S₁, S₂, extra sounds

DOCUMENT

Document your findings and develop your plan of care.


S U M M A R Y

- A thorough cardiovascular assessment includes a health history and physical examination and provides invaluable data about the patient's overall health status.
- Before you begin, visualize the underlying structures and review expected normal findings. Understanding normal cardiovascular functioning is crucial to interpreting your findings.
- As you work through the assessment, systematically look for cardiovascular changes in every system.
- Let your patient's current health status direct your assessment. If she or he has an acute problem, perform a focused assessment.
- After you complete your assessment, document your findings. Your plan of care should flow from the assessment.
- Share your findings with your patient—this is an ideal teaching opportunity. Involving the patient in developing the plan of care allows him or her to actively participate in his or her healthcare and to make decisions that promote a healthy cardiovascular system.

R E V I E W Q U E S T I O N S

1. Tom Roberts, age 59, presents in the ER with chief complaint of "chest pain." Which description of chest pain would least likely be cardiac in origin?
 - a. Chest pain feels like a burning sensation.
 - b. Chest pain increases with palpation of the chest.
 - c. Chest pain radiates to the jaw.
 - d. Chest pain radiates to the left arm.
2. There are many different causes of chest pain. Which area should be addressed first when your patient has chest pain?
 - a. Respiratory
 - b. GI
 - c. Cardiac
 - d. Musculoskeletal

3. Part of a cardiac assessment includes palpating the chest for a PMI. Which cardiac site would be best for assessing the PMI?
 - a. Left base
 - b. Right base
 - c. Apex
 - d. LLSB
4. While performing a cardiovascular assessment, you palpate a thrill at base right. Considering this finding, what might you expect to hear upon cardiac auscultation?
 - a. Murmur
 - b. S₃
 - c. S₄
 - d. Ejection click
5. During cardiac auscultation, you hear a high-pitched, early diastolic sound at the apex. Identify the sound.
 - a. S₁
 - b. S₂
 - c. Ejection click
 - d. Opening snap
6. Your patient has HTN. While performing a cardiovascular assessment, you hear a low-pitch, late diastolic sound at the apex. Identify the sound.
 - a. S₁
 - b. S₂
 - c. S₃
 - d. S₄
7. Your patient has CHF. While performing a cardiovascular assessment, you hear a low-pitch, early diastolic sound at the apex. Identify the sound.
 - a. S₁
 - b. S₂
 - c. S₃
 - d. S₄
8. While performing a cardiovascular assessment, you palpate a carotid thrill. What sound would you expect to hear when auscultating the carotids?
 - a. Bruit
 - b. No sound
 - c. Murmur
 - d. Venous hum
9. Assessing the jugular venous system is part of the cardiovascular system. What patient position would best facilitate assessment of the jugular veins?
 - a. Supine
 - b. Standing
 - c. Sitting
 - d. Semi-Fowler's
10. While measuring your patient's JVP, you elevate your patient's head of the bed. What level is best for obtaining a JVP measurement?
 - a. 10 degrees
 - b. 45 degrees
 - c. 90 degrees
 - d. 180 degrees



Assessing the Peripheral-Vascular and Lymphatic Systems

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent peripheral-vascular/lymphatic history questions
- Obtain a peripheral-vascular/lymphatic history
- Perform a peripheral-vascular/lymphatic physical assessment
- Document peripheral-vascular/lymphatic findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

The **peripheral-vascular (PV) system** is a branching network of vessels that transports oxygenated blood to all body organs and tissues and then returns it to the heart for reoxygenation in the lungs. A disruption in the PV system can cause significant pain, loss of limb, or even death. The lymphatic system helps the heart and peripheral vasculature maintain adequate circulation. The function of the lymphatic system is to collect and drain excess tissue fluid that accumulates from the cardiovascular system and return this fluid to the heart. Lymphatic dysfunction may result in reduced range of motion, permanent disfigurement, and susceptibility to infection.

When assessing these two systems, your goal is first to identify any acute changes that require immediate intervention and then to identify any unhealthy behaviors that may affect the patient's health. A complete, accurate assessment is the first step in planning care, implementing therapeutic measures, and developing appropriate teaching strategies.

Anatomy and Physiology Review

Before you begin your assessment, you must understand how the PV and lymphatic systems work.

Peripheral-Vascular System

The primary function of the PV system is to deliver blood to all areas of the body. Blood flows through arteries, veins, and capillaries and is measured by pulse and pressure. Adequate perfusion to the vital organs and extremities is essential to life. Figure 15.1 and the box that follows it illustrate and describe the arterial system, and Figure 15.2 and the box that follows it illustrate and describe the venous system. (For more information on the function of arteries, veins, and capillaries, see Chapter 14, Assessing the Cardiovascular System.)

Lymphatic System

The **lymphatic system** complements the function of the vascular system. It is made up of lymphatic capillaries, lymph fluid, collecting ducts, and various tissues including the lymph nodes, spleen, thymus, tonsils, adenoids, and Peyer patches. The collecting ducts are thin, small, veinlike vessels that start at the capillaries and branch into their own circulation. Lymphatic vessels tend to lie near veins and drain into larger lymphatic vessels. **Lymph capillaries** drain the tissue spaces of lymph and tissue fluid. The lymph flows through small, oval bodies called **lymph nodes** before entering the bloodstream. As the lymph is returned to the central circulation, collec-




FIGURE 15.1. Peripheral arterial system. Pulse sites in red.

tions of lymph nodes filter large molecules and debris. The lymphatic system drains toward the center of the body. All lymphatic vessels eventually flow into two main vessels, the **thoracic duct** and the **right lymphatic duct**. The drainage point for the right upper body is a lymphatic trunk that empties into the right subclavian vein. The **thoracic duct**, the major vessel of the lymph system, drains lymph from the rest of the body into the left subclavian vein. It returns the various fluids and proteins to the cardiovascular system, forming a closed, porous circle.

Important functions of the lymphatic system include:

- Movement of lymph fluid in a closed circuit with the cardiovascular system.
- Development and maintenance of the immune system.
- Reabsorption of fat and fat-soluble substances from the small intestine.

Figure 15.3 and *Structures and Function of the Lymphatic System* that follows it illustrate and describe the lymphatic system.

Peripheral Arterial Pulse Sites and Areas Supplied

ARTERY	AREAS SUPPLIED
Carotid	
■ Internal	■ Branches into the anterior and middle cerebral arteries and supplies the brain.
■ External	■ Branches into the superior thyroid, inguinal, facial, occipital posterior auricular, superficial temporal, and maxillary arteries and supplies the thyroid, head, and mouth.
Brachial	■ Supplies the humerus and the muscles and skin of the upper arm.
Radial	■ Supplies the forearm and hand on the radial side.
Ulnar	■ Supplies the forearm and hand on the ulnar side.
Femoral	■ Supplies the thighs.
Popliteal	■ Supplies the knees; the posterior femoral, gastrocnemius, and soleus muscles; and the skin on the back of the leg.
Posterior Tibialis	■ Supplies the back of the leg and the ankle.
Dorsalis Pedis	■ Supplies the feet.




FIGURE 15.2. Peripheral venous system.

Peripheral Venous System and Areas Drained

VEIN	AREAS DRAINED
Jugular Veins	■ Head, face, neck, and brain
■ Internal	
■ External	
Deep Veins	■ Arms
■ Brachial	
■ Axillary	
■ Subclavian	
Superficial Veins	■ Elbow, forearm, and hand
■ Dorsal venous arch	
■ Basilic	
■ Cephalic	
■ Medial cubital	
Common Iliac Vein	■ Lower limbs, pelvis, and genitourinary structures
Inferior Vena Cava	■ Renal, gonadal, and hepatic veins
Portal Veins	■ Spleen and small and large intestines.
Azygos Vein	■ Chest and bronchi




FIGURE 15.3. Lymphatic system.

Interaction With Other Body Systems

The PV system has a complex mechanism to control blood flow to different parts of the body. There are three major types of control. First, local control of blood flow in each individual tissue is delivered in proportion to that tissue's need for blood perfusion. Second, nervous system control of the blood flow affects large segments of the systemic circulation, for example, by shifting blood flow from nonmuscular vascular beds to muscular beds during exercise or by altering blood flow in the skin to control body temperature. Last, the endocrine system, through hormones, ions, and other chemicals, can cause local increases, decreases, and generalized changes in tissue flow.

The Cardiovascular System

The lymphatic system has no pumping mechanism of its own, so it depends on the cardiovascular system, contraction of smooth muscles within lymphatic vessels, and skeletal muscle contraction to pump lymph throughout the body. Like the veins, the lymphatic vessels have one-way valves to prevent backflow. As lymph volume in-

creases, the lymph flows faster in response to an increasing capillary pressure and greater permeability of the capillary walls of the cardiovascular system. Increased metabolic activity and massage also contribute to lymph movement. Mechanical obstruction slows or stops the movement of lymph, dilating the system. If the flow of the lymphatic system is obstructed, lymph may diffuse into the vascular system or collateral channels may develop. For more information on how the PV and lymphatic systems interact with other systems, see Chapter 14, Assessing the Cardiovascular System.

Developmental, Cultural, and Ethnic Variations

Infants and Children

The lymphatic system begins developing at about 20 weeks' gestation. During the first few months after birth, an infant's ability to produce antibodies is still immature, so he or she is more vulnerable to infection and relies on the mother's immunity. Lymphoid tissue is abundant in infants, increases especially between ages 6 and 10, and then regresses to adult levels by puberty. The thymus grows rapidly from birth to age 2, and then slows down, reaching its greatest weight at puberty. Afterward, involution begins to occur until the thymus becomes a rudimentary organ in adulthood. The palatine tonsils are larger during early childhood than after puberty. The lymph nodes have the same distribution in children as in adults.

Like the cardiovascular system, the PV system becomes very similar to an adult's early in fetal life. The fetal circulation compensates for the nonfunctional fetal lung, with the right ventricle pumping blood through the patent ductus arteriosus rather than into the lungs. In the fetus, both ventricles are equal in weight because they both pump blood into the systemic circulation.

At birth, both the ductus arteriosus and the interatrial foramen ovale close, causing pressure to rise in the left atrium. Demand on the right ventricle changes as the pulmonary circulation becomes established. Demand also changes on the left ventricle, which now assumes total responsibility for the systemic circulation. These changes increase the size of the left ventricle, and by age 1 year, the relative size of both ventricles approximates the adult ratio of 2:1. The arterial pressure on the first day after birth averages about 70/50 mm Hg, increasing slowly over the next several months to about 90/60 mm Hg. Then it rises very slowly until the adult pressure of 120/80 mm Hg is reached at adolescence.

Pregnant Women

The immunologic processes that allow a woman to tolerate the implantation and development of a fetus are not fully understood. For unknown reasons, during the first

Structures and Functions of the Lymphatic System

STRUCTURE	FUNCTION
Lymph	<ul style="list-style-type: none"> Excessive fluid from body tissue that has drained into lymphatic capillaries. Responsible for 1%–3% of body weight. Contains leukocytes.
Lymphatic Capillaries	<ul style="list-style-type: none"> Collects fluid from interstitial space and surrounding tissue.
Lymphatic vessels	<ul style="list-style-type: none"> Formed by merging of lymphatic capillaries. Drain lymph back to right lymphatic duct and thoracic duct.
Lymph Nodes	<ul style="list-style-type: none"> Filter microorganisms and foreign substances from lymph.
<ul style="list-style-type: none"> Cervical Axillary Epitrochlear Intestinal Iliac Inguinal Lumbar 	<ul style="list-style-type: none"> Drains head, face, mouth, and neck. Drains arms, thoracic, breast, and upper abdominal wall. Drains hands and forearms. Drains abdominal viscera. Drains pelvic viscera. Drains legs, external genitalia, and lower abdominal wall. Drains pelvic viscera.
Tonsils	<ul style="list-style-type: none"> Lymphatic tissue that destroys microorganisms and foreign substances at beginning of digestive and respiratory tracts.
Thymus	<ul style="list-style-type: none"> Lymphatic tissue in thorax that forms antibodies in newborn and developing immune system and secretes thymosin, which helps with T-cell differentiation.
Spleen	<ul style="list-style-type: none"> Lymphatic tissue that filters blood and produces lymphocytes and monocytes.
Peyer's Patches	<ul style="list-style-type: none"> Clusters of lymphoid tissue found in small intestines. Respond to antigens by producing antibodies.

and second trimesters of pregnancy, the leukocyte count gradually increases and immunoglobulin G (IgG) concentrations decrease slightly. Maternal host defenses are altered because of reduced chemotaxis (the movement of neutrophils toward the infection site) and cause a delay in initial maternal responses to infection.

Systemic vascular resistance decreases with pregnancy, and peripheral vasodilatation occurs. This may lead to **palmar erythema**, a redness of the palms, and to **spider telangiectasis**, a spider-like image on the skin caused by a branched group of dilated capillary blood vessels. Blood pressure (BP) decreases during the second trimester and may rise thereafter to the prepregnancy level.



ALERT

Generally, an increase in BP of more than 30 mm Hg systolic or 15 mm Hg diastolic over prepregnancy levels is cause for concern and may indicate **preeclampsia**, a hypertensive condition of pregnancy involving many systems and affecting primarily the vasculature, kidneys, liver, and brain.


Older Adults

The number and size of lymph nodes decrease with advancing age, and some of the lymphoid elements are lost. Typically, the nodes of older patients are more likely to be fibrotic and fatty and result in impaired ability to resist infection. Degenerative changes in the vascular system occur as part of the normal aging process. Fibrosis causes increased thickness in the intimal wall. Wall stiffness is caused by an accumulation of collagen and calcium in the intima and media. Elastic fibers of the media become thin and calcified. These changes dramatically decrease the flexibility and elasticity of the vessels and increase PV resistance. Decreased venous elasticity increases the risk of varicosities, and increased PV resistance usually elevates BP. Throughout life, damage to the endothelium reduces its ability to protect the PV system and leads to the development of atherosclerotic plaques.

People of Different Cultures or Ethnic Groups

Although socioeconomic and environmental factors may play a part, certain diseases of the PV and lymphatic systems are more prevalent in certain cultural groups.

- Hypertension (HTN) is a major health problem for African Americans. There is also a high incidence of



Relationship of Lymphatic System to Other Systems

HTN in Puerto Ricans, Cubans, and Mexican Americans. One-sixth of the Iranian population has HTN, with stress as a major contributing factor.

- Navajo Native Americans have a high incidence of severe combined immunodeficiency syndrome (SCIDS), failure of antibody response, and cell-mediated immunity, unrelated to acquired immunodeficiency syndrome (AIDS).
- African Americans account for 13 percent of the population in the United States, but they account for 30 percent of AIDS cases.

Introducing the Case Study



Tasha Jordan is a 38-year-old African American woman who reports increasing pain and tenderness in her left leg since she delivered a premature baby 1 week ago. She describes the pain as dull and throbbing, and says that it gets worse when she stands and walks and gets better with rest and elevation. She also has noticed swelling in the leg. Aspirin has not relieved her symptoms. She has felt more tired than usual. Her past medical history is unremarkable. She is a smoker with a family history of cardiac disease. She had prolonged bedrest before delivery.

Performing the Peripheral-Vascular and Lymphatic Systems Assessment

Health History

Assessment of the PV and lymphatic systems includes a comprehensive health history that provides the framework for an accurate physical examination. If the patient's medical records are available, review them before starting the interview. Medical records can help you identify areas that need further exploration, but they are never a substitute for obtaining a health history directly from the patient. If time is an issue and you are unable to perform a complete health history, perform focused histories on the PV and lymphatic systems.

Biographical Data

The incidence of PV disease increases with age. Most patients are over age 45. The incidence of HTN also increases with age. Risk factors for vascular disease may differ according to sex and race. In men, half of all new cases occur in the fifth decade of life, but in women, the disease occurs much later. By the seventh decade, prevalence is similar in both sexes. White women have a slightly higher incidence of vascular disease than African American women. Infection in the lymphatic system can occur at any age.

Focused Peripheral-Vascular History

Take a focused history if your patient's condition or time restraints prohibit a detailed one. Arterial occlusion of an extremity can become a situation requiring rapid assessment, so be sure to ask the following questions:

- Have you noticed pain, pallor, pulselessness, polar sensation (coldness), paresthasias, or paralysis in an extremity (the "six Ps" of acute occlusion)?
- Do you have aching, heaviness, throbbing or burning pain, itching, or cramping in your legs?
- Are your ankles swollen? Is it difficult to fit into your shoes or wear your wedding band lately?
- Do you have leg pain when walking or at rest? What makes the leg pain better?
- Have you noticed any sores or ulcers on your feet or legs? How long have they been there? What have you used to treat them?
- Do you have a history of high BP, high cholesterol, cardiovascular or PV disease, or diabetes mellitus?
- Do you smoke? If so, how long and how much?

Current Health Status

Begin your assessment by asking for the person's chief complaint. Because the assessment of the PV and lymphatic systems covers a wide range of possible symptoms, you may need to ask which symptom is the most bothersome. Major vascular symptoms to watch for are swelling, limb pain, and changes in sensation of an

Focused Lymphatic History

Generally, there is no need for rapid assessment of this system. However, these questions will help you narrow your assessment focus:

- Have you noticed any swelling in your neck, armpits, or groin? If so, are the swollen areas sore, hard, or red? Do they appear on both sides of your body?
- Are you unusually tired? If so, are you tired all the time or only after exertion? Do you need frequent naps, or do you sleep an unusually long time at night?
- Have you had a fever recently? If so, how high was it? Was it constant or intermittent? Did it follow a pattern?
- Do you ever have joint pain? If so, which joints are affected? Does swelling, redness, or warmth accompany the pain? Do your bones ache?
- Have you noticed any sores that heal slowly?
- Do you have a history of blood transfusions?
- Have you ever been diagnosed with a chronic infection?

extremity. Always ask about fever, fatigue, and lumps when evaluating the lymphatic system.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Swelling

Swelling (edema) can be generalized or localized, acute or chronic. It occurs when fluid accumulates in tissue because of increased cellular permeability. This results in increased capillary pressure and can be caused by venous obstruction, heart failure, renal disease, lymphatic obstruction, decreased plasma proteins, inflammatory conditions, fluid and electrolyte imbalance (especially an excess of sodium), malnutrition, bacterial toxins, venoms, caustic substances, and histamine.

Limb Pain

Limb pain associated with vascular or lymphatic problems may be associated with a chronic condition or be a signal of a more serious acute problem that requires immediate medical attention. Pain associated with venous

or arterial vascular problems is often accompanied by other signs and symptoms.

Changes in Sensation

Paresthesia is a change in sensation such as numbness, tingling, pins and needles, or burning. It may be neurological in nature, but it can also be caused by vascular problems. Paresthesia can be acute or chronic, temporary or permanent.

Fatigue

Fatigue is the seventh most common complaint in primary care. Fatigue is a normal response to overexertion, but it can also be the presenting symptom of almost any disease. A thorough history and physical examination will establish the cause in about 85 percent of all patients.

Past Health History

The past health history can provide direct links to your patient's present health status. Look for data that may affect the PV and lymphatic systems.

Past Health History

RISK FACTORS/QUESTIONS TO ASK

Childhood Illnesses

- Did you ever have Hodgkin's lymphoma, acute lymphocytic leukemia, or sarcoma?

Surgery

- Have you ever had surgery? If so, when and what type?

Hospitalizations/Diagnostic Procedures

- Were you ever hospitalized for PV disease or lymphatic disorders?
- Did you receive a blood transfusion before 1985?

Serious Injuries

- Have you ever had trauma to your spleen?

Serious/Chronic Illnesses

- Do you have any other medical problems, such as cardiac or renal disease, cancer, hypertension, diabetes, AIDS, or autoimmune diseases such as systemic lupus erythematosus (SLE) or rheumatoid arthritis (RA)?

RATIONALE/SIGNIFICANCE

- These cancers frequently occur in children, who often receive bone marrow suppression drugs or radiation.

- May explain physical findings.
- Surgery exerts a negative effect on immune system. Splenectomy increases risk for disseminated infection. Mastectomy with lymph node removal can cause lymphedema.

- Use previous hospitalizations as a comparative.
- Blood was not routinely screened for human immunodeficiency virus (HIV) before 1985.
- Transfusions can also cause other infectious disorders such as hepatitis B.

- The spleen is very vulnerable to blunt abdominal trauma and may rupture.

- All these diseases may have PV or lymphatic symptoms.
- *Hypertension*: Doubles risk of symptomatic peripheral artery occlusive disease.

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Immunizations</p> <ul style="list-style-type: none"> ■ What immunizations have you had? Are they up to date? <p>Allergies</p> <ul style="list-style-type: none"> ■ Do you have allergies or asthma? <p>Medications</p> <ul style="list-style-type: none"> ■ Are you taking any medications? <ul style="list-style-type: none"> ■ Are you taking birth control pills? ■ Are you on chemotherapy? <p>Recent Travel/Military Service/Exposure to Infectious or Contagious Diseases</p> <ul style="list-style-type: none"> ■ Have you been in the military service? 	<ul style="list-style-type: none"> ■ <i>Diabetes mellitus</i>: Damages blood vessels and accelerates atherogenesis. ■ <i>Autoimmune diseases</i>: Increase risk for other diseases. <ul style="list-style-type: none"> ■ Up-to-date immunizations can prevent many communicable diseases. <ul style="list-style-type: none"> ■ Allergies reflect immune system function. ■ Asthma may indicate immunopathology. <ul style="list-style-type: none"> ■ <i>Antihypertensives, anticoagulants</i>: Determines compliance with treatment; therapeutic vs. toxic effects. ■ <i>Over-the-counter (OTC) medications</i>: May alter affects of prescribed medications. ■ <i>Steroids</i>: Cause immunosuppression. ■ <i>Birth control pills</i>: Increase risk for thrombus formation. ■ <i>Chemotherapy</i>: May suppress the immune system and lower platelet counts. ■ See <i>Drugs That Adversely Affect the Lymphatic System</i>. <ul style="list-style-type: none"> ■ Exposure to certain chemicals or toxins may affect the immune system.

Drugs That Adversely Affect the Lymphatic System

For information on drugs that affect the cardiovascular system, see Chapter 14, Assessing the Cardiovascular System.

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Antiarrhythmics	Procainamide Quinidine	Positive antinuclear antibody (ANA) titer; SLE-like syndrome Thrombocytopenia, hemolytic anemia, agranulocytosis
Anticonvulsants	Carbamazepine	Aplastic anemia (characterized by decreased levels of all formed elements of blood), leukopenia (decreased leukocyte levels), agranulocytosis (severely decreased agranulocyte levels), eosinophilia (increased eosinophil levels), leukocytosis (increased leukocyte levels), thrombocytopenia (decreased thrombocyte levels)
	Phenytoin	Thrombocytopenia, leukopenia, granulocytosis, pancytopenia (decreased levels of all blood cells), megaloblastic anemia (characterized by increased, immature, enlarged erythrocytes)
Antidiabetics	Acetohexamide, chlorpropamide, glipizide, glyburide, tolbutamide	Leukopenia, thrombocytopenia, pancytopenia, agranulocytosis, aplastic anemia, hemolytic anemia (characterized by premature erythrocyte destruction)

(continued)

Drugs That Adversely Affect the Lymphatic System (continued)		
DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Antihypertensives	Captopril	Neutropenia (decreased neutrophil levels), agranulocytosis
	Hydralazine	Positive ANA titer that occurs when the immune system creates antibodies against some of the body's own cells; SLE-like syndrome
	Methyldopa	Positive Coombs' test, indicating that some type of immunoglobulin coats the red blood cells
Anti-infectives	Cephalosporins	Positive Coombs' test, hypothermia (decreased thrombin levels), with or without bleeding
	Chloramphenicol	Bone marrow depression, pancytopenia, aplastic anemia
	Penicillins	Eosinophilia, hemolytic anemia, leukopenia, neutropenia, thrombocytopenia, positive Coombs' test
	Pentamidine isethionate Sulfonamides	Leukopenia, thrombocytopenia Granulocytopenia, leukopenia, eosinophilia, hemolytic anemia, aplastic anemia, thrombocytopenia, methemoglobinemia (increased levels of an oxidative compound of hemoglobin)
Antineoplastics	Busulfan	Severe leukopenia, anemia, severe thrombocytopenia
	Chlorambucil	Leukopenia, thrombocytopenia, anemia
	Cisplatin, cyclophosphamide, doxorubicin	Leukopenia, granulocytopenia, thrombocytopenia, anemia
	Methotrexate	Leukopenia, thrombocytopenia, anemia
	Mitomycin	Anemia
Antipsychotic agents	Chlorpromazine, thioridazine	Agranulocytosis, mild leukopenia
Gold compounds	Auranofin, gold sodium thiomalate	Leukopenia, thrombocytopenia, anemia, eosinophilia, aplastic anemia
Nonsteroidal anti-inflammatory agents	Ibuprofen	Neutropenia, agranulocytosis, aplastic anemia, thrombocytopenia, decreased hemoglobin and hematocrit
Miscellaneous agents	Furosemide	Anemia, leukopenia, thrombocytopenia
	Lithium	Leukocytosis

Family History

The family history identifies any causative factors that may lead to problems with the PV or lymphatic system.

The cause of lymphomas, lymphosarcomas, and cancer of the lymph glands is unknown.

Family History

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Familial/Genetically Linked PV Disorders <ul style="list-style-type: none"> Does anyone in your family have heart disease, arteriosclerosis, stroke, HTN, or hyperlipidemia? 	<ul style="list-style-type: none"> Increases risk for PV disease. Indicates need for risk factor modification.
Diabetes Mellitus <ul style="list-style-type: none"> Does anyone in your family have diabetes? If so, how old was she or he when it was diagnosed? 	<ul style="list-style-type: none"> Diabetics are more susceptible to PV disease.
Familial/Genetically Linked Lymphatic Disorders <ul style="list-style-type: none"> Is there any family history of lymphomas or leukemias? 	<ul style="list-style-type: none"> Positive family history increases risk for malignancies.

Review of Systems

The review of systems (ROS) assesses the interrelationship of the PV and lymphatic systems to every other

body system. Because signs and symptoms of lymphatic disorders are typically nonspecific, a complete ROS is an essential component of your assessment.

Review of Systems

SYSTEM/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- How have you been feeling?
- Have you had any changes in your energy level?
- Have you had any fevers or weight changes?

- *Fatigue/activity intolerance*: PV changes, infection.
- *Unexplained fever, night sweats, weight loss*: AIDS or cancer involving lymphatic system.

Integumentary

- Have you noticed changes in the color of your skin, hair, or nails? Hair loss?
- Ankle swelling or tight shoes?

- *Color changes/hair loss*: Chronic arterial or venous insufficiency.
- *Ankle swelling/tight shoes*: Fluid retention associated with PV disease or lymphadenopathy.

HEENT

- Do you have any lumps in your neck?
- Sore throat?
- Vision problems?

- *Neck swelling or masses*: Enlarged lymph nodes associated with infection or malignancy.
- *Sore throat/difficulty swallowing*: Bacterial or viral infection of lymphoid tissue.
- *Vision problems*: PV disease such as HTN.

Respiratory

- Do you have breathing problems?

- *Dyspnea with sputum production*: Respiratory infection such as pneumonia.
- *Dyspnea* is also associated with cardiovascular problems.

Cardiovascular

- Do you have cardiovascular problems?

- *Color changes in fingers and toes, leg pain, ankle swelling*: PV disease.
- *Chest pain/dyspnea*: Cardiac involvement.

Gastrointestinal

- Do you have gastrointestinal complaints? Bowel changes?

- *Diarrhea, abdominal tenderness and pain, weight loss*: Infections occurring with immunodeficiency disorders.

Genitourinary

- Do you have to urinate frequently, especially at night?

- *Nocturia*: CHF.

Reproductive

- Do you have reproductive/sexual performance problems?
- Do you practice safe sex?

- *Impotence*: PV disease.
- *Unprotected sexual activity*: Increased risk for HIV.

Musculoskeletal

- Do you have limb pain on activity?
- Limb tenderness?

- *Exercise-induced limb pain*: PAD (intermittent claudication).
- *Bone tenderness*: Leukemic or immunoproliferative disorder.

Neurological

- Do you have numbness or tingling anywhere in the body?

- *Paresthesia*: Arterial insufficiency.

Endocrine

- Do you have a history of diabetes or thyroid disease?

- *Diabetes* can cause vascular changes. Hypothyroidism can cause edema.

Psychosocial Profile

The psychosocial profile reveals patterns in the patient's life that may affect the health of her or his PV and lym-

phatic systems. It may also help determine the cause of a particular health concern. Assessing the lifestyle may identify specific health maintenance educational needs.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Health Practices and Beliefs/Self-Care Activities <ul style="list-style-type: none"> ■ Do you have regular health checkups? ■ Do you know the risk factors for PV disease (e.g., high-fat diet, protein-deficient diet, smoking, lack of exercise)? 	<ul style="list-style-type: none"> ■ May identify screening for risk factors and teaching needs.
Typical Day <ul style="list-style-type: none"> ■ Can you describe your typical day? 	<ul style="list-style-type: none"> ■ Fatigue and limb pain may limit ability to perform activities of daily living (ADLs).
Nutritional Patterns <ul style="list-style-type: none"> ■ What did you eat in the last 24 hours? ■ Has your weight changed recently? 	<ul style="list-style-type: none"> ■ Protein-deficient diets can cause edema. ■ High-fat diet is risk factor for PV disease. ■ Weight loss seen with cancer and AIDS; weight gain seen with edema. ■ Obesity is risk factor for cardiovascular disease.
Activity/Exercise Patterns <ul style="list-style-type: none"> ■ Do you exercise? What type of activity? 	<ul style="list-style-type: none"> ■ Exercise decreases risk for cardiovascular disease. ■ Arterial insufficiency can cause intermittent claudication with exercise.
Pets/Hobbies <ul style="list-style-type: none"> ■ Do you have any pets? ■ Do any of your hobbies involve using chemicals? 	<ul style="list-style-type: none"> ■ Pets and hobbies can enhance mental well-being if they do not compromise immunocompetence.
Sleep/Rest Patterns <ul style="list-style-type: none"> ■ How many hours of sleep do you get a night? 	<ul style="list-style-type: none"> ■ Pain can interrupt sleep.
Personal Habits <ul style="list-style-type: none"> ■ Do you or did you ever use tobacco? If so, in what form and how much? ■ Do you drink alcohol? ■ Do you use street drugs? 	<ul style="list-style-type: none"> ■ Smoking is a major risk factor for vascular disease. ■ Regular consumption of three or more drinks a day increases risk of HTN and decreases immunocompetence. ■ Intravenous drug use may transmit infectious diseases like hepatitis and HIV.
Occupational Health Patterns <ul style="list-style-type: none"> ■ What type of work do you do? ■ Are you exposed to chemicals or toxins? 	<ul style="list-style-type: none"> ■ Prolonged sitting or standing and exposure to temperature extremes may contribute to circulatory problems. ■ Chemicals or toxins may be carcinogenic and affect the immune system.
Environmental Health Patterns <ul style="list-style-type: none"> ■ Are you exposed to environmental chemicals? Are there any safety hazards in your household? 	<ul style="list-style-type: none"> ■ Chemicals may decrease immunocompetence. ■ Household hazards increases risk of falls. Protective shoes are essential for safety and proper healing of venous/arterial ulcers.
Roles/Relationships/Self-Concept <ul style="list-style-type: none"> ■ How do you see yourself? What is your role in the family? 	<ul style="list-style-type: none"> ■ Chronic vascular disease may affect role performance.
Religious/Spiritual/Cultural Influences <ul style="list-style-type: none"> ■ Do you have any religious practices or beliefs or cultural influences that may affect your healthcare practices? 	<ul style="list-style-type: none"> ■ It is important to identify any religious practices or beliefs and cultural influences that may affect the person's healthcare practices.

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Sexuality Patterns ■ Can you describe your sexual activity?	■ Multiple sex partners and anal sex are risk factors for HIV. ■ Impotence is associated with PVD.
Social Supports ■ Who are your supports?	■ Identifies supports within family and community. May indicate need for referral. ■ Limited mobility from PV disease may lead to social isolation.
Stress and Coping ■ How do you deal with stress?	■ Stress is risk factor for cardiovascular disease. ■ Stress decreases immunocompetence and increases BP.



Case Study Findings

Reviewing Mrs. Jordan's

Biographical data:

- 38-year-old African American female, married, one child.
- Works full-time as a secretary.
- Born and raised in the United States; Baptist religion.
- High school graduate, interested in attending community college when her baby gets older.
- Healthcare insurance, PPO, through husband's work plan.
- Husband is a firefighter.
- *Referral:* Private practice, internal medicine office.
- *Source:* Self, seems nervous but reliable.

Current health status:

- 6-day history of dull, aching pain and tenderness in left leg that is progressively getting worse.
- Pain is worse when standing and walking and is minimally relieved by rest and elevation.
- Aspirin has not helped.
- Denies chest pain or shortness of breath.

Past health history:

- Delivered premature baby 1 week ago.
- Prolonged bedrest (2 months) before delivery.
- No history of surgery. Hospitalized only for childbirth. Uncomplicated vaginal delivery; baby's gestational age is 31 weeks.
- Gained 50 lb during pregnancy.
- No history of chronic illness.
- Denies prescribed drugs.
- Has smoked 1 pack of cigarettes a day for 22 years (22 pack-years).

Family history:

- Mother, age 65, has HTN and is obese.
- Father, age 67, had myocardial infarction (MI) at age 50; quit smoking 17 years ago.
- All other known relatives alive and well.

Review of systems:

- *General Health Survey:* General health good, feeling tired since delivery of baby.
- *Integumentary:* Inflammation and swelling of left calf.
- *Head, Eyes, Ears, Nose, and Throat (HEENT):* No swelling or masses, no sore throat.
- *Respiratory:* No breathing problems.
- *Cardiovascular:* No chest pain.
- *Gastrointestinal:* No complaints, daily bowel movement.
- *Genitourinary:* No nocturia since delivery of baby.
- *Reproductive:* Has not been sexually active for about 3 months because of complications of pregnancy. No problems before this. Plans to bottle-feed baby.
- *Musculoskeletal:* Tenderness in left calf.
- *Neurological:* No numbness, tingling, or loss of sensation.
- *Endocrine:* No history of thyroid disease or diabetes.

Psychosocial profile:

- Has Pap smear by gynecologist every year. Last complete physical exam 5 years ago.
- Typical day consists of arising at 7 A.M., showering and dressing, and having breakfast with husband. Takes the train to work. Works 8 A.M. to 5 P.M. 5 days a week as a secretary,

(continued on page 504)



Case Study Findings (continued)

which involves mostly sitting at desk with computer. Spends evenings watching TV, reading, going out socially with husband. This routine was dramatically altered 2 months ago because of premature labor and bedrest.

- Usually eats lots of fast foods but worked hard to maintain a healthy diet during pregnancy. Gained 50 lb during pregnancy, which she attributes primarily to 2 months on bedrest. Would like to lose weight.
- Likes to walk and continued this activity throughout pregnancy until she was put on bedrest. Before pregnancy, played tennis once a week with husband. Understands need to resume previous exercise regimen but worries that new baby and current leg problem might prevent this.
- Normally sleeps about 8 hours nightly without difficulty. During pregnancy, sleep was interrupted because of need to void.
- No pets. Hobbies include walking, tennis, sewing, and reading.
- Has smoked 1 pack of filtered cigarettes a day for past 22 years (22 pack-years). Has tried behavior modification, cold turkey, and nicotine patches without success. Understands potential health effects of smoking on self and others and would like to quit. Drinks wine with dinner on weekends, but none since pregnancy. Denies recreational drug use.
- Sits for long hours in front of computer at work. Denies other occupational hazards.
- Lives in ranch home in suburbs with ample living space, air conditioning, and hot water heating. Denies travel outside state.
- Has supportive, caring husband who worries about premature baby who will be coming home from hospital shortly. Husband visits baby daily and is concerned about patient's ability to care for baby with her leg problem. Would like patient to quit smoking before baby comes home from hospital.
- Worries about caring for premature baby, especially in view of her own medical problems.
- Denies religious or cultural influences on health. Denies any sexual problems. Had sexual relations with husband until bedrest was ordered.
- Has no family close by but has many friends at work who will help out once baby comes home. Plans to return to work as soon as she can hire a reliable babysitter.
- Copes with stress by smoking and talking with husband and friends.



Case Study Evaluation

Before you proceed with the physical examination, document the PV and lymphatic history related to Mrs. Jordan.



CRITICAL THINKING ACTIVITY #1

Based on her health history, what circulatory problem do you think Mrs. Jordan might have? What factors contributed to this situation?



CRITICAL THINKING ACTIVITY #2

What family concerns will arise if Mrs. Jordan requires continued hospitalization?

Anatomical Landmarks

Before you can examine the PV and lymphatic systems effectively, review the location of pulse sites, and remember that the lymphatic system drains toward the center of the body (see Figs. 15.1 through 15.3).

Physical Assessment

Now that you have collected the subjective data, it is time to collect objective data during the physical examination. Keep key history findings—especially the chief complaint—in mind as you assess.

Approach

You may be able to combine the PV and cardiovascular physical assessments. However, if the patient has known or suspected PV disease or is at risk for circulatory complications, conduct the PV examination separately. The techniques used include inspection (of the skin for edema, ulcerations, and color or temperature changes), palpation (of the major pulse points of the body), and auscultation (of BP).

Physical assessment of the lymphatic system is usually integrated into the assessment of other body systems. For example, you can observe the tonsils along with the pharynx during the head and neck assessment, and evaluate the regional lymph nodes as you assess each body region. The techniques of inspection and palpation are used. A head-to-toe approach is suggested to incorporate these two systems. As you assess, be sure to compare the contralateral side of the body. Also ensure your patient's privacy and comfort.

**TOOLBOX**

The tools for examination of the PV and lymphatic systems include a stethoscope, sphygmomanometer, flashlight, ruler, and nonstretchable tape measure.

Performing a General Survey

Before examining specific areas, scan the patient from head to toe, looking for general appearance and signs of PV and lymphatic system dysfunction. For example:

- Be alert for signs of an acute illness, such as grimacing, pallor, or diaphoresis, or of a chronic problem, such as wasting or listlessness.
- Check appearance against chronological age. Does the person look older than his or her stated age? Chronic autoimmune diseases can “age” a person.
- Observe for edema, which is associated with PV or lymphatic disease.
- Observe posture. Abnormal posture, gait, and movements may signal an autoimmune disease such as rheumatoid arthritis.

- Check vital signs. Elevated temperature may indicate an infection; changes in BP and pulse may indicate a cardiovascular problem.

Performing a Head-to-Toe Physical Assessment

Next, look for changes in every system that might signal a PV or lymphatic system problem.

Performing a Physical Assessment of the Peripheral-Vascular and Lymphatic Systems

Now that you have looked at the relationship of the PV and lymphatic systems to all other systems, focus on the specifics of the examination. Begin with inspection of the veins and lymph nodes, and then proceed to palpation and auscultation. The patient should be sitting for most of the assessment and be supine only for palpation of the pulses of the lower extremities. Palpation of the lymph nodes and pulses is essential. For best results, use the pads of your index and middle fingers. Always palpate gently, beginning with light pressure and gradually increasing it.

Performing a Head-to-Toe Physical Assessment**SYSTEM/NORMAL FINDINGS****ABNORMAL FINDINGS/RATIONALE****General Survey***INSPECT*

- Awake, alert, and oriented to time, place, and person.
- Affect appropriate.

- Changes in mental status may indicate impaired cerebral circulation or an autoimmune disease such as SLE, which can cause mental status changes, including depression or even psychosis.

Integumentary*INSPECT/PALPATE***Skin**

- Skin color varies with client's ethnicity. Texture smooth and skin intact with no lesions.

- *Cyanosis*: Impaired circulation.
- *Pallor or rubor*: Vascular problems.
- *Thin, shiny skin*: Arterial insufficiency
- *Thick, leathery skin*: Venous insufficiency.
- *Rash*: Immune disorders such as the butterfly rash of SLE.
- *Poor wound healing*: Vascular insufficiency or immune deficiency.

Hair

- Even hair distribution.

- *Alopecia and short broken hair above forehead*: SLE.
- *Decreased hair on extremities*: Arterial insufficiency.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p>HEENT <i>INSPECT, PALPATE</i></p> <ul style="list-style-type: none"> ■ Oral and nasal mucosa pink, moist, and intact without lesions. ■ Tonsils pink; gum pink and intact with no bleeding. ■ No masses or nodes. ■ Thyroid not palpable. ■ Eye clear and bright, vision and hearing intact 	<ul style="list-style-type: none"> ■ <i>Pale nasal mucosa, boggy turbinates:</i> Chronic allergies. ■ <i>Enlarged, red, inflamed tonsils with exudates:</i> Infection. ■ <i>Oral lesions (e.g., lacy leukoplakia):</i> AIDS. ■ <i>Bleeding gums:</i> Leukemia. ■ <i>Enlarged nodes:</i> Infection or malignancy. ■ <i>Enlarged thyroid:</i> Chronic thyroiditis (Hashimoto's), an autoimmune disorder. ■ <i>Ptosis, extraocular movement deficits:</i> Myasthenia gravis. ■ <i>Decreased auditory function:</i> May occur with Hashimoto's.
<p>Respiratory <i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Lungs clear. 	<ul style="list-style-type: none"> ■ Adventitious sounds such as wheezes associated with asthma or allergic responses.
<p>Cardiovascular <i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Regular heart rate/rhythm. PMI 1 cm at apex. No extra sounds. 	<ul style="list-style-type: none"> ■ Enlarged, displaced PMI associated with ventricular hypertrophy such as occurs with HTN.
<p>Gastrointestinal <i>INSPECT/PALPATE/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Positive bowel sounds; abdomen soft, nontender, no organomegaly. 	<ul style="list-style-type: none"> ■ <i>Hyperactive bowel sounds:</i> Ulcerative colitis, an autoimmune disorder. ■ <i>Hypoactive bowel sounds:</i> Scleroderma, an autoimmune disorder. ■ <i>Enlarged liver:</i> Infection, cardiovascular problem, immune disorder, malignancy. ■ <i>Enlarged spleen:</i> Infection or cancer.
<p>Genitourinary <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ External genitalia pink, intact, no lesions 	<ul style="list-style-type: none"> ■ <i>Lesions:</i> Associated with STDs and may cause lymphadenopathy.
<p>Musculoskeletal <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ Full ROM in joints, no deformities or tenderness, +5 muscle strength 	<ul style="list-style-type: none"> ■ <i>Joint pain, deformity, limited ROM, muscle weakness:</i> PV disease multiple sclerosis, or autoimmune disorder such as rheumatoid arthritis.
<p>Neurological <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ Sensory intact, +2 DTR 	<ul style="list-style-type: none"> ■ <i>Decreased sensation:</i> Vascular insufficiency. ■ <i>Decreased DTR:</i> Peripheral neuropathies.

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> Fatigue Fevers Weight changes	General	<i>Inspect:</i> Signs of distress <i>Measure:</i> Vital signs Weight
Integumentary	<i>Ask about:</i> Skin lesions, rashes Hair loss Tight shoes	Integumentary	<i>Inspect:</i> Skin lesions Hair distribution Edema
HEENT	<i>Ask about:</i> Mouth lesions Bleeding gums Sore throat	HEENT	<i>Inspect:</i> Oral mucosa, gums, tonsils Nasal mucosa
Respiratory	<i>Ask about:</i> History of asthma, allergies	Respiratory	<i>Auscultate:</i> Adventitious breath sounds
Cardiovascular	<i>Ask about:</i> Chest pain History of murmurs	Cardiovascular	<i>Auscultate:</i> Heart sounds Murmurs, rubs
Gastrointestinal	<i>Ask about:</i> GI complaints History of abdominal trauma	Gastrointestinal	<i>Palpate:</i> Organs for enlargement
Genitourinary/Reproductive	<i>Ask about:</i> UTIs Safe sex practices STDs	Genitourinary/Reproductive	<i>Inspect:</i> Discharge STDs, lesions
Musculoskeletal	<i>Ask about:</i> Weakness Joint pain	Musculoskeletal	<i>Palpate:</i> Muscle strength Joints
Neurological	<i>Ask about:</i> Changes in mental status Loss of sensation	Neurological	<i>Test:</i> Mental status Sensation

Assessment of the Lymphatic System's Relationship to Other Systems

Inspection

Begin with inspection. Work from head to toe, inspecting the head and neck, upper extremities, abdomen, and

lower extremities. For information on inspecting the neck, see Chapter 11, Assessing the Head, Face, and Neck, and Chapter 14, Assessing the Cardiovascular System.

INSPECTING THE VEINS AND LYMPH NODES

1. Upper Extremities

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color, lesions, edema.
- Test capillary refill: press nail till it blanches; release and note time for color return.
- *Skin color uniform.*
- *Fingernails of equal thickness.*
- *Positive brisk capillary refill less than 2 to 3 seconds.*
- *No edema, erythema, red streaks, or skin lesions.*



Inspecting the hands



Checking capillary refill



Positive capillary refill

ABNORMAL FINDINGS/RATIONALE

- *Delayed capillary refill time:* Arterial occlusion, hypovolemic shock, hypothermia.
- Capillary refill time may also be delayed by environmental influences such as decreased ambient temperature, suggesting a problem that may not exist.
- *Edema:* Cellulitis, lymphedema, venous obstruction (thrombophlebitis)
- Lymphedema secondary to mastectomy.
- *Intermittent pallor and cyanosis of hands and fingers:* Episodic constriction of peripheral small arteries or arterioles caused by Raynaud's phenomenon or Raynaud's disease (a condition in which digital arteries respond excessively to vasospastic stimuli). Constriction causes hyperemia and rubor (redness).
- *Ischemic changes and gangrene of hands and fingers:* Buerger's disease (thromboangiitis obliterans).
- *Streaky redness, tenderness, warmth along course of a vein:* Thrombophlebitis.

2. Abdomen

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect arterial pulsations over the abdominal aorta.
- Note abdominal venous pattern and flow.
- *Abdominal contour flat, concave, or round.*
- *Abdominal veins barely visible. Veins flowing upward above and downward below umbilicus.*
- *Arterial pulsations at midline over aorta in thin adults.*



Inspecting the abdomen

ABNORMAL FINDINGS/RATIONALE

- *Tense, shiny abdominal skin:* Ascites or edema.
- *Upward or centrifugal venous flow:* Inferior vena cava obstruction or portal HTN.
- *Visible, large, diffuse pulsations:* Aneurysm (for more information, see Chapter 17, Assessing the Abdomen).

INSPECTING THE VEINS AND LYMPH NODES

3. Lower Extremities

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color, lesions, edema, hair distribution, and condition of nails.



Inspecting the legs

- *No varicosities or only superficial ones.*
- *No swelling or edema.*
- *No lesions or ulcers.*



Superficial varicosities

- *Leg hair distributed evenly. Older patients may have thinner, drier skin with less leg hair and altered pigmentation.*



HELPFUL HINT

Have patient stand when assessing for varicosities so that veins are more prominent.

3. Lower Extremities

ABNORMAL FINDINGS/RATIONALE

- *Hair loss, thin shiny skin, and thick nails:* Arterial insufficiency.
- *Eczema, stasis dermatitis:* Chronic venous insufficiency.
- *Prominent leg veins, possibly with ropelike, dilated appearance or purplish spider-like appearance:* Chronic venous insufficiency.
- *Edema:* Injury, cellulitis, venous/lymph obstruction, thrombophlebitis, varicosities.
- *Skin ulcers:* Trauma or venous/arterial insufficiency.



Lymphedema




Venous stasis ulcer



Arterial insufficiency

Assessing Edema

To assess edema, press your index finger over the bony prominence of the tibia or medial malleolus. Orthostatic (pitting) edema results in a depression that does not rapidly refill and resume its original contour. It is not usually accompanied by thickening or pigmentation of the overlying skin. The severity of edema can be graded on a scale of +1 to +4.



- +1: Slight pitting with about 2 mm depression that disappears rapidly. No visible distortion of extremity.
- +2: Deeper pitting with about 4 mm depression that disappears in 10 to 15 seconds. No visible distortion of extremity.
- +3: Depression of about 6 mm that lasts more than a minute. Dependent extremity looks swollen.
- +4: Very deep pitting with about 8 mm depression that lasts 2 to 3 minutes. Dependent extremity is grossly distorted.

Palpation

Certain parts of your hands and fingers are better than others for specific types of palpation. When evaluating pulses, use the distal pads of your second and third fingers. Palpate firmly but not so hard that you occlude the artery. If you have trouble finding a pulse, vary your pressure, feeling carefully at the correct anatomical location. Be sure you are not feeling your own pulse. Assess pulses for rate, rhythm, equality, amplitude, and elasticity. Pulse amplitude should be described on a

Measuring Pulse Amplitude

0 to 3 Scale

- 0 = Absent, not palpable
- 1 = Diminished
- 2 = Normal, expected
- 3 = Bounding

0 to 4 Scale

- 0 = Absent, not palpable
- 1 = Diminished, barely palpable
- 2 = Normal, expected
- 3 = Full, increased
- 4 = Bounding

PALPATING THE PULSES AND LYMPH NODES

1. Head and Neck Pulses

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note rate, rhythm, amplitude, contour, symmetry, thrills.
- *Temporal pulse:* Gently palpate next to eyes over temporal area.
- *Carotid pulse:* Gently palpate along sternocleidomastoid muscle; always palpate carotids separately.
- *Temporal and carotid arteries are regular, smooth, and +2 bilaterally.*



Palpating the temporal artery



Palpating the carotid artery

PALPATING THE PULSES AND LYMPH NODES

1. Head and Neck Pulses (continued)

ABNORMAL FINDINGS/RATIONALE

- *Alterations in pulse rate/rhythm:* Cardiac arrhythmia.
- *Absent pulse:* Arterial occlusion.

2. Neck Lymph Nodes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note size, shape, tenderness, mobility, consistency.
- Gently palpate lymph nodes of head and neck. (For more information on assessing lymph nodes in the head and neck, see Chapter 11, Assessing the Head, Face, and Neck.)
- *Cervical nodes are nonpalpable.*



Palpating the cervical lymph nodes

ABNORMAL FINDINGS/RATIONALE

- *Tender, palpable lymph nodes:* Recent infection.
- *Large, well-defined, tender nodes:* Acute infection.
- *Less-defined node borders:* Chronic infection.
- *Hard, enlarged, nontender, immobile nodes:* Malignancy.
- *Involvement of three or more node groups (generalized lymphadenopathy):* Autoimmune disease or neoplasm.

3. Upper Extremity Pulses

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Brachial pulse:* Gently palpate in the antecubital space along the medial side of the arm.
- *Radial pulse:* Run your fingers from the thumb to the wrist on the radial side and gently palpate pulse.
- *Ulnar pulse:* Run your fingers from the little finger to the wrist on the ulnar side and gently palpate pulse.
- *Skin temperature is warm bilaterally.*



Palpating the brachial pulse



Palpating the radial pulse



Palpating the ulnar pulse

- *Brachial, radial, ulnar pulses are easily palpated and equal in strength and amplitude bilaterally + 2.*
- *Ulnar artery may be difficult to palpate.*

(continued)

PALPATING THE PULSES AND LYMPH NODES

3. Upper Extremity Pulses (continued)

ABNORMAL FINDINGS/RATIONALE

- *Cool extremities*: Decreased circulation, vasoconstriction, Raynaud's disease, Buerger's disease, response to cold external temperature.
- *Absent pulse, pallor, pain, or paresthesias of distal upper extremities*: Arterial occlusion.
- *Unequal pulses*: Arterial narrowing or obstruction on one side.

4. Upper Extremity Lymph Nodes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Epitrochlear lymph nodes*: Gently palpate about ½ to 1 inch above elbow on the outer aspect of the arm.
- *Infraclavicular, axillary, and epitrochlear nodes are either nonpalpable or small, soft, mobile, and nontender*. (See Chapter 16, *Assessing the Breasts*.)



Palpating the epitrochlear nodes

ABNORMAL FINDINGS/RATIONALE

- *Extremity swelling, along with pallor or cyanosis and dilation of superficial veins*: Thrombosis of axillary or subclavian veins.
- *Diminished or absent pulses*: Arterial spasm.
- *Painful reddish streaks beneath skin; tender, enlarged nodes*: Inflamed lymphatic vessels caused by bacterial infection.

5. Lower Extremity Pulses

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Femoral Pulse

- Place patient supine and feel below the inguinal ligament, halfway between the symphysis pubis and the anterior-superior iliac spine.
- Palpate in the groin crease halfway between the symphysis pubis and the anterior-superior iliac spine.

HELPFUL HINT

- If you are unable to palpate a pulse, use a Doppler if available.



Palpating the femoral pulse

PALPATING THE PULSES AND LYMPH NODES

5. Lower Extremity Pulses (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Popliteal Pulse

- Flex the patient's knee so that his or her foot rests on the examination table.
- Place one hand on each side of the knee with your thumbs near the front of the patella.
- Curl your fingers around the knee and rest them in the popliteal fossa.



HELPFUL HINT

If the popliteal pulse is difficult to feel, try slightly straightening the patient's leg.



Palpating the popliteal pulse

Dorsalis Pedis

- Place your fingers between the patient's great toe and the first toe and slowly move away from the toes between the extensor tendons until you feel pulse.



HELPFUL HINT

- Plantarflexing the foot may make pedal pulse easier to palpate.
- Pedal pulse may be congenitally absent or branch high in the ankle.



Palpating the dorsalis pedis

Posterior Tibialis

- Place your fingertips in the groove between the medial malleolus and the Achilles tendon, and feel for pulse.



HELPFUL HINT

- Passive dorsiflexion of the foot makes the posterior tibial pulse easier to palpate.



Palpating the posterior tibialis.

- *Femoral, popliteal, posterior tibial, and dorsalis pedis pulses are easily palpated and bilaterally equal in strength and amplitude +2.*



Using a Doppler

(continued)

PALPATING THE PULSES AND LYMPH NODES

5. Lower Extremity Pulses (continued)

ABNORMAL FINDINGS/RATIONALE

- *Diminished or absent pulses:* Partial or complete arterial occlusion of proximal vessel, often caused by arteriosclerosis obliterans.
- *Absent posterior tibial pulse with signs and symptoms of arterial insufficiency:* Acute occlusion by thrombosis or embolus.
- *Increased, widened popliteal pulses:* Aneurysm.
- Popliteal occlusion associated with diabetes mellitus.

6. Skin Temperature

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Compare bilaterally, use light palpation and the dorsal portion of your hand to assess skin temperature, and lightly palpate texture.
- If edema present, grade degree of edema.
- *Temperature is cool to warm.*



ALERT

Sudden loss of peripheral pulse with cold, mottled extremity suggests arterial occlusion, a medical emergency.

- *Cold feet:* Arterial insufficiency, especially if unilateral.



Checking skin temperature

7. Calf Circumference

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Measure leg circumference to monitor edema at forefoot, above ankle, at calf, and at midhigh. Measure in centimeters.
- *Leg circumferences at forefoot, above ankle, at calf, and at midhigh are equal bilaterally.*



Measuring calf circumference

ABNORMAL FINDINGS/RATIONALE

- *Difference in leg circumference over 1 cm above ankle or 2 cm at calf:* Edema.

8. Calf Pain

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Support patient's leg while dorsiflexing foot.
- Note pain with dorsiflexion of foot.
- *No calf pain is present on dorsiflexion.*



Assessing for Homans' sign

PALPATING THE PULSES AND LYMPH NODES

8. Calf Pain (continued)

ABNORMAL FINDINGS/RATIONALE

- *Pain on dorsiflexion (Homans' sign)*: Deep vein thrombosis (in 50% of patients). Negative sign does not rule out deep vein thrombosis.

9. Inguinal Lymph Nodes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Inguinal lymph nodes*: Gently palpate horizontally in the groin and vertically in the inner aspect of the thigh.
- *Horizontal and vertical nodes are either nonpalpable or small (0.5–1 cm), soft, mobile, and nontender.*



Palpating the horizontal lymph nodes



Palpating the vertical lymph nodes

ABNORMAL FINDINGS/RATIONALE

- *Enlarged or tender nodes*: Current or recent infection in abdominal or genital area.
- *Difference in leg circumference*: Infection.
- *Nodes that run together (confluent)*: Chronic infection.
- *Unilateral discrete, nontender, firm or hard, fixed nodes*: Metastasized cancer.
- *Generalized lymphadenopathy (of three or more node groups)*: Autoimmune, infectious, or neoplastic disorder.

scale of 0 to 3 or 0 to 4. For more information, see Chapter 14, Assessing the Cardiovascular System.

To palpate lymph nodes, use the pads of your second, third, and fourth fingers, and gently palpate for superficial nodes. Be sure to note the size, shape, symmetry, consistency, delineation, mobility, tenderness, sensation, and condition of overlying skin. In areas in which the skin is more mobile, move it over the area of the nodes. Press gently at first, adding pressure gradually. If you find enlarged lymph nodes, explore adjacent areas and regions drained by those nodes for signs of possible infection or malignancy. The following section shows how to palpate the head and neck and upper and lower extremities. For more information on palpation, see Chapter 11, Assessing the Head, Face, and Neck, Chapter 16, Assessing the Breasts, and Chapter 17, Assessing the Abdomen.

Auscultation

Auscultate the pulses in the head, neck, upper extremities, and abdomen, and check BP.



HELPFUL HINTS

- If edema is unilateral, suspect occlusion of a major vein. If edema occurs without pitting, suspect arterial disease and occlusion. Arterial insufficiency may occur without edema.
- Because arteries are elastic, if you push them into soft tissue, you will have trouble palpating the pulse. So try to palpate arteries over bone.
- When assessing for edema, check the most dependent part of the body, so if the patient is supine, check the sacral area and scrotum in men.

AUSCULTATING THE PULSES

1. Head and Neck Pulses

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Auscultating bruits:* Listen with the bell portion of the stethoscope for bruits.
- *Auscultating carotid arteries:* Place bell of stethoscope over each artery and ask patient to hold her or his breath.
- *Heart sounds may normally be heard in neck. No other audible sounds when auscultating temporal and carotid arteries.*



Auscultating for bruits

ABNORMAL FINDINGS/RATIONALE

- *Soft, low-pitched, rushing sound during cardiac cycle:* Bruit in temporal or carotid artery signifies narrowing of artery.

2. Blood Pressure in Upper Extremities

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Refer to Chapter 3, Approach to the Physical Assessment and *Steps for Obtaining an Accurate Blood Pressure Reading*, for assessment techniques for measuring BP.
- *Systolic BP, < 120 mm Hg bilaterally in adults.*
- *Diastolic BP, < 80 mm Hg bilaterally in adults.*
- *Systolic BP dropping less than 20 mm Hg when patient rises from sitting position.*
- *Pulse pressure (difference between systolic and diastolic pressure) one-third of systolic pressure. BP rises slightly with age.*

ABNORMAL FINDINGS/RATIONALE

- *Below normal BP (hypotension):* Heart failure, dehydration, endocrine disorders (hypothyroidism), neurogenic vena cava obstruction, cardiac tamponade.
- *Decrease in systolic BP of 20 mm Hg and drop in diastolic BP on standing accompanied by rise in pulse rate:* Orthostatic hypotension caused by antihypertensive medications, volume depletion, peripheral neurovascular disease, or bedrest.
- *Three consecutive BP readings above*
A systolic reading 120 to 139 mm Hg and a diastolic reading 80 to 89 mm Hg is considered prehypertension; a systolic 140 to 159 with a diastolic 90 to 99, stage 1 HTN; a systolic \geq 160 with a diastolic \geq 100, stage 2 HTN.
- *Auscultatory gap (silent interval between systolic and diastolic BP):* Normal variation or sign of systolic HTN.
- *Korotkoff's sounds down to zero:* Cardiac valve replacement, hyperkinetic states, severe anemia, thyrotoxicosis, or following strenuous exercise.
- *Difference of over 10 mm Hg between arms:* Arterial compression on side of lower reading, aortic dissection, coarctation of aorta.
- *Widened pulse pressure with increased systolic BP:* Exercise, arteriosclerosis, severe anemia, thyrotoxicosis, increased intracranial pressure.
- *Narrowed pulse pressure with decreased systolic BP:* Shock, cardiac failure, pulmonary embolus.

3. Abdominal Aorta

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Auscultate in the epigastric area over the abdominal aorta for bruits with the bell. Can also auscultate the renal, iliac, and femoral arteries (see Chapter 17, Assessing the Abdomen, for abdominal vascular sites).
- *No vascular sounds heard on auscultation.*

ABNORMAL FINDINGS/RATIONALE

- Abdominal bruits reflect turbulent blood flow associated with partial occlusion.
- Bruits heard over epigastrium and radiating laterally, especially with HTN, suggest renal artery stenosis or aneurysm.
- Bruits heard in lower abdominal quadrants suggest partial occlusion of iliac arteries.

Steps for Obtaining an Accurate Blood Pressure Reading

- Tell patient to rest for 5 minutes before you take his or her BP and to wait 30 minutes if he or she has smoked or ingested caffeine.
- Have patient sit comfortably with arm resting on a flat surface at heart level.
- Palpate brachial artery. Center cuff bladder over this artery with bottom edge of cuff 1 inch above antecubital space.
- Estimate systolic BP by applying cuff and palpating radial pulse. Then inflate cuff until pulse disappears—this point is estimated systolic pressure. Deflate cuff and wait 1 minute before taking BP.
- Wrap cuff evenly and snugly around upper arm and place manometer at eye level. Place bell of stethoscope over brachial artery. Apply light pressure with good skin contact; heavy pressure may distort sounds. The bell transmits low-frequency sounds, such as the Korotkoff's sounds, better than the diaphragm.
- Close valve of pressure bulb clockwise until tight. Rapidly and steadily inflate cuff 30 mm Hg higher than patient's estimated systolic BP.
- Release valve at rate to 2 to 3 mm Hg/sec.
- Note point at which first sound is heard (phase I Korotkoff's sound). This is systolic pressure. Deflate cuff and note point at which sound disappears (phase V Korotkoff's sound).
- In adults, phase V is diastolic pressure.
- Listen 10 to 20 mm Hg below last sound and then deflate cuff rapidly and remove it from patient's arm. Wait 2 to 3 minutes before repeating procedure.

Additional Tests to Assess Peripheral-Vascular Flow

Allen Test

This test assesses arterial flow in the hands. Follow these steps:

- Have patient make a tight fist.
- Compress radial and ulnar arteries.
- Tell patient to open hand to a slightly flexed position.
- Observe for pallor.
- Release ulnar artery and watch for return of pink color in 3 to 5 seconds.



Compressing the radial and ulnar arteries



Observing for pallor



Releasing pressure and observing for return of normal color.

- Repeat procedure on radial artery.

Persistent pallor with the Allen's test suggests ulnar artery occlusion. Perform this test before obtaining an arterial blood specimen from the wrist. If both ulnar and radial arteries are not patent, the site cannot be used to obtain an arterial specimen.

Additional Tests to Assess Peripheral-Vascular Flow (continued)

Ankle-Brachial Index

This test assesses circulation to the feet. Measure ankle BP using a Doppler at the posterior tibialis or dorsalis pedis pulse site; then measure brachial BP. Normally, leg pressures are equal to or slightly greater than arm pressures. This test has limited use in diabetics with vascular calcification because the findings may be falsely high.

Calculate the ABI by:

Ankle systolic pressure divided by brachial systolic pressure = ABI

For example, if ankle pressure is 140 and brachial pressure is 110, ABI is 1.25.

ABI findings

- Normal = 1 or greater.
- Minimal disease = 0.8–0.95.
- Moderate disease = 0.4–0.8.
- Severe disease = 0–0.4.



Manual Compression Test

This test checks venous valve competence in patients with varicose veins. Have the patient stand and then compress first the distal portion of the vein and then the proximal portion. Normally, no backflow should occur, and you should not feel a wave with your distal hand. If the valves are incompetent, you will feel a wave with your lower hand as a result of backflow.

Trendelenburg Test

This test also evaluates venous valve competence. With the patient supine, elevate the leg to promote venous return. Then place a tourniquet around the thigh and have the patient stand. The veins should fill slowly from the lower leg up. After 30 seconds, remove the tourniquet. If the veins fill rapidly from the upper leg down, the saphenous valves are incompetent.

Color Change Test

This test evaluates arterial insufficiency. With the patient supine, elevate the legs to increase venous return, and then have the patient sit and dangle the legs. Note color changes. Normally, color returns to the feet within 10 seconds. Elevation pallor and dependent rubor are associated with arterial insufficiency.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Jordan have concluded. The following are your findings:

■ General Health Survey

- Well-developed 41-year-old female, appears stated age.
- Posture relaxed, sitting upright.
- Guarding left leg.
- Alert, cooperative, responding appropriately.
- Head-to-toe scan reveals decreased range of motion in left lower extremity, calf warm, tender, edematous. No other signs of problems in other systems/areas that would affect the peripheral-vascular/lymphatic systems.

■ Vital Signs

- Temperature, 99.8°F
- Pulse, 90 beats per minute (BPM), strong and regular.
- Respirations, 20/min, unlabored.
- BP, 140/70.
- Height, 5' 1".
- Weight, 160 lb.

■ Peripheral-Vascular and Lymphatic Systems

- Skin moist and of uniform color throughout except for slight cyanosis of left leg.
- Capillary refill, 2 seconds.
- Leg hair distributed evenly.

- No prominent leg veins or skin lesions.
- Left leg warm to touch, markedly tender on palpation, and edematous from popliteal space to ankle.
- Femoral and popliteal pulses +2 bilaterally; left posterior and pedal pulses difficult to locate.
- Calf circumference: right leg, 38 cm; left leg, 41 cm.
- Positive Homans' sign.
- No palpable lymph nodes.



Case Study Analysis and Plan

Now that your examination of Mrs. Jordan is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Now, from the data collected, develop nursing diagnoses for Mrs. Jordan. The following are some possible nursing diagnoses. Cluster the supporting data for each.

1. Chronic Pain related to interruption of venous blood flow and inflammation in left leg
2. Anxiety, related to unexpected hospitalization
3. Ineffective Role Performance related to being a new mother
4. Impaired Walking related to pain and edema in left leg
5. Health-Seeking Behaviors
6. Imbalanced Nutrition: more than body requirements, related to prolonged period of bedrest during pregnancy

Identify any additional nursing diagnoses and any collaborative nursing diagnoses.



CRITICAL THINKING ACTIVITY #3

Now that you have identified some nursing diagnoses for Mrs. Jordan, select one and develop a nursing care plan and a brief teaching plan for her, including learning objectives and teaching strategies.

RESEARCH TELLS US

Peripheral arterial disease (PAD) is the most common manifestation of systemic atherosclerosis, but it is also the most underdiagnosed and undertreated atherosclerotic disease. Leg pain complaints in older patients are often attributed to age or musculoskeletal problems. Histories routinely screen for cardiac disease (92%), but only 37 percent of histories screen for PAD (Mohler, 2003). Physical exam routinely includes heart and lung exam (95%), with 60 percent including pedal pulse assessment, and only 8 percent ABI assessment (Mohler, 2003).

Screening for PAD begins with a history and physical examination. Patients with PAD may experience intermittent claudication (characterized by cramping, tightness, tiredness, or aching of lower extremities during exercise/activity, relieved with rest) or may be asymptomatic. Because other conditions such as spinal stenosis may also result in lower extremity pain, a thorough analysis of the pain is important. A description of the pain may give clues as to arterial involvement, as calf pain may indicate involvement of the superficial femoral artery and hip, thigh, or buttock pain may indicate involvement of the aortoiliac artery.

The physical exam should include lower extremity pulse assessment, auscultation for bruits, and a skin assessment. Absence of the posterior tibialis pulse is always abnormal. Changes in skin associated with

PAD include dry, cool skin; atrophy and edema; absence of hair and perspiration; thick nails; and lesions such as ulcers or gangrene.

Although the history and physical are helpful in determining PAD, further evaluation is needed for patients with suspected PAD. The ABI is a noninvasive, inexpensive assessment test that correlates with angiographic findings. Normally, the systolic pressure in the lower extremities is slightly higher or equal to the arm pressures. The normal ABI is 1.0 to 1.3. However, patients with PAD will have lower pressures in the lower extremities than in the upper as a result of the occlusion, with an ABI <1.0. An ABI of <0.90 is diagnostic of PAD. The ABI can also be used to monitor progression of PAD. However, a major limitation of ABI is that diabetic patients with calcified tibial peroneal arteries may have falsely high ABI, thereby making the test unreliable among this patient population.

The presence of PV disease also reflects the severity of atherosclerosis in the vascular system and the presence of other vascular disorders such as CAD or cerebrovascular disease (CVD).

Because a low ABI correlates with greater risk of atherothrombotic events within the vascular system, screening for PAD including use of ABI should be routine in those at risk.

HEALTH CONCERNS

Facts About Peripheral Arterial Disease and Hypertension

- The “big-three” modifiable risk factors for coronary heart disease are cigarette smoking, dyslipidemia, and elevated BP.
- The prevalence of PAD increases with age, and rates are higher among men than among women.
- Rates of systolic BP are higher in older adults and in men.
- Other important risk factors include diabetes, obesity, and physical inactivity.
- Because each of the above risk factors is thought to influence atherosclerosis, they should be related to PAD as well as coronary heart disease (CHD).
- Cigarette smoking and diabetes appear to be the most important risk factors for PAD.

How to Manage Patients With Lymphedema

- Support and compress the affected limb, using multilayer bandaging, which gives a nonrigid outer casing of pressures up to 40 mm Hg.
- Use a combination of spiral and figure-8 bandage techniques to give an increased massage effect at rest and during exercise.
- Position wadding and foam of different densities along limb at pressure points to protect delicate areas, encourage fibrotic breakdown, and give an even compression once the entire bandage is in place.
- Change bandages daily because compression loosens with movement and bandages may slip.
- Custom-fitted pressure sleeves can also be used to control lymphedema.
- An inflatable sleeve (pneumatic pump) can also be used. An effective pump should have both gradient pressure (with the strongest pressure being distal [the hand] with less on the upper arm), and sequential pressure (in which pressure moves up the arm like a “milking technique”).
- Manual lymphatic drainage (complex decongestive physiotherapy) can be used to control lymphedema. This involves gentle stimulation of the skin and underlying lymphatic vessels by hand. Manual lymphatic drainage is usually performed by a physical, occupational, or massage therapist.

Do's and Don'ts of Preventing Lymphedema

Do's

- Do recognize early signs of infection: fever, redness, swelling, warmth, and tenderness in the arm at risk.
- Do moisturize skin.
- Do keep at-risk hand and arm extra clean. Use mild soap such as Dove.
- Do use rubber gloves when washing dishes or clothes.
- Do wear gloves when gardening.
- Do take frequent breaks when doing vigorous or repetitive activities, such as mopping, or when your arm feels tired, heavy, or achy.
- Do use potholders or oven mitts when handling hot foods.
- Do use an electric razor for shaving when shaving underarms.
- Do use insect repellent, such as Avon's “Skin So Soft.” Avoid using anything with alcohol in it because it dries the skin.
- Do apply an antibacterial ointment to any break in the skin, such as an insect bite or torn cuticle.
- Do use sunscreen that is at least SPF 15.
- Do use a thimble when sewing.
- Do elevate and rest your arm.
- Do control blood sugars if you have diabetes.
- Do wear a compression bandage or sleeve when flying in planes if you have any arm swelling.

Don'ts

- Don't bathe with hot water.
- Don't go from extreme temperatures (hot and cold).
- Don't use hot tubs, saunas, or steam baths.
- Don't use heating pads or hot compresses on the arm, neck, and back of affected side.
- Don't allow heat-producing treatment or vigorous massage of the affected side.
- Don't carry heavy objects on the affected side.
- Don't carry heavy shoulder bags on affected side.
- Don't wear tight or restrictive clothes.
- Don't wear watches or jewelry on affected arm.
- Don't wear a heavy breast prosthesis.
- Don't drink alcohol excessively.
- Don't smoke.
- Don't cut nails or cuticles.
- Don't use affected arm for BP readings.
- Don't allow any piercing of the affected arm, such as injections or blood drawing.

Common Abnormalities

ABNORMALITY/AREA

ASSESSMENT FINDINGS

Acute Lymphangitis

An inflammation of one or more lymphatic vessels. The cause is usually a bacterial infection that has spread from its initial site (e.g., skin cut) to nearby lymph glands. Area affected depends on site of inflammation.

- Chills and fever.
- Swelling of lymph glands.
- Red lines on arms or legs if lymphatics are inflamed, indicating concentration of inflammation along lymph channels.
- Inflammation may be slightly indurated and palpable to gentle touch.
- Infection may occur distal to inflammation, particularly interdigitally.

Aneurysm

An abnormal dilation of a blood vessel, commonly at the site of a weakness or tear in the vessel wall. The following are possible areas affected:

- Thoracic aorta

- Pain in back, neck, and substernal areas.
- Dyspnea, stridor, or cough if pressing on trachea.
- Asymptomatic—first sign may be rupture.
- Hoarseness and dysphagia if pressing on esophagus or laryngeal nerve.
- Pulsating abdominal mass.
- Mild midabdominal or lumbar pain to severe abdominal and back pain.
- Peripheral emboli.
- Cool, cyanotic extremities if renal, iliac, or mesenteric arteries are involved.
- Possible claudication, depending on which vessels are affected.
- Abrupt, extreme, tearing pain in area of aneurysm.
- BP markedly elevated initially; then drops.
- Radial pulses may be unobtainable as disorder progresses.
- Syncope, hemiplegia, or paralysis of lower extremities.
- Possible heart failure.

- Abdominal aorta

- Dissecting aorta

Arterial Embolism

Occlusion of blood flow by a foreign object or blood clot within the vessel. Area affected varies because embolism moves.

- Sudden or insidious pain in extremity.
- Numbness in extremity.
- Coldness of extremity.
- Tingling in extremity.
- Pulselessness distal to blockage.
- Pallor or mottling of extremity.
- Muscle weakness.
- Muscle spasm.
- Paralysis.
- A line of demarcation, with pallor, cyanosis, and cooler skin distal to blockage.
- Pain in region of affected vessel.
- Numbness in affected vessel.

(continued)

Common Abnormalities (continued)

ABNORMALITY/AREA	ASSESSMENT FINDINGS
<p>Arterial Thrombosis (Blood Clot) An aggregation of blood and other cells that adhere to the vessel wall. Area affected depends on site of blockage.</p>	<ul style="list-style-type: none"> ■ Pallor or mottling of skin in affected area. ■ Muscle spasms. ■ Pulselessness distal to blockage. ■ Possible paralysis.
<p>Buerger's Disease A vasculitis confined to the intermediate and small arteries and veins of the extremities, which are the areas affected.</p>	<ul style="list-style-type: none"> ■ Claudication. ■ Migrating thrombophlebitis. ■ Numbness, pain, and ulceration of toes. ■ Pain in arch of foot that may occur at rest. ■ Paresthesias. ■ Sensitivity to cold. ■ Diminished peripheral pulses. ■ Stasis dermatitis and trophic nail changes. ■ Ulceration and gangrene. ■ Nicotine use is strong risk factor.
<p>Chronic Arterial Insufficiency Reduced blood flow to the lower extremities (area affected), usually caused by an atherosclerotic process.</p>	<ul style="list-style-type: none"> ■ May be asymptomatic or present as intermittent claudication. ■ Pulses decreased or absent. ■ Pale color, especially when elevated; dusky red when dependent. ■ Temperature of extremity is cool. ■ Edema may be absent or mild. ■ Skin is thin, shiny, and atrophic. ■ Loss of hair is seen over foot and toes. ■ Nails are thickened and ridged. ■ Ulcers, if present, involve toes or points of trauma on feet. ■ Skin around ulcer has no excess pigment. ■ Pain is often severe unless neuropathy masks it. ■ Gangrene may develop.
<p>Chronic Venous Insufficiency Stasis of blood in lower extremities (area affected), resulting from chronic lower leg edema caused by thrombophlebitis or valvular incompetence.</p>	<ul style="list-style-type: none"> ■ Prominent leg veins, may appear ropelike and dilated or purplish and spider-like. ■ Lower leg edema may extend to knee of affected extremity. ■ Affected leg hard and leathery to touch. ■ Pulses normal, but may be difficult to feel through edema. ■ Normal temperature. ■ Brownish skin pigmentation. ■ Gangrene does not occur. ■ Ulcers may occur at sides of ankles. ■ Skin surrounding ulcers pigmented and sometimes fibrotic. ■ Pain is not severe. ■ Eczema or stasis dermatitis.
<p>Deep Vein Thrombosis A thrombus formation that obstructs venous flow. Area affected may be anywhere in the body.</p>	<ul style="list-style-type: none"> ■ Local tenderness and pain. ■ Increased skin turgor. ■ Swelling.

ABNORMALITY/AREA	ASSESSMENT FINDINGS
<p>Lymphedema A blockage or dysfunction of lymph vessels or nodes. It is often associated with lymphatic disorders following surgery for breast cancer or prostate cancer. Congenital lymphedema (Meige-Milroy's disease) is chronic edema of the lower extremities, resulting in swelling and often grotesque distortion. Areas affected are extremities.</p> <p>Peripheral Atherosclerotic Disease (Arteriosclerosis Obliterans) Occlusion of the blood supply by atherosclerotic plaque. The following areas are affected:</p> <ul style="list-style-type: none"> ■ Lower extremities. ■ Upper extremities. 	<ul style="list-style-type: none"> ■ Increased skin temperature. ■ Possible positive Homans' sign. ■ Swelling of arms or legs. ■ Overlying skin eventually thickens and feels tougher than usual. ■ Usually painless. ■ Weak or absent peripheral pulses. ■ Color changes associated with position changes (blanching when raised above heart level; rubor when dependent). ■ Filling of dorsal veins. ■ Ulceration and gangrene. ■ Aching, fatigue, weakness in upper arm. ■ Unilateral coolness. ■ Pallor. ■ Weak or absent radial or ulnar pulses. ■ Unilateral and focal filling abnormalities.
<p>Raynaud's Phenomenon/Raynaud's Disease Involves enhanced response to cold or stress. Idiopathic, intermittent spasm of the arterioles in the digits (Raynaud's disease) occurs most frequently in young, otherwise healthy women. Raynaud's phenomenon is secondary to connective tissue diseases, neurogenic lesions, and trauma. Areas affected are hands and fingers.</p>	<ul style="list-style-type: none"> ■ Pallor of fingertips with cold exposure, followed by cyanosis, then redness and pain with warming. ■ Ulceration of finger pads, progressing to autoamputation in severe, prolonged cases. ■ Raynaud's phenomenon is almost always bilateral. ■ Physical exam is normal when patient is not having symptoms. ■ May see pitting digital scars.
<p>Thrombophlebitis Inflammation of a vein associated with thrombus formation. Area affected may be anywhere in the body.</p>	<ul style="list-style-type: none"> ■ Redness and cyanosis of affected extremity. ■ Warmth and tenderness to touch. ■ Increased size of affected extremity caused by swelling. ■ Palpable cordlike structures (if superficial). ■ Possible positive Homans' sign. ■ Decreased or absent pedal pulse on affected extremity. ■ Slowed capillary refill.
<p>Varicose Veins Irregular, tortuous veins with incompetent valves. Areas affected are lower extremities.</p>	<ul style="list-style-type: none"> ■ Severe, aching pain in leg. ■ Leg fatigue, heaviness. ■ Itching over affected leg (stasis dermatitis). ■ Feelings of heat in leg. ■ Visibly dilated veins.

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the peripheral-vascular and lymphatic systems. Ask the patient the following:

- Do you have:
 - Swelling?
 - Limb pain?
 - Changes in sensation?
 - Fatigue?
- Have you noticed pain, pallor, pulselessness, polar sensation (coldness), paresthesias, or paralysis in an extremity (the “six Ps” of acute occlusion)?
- Do you have aching, heaviness, throbbing or burning pain, itching, or cramping in your legs?
- Are your ankles swollen? Is it difficult to fit into your shoes or wear your wedding band lately?
- Do you have leg pain when walking or at rest? What makes the leg pain better?
- Have you noticed any sores or ulcers on your feet or legs? How long have they been there? What have you used to treat them?
- Do you have a history of high BP, high cholesterol, cardiovascular or PV disease, or diabetes mellitus?
- Do you smoke? If so, how long and how much?
- Have you noticed any swelling in your neck, armpits, or groin? If so, are the swollen areas sore, hard, or red? Do they appear on both sides of your body?
- Are you unusually tired? If so, are you tired all the time or only after exertion? Do you need frequent naps, or do you sleep an unusually long time at night?
- Have you had a fever recently? If so, how high was it? Was it constant or intermittent? Did it follow a pattern?
- Do you ever have joint pain? If so, which joints are affected? Do swelling, redness, or warmth accompany the pain? Do your bones ache?
- Have you noticed any sores that heal slowly?
- Do you have a history of blood transfusions?
- Have you ever been diagnosed with a chronic infection?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam.

Stethoscope with the ability to detect high and low pitch sounds

Sphygmomanometer
Marker
Ruler

Position

Supine, sitting, standing

Technique

Remember to scan your patient from head-to-toe looking for signs related to the PV/lymphatic systems.

Inspection, Palpation, Auscultation

Inspect

Upper Extremities

- Color, condition of skin
- Capillary refill

Abdomen

- Shape
- Pulsations
- Veins

Lower Extremities

- Color, condition of skin, hair growth
- Veins
- Edema

Palpate

Pulses: Note rate, rhythm, contour, elasticity, equality amplitude

- Temporal
- Carotid
- Brachial
- Radial
- Ulnar
- Femoral
- Popliteal
- Dorsalis pedis
- Posterior tibialis

Lymph Nodes

- Cervical
- Epitrochlear
- Inguinal

Auscultate

- Arteries for bruits
- BP

DOCUMENT


Document your findings and develop your plan of care.

S U M M A R Y

- The incidence of PV disease continues to rise.
- Lower-extremity edema is common and represents a variety of possible disease processes. Therefore, it is essential that nurses understand the arterial, venous, and lymphatic circulatory mechanisms and the pathophysiological alterations that accompany common diseases of these systems.
- A thorough patient history and physical examination are the mainstays of assessing for PV and lymphatic disorders and the cornerstones for further diagnostic and laboratory evaluation.
- Providing an initial and ongoing assessment to obtain a diagnosis and monitor disease progression is a key task for the nurse and enhances the preservation of function, life, and limb in patients with these disorders.

R E V I E W Q U E S T I O N S

1. You are performing a PV exam. Which pulse is located at the lateral side of the wrist?
 - a. Radial
 - b. Ulnar
 - c. Brachial
 - d. Temporal
2. You are performing a PV exam. Which pulse is located at the lateral side of the head?
 - a. Radial
 - b. Ulnar
 - c. Brachial
 - d. Temporal
3. You are performing a PV exam. Which pulse is located behind the knee?
 - a. Ulnar
 - b. Brachial
 - c. Temporal
 - d. Popliteal
4. You are performing a PV exam. Which pulse is located on the top of the foot?
 - a. Posterior tibialis
 - b. Dorsalis pedis
 - c. Popliteal
 - d. Femoral
5. As part of Mr. Craig's vascular assessment, you measure his ABI. Which finding would indicate PV disease?
 - a. 0.5
 - b. 1.0
 - c. 1.5
 - d. 2.0
6. Arterial blood gases are ordered for your patient. Which test should be performed prior to obtaining an arterial blood specimen from the wrist?
 - a. Allen test
 - b. Manual compression test
 - c. Trendelenburg test
 - d. Color change test
7. You are assessing your patient's lower extremities for arterial insufficiency. Which test is best for assessing arterial flow?
 - a. Allen test
 - b. Manual compression test
 - c. Trendelenburg test
 - d. Color change test
8. Robert Mooney, age 61, has had a total right knee replacement. As part of your vascular exam, which test would be best to assess for deep vein thrombosis?
 - a. Homans' sign
 - b. Trendelenburg test
 - c. ABI
 - d. Color change test
9. You are palpating your patient's lymph nodes about 1 inch above the elbow. This site is best for assessing the:
 - a. Epitrochlear lymph node.
 - b. Horizontal lymph nodes.
 - c. Vertical lymph nodes.
 - d. Cervical lymph nodes.
10. You are palpating your patient's lymph nodes in the groin. This site is best for assessing the:
 - a. Epitrochlear lymph node.
 - b. Horizontal lymph nodes.
 - c. Vertical lymph nodes.
 - d. Cervical lymph nodes.



Assessing the Breasts

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent breast history questions
- Obtain a breast history
- Perform a physical assessment of the breast
- Document breast findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

Assessment of the breasts enables you to identify abnormalities and obtain valuable data, especially information that can help in early detection of disease. Breast cancer is the second leading cause of death in women, and successful treatment depends on early diagnosis. Knowing how to perform an accurate and thorough breast examination is crucial. Because most breast masses are detected by self-examination, teaching your patients how to examine their breasts is just as critical.

Anatomy and Physiology Review

To perform the assessment accurately, you must understand the anatomy and physiology of the breast. You need to be able to identify normal structures before you can identify abnormal findings. Understanding the physiology will also guide your assessment and enable you to interpret your findings.

Structures and Functions

The breast contains the mammary gland. The primary function of the breasts in female mammals is to produce nourishment to feed their offspring. In many countries, including the United States, the breasts are also associated with female sexuality. The breast lies between the sec-

ond and the sixth ribs and between the sternal edge and the midaxillary line. The tail of the breast extends into the axilla. Approximately two-thirds of the breast lies over the pectoralis major muscle, and the remaining one-third is superficial to the serratus anterior muscle.

Breasts are conical in shape and often unequal in size. The primary structures of the breast include lobes, lobules, milk (lactiferous) ducts, milk (lactiferous) sinuses, acini cells, areola, nipple, ligaments, blood vessels, lymphatic tissue, and supporting muscular tissue. The breast is composed of glandular, connective, and adipose tissues, as well as smooth muscle and nerve fibers. Figure 16.1 shows external breast structures and quadrants. Figure 16.2 is a cross-section of the breast, and Figure 16.3 shows the breast lymph nodes.

Although men's breasts are different from women's, men do have a few ducts surrounded by breast and other tissue. However, men generally do not secrete the same amounts of hormones (such as estrogen) that cause breasts to develop. Although the incidence is very rare (about 10 in 1 million), men can develop breast cancer. Therefore, you need to carefully examine male patients' breasts also.

Interaction With Other Body Systems

The breasts are not a system in and of themselves but are a part of the reproductive system. However, the breasts have a specific relationship to other systems.




FIGURE 16.1. External breast structures and quadrants.




FIGURE 16.2. Cross-section of breast.




FIGURE 16.3. Lymph nodes.

Structures and Functions of the Breast

STRUCTURE	FUNCTION
<p>Mammary Gland</p> <ul style="list-style-type: none"> Composed of 15 to 20 lobes. Each lobe has 50 to 75 lobules. <p>Lactiferous Ducts</p> <p>Lactiferous Sinuses</p> <p>Areola</p>	<ul style="list-style-type: none"> Lobules contain acini cells, which produce milk. Carry milk from glands to lactiferous sinuses. Store milk. Darker tissue surrounding nipple. Contains occasional hair follicles and small glands that lubricate nipple during breastfeeding. Sebaceous glands.
<p>Montgomery's Glands</p> <p>Nipple</p> <ul style="list-style-type: none"> Composed of connective tissue, smooth muscle, blood vessels, and nerve endings. 	<ul style="list-style-type: none"> Contraction of smooth muscles makes nipple firm and erect. Sensitive nerve endings are important area for sexual stimulation.
<p>Cooper's Ligament</p> <ul style="list-style-type: none"> Composed of bands of breast tissue fused with outer layers of superficial fascia. <p>Pectoralis Major and Serratus Anterior Muscles</p> <p>Internal Mammary Artery</p> <p>Lymph Nodes of Breasts</p> <ul style="list-style-type: none"> Anterior (pectoral) Central (midaxillary) Lateral (brachial) Posterior (subscapular) 	<ul style="list-style-type: none"> Suspensory ligaments attached to chest wall musculature that supports breasts. Breast lies over these chest muscles. Supplies blood to breast.
<ul style="list-style-type: none"> Internal mammary Epitrochlear Supraclavicular Infraclavicular 	<ul style="list-style-type: none"> On lower border of pectoralis major in anterior axillary fold; drains most of breast and anterior chest wall. High in axilla; pectoral, brachial, and subscapular drain into central nodes. High in axilla on inner aspect of humerus; drains most of arm. High in axilla on lateral scapular border deep in posterior axillary fold; drains part of arm and posterior chest wall. Deep in anterior chest; drains mammary glands. In depression above and posterior to medial area of elbow; drains arm. Above clavicle; often site for metastatic breast cancer. Below clavicle; often site for metastatic breast cancer.

The Neuroendocrine System

Sucking stimulates impulses to the hypothalamus. The hypothalamus stimulates the production of prolactin by the anterior pituitary, which stimulates milk production. The hypothalamus also stimulates the production of oxytocin by the posterior pituitary gland, which stimulates muscle cells surrounding the glandular tissue to contract and force the milk into the excretory ducts. Estrogen and progesterone are hormones that influence breast growth.

The Reproductive System


Breast development and function depend on estrogen and progesterone, which are produced in the ovaries. Estrogen elongates the excretory ducts and causes them

to create tributaries. Progesterone increases the number and size of the lobules to prepare the breasts for milk production and storage for nourishing a baby. Progesterone increases breast cell growth and dilation of blood vessels after ovulation. This engorgement is the reason why the breasts often become tender and swollen 5 to 7 days before menstruation, increasing in size, nodularity, density, and sensitivity.

Developmental, Cultural, and Ethnic Variations

Infants

At birth, the infant's nipple is elevated, and a slight secretion of milky material, "witch's milk," may occur for 5 to 7 days after birth. Palpable breast tissue is normal in



Relationship of Breasts to Other Systems

infancy, but not beyond this period. Some newborns have supernumerary nipples, a normal finding.

Children and Adolescents

A female child's breast is underdeveloped compared with an adult's. Growth of breast tissue generally begins in the prepubertal period. Physiological changes occurring in the breasts are the result of growth. Tanner staging (see Stages of Breast Development) is used to track breast development in females during puberty. (For more information, see Chapter 26, Assessing the School-Age Child and Adolescent.)

Pregnant Women

Breasts become fuller and firmer, and the areola and nipples darken and enlarge. In the third trimester, colostrum, a yellow secretion, occurs. Colostrum continues to be secreted after the birth of a baby until milk is produced.

(For more information, see Chapter 23, Assessing the Mother-to-Be.)

Older Adults


As women age, glandular tissue is replaced with adipose tissue and the ducts become more fibrous. These changes, along with a general reduction of muscle mass and tone, cause breasts to be less firm and more pendulous. Arthritic changes in the joints and hands make breast self-examination (BSE) a little more challenging. (For more information, see Chapter 27, Assessing the Older Adult.)

People of Different Cultures and Ethnic Groups


Some health problems are culturally or racially bound. That is, diseases either are associated exclusively with a certain race or ethnic group or occur more frequently in that group. Ashkenazi Jews have a greater

Stages of Breast Development


Stage 1: Prepuberty
Elevation of papilla




Stage 2: Breast bud
Elevation of breast and nipple and increased diameter of areola




Stage 3: Areola deepens in color and enlarges further. Glandular tissue begins to develop beneath areola.



Stage 4: Areola appears as a mound; breast appears as a mound.



Stage 5: Mature breast
Areola recesses to general contour of breast; nipple projects forward.





HELPFUL HINTS

- Pregnancy is not an ideal time for a woman to learn how to do BSE. Getting to know what is normal for her will be difficult until pregnancy and lactation are completed.
- Older women may be reluctant to examine their breasts because they have the idea that “touching yourself” is taboo.
- Older women may have trouble assuming some positions necessary for the physical examination. Besides being sensitive to their needs, adapt technique to meet patient’s needs.
- Advise older women to sprinkle talcum powder on their hands so that they will glide more smoothly during BSE.

incidence of breast cancer than other ethnic groups because they are believed to have a gene that predisposes them to familial breast cancer. White women have a higher incidence of breast cancer than non-white women, but African American women have lower survival rates than white women. African American men appear to be at greater risk for breast cancer than white men. In addition, some cultural practices may delay early diagnosis or treatment. For instance, African American women in the southern United States have been known to treat breast lesions with home remedies.

Introducing the Case Study



Lynn Kobrynski is a childless 46-year-old married woman who visits her nurse practitioner for an annual physical exam. She states: “My husband thinks I have a lump in my right breast.” Mrs. Kobrynski has been active and healthy and has never been pregnant. She has a family history of breast cancer. Considering her complaint and the fact that she has not had a complete physical in 5 years, a complete health history and physical examination are warranted.

Performing the Breast Assessment

A comprehensive assessment includes taking the patient’s health history and performing a physical examination of the breasts and axillae.

Focused Breast History

If you are unable to perform a detailed history, ask key questions that focus on a history of breast problems, presence of symptoms, identification of risk factors, and health promotion activities. The following questions focus on the warning signs of breast cancer:

- Do you have a lump or thickening in or near your breast or under your arm that persists through the menstrual cycle?
- Is the skin on your breast or nipple red, dimpled, puckered, scaly, or inflamed?
- Do you have nipple changes? For example, a change in the direction in which one nipple points, inversion, eversion, or discharge?
- Has your breast changed in size, shape, or contour?

Comprehensive Breast Assessment Health History

The health history involves interviewing the patient regarding her perception of her health status. It includes a broad range of questions so that possible problems associated with breast cancer may be identified. Keep in mind that information collected as part of the health history may uncover problems related to systems outside the breast (e.g., metastatic disease). If time is an issue and you are unable to perform a complete health history, perform a focused history on the breasts.

Biographical Data

Biographical information can provide valuable insights about the health of your patient’s breasts. Begin by asking your patient’s age—it influences the questions you ask during the rest of the health history. Breast development varies according to the age of your patient, and various breast problems are age related. For example, the risk for breast cancer increases with age; fibroid adenomas usually occur before age 30; the incidence of intraductal papilloma, a benign lesion of the lactiferous duct, peaks at age 40; fibrocystic breast changes usually occur between ages 35 and 50; and ductal ectasia occurs most commonly in perimenopausal women in their 50s.

Also ask about your patient’s ethnic background because it may influence the risk for breast disease. For example, the incidence of breast cancer is greater in white women than in nonwhite women, and fibroadenomas most commonly occur in African Americans.

Current Health Status

If your patient’s chief complaint is about her breast, investigate this first; then perform a symptom analysis. Common chief complaints involving the breast include

lumps or masses, breast pain or tenderness, and nipple discharge. Related complaints include changes in the venous pattern of the breast (increased vascularity), dimpling or puckering of breast skin, redness or scaliness of breast skin, and change in the direction of the nipple.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Lump or Mass

The most common complaint related to the early detection of breast cancer, a lump or mass, may be a normal physiological nodularity or may indicate a neoplasm. Careful analysis of this symptom can provide data that will help you distinguish between normal and abnormal breast changes.



HELPFUL HINTS

- If your patient is premenopausal, be sure to determine the relationship of symptoms to the menstrual cycle.
- When assessing precipitating or palliative factors, explore the patient's use of culturally related health practices or remedies. Use of these remedies may delay the initiation of traditional Western treatment, which can be deleterious to the patient's health.
- Give special attention to adolescent girls who are sensitive about body image. Ask about recent breast trauma.

Pain or Tenderness

Breast pain or tenderness (mastalgia) is usually related to the normal menstrual cycle and is generally not charac-

teristic of a malignant lesion. However, it may be associated with extensive or metastatic disease and therefore needs to be evaluated. Pain can also be caused by a rapidly enlarging cyst, an infection or abscess, mastitis, hematoma, trauma, or fibrocystic disease. Some disorders have classic signs located in the breast that are diagnostically significant. For instance, redness and pain may indicate inflammatory breast cancer, trauma, or an abscess.



CRITICAL THINKING ACTIVITY # 1

How would you explain pain or tenderness associated with the menstrual cycle?

Nipple Discharge

Nipple discharge may be normal or abnormal. Discharge may occur spontaneously or be manually expressed and may be white, milky, serous, clear, bloody, gray, brown, or green. A thorough evaluation is necessary to determine the cause. Unless the patient is lactating, spontaneous discharge is usually not considered normal and warrants further investigation.



ALERT

Any spontaneous discharge from a nonpregnant or lactating woman's breast is an abnormal, potentially serious finding and must be followed up.

Past Health History

The past health history compares the patient's past health and present health and uncovers risk factors that might predispose her to breast disease.

Past Health History

RISK FACTORS/QUESTIONS TO ASK

Childhood Illnesses

- Do you have a history of breast lumps as an adolescent or young adult?
- When did you begin menstruation?

Surgery

- Have you ever had surgery or other treatments involving the breasts?

Hospitalizations/Diagnostic Procedures

- Have you ever been treated for a breast problem?
- How often do you have a mammogram?

RATIONALE/SIGNIFICANCE

- Childhood illnesses may cause permanent effects.
- Fibroadenomas usually occur between ages 15 and 25.
- Early menarche (<age 12) is a risk factor for breast cancer because breasts are under hormonal influence longer.
- Provides information on possible reasons for symptoms (e.g., breast implants).
- Helps identify potential or recurring problems or may explain physical findings.
- Identifies current health practices.

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Serious Injuries</p> <ul style="list-style-type: none"> ■ What serious injuries have you had? ■ What treatment did you receive? ■ Do you have any residual effects from injuries or treatments? <p>Serious/Chronic Illnesses</p> <ul style="list-style-type: none"> ■ Do you have any other medical problems? For example, cancer in any other area? ■ Recent infection? ■ Have you recently stopped nursing a child? <p>Allergies</p> <ul style="list-style-type: none"> ■ Do you have any allergies (e.g., to perfume or deodorant)? <p>Medications</p> <ul style="list-style-type: none"> ■ Are you taking any prescribed or over-the-counter (OTC) medications? Hormones or HRT? BCPs? Other medications? 	<ul style="list-style-type: none"> ■ Data on past breast injuries provide baseline data for physical assessment. ■ Women who received radiation therapy for breast disorders may have redness, scaliness, and other symptoms that may appear to be a malignancy. ■ Liver disease, testicular tumors, hyperthyroidism, and adrenal and pituitary tumors can cause gynecomastia in men. ■ Cancer from another area may have local spread involving the lymph nodes. ■ Stopping nursing abruptly can cause breasts to become engorged, red, and hard. ■ May cause redness, pruritus, and scaliness in breasts and axillae. ■ HRT is associated with increased risk for breast cancer. ■ Unopposed use of estrogen may increase risk of breast cancer. ■ Digoxin can cause gynecomastia in men. ■ See <i>Drugs That Adversely Affect the Breasts</i>.

Drugs That Adversely Affect the Breasts

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Androgens	Danazol, fluoxymesterone, methyltestosterone, testosterone	<i>Female:</i> Decreased breast size. <i>Male:</i> Gynecomastia.
Antidepressants	Tricyclic antidepressants	<i>Female:</i> Breast engorgement and galactorrhea (spontaneous milk flow). <i>Male:</i> Gynecomastia.
Antipsychotics	Chlorpromazine hydrochloride, fluphenazine, haloperidol, perphenazine, prochlorperazine maleate, thioridazine, thiothixene	<i>Female:</i> Galactorrhea, moderate engorgement of the breast with lactation, mastalgia (breast pain). <i>Male:</i> Gynecomastia, mastalgia.
Cardiac glycosides	Digitoxin, digoxin	<i>Male:</i> Gynecomastia.
Estrogens	Conjugated estrogens, esterified estrogens, estradiol, estrone, ethinyl estradiol	<i>Female:</i> Breast changes, tenderness, enlargement, secretions. <i>Male:</i> Breast changes, tenderness, gynecomastia.
Oral contraceptives	Dienestrol Estrogen-progesterone combinations	<i>Female:</i> Breast tenderness. <i>Female:</i> Breast changes, tenderness, enlargement, secretions.
Progestins	Medroxyprogesterone acetate, norethindrone, norgestrel, progesterone	<i>Female:</i> Breast tenderness or galactorrhea.

(continued)

Drugs That Adversely Affect the Breasts (continued)

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Miscellaneous agents	Isoniazid	<i>Male:</i> Gynecomastia.
	Reserpine	<i>Female:</i> Breast engorgement, galactorrhea. <i>Male:</i> Gynecomastia.
	Spironolactone	<i>Female:</i> Breast tenderness. <i>Male:</i> Painful gynecomastia.
	Cimetidine	<i>Female:</i> Galactorrhea. <i>Male:</i> Gynecomastia.

Family History

The family history identifies familial tendencies toward

breast disease and disorders. The following questions may help you elicit this information.

Family History

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Familial or Genetically Linked Breast Problems</p> <ul style="list-style-type: none"> ■ Do you have a family history of breast cancer? ■ If yes, what family members were afflicted? What type of breast cancer? ■ How old were they when diagnosed? ■ Do you have a family history of other types of cancer? What types? 	<ul style="list-style-type: none"> ■ Women with two or more first-line relatives (mother or sister) with breast cancer have increased risk. ■ Number of first-line relatives with breast cancer may raise questions about genetic alterations leading to predisposition for breast cancer. ■ Women with altered or mutated BRCA1 or BRCA2 gene have an 85 to 95 percent lifetime chance of developing breast cancer. ■ Positive family history for cancer increases risk to patient. ■ Risk of cancer increases with age. ■ If any family member had premenopausal breast cancer, need to investigate possibility of genetic alteration. ■ May indicate hereditary family syndromes.

Review of Systems

The review of systems (ROS) enables you to relate

the breast to other systems and often reveals information that your patient might have omitted earlier.

Review of Systems

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>General Health Survey</p> <ul style="list-style-type: none"> ■ How have you been feeling? ■ Do you have a fever? ■ Have you lost weight? <p>Integumentary</p> <p>Skin on Breasts and Axillae</p> <ul style="list-style-type: none"> ■ Do you have any dimpling, redness, rashes, lesions, or lumps? 	<ul style="list-style-type: none"> ■ Provides an indicator of general overall health. ■ Associated with mastitis, abscess, infection, and sometimes, metastatic disease. ■ Unexplained weight loss may be associated with malignancy. ■ <i>Dimpling:</i> Malignancy.

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Respiratory</p> <ul style="list-style-type: none"> ■ <i>If patient has breast cancer:</i> Any breathing difficulty? <p>Cardiovascular</p> <ul style="list-style-type: none"> ■ Are you on any cardiovascular medications? <p>Gastrointestinal</p> <ul style="list-style-type: none"> ■ <i>If patient has breast cancer:</i> Any increase in abdominal size? ■ Liver disease? <p>Reproductive</p> <ul style="list-style-type: none"> ■ At what age did your menstrual cycle begin? ■ Have you ever taken BCP? ■ Have you ever been pregnant? If so, at what age(s)? ■ Do you have a history of infertility? ■ Did you breastfeed? ■ <i>If postmenopausal:</i> At what age did your menstrual cycles cease? ■ Have you ever been on HRT? If so, what type and for how long? <p>Endocrine</p> <ul style="list-style-type: none"> ■ Do you have a history of hyperthyroidism? 	<ul style="list-style-type: none"> ■ <i>Scaly erosion of areola and nipple:</i> Paget's disease (breast malignancy of areola and nipple). ■ <i>Redness over affected area:</i> Mastitis. ■ Lung is common site for metastasis. ■ Some medications such as digoxin have side effect of gynecomastia. ■ Liver is common site for metastasis. ■ Can cause gynecomastia in men. ■ Early menarche and late menopause are risk factors for breast cancer. ■ Can cause breast changes, such as tenderness, enlargement, and secretions. ■ Nulliparity, first pregnancy after age 30, and infertility have been linked to breast cancer. ■ Breastfeeding decreases risk for breast cancer. ■ HRT has been linked with increased incidence of breast cancer. ■ Can cause gynecomastia in men.

Psychosocial Profile

The psychosocial profile can reveal patterns in your

patient's life that place her at risk for breast disease. You may also identify teaching needs.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Health Practices and Beliefs/Self-Care Activities</p> <ul style="list-style-type: none"> ■ How do you promote breast health? ■ Do you get regular CBEs? ■ Do you get regular mammograms? ■ Do you perform monthly BSEs? <div style="border: 1px solid #ccc; padding: 5px; margin-top: 10px;">  <p>HELPFUL HINT</p> <p>Ask patient to demonstrate BSE. Do not rely on self-report to ascertain adequate knowledge of technique.</p> </div> <p>Typical Day</p> <ul style="list-style-type: none"> ■ Do you have trouble performing activities of daily living (ADLs)? ■ Is your range of motion (ROM) restricted? 	<ul style="list-style-type: none"> ■ Helps identify activities that reduce breast cancer risk. ■ Early detection decreases morbidity and mortality. ■ American Cancer Society (ACS) screening guidelines include: <ul style="list-style-type: none"> ■ Monthly BSE beginning at age 20. ■ Annual screening mammography beginning at age 40. ■ CBE by medical professional every 3 years from age 20 to 40; then yearly. ■ Helps identify if patient will have trouble caring for herself and performing BSE.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ What did you eat in the last 24 hours? ■ Have you recently lost or gained weight? ■ Does your menstrual cycle affect your appetite and the type of food you eat? ■ Do you have PMS? ■ Do you eat a lot of high-fat foods? Salty foods? 	<ul style="list-style-type: none"> ■ May provide baseline information on dietary habits. ■ Weight gain may increase breast size, causing tenderness. ■ May need dietary consultation to discuss foods to decrease premenstrual syndrome (PMS) symptoms (low sodium, low fat). ■ Cancer, especially of the breast, is associated with a high-fat diet. ACS recommends <30 g of fat/day. ■ Diets high in sodium and methylxanthines, such as chocolate and coffee, can aggravate fibrocystic breast disease.
<p>Activity and Exercise Patterns</p> <ul style="list-style-type: none"> ■ Do you exercise? If so, how much and what type? 	<ul style="list-style-type: none"> ■ Sedentary lifestyle increases risk for cancer. ■ Exercise is linked with decreasing cancer risks overall.
<p>Sleep and Rest Patterns</p> <ul style="list-style-type: none"> ■ Do you have trouble sleeping at night? 	<ul style="list-style-type: none"> ■ Patients with breast swelling or pain/tenderness may have trouble finding a comfortable position for sleep.
<p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you drink alcohol? If so, how much? ■ Do you drink caffeinated beverages? ■ Do you use heroin or marijuana? 	<ul style="list-style-type: none"> ■ High intake of alcohol may be linked with cancer. ■ Can worsen fibrocystic breast disease. ■ Can cause gynecomastia in men.
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ What type of work do you do? Does it involve heavy lifting? 	<ul style="list-style-type: none"> ■ Diagnosis or treatment of breast disorder may affect whether patient can return to work at same job.
<p>Roles/Relationships/Self-Concept</p> <ul style="list-style-type: none"> ■ How important are your breasts to a positive view of yourself? ■ Do you have a difficult time discussing your problem? 	<ul style="list-style-type: none"> ■ Breast cancer can cause body image disturbances and have a major impact on a woman's self-image. Fear of losing breast may delay treatment.
<p>Religious/Cultural Influences</p> <ul style="list-style-type: none"> ■ What are your religious practices/beliefs or cultural influences in regard to healthcare? 	<ul style="list-style-type: none"> ■ May affect patient's healthcare practices.
<p>Sexuality Patterns</p> <ul style="list-style-type: none"> ■ Has your breast disorder affected your sex life? How? 	<ul style="list-style-type: none"> ■ Breast pain or tenderness (mastalgia) can interfere with sexual activity. ■ Loss of a breast to cancer can make a woman avoid sex because she feels less attractive.
<p>Social Supports</p> <ul style="list-style-type: none"> ■ Who are your main sources of support? Have you seen a change in the way your support person or spouse sees you since you developed a breast problem? 	<ul style="list-style-type: none"> ■ Identifying patient's support system is essential in planning care.



Case Study Findings

Reviewing Mrs. Kobrynski's

Biographical data:

- 46-year-old married woman.
- No pregnancies, no children.
- Caucasian, Catholic religion, Polish descent.
- College graduate; employed as high school teacher.
- *Contact person:* Husband, Edward Kobrynski, a steelworker.

- Insurance from full-time work.
- *Source of information:* Self, seems reliable.

Current health status:

- Chief complaint is that her husband thinks she has a lump.
- Does not perform monthly BSE.

Past health history:

- Menarche at age 12; menstrual cycles have always been regular—every 28 days and lasting 3 days.

- Had lumpectomy for benign breast lesion; otherwise negative.
- No pregnancies; no history of infertility.
- No known food, drug, perfume, or deodorant allergies.
- Does not take any medications other than occasional Tylenol for headache. No history of birth control pills (BCP) or hormone replacement therapy (HRT).
- No other medical problems. Does not examine breasts, and has never had a complete breast examination (CBE) or mammogram.

Family history:

- Positive for ductal breast cancer.
- Maternal grandmother died at age 87; diagnosed with ductal breast cancer at age 85.
- Mother had benign breast lesion at age 57.
- Patient has two younger sisters, both alive and well.
- One maternal aunt, alive and well.
- No other cancer history in family.

Review of systems:

- *General Health Survey:* Usual state of health good.
- *Integumentary:* No dimpling, redness, or rashes on breasts or axillae.
- *Cardiovascular:* Reports no problems or medications.
- *Reproductive:* Menarche at age 12; menstrual cycle regular every 28 days, lasting 3 days; no history of BCP or HRT.

Psychosocial profile:

- Has not had physical exam in 5 years; has never had a mammogram; does not perform BSE; has not had CBE in 5 years.
- Typical day consists of awakening at 6 A.M., showering and dressing, eating breakfast with husband, and driving 30 minutes to school. Works 8 A.M. to 3:30 P.M. Monday through Friday. Returns home by 4 P.M., takes a walk or relaxes, fixes dinner. After dinner, corrects papers, watches TV, or reads. Spends weekends catching up on housework, shopping and running errands, and socializing with friends.
- Usually eats well, although diet premenstrually tends to be high in fats and salt. Temporarily gains 5 lb before she gets her period; otherwise, weight stable for last 10 years.
- Walks 3 miles, three to four times a week.
- Usually gets 8 hours sleep; no difficulty sleeping.
- Denies use of alcohol and caffeine.
- Employed full time, likes job, which does not require lifting.
- Satisfied with sex life.

- Married 15 years, husband and friends are supports.
- Goes to church weekly; religion is an important aspect of her life.



Case Study Evaluation

Before you begin the physical examination, document the findings from your interview with Mrs. Kobrynski.



CRITICAL THINKING ACTIVITY #2

What additional questions would you ask Mrs. Kobrynski to further assess her chief complaint?



CRITICAL THINKING ACTIVITY #3

Based on the history you obtained from Mrs. Kobrynski, what breast problem is she at risk for? What are the risk factors?

Physical Assessment

Now that you have obtained the history, turn your attention to collecting objective data by performing the physical examination. As you assess your patient's breasts, remember that this is an ideal opportunity to teach BSE.

Anatomical Landmarks

The breast examination should include the areas above the clavicle to the sixth or seventh rib and from the sternum to the midaxillary line (Fig. 16.4).

Approach

You will use the techniques of inspection and palpation to examine the breasts. The examination is performed in several positions:

- Sitting, arms at side.
- Sitting, arms over head.
- Sitting, hands on hips.
- Sitting, leaning forward.
- Supine with pillow under shoulder of breast being examined.

The hands over head and hands pressed on hips positions are helpful in detecting dimpling or retraction of the breast tissue. The sitting, leaning-forward position is helpful when assessing large, pendulous breasts. With the supine position, have the woman lie on the examination table and place a small pillow or rolled towel under the shoulder of the breast you will examine first. The pillow or towel helps spread the breast tissue over the chest wall.




FIGURE 16.4. Anatomical landmarks.

Examination in Breast Reconstruction

Today, many women have breast reconstruction for cancer or for cosmetic reasons. Breast examination in these women is done exactly the same way as for natural breasts and must not be omitted. If an implant was used for the reconstruction, press firmly inward at the edges of the implant to feel the ribs beneath.



TOOLBOX

Assemble the following equipment for a routine breast examination: Small pillow or towel, mirror, centimeter ruler, nonsterile gloves, specimen slide, and specimen culture swab.

Performing a General Survey

Before examining specific areas, scan your patient from head to toe, looking at her general appearance. Then take vital signs and document them. A temperature elevation may be associated with mastitis.

Performing a Head-to-Toe Physical Assessment

Now, look for changes in every system that may relate to the breast. Because the breasts are not a system, there may not be an obvious relationship to other systems. However, if your patient has breast cancer, be alert for changes in other systems that reflect metastatic disease.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

General Health Survey

INSPECT

- No acute distress; AAO × 3; responses clear and appropriate; affect pleasant; well-groomed.

Integumentary

INSPECT

- No rashes or lesions on breasts or axillae.

ABNORMAL FINDINGS/RATIONALE

- *If client has breast cancer:* Changes in level of consciousness (LOC), dysphasia, or hemiparesis may indicate brain metastasis.
- *If client has breast lump,* mood may be depressed or anxious.
- *Axillary rashes:* Allergies.
- *Breast lesions/redness:* Malignancy or mastitis.

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p>HEENT <i>Eyes</i> <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ Vision intact; no papilledema. <p><i>Neck</i></p> <ul style="list-style-type: none"> ■ No palpable nodes. ■ Thyroid nonpalpable. <p>Respiratory <i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Respiratory rate 18/min. ■ Lungs clear. <p>Cardiovascular <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ No increased venous pattern on chest. <p>Gastrointestinal <i>INSPECT/PALPATE/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Swallowing intact; positive bowel sounds; abdomen soft, nontender, no hepatomegaly. <p>Genitourinary/Reproductive <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ Normal pelvic exam. <p>Musculoskeletal <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ Muscle strength +5. ■ Movements smooth and coordinated. ■ Gait balanced, no ataxia. <p>Neurological <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ No sensory loss. 	<ul style="list-style-type: none"> ■ <i>If client has breast cancer:</i> Visual changes and papilledema may indicate brain metastasis. ■ <i>Enlarged lymph nodes:</i> Infection or metastasis. ■ <i>Palpable thyroid:</i> Hyperthyroidism, which can cause gynecomastia in men. ■ <i>If client has breast cancer:</i> Adventitious breath sounds, decreased chest excursion, and dullness to percussion seen with pleural effusion and may indicate metastasis to lungs. ■ <i>Increased venous pattern, especially asymmetrical:</i> Breast cancer. ■ <i>If client has breast cancer:</i> Ascites and hepatomegaly may indicate metastasis to liver. ■ Endometrial and ovarian cancer increase risk for breast cancer. ■ <i>If client has breast cancer:</i> Weakness and ataxia may indicate metastasis to bone and brain. ■ <i>If client has breast cancer:</i> Unilateral sensory loss and hemiparesis may indicate metastasis.

Performing a Physical Assessment of the Breast

Now that you have looked at the relationship of the breasts to other systems, focus on the specifics of the examination. First inspect the breasts and axillae; then palpate them.

Inspection

Inspect the breasts with the patient in all five positions, as described in the Approach section. Inspect the breasts for size, shape, symmetry, skin condition, and venous pattern. Inspect the nipple and areola for nipple inversion or eversion, nipple direction, and nipple discharge. (See *Inspecting the Breasts*.)

Figure 16.5 shows milk lines, mentioned under Nipples and Areolae.

Palpation

Palpation enables you to feel for lumps, masses, and other abnormalities. Palpate the entire breast and axilla on both sides, making three small circles with the fingerpads of your middle three fingers. Use different pressure for each circle—light, medium, and deep. Light pressure allows you to palpate lesions beneath the skin; deep pressure allows you to palpate lesions that are deeper and near the ribs. Feel for lumps, thickening, or other changes. Because the breast extends from the midaxillary line to the sternal notch and the second to the sixth ribs, you must palpate all these areas. Put another way, you must palpate from the bottom of the bra line (sixth or seventh intercostal space) to the collarbone (clavicle) and from

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	Ask about: General health Fever, weight changes	General	Observe: LOC Affect Measure: Vital signs Weight
Integumentary	Ask about: Rashes Lesions Allergies	Integumentary	Inspect: Rashes Lesions Color changes
HEENT	If patient has breast cancer, ask about: Headaches Changes in mentation	HEENT	If patient has breast cancer: Test vision.
Respiratory	If patient has breast cancer, ask about: SOB Cough Breathing difficulty	Respiratory	If patient has breast cancer: Auscultate and percuss lungs
Cardiovascular	Ask about: CV medications	Cardiovascular	Inspect: Increased vascularity of breasts
Gastrointestinal	If patient has breast cancer, ask about: Increase in size of abdominal girth.	Gastrointestinal	If patient has breast cancer: Palpate abdomen.
Genitourinary/ Reproductive	Ask about: Last menstrual period, whether patient is pregnant, number of pregnancies, year of first pregnancy, whether patient is pre- or postmenopausal. Medications (BCP, HRT)	Genitourinary/ Reproductive	Inspect: Perform pelvic exam
Musculoskeletal	If patient has breast cancer, ask about: Weakness Bone pain Fractures	Musculoskeletal	If patient has breast cancer: Test muscle strength
Endocrine	Ask about: History of thyroid disease	Neurological	If patient has breast cancer: Test LOC, sensory and motor function
Lymphatic/ Hematologic	Ask about: Infection or malignancy	Endocrine	Palpate thyroid
		Lymphatic/ Hematologic	Palpate lymph nodes

Assessment of the Breasts' Relationship to Other Systems




FIGURE 16.5. Milk lines.

the line down the middle of the breastbone (sternum) to the line down the middle of the armpit (midaxillary line), including the axilla.

**HELPFUL HINTS**

- Do not remove your fingerpads from the skin surface or jump from area to area.
- Most breast lesions in women are found in the upper outer quadrant. This area includes the tail of Spence and contains more breast tissue than the other quadrants.
- Although only about 1 percent of men get breast cancer, you still need to assess their breasts. Breast cancer in men most frequently occurs in the areola.

Palpating the Breasts

There are three techniques of breast palpation—the vertical strip method, the pie wedge method, and the concentric circle method. When demonstrating these methods for your patient, stress that it doesn't matter which method she uses during BSE, as long as she uses the same method consistently and examines the entire breast area and axillae.




FIGURE 16.6. Vertical strip method.

Vertical Strip Method

With this method (Fig. 16.6), you start at the sternal edge and palpate the breast in parallel lines until you reach the midaxillary line. This is like “mowing the grass,” in which you go up one area and down the adjacent strip. Be sure to palpate the tail of Spence as well. End by examining and palpating the nipple. Note its elasticity, and squeeze it between your thumb and your index finger to see if there is a discharge. Because of the possibility of discharge, be sure to wear nonsterile gloves when palpating the nipple. If the patient is supine, shift the pillow or rolled towel to the opposite side and repeat the procedure with the other breast.

**HELPFUL HINT**

If you notice an open lesion on the breast or axillae, wear nonsterile gloves to prevent contact with body fluids.

Pie Wedge Method

This method (Fig. 16.7) examines the breast in wedges or “pie slices.” Once one wedge is examined, move to the adjacent wedge until you examine the entire breast and the tail of Spence. Then examine the nipple as you did in the vertical strip method. Repeat with the opposite breast.



FIGURE 16.7. Pie wedge method.

Concentric Circles Method

This method (Fig. 16.8) uses concentric circles to examine the entire breast area. Start in the outermost area (or largest circle) at 12 o'clock; then move to 1 o'clock, 2 o'clock, and so on until the first circle is completed. When you complete the circle and return to 12 o'clock, move your fingers two fingerbreadths inward and examine another concentric circle. Repeat this procedure until you reach the nipple area. Examine the nipple in the same fashion as the previous two methods. Also examine the tail of Spence. Repeat with the other breast.



FIGURE 16.8. Concentric circles method.

INSPECTING THE BREASTS

1. Positions for Breast Examination

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect breast with the patient in various positions:
- Sitting, arms at side.



- Sitting, arms over head.



- Sitting, hands on hips.



- Sitting, leaning forward.



- Supine, with pillow under shoulder of breast being examined.



- Sitting, pushing hands together.



HELPFUL HINTS

- Large, pendulous breasts are easier to visualize if woman leans forward so they hang free.
- Note patient's dominant side—it may be more developed.

HELPFUL HINT

To accentuate dimpling or retraction, have patient press hands on hip or push hands together.

- Note size, shape, symmetry, lesions, discharge, venous pattern, dimpling, or retraction.
- *Breast conical, symmetrical, or slightly asymmetrical.*
- *Skin color lighter than in exposed areas; no lesions, redness, or edema; texture even.*
- *Striae often seen with breast enlargement during pregnancy.*
- *No dimpling or retraction.*
- *No increase in venous pattern unless patient is pregnant. Then symmetrical increase is normal.*

INSPECTING THE BREASTS

1. Positions for Breast Examination (continued)

ABNORMAL FINDINGS/RATIONALE

- Change in symmetry warrants further investigation.
- *Edema and peau d'orange (orange skin appearance)*: May be related to lymphatic obstruction.
- *Erythema*: Infection, abscess, or inflammatory carcinoma of breast.
- *Dimpling/puckering*: Sign of retraction phenomena or abnormal traction on Cooper's ligaments caused by neoplasm. Attachment of tumor to fascia and pectoralis muscle pulls on skin and produces dimpling.
- *Lesions/asymmetrical increased venous pattern*: Signs of breast cancer.

2. Nipples and Areolae

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect position, direction, discharge, or lesions.
- Note presence of supernumerary nipples.
- *Nipples everted, pointing in same direction, no discharge or lesions.*
- *Spontaneous discharge normal during pregnancy and lactation.*
- *Areola and nipple darker than breast tissue. Become even darker during pregnancy.*
- *Supernumerary breasts or nipples are congenital anomalies in which small, palpable masses or nipples are present along milk lines, embryonic ridges that extend from axilla to groin. They usually atrophy during fetal development, except where the breasts develop. Cracks, redness of nipple can occur with nursing.*

ABNORMAL FINDINGS/RATIONALE

- Inverted nipples may make breastfeeding difficult.
- Nipple changes—eversion to inversion or changes in direction they are pointing—may indicate an underlying mass.
- Flattened or inverted nipples are caused by shortening of mammary ducts.
- Spontaneous discharge not associated with pregnancy or breastfeeding warrants follow-up. Obtain a specimen for evaluation.
- *Lesions/erosion/ulceration of areola and nipple*: Paget's disease.
- Discoloration of areola and nipple that is not associated with pregnancy warrants follow-up.

3. Axillae

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect color, lesions.
- *Skin intact. No rashes or lesions.*



Inspecting the axillae

ABNORMAL FINDINGS/RATIONALE

- *Rashes/redness/unusual pigmentation*: Infection, allergy.
- *Dark pigmentation/velvety skin texture of axilla*: Malignant acanthosis nigricans, a rare cancer.

Palpating the Axillae and Lymph Nodes

Using your fingerpads, move your fingers in circular fashion to lightly palpate the axillary and clavicular nodes. Palpate the axillae with the patient supine or sitting up with her arms at her sides. Palpate the supraclavicular, infraclavicular, central, lateral, anterior, posterior, and epitrochlear nodes. (For more information, see Chapter 15, Assessing the Peripheral-Vascular and Lymphatic Systems.)



HELPFUL HINT

Older women may have trouble changing positions, so examine the axillae after examining the breasts, with the patient in the same position.

PALPATING THE BREASTS, AXILLAE, AND LYMPH NODES

1. Breasts

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Position patient supine with arm over head and small pillow under shoulder of breast being examined.
- Use three levels of palpation: light, medium, and deeper.
- Using your finger pads, move in circular fashion across breast using vertical strip method, pie edge method, or concentric circle method.
- Be sure to cover entire breast area from the sternum to the midaxillary line and from the clavicle to the sixth or seventh intercostal space.
- Wear gloves if open lesions or discharge present.
- *Breasts soft, nontender.*
- **Consistency is age dependent:** *More firm and elastic in premenopausal women; less firm and elastic with ducts that may feel stringy or cordlike in postmenopausal women.*



HELPFUL HINTS

- In older adults, the inframammary ridge below the breast thickens and can be mistaken for a mass.
- In pregnancy, the breast may feel firm; during lactation, it may feel hard if engorged.



Palpating the breast

ABNORMAL FINDINGS/RATIONALE

Benign Lumps

- *Fibroadenoma:* Smooth, firm, round, movable, nontender, 1 to 5 cm in size.
- *Fibrocystic breast disease:* Nodular, tender, movable, soft to firm, in postmenopausal women.

Malignant Breast Lumps

- *Breast cancer:* Irregular shape; irregular, poorly defined borders; nontender, immovable; increase in size as disease progresses.
- *Breast warm and indurated (hard):* Mastitis; *Staphylococcus aureus* is most common cause.

2. Nipples

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Gently palpate the nipple.
- Note elasticity, tenderness, discharge.
- *Nipples elastic, nontender. No discharge or white, sebaceous secretion.*



Palpating the nipple

PALPATING THE BREASTS, AXILLAE, AND LYMPH NODES

2. Nipples (continued)

ABNORMAL FINDINGS/RATIONALE

- Loss of elasticity may indicate underlying malignancy.
- *Bloody, purulent discharge*: Infection
- *Serous, serosanguineous, or bloody drainage*: Intraductal papilloma.
- *Thick, gray drainage, fixation of nipple*: Ductal ectasia.

3. Axilla and Clavicular Nodes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- If nodes palpable, note size, shape, tenderness, mobility, consistency.
- *Supraclavicular nodes*: Gently palpate above the clavicle.



Palpating supraclavicular nodes

- *Infraclavicular nodes*: Gently palpate below the clavicle.



Palpating infraclavicular nodes

- *Central nodes*: With the patient sitting, support arm on side being examined, reach high into the axilla and gently palpate.



Palpating central nodes

- *Lateral nodes*: Support patient's arm, and turn your hands toward the humerus. Gently palpate the inner aspect of the arm.



Palpating lateral nodes

- *Posterior or scapular nodes*: Support patient's arm, place your hand toward the back at the scapula, and gently palpate.



Palpating posterior nodes

- *Anterior or pectoral nodes*: Support the patient's arm, and place your hand forward toward the anterior axillary fold.



Palpating anterior nodes

- *Epitrochlear nodes*: Gently palpate about ½ to 1 inch above the elbow on the outer aspect of the arm.
- *Nonpalpable*



Palpating epitrochlear nodes

(continued)

PALPATING THE BREASTS, AXILLAE, AND LYMPH NODES

3. Axilla and Clavicular Nodes (continued)

ABNORMAL FINDINGS/RATIONALE

- *Palpable nodes:* Infection or metastatic disease.
- Enlarged lymph nodes caused by infection are usually tender; those caused by malignancy are nontender.




FIGURE 16.9. Describing the location of abnormalities.

Documenting Your Findings

If you palpate a mass or lump, describe its size, shape, symmetry, mobility, delimitation, tenderness, consistency, temperature, and degree of redness. To document the location of breast masses, divide the breast into four quadrants by horizontal and vertical lines. Then visualize the breast as a clock face with the nipple as the center. Locate the pain or lump by the time on the clock (e.g., 2 o'clock), and measure the distance in centimeters from the nipple (Fig. 16.9).



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Kobrynski have concluded. The following are your findings:

- **General Health Survey**
 - No acute distress.
 - Awake, alert, and oriented x 3.
 - Responses clear and appropriate.

- Affect pleasant, but slightly anxious.
- Well-groomed, looks younger than stated age.
- Head-to-toe scan revealed no pertinent findings related to breast.

Vital Signs

- Temperature, 98.4° F
- Pulse, 72 beats per minute (BPM).
- Respirations, 18/min.
- Blood pressure (BP), 110/68.
- Height, 5'4".
- Weight, 116 lb.

Breasts

- Right breast slightly larger than left.
- Uniform and symmetrical.
- Ovoid in shape.
- No skin lesions noted.
- No redness or dimpling.
- Small, pea-sized (0.5 cm), easily movable, rubbery, smooth-edged lesion palpated in right breast at 2 o'clock; 4 cm from areola in the right upper outer quadrant.
- No lesions palpated in left breast.

Nipple

- No lesions observed or palpated.
- Nipples erect and point in same direction.
- No inversion or eversion.
- Denies discharge from nipple.
- Unable to elicit discharge from either nipple; no specimens sent.

Axillae

- No lesions or thickening noted.



Case Study Analysis and Plan

Now that you have completed a thorough assessment of Mrs. Kobrynski, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Your next step is to analyze the data from Mrs. Kobrynski's assessment and develop nursing diagnoses. The following are possible nursing diagnoses for Mrs. Kobrynski. Cluster the supporting data for each.

1. Knowledge deficit related to BSE
2. Ineffective Health Maintenance related to need for incorporation of preventive behaviors

3. Fear related to presence of lump and need for mammogram/potential for life-threatening disease
4. Potential disturbed body image



CRITICAL THINKING ACTIVITY # 4

Now that you have identified some nursing diagnoses for Mrs. Kobrynski, select one diagnosis and develop a nursing care plan and a related teaching plan, including learning outcomes and teaching strategies.

RESEARCH TELLS US

Fatty diets have been an identifiable risk factor for many types of cancers, including breast cancer. Metabolic syndrome is a condition characterized by obesity, glucose intolerance, low serum high-density lipoprotein cholesterol (HDL-C), high serum triglycerides, and hypertension. The incidence of metabolic syndrome is increasing and is parallel with increasing rates of breast cancer worldwide.

Furberg, Veierod, Wilsgaard, Bernstein, and Thune (2004) studied the relationship between HDL-C as a component of metabolic syndrome and risk for breast cancer. The participants in this study were based on a follow-up of 38,823 Norwegian women with repeated assessments of serum lipids, height, weight, diet, and lifestyle factors. The initial screening occurred from 1977 through 1983 and a second screening took place from 1985 to 1987. The women in the study were between 17 and 54 years of age. During the median 17.2 years follow-up, 708

cases of invasive breast cancer were identified. Multivariable analysis revealed that the risk of breast cancer was inversely related to HDL-C levels. The study showed that low HDL-C levels are associated with increased levels of cancer-promoting hormones, such as androgens, estrogens, insulin and insulin-like growth factor (IGF-I). It also showed that low HDL-C levels as part of metabolic syndrome are associated with an increased risk of postmenopausal breast cancer.

These findings may prove to be an important marker at identifying those at risk for developing breast cancer. When screening for risk factors of breast cancer, nurses should consider the presence of metabolic syndrome with low HDL-C levels. Furthermore, if metabolic syndrome is identified, clinicians should include weight management, diet, BP control, and activity and exercise programs into their health promotion teaching.

Source: Furberg, A.S., Veierod, M.B., Wilsgaard, T., Bernstein, L., Thune, I. (2004). Serum high-density lipoprotein cholesterol, metabolic profile, and breast cancer risk. *Journal of the National Cancer Institute* 96(15): 1152-1160.

HEALTH CONCERNS

BSE is now recognized as a means for women to become familiar with their breasts and notice changes (American Cancer Society, 2005). The American Cancer Society's current guidelines for early detection are age dependent.

Current Recommendations

- **Women ages 40 and older:** Annual mammogram, annual CBE (CBE by healthcare professional), monthly BSE optional.
- **Women ages 20 to 39:** CBE every 3 years, monthly BSE optional.

Mammography

Mammography is the only screening method found to reduce breast cancer deaths. It is the best method of early detection because it detects breast cancers before they can be felt. However, monthly BSE and regular clinical examinations by a trained medical professional still have value, as women generally don't have yearly mammography screenings until age 40 and because there are women who are without medical insurance or the financial means to have mammograms. In addition, a small percentage of breast cancers may not show up on mammogram.


(continued)

HEALTH CONCERNS (continued)

The American Cancer Society recommends that healthcare professionals inform women of the benefits and limitations of BSE and provide instruction if requested. If your patient decides to perform BSE, it should be performed monthly. The purpose of BSE is to allow women to detect breast changes from month to month. At first, a woman may not know how to interpret what she feels, but she will soon become familiar with what is normal for her. In premenopausal women, examination should occur 5 to 7 days after the menstrual cycle begins, or 3 to 5 days after it ends. In postmenopausal women, BSE should be performed on the same day each month—such as the first day of the month, the patient's birthday, or another meaningful day—so that the patient will be reminded. The patient should report unusual findings to her healthcare provider.

Follow these four steps when teaching BSE:

- **Step 1.** First, teach the patient how to look at herself in a mirror and, with her arms at her sides, check for any visible abnormalities. She should observe for dimpling, retraction, or breast flattening as she first elevates her arms slowly, then presses her hands against her hips, and finally, bends forward.



- **Step 2.** Next, by placing your hand over the patient's, show her how to use the pads of the middle three fingers of the opposite hand to palpate the breast systematically by compressing the breast tissue against the chest wall. She should palpate all portions of the breast, areola, nipple, tail of Spence, and axilla when she is in the shower or standing before a mirror. She should repeat the procedure lying down with a pillow or folded towel under the shoulder of the side she is examining.



- **Step 3.** Next, show her how to compress the nipple gently between the thumb and the index finger as she observes any discharge.



- **Step 4.** Finally, explain that she should report any redness or inflammation, swelling, masses, flattening, puckering, dimpling, retraction, sunken areas, asymmetrical nipple direction, discharge, bleeding, lesions, or eczematous nipple changes to her physician.



HELPFUL HINTS

- Some women do not feel comfortable touching their own breasts. Before teaching BSE, ascertain a woman's comfort level regarding touching herself.
- Suggest that the patient stand in front of a mirror to inspect her breasts.

Risk Factors for Breast Cancer

Breast cancer risk factors can be either modifiable or unmodifiable. As a nurse, you are in an ideal position to educate your patient as to the benefits of changing modifiable risk factors through healthy lifestyle changes.

Unmodifiable Factors

- **Age:** Risk increases with age.
 - **Race/ethnicity:** Prior to age 35, African American women have a higher incidence of breast cancer and a higher mortality rate from breast cancer than white women. After age 35, white women have higher incidence of breast cancer than African American women.
- **Personal or family history of breast cancer:** Women who have a first-degree relative (mother, sister, daughter) with a history of breast cancer have an increased risk. Risk increases when more than one family member has a history of breast cancer. Risk increases the younger the relative was at

the time of diagnosis. Family history of ovarian cancer, or personal history of endometrium, ovary, or colon cancer also increases risk. Being a carrier of either of two familial breast cancer genes called BRCA1 or BRCA2 increases the risk. Five to 10 percent of breast cancer cases result from BRCA1 or BRCA2 mutated genes.

- **Hormonal factors:** Women who are nulliparous or have their first child after age 30 are at increased risk.
 - Women who were never breast fed, experienced early menarche (earlier than age 12 years) or late menopause (after age 50 years), or suffered unopposed estrogen stimulation of breasts by BCP or HRT are at increased risk.

Modifiable Factors

- High-fat diet.
- Postmenopausal obesity.
- Use of postmenopausal hormones.
- Alcohol consumption.
- Physical inactivity.

Source: American Cancer Society, 2005 <http://www.cancer.org>

Common Abnormalities

ABNORMALITY

ASSESSMENT FINDINGS

Breast Cancer

- Nipple erosion; retraction; or discharge, usually bloody.
- Enlarged, shrunken, or dimpled breast with no pain.
- Nontender, firm or hard lump, irregularly shaped and fixed to skin or underlying tissues.
- Enlarged surrounding lymph nodes.
- Usually seen in white women over age 35 with family history of breast cancer; personal history of long menstrual cycles, early menarche, late menopause, or first pregnancy after 30.

Fibroadenoma of Breast

- Possible history of endometrial or ovarian cancer.
- Well-defined mass or masses in breast, no pain.
- Mass is round, firm, discrete, movable, $\frac{3}{8}$ to 2 inches (1 to 5 cm) in diameter.
- Usually solitary but may be multiple and bilateral.
- Patient usually in teens or early 20s.

Fibrocystic Disease

Includes mammary dysplasia, cystic adenosis, chronic cystic disease, cystic mastitis.

- Thickened, nodular areas in breast (usually bilateral); may be slight; pain and tenderness possible, especially premenstrually.
- May be single or multiple; mobile, well defined, tender; in upper outer quadrant.
- Exacerbated by caffeine intake.
- Cystic fluid (typically gray-green) from aspirated cysts.
- Patient in childbearing years.

Interductal Papilloma

- Serous or serosanguineous discharge from one nipple duct unilaterally, moderate pain.
- No palpable tumor or mass, or soft mass difficult to distinguish from surrounding tissue.
- Patient usually aged 35 to 55.

(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
Mammary Duct Ectasia (Dilatation)	<ul style="list-style-type: none"> ■ Nipple discharge and retraction, pain in affected areas, itching around nipple. ■ Subareolar ducts feel like rubbery lesions filled with pastelike material. ■ Enlarged regional lymph nodes possible.
Mastitis	<ul style="list-style-type: none"> ■ Patient usually in early-stage menopause. ■ Pain accompanied by breast tenderness. ■ Firm, tender, warm, and reddened area in affected breast. ■ History of cracked nipples. ■ Breast abscess. ■ Painful, enlarged axillary lymph nodes. ■ Patient usually third or fourth week postpartum.

ASSESSMENT

CHECKLIST

HISTORY

Remember to look at each history component as it relates to the breasts.

Ask the patient the following:

- Do you have:
 - Lump or mass?
 - Pain or tenderness?
 - Nipple discharge?
- Do you have a lump or thickening in or near your breast or under your arm that persists through the menstrual cycle?
- Is the skin on your breast or nipple red, dimpled, puckered, scaly, or inflamed?
- Do you have nipple changes? For example, a change in the direction in which one nipple points, inversion, eversion, or discharge?
- Has your breast changed in size, shape, or contour?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam.

- Small pillow
- Mirror
- Gloves
- Ruler
- Specimen slide and culture swab

Position

Supine with small pillow under shoulder of breast being examined.
Sitting arms at side, arms over head, hands on hips, leaning forward.

Technique

Remember to scan your patient from head-to-toe looking for signs related to the breasts.

Inspection, Palpation

Inspect (Different Positions)

Breasts

- Size, shape, symmetry
- Condition of skin, lesions, venous pattern

Nipple

- Inversion/eversion
- Direction
- Discharge

Axilla

- Condition of skin
- Lesions

Palpate

Breasts

(vertical strip, pie wedge, or concentric circles methods)

- Consistency
- Tenderness
- Masses

Nipple

- Elasticity
- Tenderness
- Discharge

Axilla and Nodes

- Palpate central, lateral, posterior, anterior, epitrochelar, and clavicular nodes

DOCUMENT


Document your findings and develop your plan of care.

S U M M A R Y

- Assessment of the breast provides invaluable information about the overall health of your patient.
- A primary goal of breast assessment is to detect changes in the breast.
- Early detection of breast disease leads to early treatment.
- For breast cancer, the earlier the diagnosis and treatment, the better the prognosis.
- Assessment of the breast also provides an ideal opportunity to teach BSE to your patient.

R E V I E W Q U E S T I O N S

1. Mary Jane McCormick, age 40, has discovered a lump in her right breast. As you obtain her health history, you identify risk factors for breast cancer. Which factor increases the risk for breast cancer?
 - a. History of fibrocystic breast disease
 - b. Multiparity
 - c. Breast fed her children
 - d. Her age
2. Mary Jane McCormick, age 40, has discovered a lump in her right breast. As you obtain her health history, you identify risk factors for breast cancer. Which diet would decrease the risk for breast cancer?
 - a. Low-calorie
 - b. Low-salt
 - c. Low-fat
 - d. Low-fiber
3. Donna Penner, age 56, is postmenopausal. You are reviewing BSE with her. At what time of the month should she perform BSE?
 - a. First of every month
 - b. Middle of the month
 - c. Last day of the month
 - d. Same time every month
4. Which is the most common site for breast cancer in men?
 - a. Upper outer quadrant in the tail of Spence
 - b. Areola
 - c. Upper inner quadrant
 - d. Lower outer quadrant
5. Part of the assessment of the breast includes assessing the lymph nodes. Which area would be best for palpating the central lymph nodes?
 - a. High in the middle of the axilla
 - b. High in the axilla on the lateral aspect of the humerus
 - c. High in the axilla toward the posterior axillary line
 - d. High in the axilla toward the anterior axillary line
6. Part of the assessment of the breast includes assessing the lymph nodes. Which area would be best for palpating the lateral lymph nodes?
 - a. High in the middle of the axilla
 - b. High in the axilla on the inner aspect of the humerus
 - c. High in the axilla toward the posterior axillary line
 - d. High in the axilla toward the anterior axillary line
7. Part of the assessment of the breast includes assessing the lymph nodes. Which area would be best for palpating the epitrochlear lymph nodes?
 - a. One inch above the olecranon process
 - b. High in the axilla on the lateral aspect of the humerus
 - c. High in the axilla toward the posterior axillary line
 - d. High in the axilla toward the anterior axillary line
8. Bill Brown, age 55, has a history of cardiac disease and is taking digoxin. How might digoxin affect his breast?
 - a. Gynecomastia
 - b. Galactorrhea
 - c. Mastalgia
 - d. Cystic breast
9. Mary Smith, age 20, has a benign fibrocystic cyst of the left breast. Which characteristic is associated with a fibrocystic breast disease?
 - a. Immobile
 - b. Hard consistency
 - c. Tender
 - d. Irregular shape
10. When inspecting the breast, which position is best for accentuating dimpling or retraction of breast tissue?
 - a. Sitting, hands at side
 - b. Sitting, hands over head
 - c. Sitting, leaning forward
 - d. Sitting, hands pressed on hips



Assessing the Abdomen

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent abdominal history questions
- Obtain a pertinent abdominal history
- Perform an abdominal physical assessment
- Document abdominal assessment findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

The abdominal assessment provides information about a variety of systems because every system, with the exception of the respiratory system, is found within the abdomen. The stomach, small and large intestines, liver, gallbladder, pancreas, spleen, kidneys, ureters, bladder, aortic vasculature, spine, uterus and ovaries, or spermatic cord are all located in the abdomen. Not only does assessment of the abdomen enable you to obtain valuable information about the functioning of the gastrointestinal (GI), cardiovascular, reproductive, neuromuscular, and genitourinary systems; it can also provide vital information about the health status of every other system.

Anatomy and Physiology Review

Before you begin your assessment, an understanding of the anatomy and physiology of abdominal structures is essential. You must be able to recognize normal structures before you can identify abnormal findings. Recognizing the structures will enable you to perform the assessment accurately, and understanding the physiology will guide your assessment and allow you to interpret your findings.

Structures and Functions

The major system assessed in the abdominal examination is the GI or digestive system. The digestive system is responsible for the ingestion and digestion of food, absorption of nutrients, and elimination of waste products.

The primary structures of the digestive system (Fig. 17.1) include the mouth, pharynx, esophagus, stomach,

small intestines (duodenum, jejunum, and ileum), large intestines (cecum, colon [ascending, transverse, descending, and sigmoid]), and rectum. These main structures of the digestive system form a hollow tube that is actually outside the internal environment of the body even though it is located inside the body. This tube, referred to as the **alimentary canal** or the **gastrointestinal tract**, begins at the mouth and ends at the anus.

The digestive system also contains accessory organs that aid in the digestion of food. The accessory organs of the digestive system include the salivary glands (parotid, submandibular, and sublingual), liver, gallbladder, and pancreas.




FIGURE 17.1. Abdominal organs and structures.

Primary Digestive Organs/Structures and Functions

ORGAN STRUCTURE	PRIMARY FUNCTION
Mouth	<ul style="list-style-type: none"> ■ Masticating (chewing) food particles and mixing them with saliva. ■ Primarily mechanical digestion and the beginning of chemical digestion.
Pharynx	<ul style="list-style-type: none"> ■ Swallowing food particles into esophagus.
Esophagus	<ul style="list-style-type: none"> ■ Propelling food downward into stomach
Stomach	<ul style="list-style-type: none"> ■ Storing food until it can be moved farther along GI tract. ■ Parietal cells secrete hydrochloric acid to aid in digestion. ■ Mucus cells secrete substances to coat the stomach. ■ Chief cells secrete pepsinogen, which is converted to pepsin, which in turn digests protein. ■ Churns food and breaks it down into small particles, then mixes them with gastric juices. ■ Secretes gastrin, which stimulates secretion of acid and pepsinogen and increases gastric motility.

ORGAN STRUCTURE	PRIMARY FUNCTION
<p>Small Intestine</p> <ul style="list-style-type: none"> ■ Duodenum: 25 cm (10 inches) long ■ Jejunum: 2.5 m (8 ft) long ■ Ileum: 3.5 m (12 ft) long 	<ul style="list-style-type: none"> ■ Secretes intrinsic factor that protects vitamin B₁₂ from stomach acids and facilitates absorption of B₁₂ across the membranes of the small intestines (parietal cells). ■ Absorbs some water, alcohol, and certain drugs. ■ Destroys some bacteria found in foods. ■ Site for both mechanical and chemical digestion. ■ Bile and pancreatic juices facilitate absorption of nutrients in small intestines. ■ Primary site for digestion, especially chemical digestion. ■ Enzymes include: <ul style="list-style-type: none"> ■ <i>Enterokinase</i>: Converts trypsinogen to active trypsin. ■ <i>Peptidases</i>: Help break down proteins. ■ <i>Maltase, lactase, and sucrase</i>: Break down carbohydrates. ■ Hormones include: <ul style="list-style-type: none"> ■ <i>Cholecystokinin</i>: Secreted from duodenal wall; stimulates gallbladder to secrete pancreatic enzymes. ■ <i>Gastric inhibitory peptide</i>: Inhibits gastric motility. ■ <i>Secretin</i>: Secreted by duodenal wall; stimulates pancreatic secretions to neutralize gastric acid.
<p>Appendix</p>	<ul style="list-style-type: none"> ■ Function is unknown. ■ Narrowest part of intestines and a frequent site for bacteria and indigestible matter to become trapped leading to inflammation (appendicitis). ■ May serve as a breeding ground for intestinal bacteria because it contains large amounts of lymphatic tissue.
<p>Large Intestine</p> <ul style="list-style-type: none"> ■ 1.5–1.8 m (5–6 ft) long 	<ul style="list-style-type: none"> ■ Absorbs salt and water and excretes waste products of digestive process (feces) from the rectum (defecation). ■ Helps synthesize vitamins B₁₂ and K.

Accessory Digestive Organs/Structures and Functions

ORGAN STRUCTURE	PRIMARY FUNCTION
<p>Salivary Glands</p> <ul style="list-style-type: none"> ■ Parotid, submandibular, sublingual 	<ul style="list-style-type: none"> ■ Produce saliva that moistens and lubricates food. ■ Secrete amylase, which converts starches to maltose.
<p>Liver</p>	<ul style="list-style-type: none"> ■ Produces and secretes bile to emulsify fats. ■ Metabolizes protein, carbohydrates, and fats. ■ Converts glucose to glycogen, which is stored in the liver. ■ Produces clotting factors and fibrinogen. ■ Produces plasma proteins such as albumin ■ Detoxifies a variety of substances such as drugs and alcohol ■ Stores vitamins A, D, E, K, and B₁₂ and minerals iron and copper ■ Converts conjugated bilirubin from blood to unconjugated bilirubin
<p>Pancreas</p> <ul style="list-style-type: none"> ■ 12–15 cm (6–9 inches) long 	<ul style="list-style-type: none"> ■ As an endocrine gland, the pancreas secretes: <ul style="list-style-type: none"> ■ Beta cells, which secrete insulin to regulate blood sugar levels. ■ Alpha cells, which secrete glucagons that store carbohydrates. ■ Delta cells, which secrete somatostatin, the hypothalamic growth-inhibiting hormone that inhibits insulin and glucagon secretion. ■ F cells, which secrete pancreatic polypeptide that regulates the release of pancreatic enzymes.

(continued)

Accessory Digestive Organs/Structures and Functions (continued)

ORGAN STRUCTURE

PRIMARY FUNCTION

Gallbladder

- 7–10 cm (3–4 inches) long and 3 cm wide.

- As an exocrine gland, the pancreas secretes the following digestive enzymes and alkaline materials:
 - Acinar units, which secrete digestive enzymes.
 - Amylase, which digests starches into maltose.
 - Lipase, which breaks down lipids into fatty acids and glycerol.
 - Pancreatic proteolytic enzymes (trypsinogen, chymotrypsinogen, and procarboxypeptidase), which are secreted in inactive form and activated in small intestine.
- Stores and concentrates bile.

The Digestive Process

The digestive process consists of **mechanical digestion**, the breakdown of food through chewing, peristalsis, and churning; and **chemical digestion**, the breakdown of food through a series of metabolic reactions with enzymes.

The digestive process begins in the mouth, where food is taken in and masticated. The bolus of food is then swallowed into the esophagus, where it is propelled slowly via peristaltic contraction to the stomach. In the stomach, the food bolus is churned, breaking it down further into smaller particles and mixing it with digestive juices and hydrochloric acid that is produced by the stomach. The food bolus becomes chyme and progresses down into the first portion of the small intestine, called the **duodenum**. In the duodenum, pancreatic juices and bile are secreted in the chyme. The food then enters the **jejunum** and **ileum**, where nutrients are absorbed into the circulatory system. Food particles that are not absorbed by the small intestines proceed into the large intestine, where they are eventually excreted as feces.

Additional Abdominal Structures

Along with the organs of the digestive system, the abdomen also contains the spleen; the urinary tract including the bladder, kidneys, and ureters; the uterus and ovaries; the aorta; and the iliac, renal, and femoral arteries. The uterus and ovaries are covered in Chapter 18, Assessing the Female Genitourinary System. The other abdominal organs are shown in Figure 17.2.

The abdominal cavity has a serous membrane called the **peritoneum**, which covers the organs and holds them in place. The peritoneum contains a parietal layer that lines the walls of the abdomen and the visceral pleura, which coats the outer surface of the organs. A small amount of fluid between these membranes allows them to move smoothly within the cavity.




FIGURE 17.2. Other abdominal organs and structures.

Interaction With Other Body Systems

The GI system requires the proper functioning of the nervous, endocrine, respiratory, cardiovascular, integumentary, and musculoskeletal systems in order to operate at its full capacity.

The Integumentary and Musculoskeletal Systems

The digestive system is protected and supported by the musculoskeletal and integumentary systems. The musculoskeletal system also assists with ingestion, mastication, deglutition (swallowing) of food, and eventual defecation of its byproducts.

Other Abdominal Organs and Structures

ORGAN STRUCTURE	PRIMARY FUNCTION
Spleen	<ul style="list-style-type: none"> ■ Stores red blood cells (RBCs) and platelets. ■ Produces RBCs and macrophages. ■ Activates B and T lymphocytes.
Urinary Tract, Bladder, Kidneys	<ul style="list-style-type: none"> ■ Maintain homeostasis of fluid-and-electrolyte and acid-base systems. ■ Control BP through secretion of renin. ■ Secrete erythropoietin to stimulate RBC production. ■ Remove waste products from body.
Spine	<ul style="list-style-type: none"> ■ Supports weight of body. ■ Protects spinal cord.
Spermatic Cord	<ul style="list-style-type: none"> ■ Protects the vas deferens, blood vessels, lymph, and nerves that run from the scrotum to the penis.
Uterus and Ovaries	<ul style="list-style-type: none"> ■ Ovaries produce ova and secrete estrogen and progesterone. ■ The uterus allows the sperm to reach the uterine tubes, where it can make contact with an ovum. If conception occurs, the uterus develops an environment in which the embryo can grow.
Vasculature of the Abdomen: Aorta and Inferior Vena Cava	<ul style="list-style-type: none"> ■ Supplies oxygenated blood to the cells and organs in the lower half of the body.

The Respiratory and Cardiovascular Systems

The respiratory and cardiovascular systems provide the oxygen needed for the digestive organs to function. The respiratory system gets oxygen for the cells of the body and rids the body of carbon dioxide. All the cells in the body, including those of the digestive system, need oxygen to function appropriately. The cardiovascular system circulates the oxygen-rich blood to all the cells in the body. Any decrease in oxygen to the cells of the digestive system affects organ function. For example, if blood flow to the bowel is disrupted, a bowel infarct can occur, causing the bowel to stop functioning.

The Neurological System

The neurological system plays an important role in digestion. When the body is in a parasympathetic response, or the “rest and repair” phase, the neurological system releases acetylcholine, the neurotransmitter for the parasympathetic system. In relation to the digestive system, acetylcholine stimulates the secretion of digestive juices and increases peristalsis.

The opposite is true for the sympathetic response. The sympathetic system is stimulated at times of physical or psychological stress. When this system is stimulated, a “fight or flight” response occurs, causing the release of norepinephrine, which produces a decrease in peristalsis and secretion of digestive juices. Therefore, the digestive system functions to its maximum capacity when it receives parasympathetic responses from the peripheral nervous system.

The Endocrine System

The secretion of digestive juices also depends on the proper functioning of the pancreas, an organ that has both endocrine and exocrine functions. The endocrine function is to release insulin, glucagon, and gastrin into the bloodstream to assist in carbohydrate metabolism. The exocrine function is to secrete bicarbonate and pancreatic enzymes into the duodenum to aid in the digestion of proteins, fats, and carbohydrates.


Developmental, Cultural, and Ethnic Variations

Infants

Some anatomical characteristics in infants are different from those in adults. It is important to note that the newborn’s bladder is located above the symphysis pubis. The liver proportionately takes up more space in the abdomen and may extend 2 cm ($\frac{3}{4}$ inch) below the rib cage. Because the infant’s abdominal muscles are weak, the abdomen normally protrudes (see Chapter 24, Assessing the Newborn and Infant).

Children

A child’s abdomen is proportionately larger than an adult’s and has a slightly protuberant appearance because of the curvature of the back. This protuberance is most obvious in toddlers and preschoolers and diminishes to adult proportions during adolescence. Abdominal respiration is common in most children. Because the abdominal



Relationship of GI System to Other Systems

muscles are underdeveloped, the organs are more easily palpated (see Chapter 26, Assessing the School-Age Child and Adolescent).

Pregnant Women

Many changes occur in the abdomen of the pregnant patient. The abdominal muscles relax, allowing the uterus to protrude during a normal pregnancy. For women having multiple pregnancies or a multiple birth, the rectus abdominis muscles, which are located medially on the anterior aspect of the abdomen, may become separated (diastasis recti abdominis). As the fetus grows, it takes up even more room in the uterus, causing the stomach to rise up and impinge on the diaphragm.

Also, bowel sounds are diminished in the pregnant patient because the bowels are compressed by the fetus. This decreased activity in the lower GI tract, along with the ingestion of prenatal vitamins containing large amounts of iron, contributes to constipation. Because of increased venous pressure in the lower abdomen, hemorrhoids may develop and cause further problems with elimination.

It is important to note that the location of the appendix in the pregnant woman is different from that in a nonpregnant woman. During pregnancy, the appendix is displaced upward and laterally to the right. This can complicate the diagnosis of appendicitis.

Pregnant patients have several particular alterations in their integumentary system. One characteristic skin change is the *linea nigra*, a darkly pigmented line that appears on the anterior abdomen from the pubis to the umbilicus in most pregnant women. Near the end of pregnancy, the umbilicus may become everted and *striae* (stretch marks) may develop on the skin of the abdomen from the increased tension of the expanding uterus (see Chapter 23, Assessing the Mother-to-Be).

Older Adults

As people age, many of their body systems slow down and become less efficient. Elderly people have changes in dentition that may affect their chewing ability and digestion. Poorly fitted dentures may result in painful mastication, which causes the patient to select foods that are easier to chew but not necessarily nutritionally balanced. In the GI tract, there is a reduction of saliva, stomach acid, gastric motility, and peristalsis that causes problems with swallowing, absorption, and digestion. These changes, along with a general reduction of muscle mass and tone, also contribute to constipation. Fat accumulates in the lower abdomen in women and around the waist in men, making physical assessment of the organs a little more challenging. The liver becomes smaller and liver function declines, making it harder to process medications (see Chapter 27, Assessing the Older Adult).



HELPFUL HINTS

- Be aware that older adults have a diminished response to painful stimuli that may mask abdominal health problems.
- Older adults may have difficulty assuming some positions necessary for the physical examination, so modify positions to meet their needs.

People of Different Cultures and Ethnic Groups

Some health problems are considered culturally or racially related. For instance, sickle cell anemia is seen almost exclusively in African Americans. Ashkenazi Jews have a greater incidence of colon cancer than other groups because they are believed to have a gene linked to the development of familial colorectal cancer.

Culturally Related Health Problems With Gastrointestinal Symptoms

CULTURAL GROUP	HEALTH PROBLEM	GASTROINTESTINAL SYMPTOMS
African Americans	<ul style="list-style-type: none"> ■ Sickle cell anemia ■ Obesity generally higher in racial/ethnic minorities than in whites. Obesity rates highest in non-Hispanic black women. ■ Lactose intolerance 	<ul style="list-style-type: none"> ■ Splenomegaly and jaundice ■ <i>Sickle cell crisis</i>: Acute abdominal pain, vomiting ■ Weight > 20 percent ideal weight ■ Abdominal cramping, diarrhea

(continued)

Culturally Related Health Problems With Gastrointestinal Symptoms (continued)

CULTURAL GROUP	HEALTH PROBLEM	GASTROINTESTINAL SYMPTOMS
Asian Americans	<ul style="list-style-type: none"> ■ GI cancer 	<ul style="list-style-type: none"> ■ Anorexia, bowel and digestion problems, pain, problems with weight loss
Jewish Americans	<ul style="list-style-type: none"> ■ Lactose intolerance ■ Crohn's disease ■ Ulcerative colitis ■ Colon cancer 	<ul style="list-style-type: none"> ■ Abdominal cramping, diarrhea ■ Abdominal pain, diarrhea ■ Abdominal pain, diarrhea, rectal bleeding ■ Changes in bowel habits, blood in stool, constipation
Mediterranean Americans (Greek and Italian Descent)	<ul style="list-style-type: none"> ■ Lactose intolerance ■ Thalassemia, anemia 	<ul style="list-style-type: none"> ■ See symptoms above ■ Jaundice, splenomegaly
Native Americans	<ul style="list-style-type: none"> ■ Alcoholism, liver disease, pancreatitis ■ Diabetes ■ Gallbladder disease 	<ul style="list-style-type: none"> ■ Jaundice, anorexia, ascites, pain, steatorrhea ■ Polyuria, thirst, weakness, weight loss, itching ■ Pain

Introducing the Case Study



Anne Robichaud is a 56-year-old divorced mother of two grown children who came to see her nurse practitioner for an annual physical examination. She is a Roman Catholic of French Canadian descent who speaks English and French and works in a local mill. Mrs. Robichaud states, "I'm constipated all the time." She has a family history of colon cancer. Considering her chief complaint and the fact that she has not had a complete physical for several years, a complete health history and physical examination are warranted.

perception of his or her health status. The health history interview includes a broad range of questions so that possible problems associated with each of the systems of the abdomen may be identified. Remember that information collected as part of the health history may uncover problems related to systems outside the abdomen (e.g., myocardial infarction [MI]). If time is an issue and you are unable to perform a complete health history, perform a focused history on the abdomen.

Biographical Data

Gathering biographical information can provide valuable insights about the patient's health status in several ways. Certain age groups are at greater risk for problems in the GI system. For example, infants and toddlers have a higher incidence of hernias than older children. Preschoolers are more likely to get parasitic infections, and teenagers may have abdominal symptoms as a result of pregnancy, sexually transmitted diseases (STDs), eating disorders like anorexia nervosa or bulimia, and infectious mononucleosis. Appendicitis occurs more frequently in children and teenagers than it does in adults. Older adults commonly develop problems with digestion, absorption, metabolism, and elimination because of changes caused by the aging process. Women aged 65 and over are commonly diagnosed with hiatal hernia, constipation, and diverticulosis.

Certain diseases occur more frequently in some races and cultures (see previous section). You will need to ask additional health history questions to determine

Performing the Abdominal Assessment

Assessment of the abdomen involves obtaining a complete health history and performing a physical examination. As you assess the patient, be watchful for signs and symptoms of actual and potential problems involving the different organs and structures in the abdomen.

Health History

The health history precedes the physical examination and involves interviewing the patient about his or her

Focused Abdominal History

If there is little time to do a complete history, a focused history can provide a baseline of information. Identify current symptoms related to digestion and elimination as well as risk factors for GI disease by asking the following questions, using the PQRST format.

- Do you have any abdominal pain?
- Have you ever had the following: stomach ulcer, hemorrhoids, hernia, bowel disease, cancer, hepatitis, cirrhosis, or appendicitis?
- Have you had abdominal surgery? If so, when, what type, and were there any subsequent problems?
- Do you have a family history of ulcer, gallbladder disease, bowel disease, or cancer?
- Do you have any problems with swallowing, heartburn, nausea, yellowing of your skin, gas, bloating, or vomiting? (Note onset, quality, and quantity.)
- Do you have any food allergies or lactose intolerance?
- Have you noticed any recent weight changes? What is your usual weight and height?
- How is your appetite? What did you eat in the last 24 hours?
- How is your health usually?
- Are you currently being treated for a health problem? If so, what?
- How often do you usually have a bowel movement? Have you noticed any changes in your bowel movements?
- Are you having problems with diarrhea, constipation, hemorrhoids, or fecal incontinence? Have you ever noticed blood in your stool or had black, tarry stools?
- How often do you urinate? Do you have incontinence or burning when you urinate?
- When was your last menstrual period?
- Do you smoke? How many packs a day? (Calculate pack-years.)
- Do you drink alcohol? If so, how often? Do you use street drugs?
- How many cups of coffee, tea, or caffeinated soda do you drink every day?
- Have you been exposed to an infectious disease recently?
- What is your occupation?
- Have you been immunized against hepatitis B?
- Have you ever had a blood transfusion? If so, when?
- Do you take any prescribed medications? What are they?
- Do you have any allergies to medications?
- What OTC medicines or herbal preparations do you use?
- Do you use antacids, laxatives, enemas, NSAIDs, or aspirin?
- What home remedies do you use?

whether symptoms of these diseases are present so that appropriate screening measures can be performed, if necessary.

The potential for exposure to environmental and occupational hazards can also be discovered in the biographical data. Where a person lives or works may raise questions about environmental hazards such as lead exposure in children (from inhalation of lead-based paint dust in older houses) or occupational health hazards such as chemical exposure (arsenic, benzene).

Current Health Status

If your patient has an abdominal complaint, investigate this first. Common chief complaints involving the body systems in the abdomen include:

- *Lymphatic:* Swelling, lymph node tenderness.
- *Digestive:* Anorexia, bruising, constipation, diarrhea, distension, dysphagia, epigastric burning, gastric reflux, indigestion, jaundice, nausea, vomiting, pain, weight changes.
- *Reproductive:* Cramping, nausea, pain, vomiting, weight gain.
- *Neurological:* Pain.
- *Cardiovascular:* Pain.
- *Urinary:* Edema, pain, problems with urination (burning, frequency).



HELPFUL HINTS

- Rapid onset of symptoms usually indicates a serious problem. Note date and time pain started.
- Women of childbearing age with abdominal complaints should always be asked about the possibility of pregnancy.

The most common abdominal complaints—pain, changes in weight, changes in bowel habits (constipation, diarrhea), indigestion, nausea, and vomiting—are analyzed in the subsequent text, using the PQRST format. The nature and intensity of the symptoms dictate the order and extent of questioning during the symptom analysis.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Abdominal Pain

The most common complaint related to the abdomen, pain is often classified as visceral, parietal, or referred. **Visceral pain** results from distension of the intestines

or stretching of the solid organs. It is often described as burning, cramping, diffuse, and poorly localized. **Parietal pain** results from inflammation of the parietal peritoneum. The pain is usually severe, localized, and aggravated by movement. **Referred pain** is felt at a site away from the site of origin. Impulses from the internal organs and structures that share nerve pathways inside the central nervous system explain the nature of referred pain.



HELPFUL HINT

When assessing severity of pain, use an accepted pain intensity rating scale consistently. Be sure to document what tool you used and the words the patient uses to describe the pain's intensity.

Acute abdominal pain (“acute abdomen”) may indicate a life-threatening abdominal condition that requires immediate medical intervention. In this situation, you should assess the patient’s vital signs to determine whether she or he is in imminent danger. Vital signs provide information about the possibility of cardiac irregularities and reveal symptoms of shock and signs of an infectious process such as peritonitis. In addition, you need to prioritize the symptom assessment questions to elicit the most essential information. The order of symptom assessment becomes RTQSP.

Pain Location

The location of the pain is often diagnostically significant. Some disorders have classic signs located in

specific regions of the abdomen. For instance, pain in the umbilical region may indicate an abdominal aortic aneurysm or early appendicitis. Abdominal problems may also cause referred pain to the chest, so chest pain can indicate either an abdominal problem or a cardiac event. Patients with a gastric ulcer can have pain in the upper epigastric region left of midline, which is also the location for angina and MI. Patients with gastroesophageal reflux disease (GERD) may have chest pain that radiates to the back, neck, or jaw, which also mimics an MI. Patients with a hiatal hernia may complain of substernal chest pain and difficulty breathing, especially after a meal. Note location of pain by quadrant or region:

- *Pain in shoulder:* Ruptured spleen, ectopic pregnancy, Pancreatitis.
- *Pain in scapula:* Cholelithiasis, MI, angina, biliary colic, pancreatitis.
- *Pain in thighs, genitals, lower back:* Renal problems, ureteral colic.
- *Pain in lower and middle back:* Abdominal aortic aneurysm.

Recognizing the relationship between the location of the pain and the possible health problem has important implications for immediate nursing assessment and care of the patient.



ALERT

If a patient presents with chest pain or indigestion, rule out cardiac disease first.

Significance of Pain by Abdominal Quadrant

Left Upper Quadrant	Right Upper Quadrant
<ul style="list-style-type: none"> ■ <i>Heart:</i> MI/ischemia ■ <i>Lungs:</i> Pulmonary embolism, pneumonia ■ <i>Pancreas:</i> Pancreatitis ■ <i>Spleen:</i> Ruptured spleen ■ <i>Stomach:</i> Gastric ulcer, esophagitis, GERD, hiatal hernia, varices 	<ul style="list-style-type: none"> ■ <i>Heart:</i> MI/ischemia ■ <i>Lungs:</i> Pneumonia ■ <i>Gallbladder:</i> Cholelithiasis, cholecystitis ■ <i>Liver:</i> Hepatitis, cancer ■ <i>Intestines:</i> Duodenal ulcer
Left Lower Quadrant	Right Lower Quadrant
<ul style="list-style-type: none"> ■ <i>Ovary/uterus:</i> Ectopic pregnancy, ovarian cyst, pelvic inflammatory disease (PID) ■ <i>Intestines:</i> Perforation, constipation, diverticulitis, hernia, ulcerative colitis ■ <i>Kidney:</i> Nephrolithiasis (kidney stones), infection 	<ul style="list-style-type: none"> ■ <i>Ovary/uterus:</i> Ectopic pregnancy, ovarian cyst, PID ■ <i>Intestines:</i> Perforation, obstruction, constipation, Crohn's disease, diverticulitis, hernia, ulcerative colitis ■ <i>Appendix:</i> Appendicitis ■ <i>Kidney:</i> Nephrolithiasis, infection

Terms Used to Describe Quality and Radiation of Pain

Quality

- Aching
- Burning
- Colicky
- Dull
- Nagging
- Piercing
- Pressure
- Cramping
- Pulsing
- Sharp
- Spasms
- Squeezing
- Stabbing
- Throbbing
- Viselike
- Waves

Radiation

- Radiating
- Shooting

How Patients With Eating Disorders Lose Weight

- *Diets*: Reducing calories by stopping or skipping meals.
- *Cathartics*: Abusing laxatives, enemas, suppositories, diuretics.
- *Exercise*: Excessive exercise to burn off calories.
- *Purgatives*: Using ipecac or self-induced vomiting.

A careful analysis of this symptom provides data that allow the nurse to distinguish between medical and behavioral problems causing the weight change. Weight changes of 2 to 3 lb (1 to 1.4 kg) within 48 hours result from fluid changes. Unexplained weight loss in an adult should raise suspicions of underlying malignancy.

Change in Bowel Patterns

Alterations in bowel movements are associated with a variety of GI disorders, such as malabsorption disorders, irritable bowel syndrome, cancer, infections (viral, bacterial, parasitic), food intolerance, and reactions to medications, as well as non-GI disorders. To determine whether a patient is having health problems that affects bowel function, first establish a baseline by asking general questions about bowel habits, such as: “How often do you have bowel movements? Do you have any problems with your bowels, such as straining, pain, constipation, or diarrhea?” Then ask more specific questions to help identify the origin of the problem.

Bowel patterns range from two movements per day to two or three per week. Identify the color of the stool:

- *Black, tarry*: Upper GI bleeding.
- *Red, bloody*: Lower GI bleeding.
- *Clay colored*: Increased bile in obstructive jaundice.



HELPFUL HINTS

- Remember to assess each new complaint as it arises in the patient history and physical examination.
- When assessing palliative or provocative factors, ask about the patient's use of culturally related remedies. These remedies may be tried first, delaying the initiation of traditional Western treatment. For some illnesses, a delay in treatment may have serious repercussions.



CRITICAL THINKING ACTIVITY #1

How would you explain the pain associated with appendicitis, beginning first in the umbilical region, then localizing to the right lower quadrant (RLQ)?

Weight Change

Weight change may indicate diseases in many body systems, reflect unhealthy behaviors, or even reveal a normal state such as pregnancy. Weight changes can be a sign of GI disease, cancer, congestive heart failure with fluid retention, metabolic or endocrine disorders, unhealthy lifestyles, major depressive disorder, and eating disorders.

Common Causes of Constipation

- Medications
- Lack of exercise
- Lack of dietary fiber
- Lack of response to urge to defecate
- Loss of abdominal muscle tone
- Cancer
- Depression
- Dehydration
- Impaction
- Lead poisoning
- Intestinal obstruction
- Hypothyroidism

Diarrhea (frequent loose bowel movements) can have a variety of causes: diet; medications; viral or bacterial infections; and medical problems such as ulcerative colitis, irritable bowel syndrome, and Crohn's disease.



HELPFUL HINTS

- If your patient has abdominal cramps, diarrhea, nausea, vomiting, and fever lasting 7 to 10 days, he or she probably has gastroenteritis caused by a virus.
- Because diarrhea and other GI disorders cause fluid and electrolyte imbalance, be sure to monitor your patient's input and output (I&O) and serum electrolytes.
- Diarrhea assessment tools such as the National Cancer Institute Scale of Severity of Diarrhea are helpful to use when assessing patients, especially those with ongoing episodes of diarrhea who are taking antineoplastic drugs.



ALERT

- Never give a cathartic to a patient with abdominal pain, nausea, and constipation who also has absent bowel sounds.
- Extended hospitalization, abdominal surgery, and antibiotics (especially cephalosporins) increase the patient's risk for developing pseudomembranous colitis. This potentially life-threatening disorder causes abdominal cramps, fever, and watery or bloody diarrhea.
- Prolonged diarrhea can cause loss of fluids and electrolytes such as potassium. Assess for dehydration by checking skin turgor, taking orthostatic blood pressure (BP) and pulse and observing for signs of hypokalemia, such as cardiac arrhythmias and muscle weakness.
- If your patient complains of indigestion, take her or his vital signs immediately. People having an MI commonly complain of indigestion.

Indigestion

Indigestion—also called *dyspepsia* or *pyrosis*—is a frequent abdominal complaint that is usually described as “heartburn.” This burning sensation is usually worse after eating a meal. Acid from the stomach flows into the lower esophagus, causing the burning sensation. GERD has heartburn as its chief symptom, but the epigastric dis-

stress occurs more frequently, lasts longer, and has more severe symptoms than indigestion. Heartburn is also a common complaint in both gastric ulcer and duodenal ulcer disease and gallbladder disease. Indigestion that increases when the person is lying flat may indicate a hiatal hernia or GERD. Indigestion associated with belching (eructation) and flatulence suggests cholecystitis.

Nausea

Nausea is caused by stresses on the stomach wall or esophagus. Distension, alterations in peristalsis, negative olfactory stimulation, inner ear problems, or medications can also cause nausea. Many GI medical conditions have nausea as an assessment finding.

Vomiting

During **vomiting**, peristalsis is reversed and the esophageal sphincter opens to allow the contents of the stomach to be ejected. The involuntary emptying of stomach contents is caused by irritation of the stomach lining caused by chemicals, trauma, or distension; stimulation of the vomiting center in the brain (medulla); and head injury. Some GI conditions that cause vomiting are intestinal obstruction, peptic ulcer, viral or bacterial infection, and appendicitis. A person with repeated vomiting is always at risk for fluid and electrolyte problems.



ALERT

Projectile vomiting without nausea (unexpected vomiting) results from central stimulation of the vomiting center (medulla) and is a sign of head injury or neurological or vascular abnormalities in the brain.



HELPFUL HINT

Coffee-ground-colored emesis indicates the presence of blood in the stomach.

Past Health History

This section of the health history involves asking questions about childhood and adult illnesses, injuries, hospitalizations, allergies, immunizations, and medications that can affect the abdominal structures. Remember to document specific dates in the patient's record.

(Text continued on page 569)

Past Health History

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Childhood Illnesses

- Did you have chickenpox (varicella)?
- Did you have digestive problems?
- Did you have malabsorption diseases?
- Did you have an eating disorder?
- Do you have sickle cell anemia?

Hospitalizations

- Have you ever been hospitalized for GI problems?
- Have you ever had any GI diagnostic tests, such as guaiac tests, colonoscopy, computed tomography (CT), magnetic resonance imaging (MRI), GI series, or barium swallow?

Surgeries

- Have you had any of the following, and if so, when?
- Abdominal surgery, such as an appendectomy?
- Blood transfusions?
- Recent insertion of GI tubes?

Serious Injuries

- Have you had recent trauma, such as auto accident, workplace injury, or sports injury?

Serious/Chronic Illnesses

- Have you had cancer?
- Do you have inflammatory bowel disease (Crohn's disease)?
- Hypertension?
- Heart disease?
- HIV or acquired immunodeficiency syndrome (AIDS)?

- Varicella always precedes herpes zoster or "shingles" (vesicular rash on lower aspect of rib cage in hypochondriac regions of the abdomen).
- May indicate a pattern of GI problems or risk for developing megacolon.
- May indicate celiac disease (impaired GI absorption related to wheat and rye ingestion).
- Eating disorders often begin during adolescence and continue into adulthood.
- Abdominal pain is associated with sickle cell crisis.
- Present symptoms may relate to previously diagnosed condition.
- Surgical procedures increase risk for adhesions, infections, obstructions, and faulty vitamin absorption.
- Rule out appendix as origin of current abdominal problem.
- Hepatitis and human immunodeficiency virus (HIV) are transmitted via blood.
- Increases risk for infection.
- Assess signs and symptoms of infection.
- Can injure organs such as the spleen.
- Symptoms may develop hours or days after an accident.
- Hemorrhage is a danger.
- Cancer in one site in the body may spread to other locations. The liver is a common site for cancer metastasis.
- Cancer of the stomach, intestines, colon, pancreas, spleen, ovaries, and uterus is often associated with abdominal complaints, including changes in bowel habits, abdominal pain, and weight loss.
- High risk for developing colorectal cancer.
- Increases risk for abdominal aortic aneurysm.
- Cardiac problems, including angina and MI, often present with indigestion, abdominal pain or pressure, and nausea and vomiting.
- Right-sided heart failure increases systemic venous pressure, which causes spleen and liver enlargement and GI complaints.
- HIV patients are at high risk for developing Kaposi's sarcoma of the abdominal organs, infections, and multiple GI problems, such as anorexia, weight loss, and cachexia.

(continued)

Past Health History (continued)

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Diabetes? <p>Immunizations</p> <ul style="list-style-type: none"> ■ Have you had hepatitis A or B series immunizations? <p>Allergies</p> <ul style="list-style-type: none"> ■ Do you have allergies to any medications or foods? 	<ul style="list-style-type: none"> ■ Antiviral agents used to treat HIV have side effects that compound these problems. ■ Diabetes can result in gastroparesis. ■ Note dates of all immunizations. ■ Discuss need for other primary prevention activities. ■ Note all allergens and responses. Milk (lactose intolerance), wheat, chocolate, peanuts, and shellfish are common food allergens. Symptoms of food allergies range from GI upset with diarrhea and vomiting to hives to life-threatening anaphylaxis.
	 ALERT
<p>Medications</p> <ul style="list-style-type: none"> ■ Are you currently taking any medication? If so, what are you taking, and what are the dosages and frequencies? ■ Are you taking any OTC medications, such as nutritional supplements, antacids, antihistamines, and laxatives? 	<ul style="list-style-type: none"> ■ Many medications have GI effects. Opioids, psychiatric medications, and iron supplements cause constipation (see <i>Drugs That Adversely Affect the Gastrointestinal System</i>). ■ NSAIDs and aspirin increase the risk for GI bleeding. ■ Overuse of laxatives may contribute to bowel problems.



ALERT

Foods do not have to be eaten to elicit an allergic reaction. Just smelling or touching the food is enough to produce an allergic reaction in some people.

Drugs That Adversely Affect the Gastrointestinal System

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Analgesics	Acetaminophen Aspirin	Hepatic necrosis with high (toxic) doses GI disturbances, GI bleeding, ulceration
Antacids	Aluminum hydroxide Calcium carbonate Magnesium hydroxide, magnesium oxide	Constipation Constipation, gastric hypersecretion, acid rebound Diarrhea
Anticholinergic agents	All anticholinergics	Nausea, vomiting, constipation, xerostomia (dry mouth), bloated feeling, paralytic ileus
Antidepressants	Amitriptyline hydrochloride, nortriptyline hydrochloride Selective serotonin reuptake inhibitors	Constipation, adynamic ileus, elevated liver enzyme concentrations Nausea, vomiting, diarrhea, constipation
Antidiabetic agents	Acetohexamide Chlorpropamide	Nausea, vomiting, diarrhea, heartburn, cholestatic jaundice Nausea, vomiting, diarrhea

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Anti-infectives	Ampicillin	Diarrhea, nausea, vomiting, pseudomembranous colitis
	Ciprofloxacin	Nausea, vomiting, diarrhea
	Clindamycin	Nausea, vomiting, diarrhea, pseudomembranous colitis
	Erythromycin	Abdominal pain and cramping, nausea, vomiting, diarrhea, hepatic dysfunction, jaundice
	Griseofulvin	Nausea, vomiting, diarrhea, flatulence
	Metronidazole	Taste disturbances, abdominal discomfort, diarrhea, nausea, vomiting
Antihypertensives	Sulfonamides	Nausea, vomiting, hepatic changes
	Tetracycline hydrochloride	Nausea, vomiting, diarrhea, stomatitis
	Clonidine	Nausea, vomiting, constipation
Antineoplastic agents	Methyldopa	Elevated liver function tests
Antituberculosis agents	All antineoplastics	Nausea, vomiting, stomatitis
	Isoniazid	Increased liver enzyme concentrations
	Rifampin	Heartburn, nausea, vomiting, diarrhea, increased liver enzyme concentrations
Biphosphonates	Alendronate	Esophagitis, heartburn
Cardiac agents	Digoxin	Nausea, vomiting, diarrhea, anorexia with toxicity
	Quinidine	Nausea, vomiting, diarrhea, abdominal cramps, colic
Narcotic analgesics	Codeine, meperidine, methadone, morphine, oxycodone	Nausea, vomiting, constipation, biliary spasm, or colic
NSAIDs	Ibuprofen, indomethacin, salicylates	Nausea, vomiting, dyspepsia, GI bleeding, peptic ulcer
Phenothiazines	Prochlorperazine, thioridazine	Constipation, dyspepsia, paralytic ileus, cholestatic jaundice (hypersensitivity reaction)
Miscellaneous agents	Acarbose	Diarrhea, flatulence, abdominal pain
	Allopurinol	Altered liver function, nausea, vomiting, diarrhea
	Aminophylline, theophylline	GI irritation, epigastric pain, nausea, vomiting, anorexia
	Barium sulfate	Cramping, diarrhea
	Colchicine	Diarrhea, nausea, vomiting, abdominal pain
	Estrogen-progestin combinations	Nausea, vomiting, diarrhea, abdominal cramps, altered liver function tests, cholestatic jaundice
	Iron preparations	Constipation, nausea, vomiting, black stools
	Gold sodium thiomalate, auranofin	Changes in bowel habits, diarrhea, abdominal cramping, nausea, vomiting
	Levodopa	Nausea, vomiting, anorexia
	Lithium	Nausea, vomiting, diarrhea
	Phenytoin sodium	Nausea, vomiting, constipation, dysphagia
	Potassium supplements	Nausea, vomiting, diarrhea, abdominal discomfort, small bowel ulceration (with enteric-coated tablets)
	Prednisone	Epigastric pain, gastric irritation, pancreatitis
	Zidovudine	Nausea, vomiting, anorexia

Family History

Questioning about diseases in the patient's family enables you to identify those that the patient may be at

risk for because of genetic predisposition. Then you can help the patient plan lifestyle changes that will help prevent those diseases and promote health.

(Text continued on page 571)

Family History

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Genetically Linked Gastrointestinal Conditions

- Do you have a family history of colorectal cancer?
- Do you have a family history of cystic fibrosis?

- Some colorectal conditions, such as familial adenomatous polyposis (FAP), increase the risk of colon cancer. Patients over age 35 with a family history of FAP should be screened with FOBT and endoscopic assessment of colon.
- Stress high-fiber, low-fat diet to minimize risk of colorectal cancer.
- Cystic fibrosis is a life-threatening, genetically transmitted disease that usually occurs in infancy and causes severe respiratory problems, pancreatic enzyme deficits, and reduced pancreatic functioning, leading to digestive difficulty. Tell patients about the availability of genetic testing.

Familial/Genetically Linked Disorders With Gastrointestinal Symptoms

- Do you have a family history of alcoholism?
- Do you have a family history of aneurysm?
- Do you have a family history of sickle cell disease?
- Do you have a family history of GI cancers?
- Do you have a family history of colorectal cancer?

- Alcoholism has severe GI consequences.
- One of several factors contributing to aneurysm formation is a genetic link.
- Assess abdominal vasculature for signs of abdominal aneurysm.
- Sickle cell disease is an inherited autosomal recessive disorder caused by a genetic aberration in hemoglobin A.
- Vaso-occlusive crisis ("sickle cell crisis") may cause severe abdominal pain.
- Family history of GI cancer increases risk of cancer.
- Teach preventive activities and recommended cancer screening procedures.
- Assess colorectal cancer in family and polyps that may precede some types of colon cancer.

Absorption Disorders

- Do you have a family history of celiac disease (also called *nontropical sprue* and *gluten-sensitive enteropathy*)?

- Genetic susceptibility exists for this sensitivity to the protein from cereal grains (e.g., wheat, rye, oats). Results in malabsorption of nutrients in small intestine.
- Teach about diet free of cereal glutes and lactose.

Inflammatory Bowel Disorders

- Do you have a family history of Crohn's disease or ulcerative colitis?

- Teach patients about increased risk for colorectal cancer with Crohn's disease and ulcerative colitis and importance of having regular cancer screening exams.

Metabolic Disorders

- Do you have a family history of diabetes mellitus?
- Do you have a family history of obesity?
- Do you have a family history of porphyria?

- Type 1 (insulin-dependent diabetes mellitus [IDDM]) is an autoimmune disorder that occurs more frequently in children whose parents, especially the father, have IDDM.
- Type 2 (non-insulin-dependent diabetes mellitus [NIDDM]) occurs even more frequently in children whose parents have the disease.
- Both genetic and environmental factors may contribute to obesity, which is an important risk factor for many serious diseases. Assess family eating and exercise patterns, explore desire for weight reduction, and teach proper nutrition and lifestyle changes to family.
- Porphyria is a relatively rare type of anemia that causes nausea, vomiting, abdominal pain, and neurological and psychological problems.

Review of Systems

A disruption in the systems contained in the abdominal cavity can cause problems in many other areas of the body. The problem in another body system depends on which organ of the abdomen is involved. For example, liver problems may cause malaise, nausea and vomiting, bruising, jaundice, and fluid in the abdomen. This is one reason why taking a careful review of systems (ROS) is so important.

Another reason is that the ROS might reveal that the primary health problem does not originate in the abdomen. Instead, you may uncover medical illnesses that have abdominal symptoms. So be sure to keep an open mind about the nature of the patient's health problem and not conclude that it lies in the GI system simply because he or she has abdominal complaints. Instead, assess each system methodically until you have collected all the data.

(Text continued on page 573)

Review of Systems

SYSTEM/NORMAL FINDINGS

General Health Survey

- Do you have changes in your energy level? Weakness? A general feeling of ill health?
- Do you have fever or chills?
- Have you been sweating?
- Do you have pain with movement or inability to perform daily activities?

Integumentary Skin

- Do you have any rashes?
- Do you have itching?
- Have you noticed any skin color changes?
- Do you have any swelling (edema)?

HEENT

Head/Neck

- Have you had a recent head injury?
- Have you noticed any neck masses, swollen nodes, or thyroid problems?

Eyes/Ears

- Have you seen any changes in the whites of your eyes?
- Do you have ear or balance problems?

Nose/Throat

- Do you have difficulty swallowing?
- Do you have a sore throat or fever?

ABNORMAL FINDINGS/RATIONALE

- Fatigue or weakness may reflect poor nutritional state, anemia related to blood loss in the GI tract, or vitamin B₁₂ deficiency.
- Malaise is commonly associated with GI diseases.
- Fever or chills are seen with infections.
- Sweating may indicate infection, female hormonal changes, or thyroid disturbances.
- Difficulty with changing positions may indicate musculoskeletal problems, pain, or an infection in the abdomen.
- Inability to carry out daily activities indicates the severity of the abdominal problem.
- Complaints of pruritus or pain along a dermatomally distributed vesicular rash located on the lower aspects of the rib cage indicates herpes zoster (shingles).
- Other rashes may reflect reactions to an antibiotic, which may also cause pain, nausea, cramping, and diarrhea.
- Associated with liver disease.
- Jaundice is associated with liver disease.
- Ascites is associated with liver disease and edema with renal disease.
- Can be related to projectile vomiting.
- Infections, such as mononucleosis, can cause spleen enlargement.
- Weight loss or gain is associated with thyroid problems.
- History of goiter (hypothyroidism) may be related to iodine deficiency in diet.
- Icteric sclera are seen with liver disease.
- Dizziness is a sign of Ménière's disease, which causes nausea.
- May be a sign of esophageal mass, muscular or neurological problems, or benign diagnoses.
- Infections, such as strep throat, often cause abdominal discomfort.

(continued)

Review of Systems (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p>Respiratory</p> <ul style="list-style-type: none"> ■ Do you have any difficulty breathing? ■ Do you have any respiratory problems? 	<ul style="list-style-type: none"> ■ Dyspnea may be caused by edema in the abdomen or large tumors of the liver, esophagus, or stomach, which impinge on respiratory structures, or may be caused by portal hypertension from liver congestion. ■ Patients with chronic obstructive pulmonary disease (COPD) often have right-sided heart failure that may cause liver enlargement.
<p>Cardiovascular</p> <ul style="list-style-type: none"> ■ Do you have cardiovascular disease such as congestive heart failure? ■ Do you have pain and pulsations in your navel or back? 	<ul style="list-style-type: none"> ■ Cardiac disease often mimics GI complaints. Congestive heart failure can cause liver enlargement and ascites. ■ Pain associated with a pulsating mass may indicate an abdominal aortic aneurysm.
<p>Genitourinary</p> <ul style="list-style-type: none"> ■ What color is your urine? ■ Do you have burning, frequency, urgency, or blood in your urine? 	<ul style="list-style-type: none"> ■ Dark yellow, orange, or brown urine can be caused by excessive breakdown of RBCs or liver problems. Blood in the urine is caused by renal calculi, renal infarct, glomerulonephritis, or pyelonephritis. ■ Urinary tract infections can cause flank pain and CVA tenderness. ■ Difficulty starting urine stream or urinary hesitancy indicates possible prostate disease in men.
<p>Female Reproductive</p> <ul style="list-style-type: none"> ■ When was your last menstrual period? ■ Do you have any vaginal discharge or itching? 	<ul style="list-style-type: none"> ■ Important to rule out pregnancy in women of childbearing age. ■ After menopause, bleeding may signal a gynecological problem. ■ Amenorrhea may indicate an eating disorder such as anorexia nervosa. ■ May indicate vaginal infection that could lead to PID.
<p>Male Reproductive</p> <ul style="list-style-type: none"> ■ Do you have a discharge from your penis? ■ Do you practice safe sex? 	<ul style="list-style-type: none"> ■ Penile discharge is related to sexually transmitted diseases (STDs). ■ Unprotected sex increases the risk for STDs for both men and women, which in turn may affect the GI system.
<p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Do you have joint pain? Have you had any broken bones? ■ Do you have rheumatoid arthritis? 	<ul style="list-style-type: none"> ■ Arthritis and osteoporosis are extraintestinal complications of ulcerative colitis. ■ Splenomegaly is associated with rheumatoid arthritis.
<p>Neurological</p> <ul style="list-style-type: none"> ■ Do you drink alcohol? How much? ■ Do you have back problems, herniated disk, numbness, or loss of bowel and bladder control? 	<ul style="list-style-type: none"> ■ A deficiency of vitamin B₁ (thiamine) may present with numbness or paresthesias in the extremities and is commonly seen in alcoholics and people with long-standing diarrhea. ■ Loss of bowel and bladder control can occur with disk problems, such as cauda equina syndrome.
<p>Endocrine</p> <ul style="list-style-type: none"> ■ Do you have diabetes or thyroid disease? 	<ul style="list-style-type: none"> ■ Diabetes can result in gastroparesis, polyuria, and polydipsia. ■ Thyroid disease can affect weight, appetite, and bowel patterns.

SYSTEM/NORMAL FINDINGS

ABNORMAL FINDINGS/RATIONALE

Immunologic/Hematologic

- Do you have food allergies?
- Do you have HIV/AIDS?
- Do you have mononucleosis?
- Do you have sickle cell anemia?

- May cause GI tract pain and changes in bowel habits.
- Patients with AIDS have problems maintaining their weight.
- Risk of an enlarged tender spleen.
- Often causes abdominal pain.

Psychosocial Profile

The psychosocial profile describes your patient's lifestyle and habits. How your patient eats, exercises, rests, and

copies with the stresses of every day has an impact on the health of the GI system.

(Text continued on page 575)

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Health Practices and Beliefs/Self-Care Activities

- When was your last dental exam?
- Have you ever had:
 - Your stool tested for occult blood?
 - An endoscopy?
 - *Women:* When was your last gynecologic examination?
 - *Men:* Have you ever had a prostate examination or prostate-specific antigen test?

- Good dentition is required for proper mastication and digestion of food. Regular flossing reduces the risk of infection.
- Recommended screenings include:
 - *Men after age 40:* Digital rectal and prostate examination annually.
 - *Women:* Gynecologic/rectal examination annually.
 - Men and women after age 50: Baseline endoscopy, then every 5 to 10 years depending on results and history.

Typical Day

- How do you usually spend your days?
- Has your daily pattern changed?

- Reflects energy expenditure and stressors that exacerbate GI problems, such as colitis and gastric ulcers.
- Pattern changes indicate how abdominal problem affects life and what adjustments patient has made.

Nutritional Patterns

- What have you eaten in the past 24 hours?
- Are you on a special diet?
- Are you concerned about your weight?
- Have you lost your appetite?
- Who prepares your meals and does the food shopping?
- Have you lost weight?
- Have you gained weight?

- Assesses usual dietary intake. Screens for eating disorders.
- May indicate cancer, ascites, depression.
- Improper food preparation and cooking puts patient at risk for salmonella and other infections.
- Weight loss is associated with long-standing diarrhea, cancer, and eating disorders such as anorexia nervosa.
- Assesses eating patterns and nutritional knowledge, stress level, fluid retention, and ascites in abdomen.



HELPFUL HINT

Take a socioeconomic history to ascertain the patient's ability to pay for food. If he or she needs financial assistance, refer him or her to the hospital social worker who can help the patient access the needed resources.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Activity and Exercise Patterns</p> <ul style="list-style-type: none"> ■ Do you have any mobility problems? ■ Do you exercise regularly? What do you do and how often? 	<ul style="list-style-type: none"> ■ Immobility contributes to constipation. ■ Exercise expends calories and strengthens abdominal muscles, which facilitates elimination.
<p>Recreation/Pets/Hobbies</p> <ul style="list-style-type: none"> ■ Do you have any pets? If so, what? ■ Are you responsible for animal care? ■ What are your hobbies? 	<ul style="list-style-type: none"> ■ Pregnant patients should avoid handling cat feces to prevent infection with toxoplasmosis. ■ Hobbies may involve toxic substances. For example, inhaling model airplane glue causes kidney problems.
<p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ What is your usual sleep pattern? ■ Do you ever awaken with indigestion? 	<ul style="list-style-type: none"> ■ Establishes baseline sleep schedule. ■ Indigestion, GERD, and hiatal hernias may interfere with falling asleep. ■ Assess whether pain or abdominal symptoms awaken patient from sleep.
<p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you drink alcohol? If yes, how much? ■ Do you use tobacco? 	<ul style="list-style-type: none"> ■ Alcohol can damage the GI tract, create vitamin deficiency states, and with chronic abuse, cause cirrhosis and esophageal varices. ■ Smoking increases risk for mouth, throat, lung, and GI cancer; chewing tobacco increases risk of mouth cancer.
<ul style="list-style-type: none"> ■ Do you use street drugs or share IV needles? ■ Do you have body piercing or tattoos? ■ Do you wash your hands after using the bathroom or preparing food? 	<ul style="list-style-type: none"> ■ IV drug use and reused tattoo or body-piercing needles put patient at risk for hepatitis and HIV. ■ Careless personal hygiene increases risk for infections, such as <i>Escherichia coli</i> and hepatitis. Handwashing is most important preventive measure.
<p>Environmental Health Patterns</p> <ul style="list-style-type: none"> ■ Where do you live? ■ Do you have public water or well water? ■ How old is your home? 	<ul style="list-style-type: none"> ■ Farmers may be exposed to insecticides or leaded gas (used in farm machines), which increase risk of GI problems. ■ Wells may be infected with microbes such as <i>Giardia</i> or <i>Cryptosporidium</i>, which cause diarrhea, vomiting, and other GI symptoms. ■ Public water may also contain chemicals that cause abdominal and neurological complaints. ■ Older homes may contain lead-based paint, putting young children at risk for lead poisoning. Chronic lead poisoning is associated with appetite loss; acute lead poisoning has symptoms of colicky abdominal pain and constipation or diarrhea. ■ Black widow spider bites cause abdominal pain. ■ Tapeworms and other parasitic intestinal worms cause abdominal symptoms.
<p>Stress and Coping</p> <ul style="list-style-type: none"> ■ How do you deal with anger and stress? 	<ul style="list-style-type: none"> ■ Explore what patient perceives as stressful and identify adaptive and maladaptive coping strategies. ■ Stress exacerbates ulcers and colitis.



Case Study Findings

Reviewing Mrs. Robichaud's

Biographical data:

- 56-year-old divorced woman, mother of two grown children.
- French-Canadian descent, speaks English and French.
- Roman Catholic.
- High school graduate and mill worker.
- Source of information: Self, reliable.

Current health status:

- No bowel movement in 4 days.
- Dull, intermittent lower abdominal pain rated 2 on a 0 to 10 scale. Walking around makes it better, and nothing makes it worse.
- Denies visible red blood in stool, black stool, or symptoms in other body systems.
- Says she has had problems with constipation for the last few years.

Past health history:

- Positive for varicella and measles.
- No history of GI problems as a child.
- Cholecystectomy in 1997; appendectomy in 1953.
- Denies having any GI assessment procedures or blood transfusions.
- Denies lactose intolerance or any allergies to substances, foods, or medications.
- Denies having any other health problems.
- Up to date on all immunizations; had hepatitis B series in 1997, pneumovax in 1997, tetanus in 1997, yearly flu shot last fall.
- Takes no prescribed medications.
- Takes no over-the-counter (OTC) medications now. Has used laxatives to relieve constipation over past few years.
- Uses no herbal preparations.

Family history:

- Mother and brother died of colon cancer.
- Father died of congestive heart failure.
- No family history of obesity or diabetes.

Review of systems:

- *General Health Survey*: No changes in energy level, just feels "bloated."
- *Integumentary*: No changes.
- *Head, Eyes, Ears, Nose, and Throat (HEENT)*: No problems with head, face, neck, eyes, ears, nose, mouth, throat; no difficulty swallowing, last dental exam 6 months ago.

- *Respiratory*: No breathing difficulties.
- *Cardiovascular*: No problems.
- *Genitourinary*: No burning, frequency, or incontinence, no history of renal disease
- *Reproductive*: Postmenopausal.
- *Neurological*: No loss of sensation, bowel and bladder control intact.
- *Musculoskeletal*: No changes, no history of arthritis.
- *Immune/Hematologic*: No allergies.
- *Endocrine*: No history of thyroid disease or diabetes.

Psychosocial profile:

- Goes for yearly physical and dental examination.
- Denies having any GI assessment procedures.
- Typical day begins at 5:30 A.M., when she bathes, eats breakfast, and gets ready for work at the textile mill. Works 7 A.M. to 3 P.M. Goes grocery shopping after work, prepares supper, does housework, calls her sister, and then relaxes by watching TV. Goes to bed at 10 P.M. every night.
- 24-hour recall reveals that Mrs. Robichaud likes "junk food," especially high-fat dishes traditional to the French culture. Breakfast usually consists of pancakes with butter and syrup; for lunch, she has a sandwich and chips; and for dinner, baked beans and brown bread. Says desserts are her weakness. She eats 1 serving of fruit; 1 serving of vegetables; 3 servings of meat; 1/2 serving of milk; 2 servings of white breads/pasta/rice/grains; and at least 5 fats a day. She reports that her usual weight is 175 pounds and she is 5'3".
- No exercise program; uses car to get to work.
- Likes to make stained glass objects in her spare time. Has no pets.
- Sleeps 7.5 hours a night, more on weekends.
- Doesn't drink alcohol or smoke cigarettes.
- Works in a textile mill, but does not know what chemicals she may be exposed to.
- Lives alone in an apartment in a part of town where she doesn't feel safe walking alone. Her apartment was built in 1985.
- Mother of two adult children, lives by herself, and calls her sister every day.
- Mill where she works has been laying off employees, and she is worried that she will be next. Has been talking with her sister and other mill workers about this concern.



Case Study Evaluation

Before you move on to the physical examination, review and summarize the key information from Mrs. Robichaud's health history. Document the abdominal history related to Mrs. Robichaud.



CRITICAL THINKING ACTIVITY #2

What risk factors for abdominal problems are evidenced in Mrs. Robichaud's history?



CRITICAL THINKING ACTIVITY #3

What factors may contribute to her chief complaint?

Anatomical Mapping

Anatomical mapping helps pinpoint the location of findings during the abdominal assessment. There are three ways to identify the location of these findings: anatomical landmarks, the four-quadrant method, and the nine regions of the abdomen.

Anatomical Landmarks

Anatomical structures are used as landmarks to help you describe abdominal findings. The following landmarks are used: xiphoid process of the sternum; costal margin; midline (down the center of the abdomen); umbilicus; anterior-superior iliac spine; inguinal ligament (Poupart's); and superior margin of the pubic bone (Figs. 17.3 and 17.4).




FIGURE 17.3. Anatomical landmarks (front).




FIGURE 17.4. Anatomical landmarks (back).

Four-Quadrant Method

Another way to mark the location of your findings is by the four-quadrant method. To use this method, draw imaginary lines separating the abdomen into four quadrants, with one line at the midline and the other horizontal at the umbilicus. These lines should intersect at the umbilicus. The aorta and the spine are located midline in the abdomen. The uterus and bladder, when enlarged, may be palpable midline in the abdomen (Fig. 17.5).




FIGURE 17.5. Four-quadrant method.

Four-Quadrant Method

Right Upper Quadrant	Left Upper Quadrant
<ul style="list-style-type: none"> ■ Liver ■ Gallbladder ■ Pylorus ■ Duodenum ■ Head of pancreas ■ Right adrenal gland ■ Portion of right kidney ■ Hepatic flexure of colon ■ Portions of ascending and transverse colon ■ Right renal artery 	<ul style="list-style-type: none"> ■ Left lobe of liver ■ Spleen ■ Stomach ■ Body of pancreas ■ Left adrenal gland ■ Portion of left kidney ■ Splenic flexure of colon ■ Portions of transverse and descending colon ■ Left renal artery
Right Lower Quadrant	Left Lower Quadrant
<ul style="list-style-type: none"> ■ Lower portion of right kidney ■ Cecum and appendix ■ Portion of ascending colon ■ Bladder (if distended) ■ Ovary and salpinx ■ Uterus (if enlarged) ■ Right spermatic cord ■ Right ureter ■ Right iliac artery ■ Right femoral artery 	<ul style="list-style-type: none"> ■ Lower section of left kidney ■ Sigmoid colon ■ Descending colon ■ Bladder (if distended) ■ Ovary and salpinx ■ Uterus (if enlarged) ■ Left spermatic cord ■ Left ureter ■ Left iliac artery ■ Left femoral artery

Nine Regions of the Abdomen

The third way to document the location of your findings is to separate the abdomen into nine regions, similar to a tic-tac-toe grid. The first two lines are vertical at the right and left midclavicular lines to the middle of the inguinal ligaments. The second two lines are horizontal beginning




FIGURE 17.6. Nine regions of the abdomen.

at the lower edge of the costal margin and at the anterior-superior iliac spine of the iliac bones (Fig. 17.6).

Physical Assessment

Now that you have completed the subjective part of your examination, proceed to the objective part. The purpose of the physical assessment is to identify normal structures and functions as well as actual and potential health problems.

Just as all the organs of the body are interrelated, so are the assessments. Assessment findings in other body areas can also indicate problems with abdominal organs. So your assessment should begin with a general survey and a head-to-toe scan to detect clues that may indicate an abdominal problem.

Approach

Perform the abdominal examination in a warm, private environment. Have your patient empty her or his bladder before the examination, so that you do not mistake a full bladder for a mass. Ask the patient to lie supine with her or his arms at the sides. Warm both your stethoscope and your hands before proceeding with the examination, and remember to work from the right side of your patient.

Once your patient is comfortable, expose the abdomen from the lower thorax to the iliac crests. Other things to remember include:

Nine Regions of the Abdomen

Right Hypochondriac	Epigastric	Left Hypochondriac
<ul style="list-style-type: none"> ■ Right lobe of liver ■ Gallbladder ■ Duodenum ■ Hepatic flexure ■ Portion of right kidney ■ Suprarenal gland 	<ul style="list-style-type: none"> ■ Pylorus ■ Duodenum ■ Head of pancreas ■ Portion of liver ■ Aorta and renal arteries 	<ul style="list-style-type: none"> ■ Stomach ■ Spleen ■ Pancreas tail ■ Splenic flexure ■ Upper portion of left kidney ■ Suprarenal gland
Right Lumbar	Umbilical	Left Lumbar
<ul style="list-style-type: none"> ■ Portion of right kidney ■ Hepatic flexure of colon ■ Ascending colon ■ Duodenum ■ Jejunum 	<ul style="list-style-type: none"> ■ Lower duodenum ■ Jejunum and ileum ■ Aorta and iliac arteries 	<ul style="list-style-type: none"> ■ Descending colon ■ Lower half of left kidney
Right Inguinal	Hypogastric	Left Inguinal
<ul style="list-style-type: none"> ■ Cecum ■ Appendix ■ Ileum ■ Right femoral artery ■ Right ureter ■ Right spermatic cord ■ Right ovary 	<ul style="list-style-type: none"> ■ Ileum ■ Bladder (if distended) ■ Uterus (if enlarged) ■ Aorta and femoral arteries 	<ul style="list-style-type: none"> ■ Sigmoid colon ■ Left ureter ■ Left spermatic cord ■ Left femoral artery ■ Left ovary

- Explain what you will be doing during the examination.
- Have adequate lighting so that you can visualize the abdomen without difficulty.
- Observe the patient's face for signs of discomfort.
- Perform the examination slowly and avoid quick movements.
- Make sure that your fingernails are short, to prevent injuring the patient during palpation.
- Distract the patient with questions or conversation.

You will use all four techniques of physical assessment to examine the abdomen. However, the sequence is inspection, auscultation, percussion, and palpation. During an abdominal examination, it is important to auscultate before percussion and palpation because the manipulation that occurs with these techniques may increase the frequency of bowel sounds.



HELPFUL HINTS

- If your patient tenses his or her muscles when you are palpating the abdomen, place a pillow under his or her knees to relax the abdominal muscles.
- If your patient complains of pain when you palpate an area, stop, and then palpate that area last.



TOOLBOX

The equipment needed for an abdominal examination includes: stethoscope (bell and diaphragm), pen, metric ruler, and reflex hammer or tongue blade to assess abdominal reflexes.

Performing a General Survey

Before physically assessing the abdomen, perform a general survey, observing the patient's overall appearance. Using your inspection skills, note nutritional status, emotional status, body habitus, and any changes that might relate to the abdomen.

Begin by taking vital signs, height, and weight. Changes in vital signs may alert you to a serious medical problem. Vital sign changes and related abdominal problems include:

- *Hypertension:* Abdominal aortic aneurysm or dissection, renal infarction, glomerulonephritis, vasculitis, or abdominal pain.
- *Orthostatic hypotension:* Hypovolemia (fluid or blood loss).

- **Fever:** GI infection, peritonitis, pelvic infection, cholangitis.
- **Pulse deficit:** Aortic dissection or aneurysm.
- **Hypotension/bradycardia:** Hypotension may indicate shock associated with ruptured abdominal aortic aneurysm. Vasovagal reaction is caused by bearing down or straining with a bowel movement. The decrease in pulse and BP is a result of decreased blood return to the heart and therefore decreased cardiac output.

In addition to taking vital signs, be alert for signs that may indicate underlying abdominal problems. For example, note:

- **Facial expression:** Is it appropriate? If your patient complains of pain, does her or his nonverbal behavior reflect this? For example, is she or he grimacing?
- **Posture:** Does your patient assume a particular posture for comfort? For example, is he or she splinting a sec-

tion of the abdomen, guarding an area of the abdomen, or drawing the knees up to his or her chest? Patients with acute appendicitis often flex their legs, because lying supine often increases the intensity of pain. Does pain seem to increase with movement?

- **Weight/nutritional status:** Is your patient malnourished and underweight or overweight? Severely thin patients may have an eating disorder. Overweight patients may have underlying cardiovascular or renal disease as a result of fluid retention. Gross abnormalities such as abdominal distension warrant further investigation.

Performing a Head-to-Toe Physical Assessment

An abdominal assessment reflects many different systems. Therefore, next examine the patient for specific changes that may indicate underlying pathology and might have an impact on the structures of the abdomen.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

General Health Survey

- Alert and oriented to time, place, and person. Remote and recent memory intact.

Integumentary

Skin

INSPECT/PALPATE

- Skin color normal and appropriate for ethnic background.
- Skin warm and moist.

ABNORMAL FINDINGS/RATIONALE

- **Diminished mental status:** Can occur with hemorrhaging.
- Confusion or other nervous system problems may relate to liver's inability to detoxify medications or toxins or kidneys' inability to rid body of harmful wastes.
- **Yellow skin:** Jaundice, associated with liver disease or biliary obstruction.
- **Dry skin and mucous membranes/poor skin turgor with tenting:** Dehydration. GI diseases may cause dehydration with fluid and electrolyte disturbances. For example, **ulcerative colitis** causes frequent, loose diarrhea and deficits in sodium, potassium, chloride, magnesium, zinc, copper, and other minerals. **Gastroenteritis** causes severe fluid and electrolyte imbalances from vomiting and diarrhea that result in loss of water, sodium, chloride, potassium, and bicarbonate.
- **Peptic ulcer disease** may cause bleeding, resulting in hematemesis and melena. Besides dehydration and fluid and electrolyte imbalances, severe blood loss (>1 L) may cause hypovolemic shock, with symptoms including hypotension, weak pulse, chills, palpitations, and cold, clammy skin.



HELPFUL HINT

Because diarrhea and other GI disorders may cause fluid and electrolyte imbalance, be sure to monitor your client's input and output (I&O) and serum electrolytes.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<ul style="list-style-type: none"> ■ No edema. ■ No lesions. 	<ul style="list-style-type: none"> ■ <i>Lower extremity edema:</i> Can be caused by iliac obstruction, pelvic mass, or renal disease. ■ <i>Linear lesions on one-half of the abdomen:</i> Shingles, an infection of herpes zoster that follows the dermatome of the nervous system.
<p>Hair/Nails</p> <ul style="list-style-type: none"> ■ Hair of normal texture. Nail shape within normal limits. 	<ul style="list-style-type: none"> ■ <i>Cachexic appearance with dry, brittle hair, thin hair, dry skin:</i> Anorexia nervosa and/or malnutrition. ■ <i>Curved fingernails:</i> Cirrhosis or some type of bowel disease.
<p>HEENT Head/Eyes/Ears/Nose <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ Normocephalic, even distribution of hair. ■ Sclera white, conjunctiva pink. ■ No periorbital edema. ■ No alterations in sight (CN IX). ■ No alterations in hearing (CN VIII). ■ No alterations in smell (CN VII). 	<ul style="list-style-type: none"> ■ <i>Icteric sclera:</i> Liver disease or biliary obstruction. ■ <i>Conjunctivitis, uveitis, episcleritis:</i> Extraintestinal complications of ulcerative colitis. ■ <i>Periorbital edema:</i> Renal disease. ■ <i>Sight alterations:</i> Can affect food intake. ■ <i>Altered hearing:</i> Can affect food intake. ■ <i>Altered smell:</i> Can affect food intake.
<p>Neck</p> <ul style="list-style-type: none"> ■ No masses or palpable lymph nodes. 	<ul style="list-style-type: none"> ■ <i>Neck masses:</i> Goiter and other masses may reflect thyroid disease and impair swallowing. ■ <i>Cervical lymphadenopathy:</i> May indicate a systemic infection or malignancy, such as mononucleosis or Hodgkin's disease, and cause splenomegaly.
<p>Mouth/Throat</p> <ul style="list-style-type: none"> ■ Lips and mucous membranes intact, pink and moist; 32 white teeth in good repair. Positive swallow and gag reflex. Tongue pink and moist without lesions and with full mobility (CN IX, CN X, CN XII intact). ■ No alterations in taste (CN I). 	<ul style="list-style-type: none"> ■ <i>Dental caries/mouth lesions (aphthous ulcers):</i> Affect nutritional intake. Mouth lesions associated with ulcerative colitis. ■ <i>Dry lips/mucous membranes:</i> Dehydration. ■ <i>Red, beefy tongue:</i> Pernicious anemia. ■ <i>Parotitis, dental problems, irritated pharynx:</i> Bulimia. ■ <i>Absence of gag/swallowing reflex:</i> Neurological diseases such as stroke may alter nutritional status. ■ <i>Taste alterations:</i> Can affect food intake.
<p>Respiratory <i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Respirations 16 with normal depth and regular rhythm. 	<ul style="list-style-type: none"> ■ <i>Hypoxia:</i> May be caused by distended abdomen, which raises diaphragm and inhibits respiration. ■ <i>Shallow respirations:</i> An attempt to avoid abdominal pain caused by movement in clients with peritonitis.
<p>Cardiovascular <i>PALPATE/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Apical and radial pulses within normal limits. ■ S₁, S₂ present and regular with no adventitious or extra heart sounds. ■ Pulses +2/3. ■ No bruits or thrills. 	<ul style="list-style-type: none"> ■ <i>Right-sided congestive heart failure:</i> Can cause ascites. ■ <i>Bruits and thrills:</i> Abdominal aortic aneurysm.

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p>Reproductive</p> <ul style="list-style-type: none"> External genitalia pink, intact, no lesions or discharge. <p>Musculoskeletal <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> 5/5 strength of upper and lower extremities. No joint deformities. <p>Neurological <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> Cranial nerves intact, positive 2 DTR 	<ul style="list-style-type: none"> STD: May cause painful oral lesions that affect client's nutritional status. AIDS: Associated with weight loss and viral and fungal infections that affect the GI tract, causing diarrhea, abdominal pain, stomatitis, esophagitis, and gastritis. Arthritis and osteoporosis: Extraintestinal complications of ulcerative colitis. Rheumatoid arthritis: Associated with splenomegaly. Muscle weakness: Lack of protein intake or absorption problems. Muscle twitching: Associated with loss of electrolytes from diarrhea. Numbness/parathesias in extremities: Vitamin B₁ deficiency, commonly seen in alcoholics and people with long-standing diarrhea. Disc problems such as cauda equina syndrome: Loss of bowel and bladder control.

Performing an Abdominal Assessment

After you have completed your general survey and head-to-toe assessment, focus on the abdomen. Begin with inspection and proceed with auscultation, percussion, and palpation. Next, examine each structure separately. As you proceed with the assessment, try to visualize the underlying structures.

Inspection

Inspect the abdomen for size, shape, and symmetry. Look at it from different angles. Check color, surface characteristics, contour, and surface movements. Look for lesions, striae, or scars. **Striae**, also known as *lineas albicantes* or *stretch marks*, are streaks of light-colored skin that occur after rapid skin stretching. Observe the location of the umbilicus and note any visible veins on the abdomen. Then, have the patient take a deep breath and bear down to assess for bulges that may indicate a hernia or organomegaly.

Assess for distension—any unusual stretching of the abdominal wall. If present, determine if it is generalized or in one area. Fluid and gas usually result in generalized, symmetrical distension, whereas anything solid, such as a fetus, mass, tumor, or stool, results in asymmetrical distension. Sometimes distension is difficult to assess, so ask the patient if her or his abdomen looks or feels any different from normal. A concrete way to

measure abdominal distention is to measure abdominal girth and compare measurements daily. Measurements should be taken at the umbilicus for consistency.

Also inspect the abdomen for any visible aortic pulsations, peristalsis, and respiratory pattern. Slight aortic pulsations and respiratory movements are readily seen in adult patients. Visualization of peristaltic waves may be seen in infants and small children, but this usually indicates a problem if seen in an adult. (See *Inspection of the Abdomen*.)

Auscultation

Begin auscultating the abdomen by placing the warmed diaphragm of the stethoscope gently in one quadrant. Proceed in an organized fashion, listening in several areas in all four quadrants. Use the diaphragm to listen for bowel sounds, which sound like high-pitched gurgles or clicks that last from 1 to several seconds. They are assessed to determine bowel motility and peristalsis. **Peristalsis** is the progressive wavelike movements of the digestive tract that move gastric contents through the tract. There will be 5 to 30 clicks per minute, or bowel sounds occurring every 5 to 15 seconds on an average adult patient. If bowel sounds are hypoactive, listen over the ileocecal valve to the right of the umbilicus.

(Text continued on page 585)

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> Weight changes Diet Fevers Dizziness	General	<i>Inspect:</i> Orientation, facial expression, posture, nutritional status <i>Measure:</i> Height, weight, vital signs
Integumentary	<i>Ask about:</i> Changes in skin, hair, and nails Rashes, itching, lesions	Integumentary	<i>Inspect:</i> Skin, hair and nails for changes in color and texture, lesions, and edema/ascites <i>Palpate:</i> Skin turgor
HEENT <i>Head and Neck</i>	<i>Ask about:</i> Thyroid disease Neck masses Recent infections	HEENT <i>Head and Neck</i>	<i>Inspect:</i> Neck masses <i>Palpate:</i> Thyroid, lymph nodes
Eyes	<i>Ask about:</i> Vision changes	Eyes	<i>Inspect:</i> Edema, color of sclera, retinal changes
Ears, Nose and Throat	<i>Ask about:</i> Trouble swallowing Sore throat Dizziness Last dental exam	Ears, Nose and Throat	<i>Inspect:</i> Mouth, throat, teeth <i>Test:</i> Cranial nerves 1, 7, 9, 10, 12
Respiratory	<i>Ask about:</i> Breathing problems, SOB History of COPD	Respiratory	<i>Measure:</i> Respiratory rate and depth <i>Auscultate:</i> Breath sounds
Cardiovascular	<i>Ask about:</i> History of CVD, HTN, CHF	Cardiovascular	<i>Palpate:</i> Pulses; check for thrills, edema <i>Auscultate:</i> Heart for extra sounds (S3), bruits
Genitourinary/Reproductive	<i>Ask about:</i> Color of urine Urinary burning, frequency, hesitancy History of STDs <i>Women:</i> LMP Vaginal discharge <i>Men:</i> Prostate problems Penile discharge	Genitourinary/Reproductive	<i>Inspect:</i> Color of urine, external genitalia for lesions or discharge <i>Palpate:</i> Bladder for distention, kidneys, prostate <i>Percuss:</i> CVA for tenderness <i>Women:</i> Perform pelvic exam <i>Men:</i> Perform rectal exam
Musculoskeletal	<i>Ask about:</i> History of fractures Joint pain Weakness	Musculoskeletal	<i>Inspect:</i> Spinal curves, joints, ROM <i>Palpate:</i> Muscle strength
Neurological	<i>Ask about:</i> Alcohol use Numbness Back problems Loss of bowel/bladder control	Neurological	<i>Test:</i> Sensation, DTR
Endocrine	<i>Ask about:</i> History of diabetes Thyroid problems	Lymphatic/Hematologic	<i>Palpate:</i> Lymph nodes, spleen
Lymphatic/Hematologic	<i>Ask about:</i> Food allergies Infection Sickle cell anemia	Lymphatic/Hematologic	

Assessment of Abdomen's Relationship to Other Systems

INSPECTION OF THE ABDOMEN

1. Inspecting the Abdomen

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Have patient empty bladder.
- Position patient supine; knees can be slightly flexed to relax the abdomen.
- Inspect the abdomen from side and foot of bed.
- Note size, shape, symmetry, movements (respirations, peristalsis, and pulsations), surface characteristics such as color, hair distribution, lesions, striae.



Inspecting the abdomen

- *Skin color should be consistent with patient's ethnicity. Color may be lighter on abdomen than on other areas of body because of lack of sun exposure. Skin color should be the same throughout the abdomen.*
- *Abdominal skin intact with no lesions or masses. Striae may be present. If new, should be pink; if old, white/silver. In a pregnant patient, striae and linea nigra may be present.*



Striae

ABNORMAL FINDINGS/RATIONALE

- *Jaundice:* Liver disease.
- *Redness:* Inflammation.
- *Cyanosis:* Hypoxia.
- *Bluish discoloration around umbilicus (Cullen's sign):* Hemorrhagic pancreatitis or intraperitoneal bleeding.
- *Discoloration of lower abdominal flanks and lower back (Grey-Turner's sign):* Pancreatitis or extraperitoneal bleeding.
- *Bruises:* Recent trauma.
- *Pink-purple striae:* Cushing's syndrome.
- *Spider angioma:* Liver failure.
- *Caput medusa* (dilated veins that extend from the umbilicus): Liver failure.
- *Hernias (protrusion of abdominal organs through opening in abdominal wall):* Caused by weakness of abdominal muscles. Seen as bulges that occur when patient bears down. Occur on old surgical incisions, around umbilicus, or in inguinal area. Epigastric or linea alba hernias occur when intestine protrudes through opening in midline of abdomen above umbilicus.
- *Diastasis recti (separation of rectus abdominis muscle that occurs in linea alba):* Occurs in pregnant women and newborns.
- *Dermatologically distributed vesicular rash on lower rib cage in upper quadrants:* Herpes zoster.

2. Umbilicus

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect umbilicus, noting position, contour, color, and discharge.
- Ask patient to raise head off bed, note any bulges (hernias)
- *Umbilicus inverted and midline.*



Lifting head to accentuate hernia

(continued)

INSPECTION OF THE ABDOMEN

2. Umbilicus (continued)

ABNORMAL FINDINGS/RATIONALE

- *Protrusion of umbilicus:* Umbilical hernia. An underlying mass may cause umbilicus to deviate from midline.

3. Symmetry

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- View abdomen from different angles.
- *Abdomen symmetrical bilaterally from costal margin to iliac crest, with umbilicus in center.*



Angle 1: Looking across abdomen



Angle 2: Looking down at abdomen


ABNORMAL FINDINGS/RATIONALE

- *Asymmetry:* Tumors, cysts, bowel obstruction, organomegaly, or scoliosis.

4. Contour and Distension

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect contour from various angles.
- *No abdominal distension. In average adult, contour should be either flat, round, or scaphoid. A flat contour is seen in a muscular patient who is physically fit. A round or convex contour is normal in infants or toddlers but indicates poor muscle tone or excessive fat deposits in adults. A scaphoid or concave contour may be seen in thin patients of all ages.*



INSPECTION OF THE ABDOMEN

4. Contour and Distension (continued)

ABNORMAL FINDINGS/RATIONALE

- *Abdominal distension*: Causes are “nine Fs”: Fat, fluid, feces, fetus, flatus, fibroid, full bladder, false pregnancy, and fatal tumor.
- The area of abdomen that is distended can pinpoint cause of distension.
- *RLQ and LLQ*: Pregnancy; ovarian or uterine tumor, bladder enlargement.
- *LUQ and RUQ*: Pancreatic cyst or tumor or gastric distension.
- *Asymmetrical*: Tumor, hernia, cyst, bowel obstruction.
- *Concave abdomen*: Can also indicate malnutrition.

5. Surface Movements

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Look for respirations, pulsations, and peristalsis.
- *On a thin patient, peristalsis and aortic pulsations may be visible. Women’s respirations are more thoracic, whereas men tend to use their abdominal muscles more with breathing.*

ABNORMAL FINDINGS/RATIONALE

- *Increased peristaltic waves*: Intestinal obstruction.
- *Reverse peristaltic waves*: Pyloric stenosis.
- *Abnormal respiratory movements (use of accessory muscles, grunting, restricted abdominal movements, labored respirations)*: May indicate respiratory distress.
- *Shallow respirations in male patients*: Abdominal pain.
- *Increased/diffuse pulsations*: Aortic aneurysm.

Listen for vascular sounds with the bell of the stethoscope. These sounds include bruits, venous hums, and friction rubs. Apply the bell of the stethoscope lightly on the abdomen. Listen over the aorta in the epigastric region, over the renal, iliac, and femoral arteries. Listen for the presence of bruits, or a swishing sound, which indicates turbulent blood flow caused by constriction or

dilation of a tortuous vessel. Also listen over the epigastric region and liver and around the umbilicus for a venous hum—a soft, low-pitched humming noise with a systolic and diastolic component. Last, listen over the liver and spleen, along the right and left costal margins, for friction rubs. These are grating sounds that increase with inspiration and indicate peritoneal irritation (Fig. 17.7).




FIGURE 17.7. Auscultating sites for vascular sounds.



HELPFUL HINTS

- If the patient identifies an area of pain, examine that area last to minimize discomfort.
- Warm your stethoscope and hands to prevent the patient from tensing the abdominal muscles.
- If the patient has a nasogastric tube attached for suction, be sure to pinch the tube off or turn off the suction before you auscultate. Otherwise, you may think the patient has bowel sounds, when what you are really hearing is gastric secretions being suctioned out into the tube.

AUSCULTATION OF THE ABDOMEN

1. Bowel Sounds

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Always auscultate prior to palpation.
- Quadrant off the abdomen with the umbilicus as the midpoint.
- Use the diaphragm portion of the stethoscope to listen for bowel sounds in all four quadrants.
- Listen for at least 5 minutes before stating patient has absent bowel sounds.
- *In an average adult patient, bowel sounds present at a rate of 5–30 clicks/min in each quadrant.*



Auscultating the abdomen



HELPFUL HINT

Listen over the ileocecal valve in the RLQ to the right of the umbilicus. This is normally a very active area connecting the small intestine to the large intestine.

ABNORMAL FINDINGS/RATIONALE

- *Bowel sounds more than 30 clicks (bowel sounds)/min:* Hyperactive bowel sounds or hyperperistalsis. Causes include irritable bowel disease, bowel infection, early bowel obstruction, diarrhea, resolving paralytic ileus, or laxative use. Types of hyperperistalsis include:
 - **Borborygmi**, which may be audible without a stethoscope and are characterized by a long stomach “gurgling.”
 - **Succussion splash**, a loud sound like splashing water, is often heard without a stethoscope as the patient moves from side to side. It occurs when the abdomen is filled with air or fluid and indicates delayed gastric emptying from an obstruction or gastric dilatation.
- *Bowel sounds less than 5 clicks/min but present:* Hypoactive or hypoperistalsis. Causes include peritonitis, medications such as opioids, bowel obstruction, or postoperative occurrence.
- Absent bowel sounds are caused by late bowel obstruction, peritonitis, or paralytic ileus after surgery during which the bowel was manipulated.

2. Arterial and Venous Vascular Sounds

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Bruits

- Use the bell portion of the stethoscope to listen for vascular sounds (bruits or venous hums).
- If indicated, listen for bruits over the abdomen at the midline in the epigastric area.
- If indicated, listen for bruits over the renal arteries above the umbilicus to the right and left of the midline in the upper quadrants.
- If indicated, listen for bruits over the iliac arteries below the umbilicus to the right and left of the midline in the lower quadrants.

AUSCULTATION OF THE ABDOMEN

2. Arterial and Venous Vascular Sounds (continued)



Auscultating the renal artery



Auscultating the iliac artery



Auscultating the abdominal aorta



Auscultating the femoral artery

- *No bruits; although bruits with only a systolic component can be a normal finding.*

Venous Hums

- If indicated, listen for a venous hum over the liver in the RUQ with the bell.



Auscultating the liver

- *No venous hums.*

Friction Rub

- If indicated, listen for friction rubs over inflamed organs or tumors with the diaphragm.
- *No friction rub.*

ABNORMAL FINDINGS/RATIONALE

- A bruit with a systolic and a diastolic component is abnormal. The etiology of the bruit depends on the area where it is heard. The following are some possible causes of bruits in these areas:
 - *Aortic:* Aortic aneurysm.
 - *Epigastric:* Renal artery stenosis.
 - *Aortic, iliac, or femoral:* Arterial insufficiency.
- *Venous hums:* Venous portal hypertension and liver disease.
- *Continuous murmur in epigastric region (Cruveilhier-Baumgarten murmur):* Portal hypertension.
- *Friction rub:* Inflammation of peritoneal surface of an organ from tumor, infection, or infarct. Liver and spleen cause friction rubs because they are in close contact with the peritoneum. Splenic friction rubs are caused by infection, abscess, infarction, or tumor and are best heard at lower rib cage in left anterior axillary line. Liver friction rubs are caused by liver cancer or abscess and can be auscultated over the lower right sternal border.




FIGURE 17.8. Percussion sites.

Percussion

Percussion is a technique used to assess the presence of fluid, air, organs, or masses in the abdominal cavity. Indirect or mediate percussion is best for assessing the abdomen. Always ask the patient if he or she has abdominal pain, and percuss painful areas last. Percuss all four quadrants, listening for tympany and dullness (Fig. 17.8). Tympany is the most common finding and indicates the presence of gas. Dullness can also be heard when percussing organs, masses, or fluid.

Percussion is also valuable for determining organ size and tenderness. The following methods are used for estimating the size of the liver, spleen, and bladder and for assessing kidney tenderness.

Assessing Liver Size

To help you locate the lower edge of the liver where it is difficult to percuss, use the **scratch test**: Place your stethoscope over the right upper quadrant (RUQ) above the liver, and with one finger of your other hand, lightly scratch the abdomen starting in the RLQ and moving up toward the liver. When the scratching sound in your stethoscope becomes magnified, you have reached the liver border.

The **liver span test** gives you an estimate of the size of the liver at the midclavicular line. To assess the upper border of the liver, start at the right midclavicular line at the third intercostal space over lung tissue and percuss down until you hear resonance change to dullness over the liver (around the fifth to seventh intercostal space). Place a mark where the dullness begins. The upper border of the liver usually begins at the fifth to seventh intercostal space. To determine the lower border of the liver, start at the right midclavicular line at the level of the umbilicus and percuss upward until tympany turns to dullness (usually at the sternal border). Mark this area with a pen. Measure the distance between the two marks—this is the liver span. The normal liver span at the midclavicular line is 6 to 12 cm. If you have a liver span




FIGURE 17.9. Percussing at lowest intercostal space at left anterior axillary line.

greater than 12 cm at the midclavicular line, you can measure the liver span at the midsternal line. The normal midsternal measurement is 4 to 8 cm.

Assessing Spleen Size

Percussion is also helpful in estimating the size of the spleen. Three methods are used. The first method is to percuss from the left midclavicular line along the costal margin to the left midaxillary line. If you hear tympany, splenomegaly is unlikely. Dullness in the area of the anterior axillary line to the midaxillary line is a sign of spleen enlargement.

A second method of assessing splenomegaly is to percuss at the lowest intercostal space at the left anterior axillary line (Fig. 17.9). Ask the patient to take a deep breath and percuss again. Tympany is normal, but with splenomegaly, the tympany turns into dullness on inspiration.

The third method is to percuss from the third to the fourth intercostal space slightly posterior to the left midaxillary line, and percuss downward until dullness is heard instead of tympany or resonance. Dullness of the normal spleen will be noted around the ninth to the eleventh rib.

Assessing Bladder Size

To percuss the bladder for distension, begin at the symphysis pubis and percuss upward to the umbilicus, noting any dullness. Normally, an empty bladder does not rise above the symphysis pubis.

Assessing Kidney Tenderness

Fist or blunt percussion can be used to assess the kidneys for tenderness. Assess the kidneys at the costovertebral angle (CVA). Posteriorly, identify the CVA where the end of the rib cage meets the spine. Place the palm of your nondominant hand over the CVA, and strike that hand with the fist of your other hand. Repeat on the other side. Tenderness upon blunt percussion at the costovertebral angle is positive CVA tenderness.

PERCUSSION OF THE ABDOMEN

1. Abdomen

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use indirect or mediate percussion.
- Percuss the abdomen in all four quadrants, percuss tender areas last.
- Note areas of dullness and tympany.
- *Tympany to dullness, depending on abdominal contents.*



Percussing the abdomen

ABNORMAL FINDINGS/RATIONALE

- *Extremely high-pitched tympanic sounds:* Distension.
- *Extensive dullness:* Organ enlargement or underlying mass.

2. Liver

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Perform the scratch test to locate the lower edge of the liver. Place diaphragm of the stethoscope over the liver in the RUQ while gently scratching the abdomen with your other hand, working from the RLQ at the midclavicular line, moving upwards toward the liver. Mark the point on the abdomen once the scratch sound is detected.



Performing the scratch test

- Use indirect or mediate percussion to determine liver size.
- Locate the upper edge of the liver by percussing downward at the midclavicular line from the fourth intercostal space from resonance to dullness, mark upper border of liver.



Percussing the upper liver edge

- Locate the lower edge of the liver by percussing upward at the midclavicular line from the RLQ from tympany to dullness; mark the lower border of liver.



Percussing the lower liver edge

(continued)

PERCUSSION OF THE ABDOMEN

2. Liver (continued)

- Measure the distance between upper and lower marks to obtain liver span.
- If the liver span is greater than 12 cm at the mid-clavicular line, a midsternal measurement can be obtained using the above technique at the midsternal line.
- *Average adult liver span at midclavicular line is 6–12 cm, 4–8 cm at midsternal line. Measurements vary according to gender and height. Men have larger liver spans than women; tall patients have larger liver spans than short patients. Liver spans in children vary with age. For example, the liver span of a 6-month-old child is around 2.5 cm, whereas that of a 24-month-old is about 3.5 cm.*



Measuring the liver

ABNORMAL FINDINGS/RATIONALE

- *Liver span greater than 12 cm:* Hepatomegaly.

3. Spleen

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Identify area of splenic dullness.
- Position patient on right side, percuss down the midaxillary line from an area of resonance over the lung to dullness over the spleen.
- *Spleen normally may not be detectable in a normal patient. If detected, the upper border should be percussed 6–8 cm above the left costal margin. The body of the spleen should be percussed between the ninth and the eleventh ribs.*



Percussing splenic dullness

ABNORMAL FINDINGS/RATIONALE

- *Upper border percussed beyond 8 cm above costal margin:* Enlarged spleen. Causes include portal hypertension, thrombosis, stenosis, atresia, deformities of splenic vein, cysts, cancer, mononucleosis, trauma, and infection as long as patient does not have a full stomach or intestines. (Dullness from full stomach and intestines is difficult to differentiate from splenic dullness.)

4. Bladder

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- If indicated, percuss for bladder dullness at the midline above the symphysis pubis.
- You should not be able to percuss the bladder above the level of pubic symphysis after the patient has voided.



Percussing the bladder

PERCUSSION OF THE ABDOMEN

4. Bladder (continued)

ABNORMAL FINDINGS/RATIONALE

- *Dullness in suprapubic area:* Full bladder (for other causes, see section on Abdominal Contour and Distension).

5. Costovertebral Angle (CVA)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Place nondominant hand over kidney.
- Make a fist with dominant hand and strike nondominant hand.
- Note any tenderness.
- Kidney tenderness is assessed at the CVA.
- *No CVA tenderness.*



Checking for CVA tenderness

ABNORMAL FINDINGS/RATIONALE

- *Severe pain or tenderness:* Kidney infection or a musculoskeletal problem.

Palpation

You will use both light and deep palpation to assess the abdomen. Begin with light palpation to put your patient at ease. Light palpation is useful in assessing surface characteristics and identifying areas of tenderness. If the patient has identified an area of pain, examine that area last. Otherwise, the patient may tense her or his muscles, affecting the accuracy of your assessment.

Perform light palpation in all four quadrants, using your fingertips. Press down 1 to 2 cm in a rotating motion, then lift your fingers and assess the next location. Palpate as much of the abdomen as possible. Observe for nonverbal signs of pain, such as grimacing or guarding. No tenderness should be noted.

To palpate for muscle guarding, perform light palpation over the rectus muscles of the abdomen. The normal response is easy palpation of the muscle. If guarding is present, determine if it is voluntary or involuntary by placing a pillow under the patient's knees and asking

him or her to take several slow, deep breaths. Palpate the rectus abdominis muscles on expiration. The patient cannot voluntarily tense this muscle during expiration, so if involuntary guarding is present, you will feel a boardlike rigidity that indicates peritonitis.

Deep palpation is used to assess organs, masses, and tenderness. It can be done using a manual or bimanual technique. To perform single-handed deep palpation, use the distal portion of your fingertips and depress 4 to 6 cm in a dipping motion in all four quadrants, assessing for masses or areas of tenderness. To perform bimanual deep palpation, place your nondominant hand on your dominant hand, then depress your hands 4 to 6 cm. Bimanual palpation is useful when palpating a large abdomen. Tenderness may be noted in a normal adult near the xiphoid or over the cecum or sigmoid colon. If you find a mass, note its location, size, shape, consistency (soft, firm, hard), tenderness, pulsation, mobility, and movement with respiration.

PALPATION OF THE ABDOMEN

1. Abdomen

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Light Palpation

- Lightly palpate about $\frac{1}{2}$ inch in each quadrant; palpating painful areas last.
- Note surface characteristics and areas of tenderness.



Light palpation



Light palpation using patient's hands

Deep Palpation

- *Single hand technique*: Use dominant hand and palpate greater than $\frac{1}{2}$ inch in each quadrant.
- *Bimanual technique*: Place dominant hand on abdomen, place nondominant hand on top, and palpate greater than $\frac{1}{2}$ inch in each quadrant.
- *Abdomen soft and nontender*.
- *No organomegaly or masses, nontender*.



Single-handed deep palpation



Bimanual deep palpation

ABNORMAL FINDINGS/RATIONALE

- *Involuntary guarding and rigidity*: Peritonitis.
- *Areas of tenderness*: Indicates underlying problem.
- *Masses*: May indicate underlying tumor, enlarged uterus, feces-filled colon.
- *Enlarged liver*: May indicate cirrhosis, tumor, hepatitis.
- *Enlarged spleen*: May indicate malignancy, infection, trauma to spleen.



HELPFUL HINTS

- If the patient is ticklish, have her or him place a hand over her or his abdomen and then place yours over it and do light palpation. When the patient starts to feel more relaxed, slip your hand underneath.
- If you detect a mass, have the patient tighten his or her abdominal muscles. If the mass is in the abdominal wall, it will become easier to palpate. If the mass is deep in the abdomen below the muscles, it will be difficult to palpate.



ALERT

- Do not palpate patients who have had an organ transplant.
- Do not palpate the abdomen of a child with suspected Wilms' tumor because it may cause the tumor to seed into the abdomen.

Assessing Abdominal Structures

While assessing the abdomen, you will need to examine specific abdominal structures. The following section describes the examination of these structures and explains the difference between normal and abnormal findings.

Abdominal Aorta

To palpate the abdominal aorta, place your fingers in the epigastric portion of the abdomen and slightly toward the patient's left midclavicular line. Palpate for aortic pulsations. You can also assess the width of the aorta by placing one hand on each side of the aorta.

Liver

To palpate the liver, place your right hand at the patient's right midclavicular line under the costal margin. Place your left hand posteriorly on the patient's right eleventh to twelfth rib and press upward to elevate the liver toward the abdominal wall. Have the patient inhale and exhale deeply while you press your right hand gently but deeply in and up during inspiration.

The hooking technique is another way to palpate the liver. Place your hands over the right costal margin and hook your fingers over the edge. Have the patient take a deep breath and feel for the liver's edge as it drops down on inspiration, then rises up over your fingers during expiration. (See *Palpating the Liver*.)



ALERT

Always percuss the spleen before palpation. Palpate gently so that you do not rupture an enlarged spleen.

Spleen

To palpate the spleen, stand on the patient's right side, place your left hand under the left CVA, and pull upward to move the spleen anteriorly. Place your right hand under the left anterior costal margin and have the patient take a deep breath in and out. During exhalation, press inward along the left costal margin and try to palpate the spleen. (See *Palpating the Spleen*.)

Kidneys

To assess the left kidney, stand on the patient's right side and place your left hand in the left CVA of her or his back. Place your right hand at the left anterior costal margin. Have the patient take a deep breath, then press your hands together to "capture" the kidney. As the patient exhales, lift your left hand and palpate the kidney with your right hand.

To assess the right kidney, remain on the patient's right side and place your right hand on the right posterior CVA and your left hand on the patient's right anterior costal margin. When the patient exhales, palpate the kidney. (See *Palpating the Kidneys*.)

Is It a Kidney or the Liver?

An enlarged liver is usually larger than an enlarged kidney. An enlarged liver extends along the right costal margin to the xiphoid, whereas an enlarged kidney does not. An enlarged liver has a sharper edge than an enlarged kidney, which is more rounded. You cannot "capture" a liver.

PALPATING THE ABDOMINAL AORTA

1. Aortic Pulsations

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Palpate in the epigastric area at the midline for abdominal aortic pulsation.



ALERT

Do not palpate a patient who has a suspected abdominal aortic aneurysm. It can cause the aneurysm to rupture, resulting in renal failure, loss of limbs, or even death.

- Aorta pulsates in an anterior direction, no diffusion. Normal width no greater than 3 cm.*



Palpating aortic pulsations

ABNORMAL FINDINGS/RATIONALE

- Lateral diffuse pulsation:* Abdominal aortic aneurysm.
- Width greater than 3 cm:* Abdominal aortic aneurysm.

PALPATING THE LIVER

1. Liver

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Stand at the right side of your patient.
- Place left hand under the patient's back at the CVA, and place right hand along the costal margin at the right midclavicular line.
- Have patient take deep breath while you press your right hand in and up, while at the same time with your left hand, press upward to elevate the liver.
- *Alternate technique (hooking technique):* Stand on right side of patient facing toward patient's feet, place your hands over the patient's right costal margin, and hook fingers over the edge. Have patient take a deep breath. Feel for the liver's edge to drop down as the patient inhales and rise up over your fingers as the patient exhales.



Palpating the liver



Hooking technique

- *The liver's edge drops down during inspiration and rises up during expiration.*
- *The liver is normally not palpable unless the patient is very thin. If it is palpable, the edge should be smooth and nontender.*

ABNORMAL FINDINGS/RATIONALE

- *Palpation below costal margin:* Congestive heart failure, hepatitis, encephalopathy, cirrhosis, cysts, or cancer. If liver is enlarged, tender, firm, and nodular or has an irregular border, suspect liver cancer.

PALPATING THE SPLEEN

1. Spleen

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Stand at the right side of your patient.
- Place right hand under the patient's back at the left CVA, and place left hand along the right costal margin.
- Have patient take a deep breath while you press inward along the costal margin.
- *The spleen is normally not palpable.*



Palpating the spleen

ABNORMAL FINDINGS/RATIONALE

- *Splenic enlargement and tenderness:* Infection, congestive heart failure, cancer, cirrhosis, or trauma.

PALPATING THE KIDNEYS

1. Kidneys

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- To palpate the right kidney, stand at right side of your patient, and place your right hand under patient at the CVA and your left hand below the right costal margin.
- “Capture” the kidney by pressing hands together as the patient breathes.
- Use the same technique to palpate the left kidney, placing the left hand under the patient at the left CVA, and the right hand below the left costal margin.
- *Kidneys are normally not palpable unless the patient is very thin or elderly (owing to loss of muscle tone), and then only the right kidney is usually palpable. If kidneys are palpable, they should be smooth and nontender.*



Palpating the kidneys

ABNORMAL FINDINGS/RATIONALE

- **Enlarged kidneys:** Hydronephrosis, neoplasm, or polycystic disease.
- **Kidney tenderness:** Trauma or infection.

Bladder

Palpate the bladder in the hypogastric area up to the umbilicus, using deep palpation. (See *Palpating the Bladder*.)

Inguinal Lymph Nodes

Using the pads of your fingers, palpate just below the inguinal ligament for the superficial superior (also called the horizontal) inguinal lymph nodes and along the inner aspect of upper thigh for the superficial inferior (also called the vertical) inguinal lymph nodes. If nodes are palpable, note size, shape, mobility, consistency, and tenderness (See *Palpating Inguinal Lymph Nodes*).

Additional Abdominal Assessment Techniques

In addition to inspection, auscultation, percussion, and palpation, several other techniques are used in assessing the abdomen. They are described in the following sections.

Assessing for Ascites

Ascites is an accumulation of fluid in the peritoneal cavity. It is usually detectable when 500 mL of fluid has accumulated. Causes of ascites include congestive heart failure, cirrhosis, nephrosis, and malignancies. There are three ways to assess for ascites: shifting dullness, fluid wave, and the puddle sign. If your patient has ascites, measure abdominal girth at the umbilicus and weigh the patient daily (Fig. 17.10).

- **Shifting dullness:** Have the patient lie supine and percuss the abdomen for dullness and tympany (Fig. 17.11). Normally, only tympany is present. In a patient with ascites, you will hear dullness laterally and tympany near the umbilical area. To further assess for ascites,

Is It an Enlarged Left Kidney or an Enlarged Spleen?

The spleen has a palpable notch on the medial edge. Percussion of tympany in the LUQ favors an enlarged kidney rather than an enlarged spleen.

have the patient lie on his or her right side and percuss again. You will note tympany on the left side of the abdomen and dullness on the right lateral aspects to the midline section. Next have the patient turn on his or her left side and percuss the abdomen (Fig. 17.12). You will hear dullness from the midline to left lateral area and tympany on the right lateral area. This shifting dullness indicates ascitic fluid of greater than 500 mL.

- **Fluid wave:** Have the patient place her or his hand vertically in the middle of her or his abdomen. Place



FIGURE 17.10. Assessing abdominal girth.




FIGURE 17.11. Percussing shifting dullness with patient supine.




FIGURE 17.12. Shifting dullness.

your hands on each side of the patient's abdomen and tap one side while palpating the other side. If ascites is present, the tap will cause a fluid wave through the abdomen and you will feel the fluid with the other hand (Fig. 17.13).

- **Puddle sign:** Ask the patient to kneel with his or her hands on the examining table. Percussing the umbilicus normally produces tympany. If the area is dull on percussion, suspect ascites because fluid pools in the dependent areas. This is the least-popular technique to assess for fluid because it may be difficult for your patient to assume the position.




FIGURE 17.13. Fluid wave test.

Abdominal Reflexes

The abdominal reflexes are superficial reflexes. To assess them, you will need a tongue blade or a reflex hammer. The upper abdominal reflexes assess T8 through T10 and are obtained by using the end of the reflex hammer or tongue blade and stroking upward and toward the umbilicus. The lower abdominal reflexes assess T10 through T12 and are obtained by stroking downward and toward the umbilicus. The normal response is movement of the umbilicus toward the stimulation. Because this is a superficial reflex, it is either positive or negative. Absent abdominal reflexes can occur when abdominal muscles are overstretched, as in pregnancy.

Ballottement

Ballottement is used to displace fluid so that a floating mass or organ can be palpated.

To perform ballottement with one hand, place your hand perpendicular to the abdomen and push in toward the mass with your fingertips. A freely movable mass will float toward your fingertips. To use the bimanual method, place one hand on the anterior abdomen and push down while the other hand is placed against the flank to push up and palpate the mass to determine the size and shape. This

PALPATING THE BLADDER

1. Bladder

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Palpate above the symphysis pubis at the midline.
- *The bladder is usually not palpable unless it is distended with urine. If palpable, it feels like a smooth, round mass.*



Palpating the bladder

ABNORMAL FINDINGS/RATIONALE

- *Nodular or asymmetrical bladder:* May indicate malignancy.

PALPATING INGUINAL LYMPH NODES

1. Inguinal Lymph Nodes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Palpate the horizontal lymph nodes in the groin.
- Palpate the vertical lymph nodes along the inner aspect of the thigh.



Palpating the horizontal inguinal lymph nodes



Palpating the vertical inguinal lymph nodes

- *If nodes are palpable, they should be less than 1 cm, smooth, mobile, and nontender.*

ABNORMAL FINDINGS/RATIONALE

- *Tender, unmovable nodes greater than 1 cm: Infection, cancer, or lymphoma.*

technique is frequently used to assess fetal position, but it may also be used to detect partially attached masses.

Kehr's Sign

Referred pain or hyperesthesia to the left shoulder is called **Kehr's sign**. It results from irritation to the hemidiaphragm caused by a splenic rupture with blood accumulating in the peritoneal cavity. Movement of the shoulder joint does not elicit pain, ruling out a musculoskeletal problem.

Ballance's Sign

Ballance's sign is present when dullness is percussed in the left upper quadrant (LUQ). This indicates peritoneal irritation or injury to the spleen.

Murphy's Sign

Murphy's sign is used to detect an inflamed gallbladder as seen in cholecystitis. Stand at the patient's right side

(Text continued on page 599)

OTHER PHYSICAL ASSESSMENT TECHNIQUES FOR THE ABDOMEN

1. Abdominal Reflexes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Using a tongue blade, stroke the abdomen upward and toward the umbilicus in each quadrant.
- *Positive abdominal reflexes. Diminished responses may be found in obese patients or those who have been pregnant. Abdominal reflexes may be absent in elderly patients.*



Assessing abdominal reflexes

(continued)

OTHER PHYSICAL ASSESSMENT TECHNIQUES FOR THE ABDOMEN

1. Abdominal Reflexes (continued)

ABNORMAL FINDINGS/RATIONALE

- *Absent abdominal reflexes*: Pyramidal tract lesion.
- *Absent upper abdominal reflexes*: Problems at spinal levels T8, T9, and T10.
- *Absent lower abdominal reflexes*: Problems at spinal levels T10, T11, and T12.

2. Ballottement

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Using bimanual technique, place one hand on the abdomen over mass and push inward toward mass. Place the other hand at the opposite end of object.
- *A partially free-floating object will move toward the other hand.*
- *In normal adults, no masses or organs should be felt. In pregnant patients, this technique is used to determine fetal position.*



Ballottement

ABNORMAL FINDINGS/RATIONALE

- *Free-floating mass in abdomen*: Malignant or benign tumor.

3. Kehr's Sign

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- With the patient supine, move patient's arm upward.
- Note referred pain to left shoulder.
- *Not normally present.*



Palpating shoulder to elicit Kehr's sign

ABNORMAL FINDINGS/RATIONALE

- *Positive Kehr's sign*: Splenic injury, renal calculi, or ectopic pregnancy.

4. Ballance's Sign

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Percuss the left upper quadrant for dullness.
- Note tenderness.
- *Not normally present.*



Percussing LUQ to elicit Ballance's sign

OTHER PHYSICAL ASSESSMENT TECHNIQUES FOR THE ABDOMEN

4. Ballance's Sign (continued)

ABNORMAL FINDINGS/RATIONALE

- *Positive Ballance's sign:* Peritoneal irritation or injury to spleen.

5. Murphy's Sign

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Stand at right side of patient, and palpate at the right midclavicular line under the costal angle as the patient takes a deep breath.
- Note pain and restricted breathing.
- *Not normally present.*



Palpating at right midclavicular line to elicit Murphy's sign

ABNORMAL FINDINGS/RATIONALE

- *Positive Murphy's sign:* Cholecystitis or carcinoma of the gallbladder.

and palpate at the right midclavicular line under the costal angle. When the patient takes a deep breath, the gallbladder moves closer to your hand and causes pain, which stops the patient from inhaling (inspiratory arrest).

The presence of pain is a positive Murphy's sign.

Assessing for Appendicitis

Various tests can be performed to assess for appendicitis. Pain during any of these tests is a positive sign.

Assessing Drains and Tubes

If your patient has an external drain, tube, or diversion (stoma), you need to evaluate how well it is functioning.

(Text continued on page 602)

ABDOMINAL SIGNS OF APPENDICITIS

1. McBurney's Sign

TEST/NORMAL VARIATIONS

- Located in RLQ, McBurney's point is one-third of the distance from anterior iliac crest to the umbilicus.
- Perform rebound tenderness test at this point, assessing for pain.
- *Pain should not occur.*



McBurney's point

ABNORMAL FINDINGS/RATIONALE

- *Pain (positive McBurney's sign):* Appendicitis.

(continued)

ABDOMINAL SIGNS OF APPENDICITIS

2. Obturator Muscle Test

TEST/NORMAL VARIATIONS

- Helps diagnose ruptured appendix or pelvic abscess.
- Patient lies supine and flexes right leg at hip and knee.
- Place one hand just above patient's knee and other hand at ankle, and rotate leg internally and externally.
- *Pain should not occur in hypogastric region.*



Obturator muscle test

ABNORMAL FINDINGS/RATIONALE

- *Pain (positive obturator muscle test):* Irritation of obturator muscle, which may be caused by a ruptured appendix or pelvic abscess.

3. Rovsing's Sign

TEST/NORMAL VARIATIONS

- Assesses peritoneal inflammation caused by appendicitis.
- Place your hand in LLQ and press deeply for 5 seconds.
- *Pain should not occur.*



Rovsing's sign

ABNORMAL FINDINGS/RATIONALE

- *RLQ pain (positive Rovsing's sign):* Peritoneal irritation and/or appendicitis.

4. Cutaneous Hypersensitivity

TEST/NORMAL VARIATIONS

- Grasp a fold of skin or touch the abdominal surface with an open safety pin to assess for pain.
- Assess entire abdominal surface.
- *Pain should not occur.*



Cutaneous hypersensitivity

ABNORMAL FINDINGS/RATIONALE

- *Cutaneous hypersensitivity:* Indicates peritoneal irritation.
- *RLQ pain:* Appendicitis.
- *Pain in middle epigastric region:* Peptic ulcer.

ABDOMINAL SIGNS OF APPENDICITIS

5. Iliopsoas Muscle Test

TEST/NORMAL VARIATIONS

- Used when an inflamed or perforated appendix is suspected.
- As patient lies supine, place your hand over her or his lower right thigh. Ask the patient to raise her or his right leg by flexing the hip while you push downward.
- Assess for pain in the RLQ.
- *Pain should not occur in RLQ.*



Iliopsoas muscle test

ABNORMAL FINDINGS/RATIONALE

- *RLQ pain (positive iliopsoas muscle test):* Iliopsoas muscle irritation caused by inflamed or perforated appendix.

6. Rebound Tenderness

TEST/NORMAL VARIATIONS

- Place your hand perpendicular to the abdomen, press firmly and slowly, and then release quickly.
- *Pain should not occur.*



HELPFUL HINT

Perform this test only if patient complains of abdominal pain.



Rebound tenderness test

ABNORMAL FINDINGS/RATIONALE

- *Rebound tenderness (pain that increases when you remove your hand):* Peritoneal irritation.
- *Rebound tenderness in RLQ (McBurney's point):* Appendicitis.

Assessing Drains and Tubes

TUBE/DRAIN/DIVERSION

ASSESS

Nasogastric Tubes

- Inserted into nose and advanced into stomach. Includes Levin's tubes and Salem sumps. These tubes are used to drain stomach contents or administer tube feedings.
- Assess skin around insertion site for signs of drainage or breakdown.
- Check tube to be sure it is secured safely to nose and then more distally to patient's gown.
- Check tube placement every shift and p.r.n. Placement can best be determined by obtaining gastric contents and testing pH and bilirubin levels.
- Assess amount, color, and consistency of fluid if tube is used for draining secretions.

(continued)

Assessing Drains and Tubes <small>(continued)</small>	
TUBE/DRAIN/DIVERSION	ASSESS
<p>Gastrostomy and Jejunostomy Tubes (G-Tubes and J-Tubes)</p> <ul style="list-style-type: none"> ■ Inserted directly into stomach or jejunum. ■ Used to drain GI secretions or deliver tube feedings. <p>Abdominal Cavity Drains (Biliary, Penrose, Jackson-Pratt, Hemovac, or Constavac Drains)</p> <ul style="list-style-type: none"> ■ Used to remove fluid or secretions from abdominal cavity. <p>Intestinal Diversions (Colostomy or Ileostomy)</p> <ul style="list-style-type: none"> ■ Provide an alternate route for evacuation of waste products out of body. Feces are not expelled through patient's rectum. <p>Urinary Diversions (Ileal Conduit, Ureteral Stent, Indwelling Catheter)</p>	<ul style="list-style-type: none"> ■ Evaluate skin around tube every shift for signs of drainage or breakdown. ■ If the patient has a percutaneous endoscopic gastrostomy (PEG) tube, rotate disk every day to prevent skin breakdown. ■ Assess amount, color, and consistency of fluid if tube is used for draining secretions. ■ Check residual volumes every 4 hours if tube is used to administer enteral feedings. ■ Assess drain for patency. ■ Evaluate color, consistency, amount, and odor of secretions. ■ Note location of drains in your nurse's notes and whether they are sutured in place. ■ Assess area surrounding drain for breakdown. ■ Assess stoma for color. Normal color is pink/red. ■ Purple stoma indicates decreased circulation to stoma and needs immediate attention. ■ Check areas surrounding stoma for skin breakdown. ■ Note color, odor, amount, and consistency of feces excreted through ostomy. ■ Record location of stoma in your nurse's notes. ■ <i>Ileal conduits:</i> Check stoma for color and skin surrounding it for areas of breakdown. ■ Note color, odor, and amount of urine from each of these urinary diversions. ■ Record location of diversions in your nurse's notes.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Robichaud have concluded. The following are your findings:

■ General Health Survey

- Well-nourished, obese 56-year-old woman who appears older than her years.
- Awake, alert, oriented × 3, memory intact.
- No distress noted.
- Posture erect, pleasant and cooperative, provides history with ease.

- In no acute distress, but mildly anxious.
- Head-to-toe scan revealed no pertinent findings related to abdomen.

■ Vital Signs

- Temperature, 98.8°F.
- Pulse, 86 and regular.
- Respirations, 22.
- BP, 168/98.
- Height, 5'2".
- Weight, 175 pounds.

■ Abdominal

- Abdomen uniformly light pink in color.
- No striae, bruises, or hernias noted; umbilicus centered; scar from previous cholecystectomy.
- Abdomen protuberant and slightly distended in lower half.
- No aortic pulsations; peristalsis noted.
- Hypoactive bowel sounds in all four quadrants.
- No bruits, friction rubs, or venous hums.
- Liver span 6.5 cm at midclavicular line.
- Unable to percuss bladder or spleen.
- No CVA tenderness.
- Abdomen soft and slightly tender in lower quadrants.
- Liver, spleen, kidneys, bladder, and inguinal nodes not palpable.
- Abdominal aorta approximately 2 cm, palpable just left of midline.
- Superficial abdominal reflexes intact.
- No ascites noted with shifting dullness test.

appropriate nursing diagnoses. Below are some possible diagnoses; cluster the supporting data for each.

1. Constipation, related to decreased activity and low-fiber diet
2. Imbalanced Nutrition: more than body requirements
3. Ineffective Health Maintenance related to dietary and exercise habits
4. Knowledge deficit related to high risk for colon cancer

Identify any additional nursing diagnoses.

Identify any collaborative nursing diagnoses.



CRITICAL THINKING ACTIVITY # 4

Now that you have identified some nursing diagnoses for Mrs. Robichaud, select one from above and develop a nursing care plan and a brief teaching plan for her, including learning outcomes and teaching strategies.



Case Study Analysis and Plan

Now that you have completed a thorough assessment of Mrs. Robichaud, document the key history and physical examination findings that will help you formulate your nursing diagnoses.



CRITICAL THINKING ACTIVITY # 5

What additional questions could you ask Mrs. Robichaud about her risk for constipation?

Nursing Diagnoses

Your next step is to analyze the data from Robichaud's assessment and group the data to determine the appro-



CRITICAL THINKING ACTIVITY # 6

What nursing interventions can you initiate to help her relieve her constipation?

RESEARCH TELLS US

Colorectal cancer is the third most common type of cancer and the second leading cause of cancer-related death in the United States. Even though colorectal screening tests have been recommended for persons 50 years or older since the mid-1990s, the screenings remain underused by the public.

The Centers for Disease Control and Prevention (CDC) analyzed colorectal cancer screening trends in the United States between 1997 and 1999, using the Behavioral Risk Factor Surveillance System (BRFSS) for fecal occult blood test (FOBT) and sigmoidoscopy/

colonoscopy. In 1997, 41 percent of the respondents reported screening within the recommended period as compared with 44 percent in 1999. Specifically, in 1997, 19.6 percent had FOBT and 30.3 percent had sigmoidoscopy/colonoscopy as compared with 20.6 percent FOBT and 33.6 percent sigmoidoscopy/colonoscopy in 1999. Although there were limitations associated with the survey, the rate of changes in testing from 1997 to 1999 were statistically significant ($p < .05$).

Even though the use of screening seems to be increasing, the proportion of the population using

(continued)

RESEARCH TELLS US (continued)

screening is low. Walsh, Posner, and Perez-Stable's (2002) report of a retrospective review of medical records from three primary care practices from 1995 to 1997 (n = 6039) to identify colon cancer screening among patients between ages 50 and 74 years revealed disturbing statistics:

- Only half of primary care patients were screened for colorectal cancer.
- Women were less likely to be screened than men.
- Only one-third of high-risk patients with family history of colon cancer were screened.

- Patients of nurse practitioners were 22 percent less likely to receive FOBT and patients of medical residents 21 percent less likely to receive FOBT than those seen by a physician/faculty.

It's important to remember that colorectal screening tests have proven effective in reducing the incidence and mortality rates associated with colorectal cancer. As health promoters, nurses are in a key position to educate the public and increase public awareness of colorectal cancer and preventative screening.

Source: Walsh, J., Posner, S., Perez-Stable, E. (2002). Colon cancer screening in the ambulatory setting. *Preventive Medicine* 35(3):209-218; and CDC, Trends in screening for colorectal cancer—United States, 1997 and 1999. www.cdc.gov/mmwr/preview/mmwrhtml/mm5009a2.htm

HEALTH CONCERNS

The major health concern for the GI system is the prevention of colorectal cancer. Risk factors for colorectal cancer include a family history of colorectal cancer, a personal history of colorectal cancer, a personal history of polyps, a personal history of inflammatory bowel disease, ethnicity (Ashkenazi Jews), aging (90 percent of patients diagnosed with colorectal cancer are over 50 years old), a diet consisting mostly of food from animal sources, and physical inactivity. Obesity, diabetes, smoking, alcohol, and night shift work have increased risk for colorectal cancer.

The American Cancer Society has identified the following preventative measures for colorectal cancer:

- **Exercise:** Exercise 30 minutes on 5 or more days a week.
- **Diet:** Consume a diet rich in fiber and low in fat, and supplement dietary intake with a multivitamin rich in folic acid, vitamin D, and calcium.
- **Medication:** Patients using nonsteroidal anti-inflammatory drugs (NSAIDs) for other conditions such as arthritis were found to have lower risk of colorectal cancer. Aspirin also has been found to prevent polyp growth. However, this therapy is not without adverse effects, so it is important to weigh benefits. (Because NSAIDs can cause severe stomach irritation and bleeding, this therapy is not recommended for people with average risk for colorectal cancer. Use of this therapy for high-risk patients is being studied.)
- **Hormone replacement therapy:** Hormone replacement therapy (HRT) in postmenopausal women

has been found to reduce risk of colorectal cancer, but women on HRT who develop colorectal cancer have a faster-growing cancer. HRT is also not without adverse effects (heart disease, blood clots, and other types of cancer); therefore, it is important to weigh benefits.

The key to successful treatment of colorectal cancer is early detection and treatment.

The American Cancer Society recommends the following screening measures:

- If you are over age 50 and have average risk for colorectal cancer, you should have one of the following five screening tests:

1. Yearly test for occult blood (FOBT) in stool or fecal immunochemical test (FIT).
2. Flexible sigmoidoscopy every 5 years.
3. FOBT or FIT every year plus a flexible sigmoidoscopy every 5 years.

OR

4. Double-contrast barium enema every 5 years.
5. Colonoscopy every 10 years.

Of the first three options, a FOBT or FIT every year plus a flexible sigmoidoscopy every 5 years is preferred.



HELPFUL HINT

If you are at high risk for colorectal cancer, undergo screening more often and earlier than other people.

Source: American Cancer Society, 2005 (<http://www.cancer.org>)

Common Abnormalities

ABNORMALITY

ASSESSMENT FINDINGS

Abdominal Aortic Aneurysm

Life-threatening outpouching of abdominal aorta, one of major blood vessels. Ruptured abdominal aortic aneurysm is a surgical emergency and must be treated immediately to prevent hemorrhages, shock, and possible death.

Appendicitis

Acute condition caused by inflammation of vermiform appendix, which may rupture, causing peritonitis.

Cancer of the Colon

Occurs most frequently in descending colon, rectum, and sigmoid colon. The colon is often the primary site of cancer in the abdomen.

Cancer of the Stomach

Difficult to detect on physical examination.

Cholecystitis

Inflammation of gallbladder, often accompanied by stone formation (cholelithiasis). Stones may obstruct bile ducts, causing bouts of colicky, acute pain in the RUQ and obstructive jaundice. Referred pain also occurs.

Cirrhosis

A degenerative disease of the liver, cirrhosis results in fibrosing and death of liver cells, which alters liver function and produces symptoms in multiple body systems. Commonly associated with alcoholism and hepatitis.

Constipation

Delayed movement of stool through large intestine, resulting in fewer bowel movements than usual. Because of length of time in intestine, more water is absorbed, causing stool to become dry, hard, and difficult to pass. Bowel movement may be painful. Fecal impaction, a

- Patient complains of boring, tearing pain; referred pain.
 - Auscultation reveals bruits, exaggerated pulsations, or a mass.
 - Diminished or diffuse femoral pulses.
 - Hypotension, tachycardia, increased respirations.
 - *Signs of shock*: pale, cool, clammy skin; decreased or absent pulses; cool extremities.
-
- Abrupt onset of constant pain unrelieved by changes in body position. Pain becomes sharp and localized.
 - Anorexia, nausea, vomiting follow complaints of pain.
 - Rigid abdomen, decreased bowel sounds.
 - Positive rebound tenderness (Blumberg's sign), positive iliopsoas sign, positive obturator muscle response.
 - Increased temperature (tympanic temperature is preferred if patient is nauseated).
-
- Symptoms include changes in bowel habits; blood in stool; and changes in quality of stool, particularly reduction in diameter.
 - Pain (late sign of rectal cancer).
-
- Associated with epigastric distress, sense of abdominal fullness, anorexia, and weight loss.
 - In late stages, ascites and lymph node enlargement are seen.
-
- Positive Murphy's sign, indigestion.
 - May be asymptomatic.
-
- Liver and spleen are enlarged and palpable; in advanced stages, liver is smaller and nodular.
 - *Other symptoms*: Ascites and caput medusa, changes in bowel sounds, anorexia, nausea, vomiting, dyspepsia, dull abdominal pain, hemorrhoids, jaundice, bruising, spider angiomas, portal hypertension, edema.
 - Encephalopathy; peripheral neuropathy with numbness and loss of sensation in extremities; asterixis (flapping hand tremor related to liver malfunction).
 - Amenorrhea in women; testicular atrophy, gynecomastia, and impotence in men.
-
- Bowel sounds on auscultation.
 - Palpation of feces in lower and possibly upper quadrants.
 - Complaints of physical discomfort.

(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>serious condition, occurs when dry, hard feces remain in rectum or colon. Prompt intervention is needed to relieve the situation and prevent life-threatening bowel obstruction.</p>	
<p>Diverticulitis Diverticula are normal structures of the colon. When they become enlarged, they fill with fecal matter, which causes inflammation (diverticulitis) and possible infection or abscess.</p>	<ul style="list-style-type: none"> ■ Pain or discomfort. ■ Diminished bowel sounds with auscultation. ■ Nausea and vomiting. ■ Constipation.
<p>Hepatitis A liver disease caused by different viruses and transmitted by blood and bodily fluids. Different types include A, B, C, D, and E. Viral hepatitis A is the least problematic. Hepatitis B can be a lengthy illness with serious complications, including death, and is a necessary precursor of Hepatitis D. Hepatitis C has been associated with complications in approximately 50 percent of patients. Hepatitis D is a long-term illness. Hepatitis E is relatively uncommon and primarily occurs during disaster situations.</p>	<ul style="list-style-type: none"> ■ Anorexia. ■ Malaise. ■ Nausea and vomiting. ■ Ascites. ■ Enlarged liver and spleen. ■ Jaundice. ■ Pruritus. ■ Amber-colored urine. ■ Gray-colored stool.
<p>Hernias Inguinal Hernia Most common type of hernia. Usually occurs in men and is caused by a muscular weakness that allows the intestine to protrude through the abdominal wall, groin, or scrotum. Usually, these protrusions can be put back in place with gentle pressure.</p>	<ul style="list-style-type: none"> ■ Discomfort worsens with prolonged standing and may diminish with rest. ■ Pain or a feeling of fullness occurs when straining for elimination.
<p>Umbilical Hernia Usually occurs in infants and children and is caused by an abdominal wall weakness around the umbilicus that allows a portion of an internal organ to protrude.</p>	<ul style="list-style-type: none"> ■ Protrusion becomes more prominent when an infant or child cries.
<p>Hiatal Hernia Causes the stomach to move through the esophageal hiatus (opening) and rise above the diaphragm.</p>	<ul style="list-style-type: none"> ■ May cause esophageal constriction, acid reflux into the esophagus, and subsequent esophageal damage.
<p>Incisional Hernia Occurs at site of a previous abdominal surgery. The most serious problem associated with hernias is the possibility of tissue death when circulation to a protruding intestine or abdominal organ becomes compromised (incarcerated or strangulated hernia).</p>	
<div style="border: 1px solid red; padding: 5px;"> <p> ALERT</p> <p>A strangulated hernia is a surgical emergency. Symptoms include severe increasing pain, vomiting, fever, and bloody stools.</p> </div>	
<p>Inflammatory Bowel Diseases Crohn's Disease Common chronic inflammatory bowel disease affecting ileum and colon that results in thickening of bowel and narrowing of intestinal lumen. Affects small intestine;</p>	<ul style="list-style-type: none"> ■ Abdominal pain. ■ Steatorrhea. ■ Diarrhea.

ABNORMALITY

therefore, problems with vitamin absorption are common. Significantly increases risk for colon cancer and arthritis.

Ulcerative Colitis

Inflammation of mucosa of rectum and sigmoid colon that causes cramping, diarrhea, and rectal bleeding and increases risk for colon cancer and bowel perforation. Has exacerbations and remissions.

Pancreatitis

Inflammation of pancreas that alters flow of enzymes to intestines and results in pancreatic retention of digestive enzymes. These enzymes cause breakdown of the pancreas.

Paralytic Ileus

Lack of peristalsis, usually in the small intestine. Serious condition requiring prompt treatment. Surgery, peritonitis, or spinal injury is a predisposing factor. Bowel necrosis and perforation are serious complications.

Peritonitis

Inflammation of the lining of the abdominal cavity (peritoneum).

Splenic Rupture

Serious abdominal condition resulting in hemorrhage. Usually follows abdominal trauma, but also may be a serious sequela of infectious mononucleosis.

ASSESSMENT FINDINGS

- Vitamin deficiency states.
 - Fatigue.
 - Weight loss.
 - Fever.
 - Arthritis.
 - Joint deformities.
-
- Watery eyes.
 - Conjunctivitis.
 - Mouth ulcers.
 - Erythema nodosum.
 - Pyoderma gangrenosum.
 - Cramping pain in lower abdomen.
 - Watery diarrhea.
 - Rectal bleeding.
 - Mucus in stool.
 - Frequent bowel movements.
 - Urgency.
 - Dehydration.
 - Fever.
 - Weight loss.
-
- Nausea and vomiting.
 - Weight loss.
 - Severe "boring" pain in LUQ.
 - Pain referred to back or shoulder.
 - Radiating pain.
 - Steatorrhea.
-
- Intermittent colicky pain.
 - Absence of bowel sounds.
 - Initially visible peristaltic waves.
 - Abdominal distension.
 - Vomiting.
-
- Assessment findings include abdominal pain of varying character, cutaneous hypersensitivity.
 - Abdominal rigidity and guarding.
 - Diminished bowel sounds.
 - Positive iliopsoas, Blumberg's, and Rovsing's signs.
 - Positive Murphy's sign.
 - Fever.
 - Nausea and vomiting.
-
- Severe LUQ pain.
 - Left shoulder pain.
 - Possible shock (drop in BP).

(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Ulcer</p> <p>Forms when gastric mucosa becomes permeable, when amount of protective mucus is reduced (in conditions such as gastritis), or when there is prolonged exposure to bile or other irritating substances (e.g., medications). GI ulcer sites include duodenum (duodenal ulcer) and stomach (gastric ulcer). Stress ulcers are erosions of the GI tract mucosa that arise in relation to trauma, burns, or severe illness.</p>	<ul style="list-style-type: none"> ■ Gastric ulcers include complaints of gnawing pain, heartburn, anorexia, vomiting (hematemesis possible), eructations, and weight loss. <div style="border: 1px solid red; padding: 5px; margin: 10px 0;"> <p> ALERT</p> <ul style="list-style-type: none"> • Suspect anorexia nervosa if your patient has a weight loss of 15 percent or more below ideal height-weight standards, says that he or she actively pursues weight loss, or expresses a distorted or almost delusion attitude about his or her body (e.g., says, “I’m fat” in spite of thin appearance). Assessment findings include starvation, decreased fluid intake, thin hair, dental erosions, dehydration, electrolyte imbalance, swollen parotid glands, decreased body temperature, dry skin, poor skin turgor, weight loss, thinness, general weakness, and constipation. • Bulimia nervosa is harder to pinpoint than anorexia. Patients may be of normal weight or slightly over- or underweight. They binge and purge, ingesting huge amounts of food during a brief period and then inducing vomiting or diarrhea. Guilt feelings are common. Suspect bulimia nervosa if your patient exercises excessively, has eroded tooth enamel (from vomiting), and enlarged parotid glands. </div> <ul style="list-style-type: none"> ■ Duodenal ulcer include intermittent RUQ pain 2 to 3 hours after meals, remissions, and exacerbations; stool may be positive for guaiac.

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the abdomen.

Ask the patient the following:

- Do you have:
 - Abdominal pain?
 - Change in bowel patterns?
 - Nausea?
 - Indigestion?
 - Vomiting?
 - Weight changes?
- Have you ever had the following: stomach ulcer, hemorrhoids, hernia, bowel disease, cancer, hepatitis, cirrhosis, or appendicitis?
- Have you had abdominal surgery? If so, when, what type, and were there any subsequent problems?
- Do you have a family history of ulcer, gallbladder disease, bowel disease, or cancer?
- Do you have any problems with swallowing, heartburn, yellowing of your skin, gas, or bloating?
- Do you have any food allergies or lactose intolerance?
- What is your usual weight and height?

ASSESSMENT CHECKLIST (continued)

- How is your appetite? What did you eat in the last 24 hours?
- How is your health usually?
- Are you currently being treated for a health problem? If so, what?
- How often do you usually have a bowel movement?
- Are you having problems with diarrhea, constipation, hemorrhoids, or fecal incontinence? Have you ever noticed blood in your stool or had black, tarry stools?
- How often do you urinate? Do you have incontinence or burning when you urinate?
- When was your last menstrual period?
- Do you smoke? How many packs a day? (Calculate pack-years.)
- Do you drink alcohol? If so, how often? Do you use street drugs?
- How many cups of coffee, tea, or caffeinated soda do you drink every day?
- Have you been exposed to an infectious disease recently?
- What is your occupation?
- Have you been immunized against hepatitis B?
- Have you ever had a blood transfusion? If so, when?
- Do you take any prescribed medications? What are they?
- Do you have any allergies to medications?
- What OTC medicines or herbal preparations do you use?
- Do you use antacids, laxatives, enemas, NSAIDs, or aspirin?
- What home remedies do you use?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam.
 Stethoscope with the ability to detect high- and low-pitched sounds
 Marker
 Ruler
 Reflex hammer or tongue blade

Position

Supine

Technique

Remember to scan your patient from head-to-toe looking for signs related to an abdominal assessment

Inspection, Auscultation, Percussion, Palpation

Inspect (From Several Different Angles)

Abdomen

- Size, shape, symmetry
- Movements (respiratory, pulsations, peristalsis)
- Condition of skin, lesions
- Hernias

Umbilicus

- Inverted/everted, position

Auscultate

Abdomen

- Bowel sounds (frequency)
- Vascular sounds: bruits, venous hums
- Friction rubs
- Scratch test

Percuss

- Abdomen
- Liver
- Spleen
- CVA tenderness

Light Palpation

- Abdomen for tenderness
- Surface characteristics
- Lymph nodes

Deep Palpation

- Abdomen for tenderness or masses
- Organs (liver, spleen, kidneys, aorta, bladder)

Additional Abdominal Assessment Techniques

- Testing for ascites
 - Shifting dullness
 - Fluid wave
 - Measuring abdominal girth and weight
- Abdominal reflexes
- Ballottement
- Kehr's sign
- Balance's sign
- Murphy's sign
- Signs for appendicitis
 - Rebound tenderness
 - Obturator test
 - Rovsing's sign
 - Cutaneous hypersensitivity
 - Iliopsoas sign

DOCUMENT

Document your findings and develop your plan of care.

S U M M A R Y


- A thorough abdominal assessment includes a detailed health history and physical examination that provides invaluable data about the patient's overall health status. Before you begin, realize that every system with the exception of the respiratory system is contained within the abdomen. As you work through the assessment, look for changes in every system.
- The comprehensive abdominal health history will direct the physical exam. The physical exam differs from other exams in that the assessment sequence

is inspection, auscultation, percussion, and then palpation.

- Cluster and analyze the data to identify actual and potential health problems. Using this information, make a prioritized list of nursing diagnoses and develop a plan of care with actions designed to restore and promote the patient's health. Share your findings with your patient. Remember: This is an ideal teaching opportunity. Involving the patient in her or his plan of care is more likely to promote positive, healthy outcomes.

R E V I E W Q U E S T I O N S

1. Johnny Fellin, age 9, is brought to the emergency room by his parents with the chief complaint of right lower quadrant pain. Which test is the best test for appendicitis?
 - a. Iliopsoas test
 - b. Shifting dullness
 - c. Kehr's sign
 - d. Murphy's sign
2. Johnny's appendix ruptures. What would you expect to hear when auscultating the abdomen?
 - a. Normal bowel sounds
 - b. Hypoactive bowel sounds
 - c. Hyperactive bowel sounds
 - d. Absent bowel sounds
3. When auscultating the abdomen for bowel sounds, how long should you listen before concluding that bowel sounds are absent?
 - a. 1 minute
 - b. 5 minutes
 - c. 10 minutes
 - d. 15 minutes
4. When assessing the abdomen, in what sequence should the assessment be performed?
 - a. Inspection, palpation, percussion, auscultation
 - b. Inspection, percussion, palpation, auscultation
 - c. Inspection, auscultation, percussion, palpation
 - d. Inspection, auscultation, palpation, percussion
5. You are assessing your patient's abdomen and are checking for hernias. What patient position is best for accentuating hernias?
 - a. Have patient relax abdomen.
 - b. Have patient flex legs.
 - c. Have patient raise head off bed.
 - d. Have patient breathe slowly.
6. You perform the scratch test on your patient. The scratch test is helpful in locating the:
 - a. Liver.
 - b. Spleen.
 - c. Kidneys.
 - d. Gallbladder.
7. You are percussing your patient's liver at the right midclavicular line. What is the normal liver span at the right midclavicular line?
 - a. 2 to 4 cm
 - b. 4 to 8 cm
 - c. 6 to 12 cm
 - d. 8 to 12 cm
8. Tom Johnson, age 56, is admitted to the hospital with the diagnosis of upper GI bleed. Considering his diagnosis, what color would you expect his stools to be?
 - a. Brown
 - b. Red
 - c. Green
 - d. Black
9. Sam Thomas, age 56, has liver disease. Which area would you most likely hear a venous hum?
 - a. RUQ
 - b. RLQ
 - c. LUQ
 - d. Left lower quadrant (LLQ)
10. You assess Mr. Thomas's abdomen for ascites. Which test is best for assessing abdominal fluid?
 - a. Shifting dullness
 - b. Scratch test
 - c. Iliopsoas test
 - d. Cullen's sign



Assessing the Female Genitourinary System

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent female genitourinary history questions
- Obtain a female genitourinary history
- Perform a physical assessment of the female genitourinary system
- Document female genitourinary findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

Assessment of the genitourinary (GU) system is essential to obtaining a picture of a woman's overall health status. The female reproductive system undergoes cyclical changes in response to hormonal levels of estrogen and progesterone throughout the life cycle. Reproductive health issues are a concern at every stage of reproductive development. Premenarchal girls may have questions about changes that occur during adolescence and menstruation, and adolescents may be concerned about sexuality and contraception. During the childbearing years, an adult woman's concerns focus on conception and pregnancy; later in life, they focus on the changes associated with menopause. Identifying a woman's reproductive stage will help you identify health concerns and differentiate your findings.

Focus your assessment on identifying any acute health problems, risks for GU problems, and health promotion teaching needs. Sexually transmitted diseases (STDs) are the most common communicable diseases in the United States. If left untreated, they can have devastating, permanent effects. GU dysfunction has not only a direct impact on the reproductive and urinary systems but also far-reaching effects on a woman's quality of life.

Anatomy and Physiology Review

Before you begin the assessment, you need to understand the anatomy and physiology of the female GU system.




FIGURE 18.1. External female genitalia.

The system consists of external and internal genitalia, which develop and function according to the hormonal influences that affect fertility and childbearing. It also consists of urinary structures.

External genitalia include the mons pubis, clitoris, vestibule, labia majora, labia minora, vaginal introitus, hymen, Bartholin's glands, Skene's glands, and the urethral meatus. Internal genitalia include the vagina, cervix, uterus, adjacent structures (adnexa), ovaries, and fallopian tubes. Internal urinary structures include the ureters, bladder, and urethra (Figs. 18.1 through 18.3).




FIGURE 18.2. Internal female genitalia and cross-section of rectum.




FIGURE 18.3. Internal structures of adnexa.

The functions of the female GU system are:

- Manufacturing and protecting ova for fertilization.
- Transporting the fertilized ovum for implantation and embryonic/fetal development.
- Housing and nourishing the developing fetus.
- Regulating hormonal production and secretion of several sex hormones.
- Providing sexual stimulation and pleasure.
- Providing a drainage route for the excretion of urine (urinary structures).

The Process of Reproduction

Mechanical Process

The process of reproduction requires that a mature ovum be fertilized by a sperm. The ovum is viable for about 24 hours and the sperm for up to 72 hours. Fertilization must occur within 24 hours of ovulation. During male ejaculation, sperm enters the vaginal canal and travels through the uterus to the fallopian tube, where fertilization usually occurs. A fertilized ovum is called a **zygote**. Once the ovum is fertilized, embryonic development begins. The fertilized ovum travels through the fallopian tube to the uterus for implantation. Once implantation has occurred, the uterus houses, nourishes, and protects the developing fetus until the time of delivery. Hormonal influences prepare the uterus monthly for implantation of a fertilized ovum. If fertilization does not occur, the endometrium sheds and the menstrual cycle begins.

Physiological Process

The hypothalamus, ovaries, and pituitary gland secrete hormones that affect the thickening and shedding of the uterine lining during the menstrual cycle. The ovarian cycle consists of two phases: the follicular phase and the luteal phase. During the **follicular phase**, the actions of the follicle-stimulating hormone (FSH) and the luteinizing hormone (LH) from the anterior pituitary stimulate the ripening of one ovarian follicle called the **graafian follicle**. The remaining follicles are suppressed by LH. Ovulation occurs when high levels of LH cause the release of the ovum from the graafian follicle. During the **luteal phase**, LH stimulates the development of the corpus luteum. The yellow pigment that fills the graafian follicle produces high levels of progesterone and low levels of estrogen. If pregnancy does not occur, the corpus luteum diminishes, hormone levels fall, and menstruation begins (Fig. 18.4).




FIGURE 18.4. Menstrual cycle.

The Urinary System

The kidneys filter metabolic waste products from the blood and excrete these wastes in excessive body water. Each kidney has a single ureter connecting the renal pelvis with the urinary bladder. Each ureter then narrows on entering the posterior wall of the urinary bladder. The bladder temporarily stores the urine. As the bladder fills with urine, neuroreceptors transmit the impulse to void. Urine then flows through the urinary meatus. Any alteration in the homeostasis or structure of the urinary tract can be life threatening.

Interaction With Other Body Systems

The Cardiovascular System

Nutrients and sex hormones are transported to the reproductive organs through the vascular system. Sexual excitement

causes vasodilation of the arterioles in the clitoris. In addition, estrogen lowers blood cholesterol levels and has positive cardiovascular effects until menopause.

The Urinary System

The close proximity of the vaginal opening to the urinary meatus increases the risk of urinary tract infections (UTIs) after frequent or vigorous sexual intercourse. Bladder continence is achieved during filling through the combination of detrusor muscle relaxation, internal sphincter muscle tone, and external sphincter contraction.

The Musculoskeletal System

The pelvic bones and pelvic floor muscles provide protection and support for some reproductive organs. Estrogen helps maintain bone mass until menopause.

Structures and Functions of the Female Genitourinary System

STRUCTURE	DESCRIPTION/PRIMARY FUNCTION
Mons Pubis	<ul style="list-style-type: none"> ■ Pad of subcutaneous fatty tissue lying over anterior symphysis pubis. ■ Protects pelvic bones during coitus.
Labia Majora	<ul style="list-style-type: none"> ■ Two longitudinal folds of adipose and connective tissue. Extend from clitoris anteriorly and gradually narrow to merge and form posterior commissure of perineum. ■ Outer surface of labia majora becomes pigmented, wrinkled, and hairy at puberty. ■ Inner surface is smoother, softer, and contains sebaceous glands. ■ Protects vulva components that it surrounds. ■ Protects urethra and vagina from infection.
Labia Minora	<ul style="list-style-type: none"> ■ Consists of two thin folds of skin that extend to form prepuce of clitoris anteriorly and a transverse fold of skin forming fourchette posteriorly. ■ Labia minora contain sebaceous glands, erectile tissue, blood vessels, and involuntary muscle tissue. ■ Secretions are bactericidal and aid in lubricating vulval skin and protecting it from urine. ■ Protects urethra and vagina from infection.
Clitoris	<ul style="list-style-type: none"> ■ Erectile body about 2.5 cm in length and 0.5 cm in diameter. ■ Contains erectile tissue and has significant supply of nerve endings. ■ Serves as primary organ of sexual stimulation.
Vestibule	<ul style="list-style-type: none"> ■ Area between two folds of labia minora. ■ Boat-shaped area containing the urethral meatus, openings of Skene's glands, hymen, openings of the Bartholin's glands and vaginal introitus.
Skene's Glands (Paraurethral Glands)	<ul style="list-style-type: none"> ■ Surround urethral meatus. Provide lubrication to protect skin.
Vaginal Introitus	<ul style="list-style-type: none"> ■ Entrance to vagina. Size and shape may vary.
Hymen	<ul style="list-style-type: none"> ■ Avascular thin fold of connective tissue surrounding vaginal introitus in women who have not had sexual intercourse.
Bartholin's Glands	<ul style="list-style-type: none"> ■ Small, pea-shaped glands deep in perineal structures. ■ Ducts are not visible. ■ Secrete clear, viscid, odorless, alkaline mucus that improves viability and motility of sperm along reproductive tract.
Perineum	<ul style="list-style-type: none"> ■ Space between fourchette and anus. ■ Composed of muscle, elastic fibers, fascia, and connective tissue.

STRUCTURE	DESCRIPTION/PRIMARY FUNCTION
Vagina	<ul style="list-style-type: none"> ■ Muscular tube from cervix to vulva. ■ Located posterior to bladder and anterior to rectum. ■ Serves as female organ of copulation, birth canal, and channel through which menstrual flow exits.
Cervix	<ul style="list-style-type: none"> ■ End of uterus that projects into vagina.
Uterus	<ul style="list-style-type: none"> ■ Pear-shaped, hollow, muscular organ between bladder neck and rectal wall. ■ Mucous membrane lining is the endometrium. Muscular layer is the mesometrium. Inferior aspect is cervix, and superior aspect is fundus. ■ Major functions include serving as implantation site of fertilized ovum and as protective sac for developing embryo and fetus.
Fallopian Tubes	<ul style="list-style-type: none"> ■ Two 7- to 10-cm-long ducts on either side of fundus of uterus. ■ Extend from uterus almost to ovaries. ■ Normally, fertilization takes place within tubes. ■ Major function includes serving as fertilization site and providing passageway for unfertilized and fertilized ova to travel to uterus.
Ovaries	<ul style="list-style-type: none"> ■ Almond-shaped glandular structures that produce ova. ■ Located beside fallopian tubes. ■ Major functions include producing ova for fertilization by sperm and producing estrogen and progesterone.
Ureters	<ul style="list-style-type: none"> ■ Hollow, tubelike structures connecting renal pelvis with urinary bladder.
Urinary Bladder	<ul style="list-style-type: none"> ■ Muscular sac behind pubic symphysis that stores urine.
Urethra	<ul style="list-style-type: none"> ■ Narrow, tubelike structure lined with mucous membrane. ■ Opens to allow flow of urine to be excreted. ■ It is 1 to 1½ inches long and located slightly posterior to clitoris and directly anterior to vagina and rectum.

The Neurological System

Neurological system impulses regulate events of the female sexual response. Hormones released by the hypothalamus of the diencephalon trigger the onset of the ovarian cycle.

Micturition is a reflex of parasympathetic control that stimulates detrusor muscle contraction with the simultaneous relaxation of the external sphincter and muscles of the pelvic floor. Urinary continence depends on the interaction of nerves that control the muscles for the bladder, bladder neck, urethra, and pelvic floor.

Developmental, Cultural, and Ethnic Variations

Infants

The female infant's genitals are enlarged at birth in response to maternal estrogen. The labia majora should cover the labia minora. The hormonal effect may also cause a pseudomenstruation. The urinary meatus and vaginal orifice should be visible. Androgens and progestins may cause masculine characteristics in female infants. If genitalia are ambiguous, a history is obtained to


determine maternal use of sex hormones during pregnancy or a family history of adrenogenital syndrome.

Children and Adolescents

Girls begin puberty changes sometime between 8 and 13 years of age. Release of estrogen initiates the changes, which are first demonstrated in the development of breast buds and growth of pubic hair, followed by menstruation. The early development of sexual characteristics allows pregnancy before the child is intellectually or emotionally prepared for it. Early maturation can also lead to anemia related to menstrual bleeding. It is essential to assess for sexual molestation. Some signs include genital trauma; depression; eating disorders; and bruising, swelling, and inflammation in the vaginal, perineal, and anal areas. (See *Maturation States in Females*.)

Pregnant Women

Pregnancy imposes a multitude of changes to the reproductive organs. The uterus, cervix, ovaries, and vagina undergo significant structural changes related to the pregnancy and the influence of hormones. The uterus



Relationship of Female GU System to Other Systems

Maturation States in Females

Stage 1



- *Preadolescent*: No pubic hair except for fine body hair similar to hair on abdomen.

Stage 2



- Sparse growth of long, slightly pigmented, downy hair, straight or only slightly curled, mostly along labia.

Stage 3



- Hair becomes darker, coarser, and curlier and spreads sparsely over pubic symphysis.

Stage 4



- Pubic hair is coarse and curly as in adults. It covers more area than in stage 3 but not as much as in adults.

Stage 5



- Quality and quantity are consistent with adult pubic hair distribution and spread over medial surfaces of thighs but not over abdomen.

becomes hypertrophied, and its capacity increases to 500 to 1000 times its nonpregnant state. The vascularity of the cervix increases and contributes to softening of the cervix. Vascular congestion creates a blue-purple blemish or change in the cervical color (Chadwick's sign). Estrogen causes the glandular cervical tissue to produce a thick mucus, which builds up and forms a mucus plug at the endocervical canal.

The vagina undergoes changes similar to those of the uterus. Hypertrophy of the vaginal epithelium occurs. The vaginal wall softens and relaxes to accommodate the movement of the infant during birth.

Older Adults

Reproductive ability in women usually peaks in the late 20s. Over time, estrogen levels begin to decline and, in response, between the ages of 46 and 55, menstrual periods become shorter and less frequent until they stop entirely. Menopause is said to have occurred when the woman has not experienced a menstrual period in over a year. Sexual organs atrophy, the clitoris becomes smaller, and vaginal secretions are not as plentiful, so intercourse may become painful. These vaginal changes increase the risk for vaginal infections. (See *Cultural and Ethnic Views of Women*.)

Women of Different Cultures or Ethnic Groups

Cultural and ethnic variations surrounding women often center on reproductive ability. (See *Cultural and Ethnic Views of Women*.)

Introducing the Case Study



Mrs. Ellen James has come for her annual gynecological examination. She is 29 years old, married, and the mother of one child. She complains of scant midcycle vaginal bleeding for the last 2 months. She is not employed outside the home.

Performing the Female Genitourinary Assessment

A comprehensive GU assessment includes taking the patient's health history and performing a physical examination of the external and internal genitalia.

Health History

Because assessment of the female GU system may be somewhat uncomfortable for both you and your patient, begin by establishing rapport. A professional, nonthreatening approach is essential in order to collect accurate data.

Cultural and Ethnic Views of Women

CULTURAL GROUP	VIEW OF WOMEN
Amish	<ul style="list-style-type: none"> High status associated to role of wife and mother.
Appalachian	<ul style="list-style-type: none"> Responsibilities include feeding, clothing, and caring for family. High status associated with motherhood. Having children associated with fulfillment. Responsible for child rearing. Older women preserve culture. Responsible for preparing herbal medicines and folk medicine.
Arab American	<ul style="list-style-type: none"> Women gain status with age. Responsible for caring for and educating children and tending to husband's needs.
Chinese American	<ul style="list-style-type: none"> Woman's role is to perpetuate male dominance.
Cuban American	<ul style="list-style-type: none"> Women expected to stay at home and care for children.
Egyptian American	<ul style="list-style-type: none"> Status and power increase with pregnancy and birthing, especially with having a son. Childbirth is expected within 1 year of marriage.
Filipino American	<ul style="list-style-type: none"> Women have role equal to that of men in health, welfare, and family finance.
Greek American	<ul style="list-style-type: none"> Pregnancy seen as a time of great respect.
Iranian American	<ul style="list-style-type: none"> Prestige associated with having children.
Jewish American	<ul style="list-style-type: none"> Woman runs home and is responsible for children.
Mexican American	<ul style="list-style-type: none"> Woman maintains home and health of family.
Native American	<ul style="list-style-type: none"> Mother is center of Native American society.
Vietnamese American	<ul style="list-style-type: none"> Women expected to be dutiful and respectful of husband and to make healthcare decisions.

Source: Purnell, L., and Paulanka, B. (2003). *Transcultural Nursing*, ed. 2. Philadelphia: F.A. Davis.

Accurate data are the keys to a successful health history. When interviewing your patient, keep these points in mind:

- Some cultures do not permit women to discuss sexual topics or be examined by a man.
- Reproductive system problems may adversely affect self-image, sexual functioning, and overall wellness.

- Consider the woman's age and choose your health history questions accordingly.

The history will provide you with data that will guide your physical examination. If time is an issue and you are unable to perform a complete health history, perform a focused history on the female GU system.

Focused Genitourinary Health History

If you are unable to perform a detailed history, ask key questions that focus on a history of GU problems, presence of symptoms, identification of risk factors, and health promotion activities. The following questions focus on the warning signs of GU problems:

- If premenopausal, when was your last menstrual period?
- Do you have any pain? Location?
- Can you describe your menstrual cycle? What was your age of menarche? What was age at menopause? Do you have any abnormal vaginal bleeding?
- What are your sexual practices? How many partners have you had? What type of sexual activity: oral, genital, anal? What is your sexual preference: heterosexual, homosexual, or bisexual? Do you practice safe sex? What do you use? Do you use contraceptives? If yes, type?
- What is your obstetric history? Have you ever been pregnant? Number of live deliveries? Number of miscarriages? Abortions? Age of first full-term pregnancy?
- Do you have any vaginal discharge? If so, can you describe it?
- Do you have a history of GU surgery, STDs, or cancer?
- Are you currently taking any medications? Hormone replacement therapy (HRT)? Birth control pills (BCP)?
- Do you have any urinary symptoms, such as burning, frequency, or dysuria?

Biographical Data

Quickly reviewing the biographical data may identify actual or potential problems. Begin with the woman's age—it influences the questions you ask. The woman's age may also help you identify risk factors such as STDs, which occur more frequently in young women, and gynecological cancers, which occur more often in older adults. If your patient is over age 50, you may want to ask questions related to menopause, rather than asking about the menstrual cycle.

Current Health Status

Reproductive health history questions relate to sexual function, fertility, sexual satisfaction, sexual practices, and self-care. Begin by asking about symptoms; leave sexuality questions to the end of the history. If the woman has a reproductive system problem, identify her chief complaint and perform a symptom analysis. Symptoms that relate to the female GU system include vaginal discharge, pain, lumps/masses, dysmenorrhea, amenorrhea, and urinary symptoms.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Vaginal Discharge

The only normal vaginal drainages are mucus and menstrual blood. Any other type of discharge may signal a problem. The description of the discharge may give clues to the underlying problem. Vaginal discharge is the most common presenting symptom in women with STDs. A white, curdlike discharge is seen with candidiasis; a gray or white, thin, malodorous discharge with bacterial vaginosis; and a yellow or green discharge with gonorrhea.

Lesions

Lesions may indicate a variety of problems. Vesicles, ulcers, warts, or rashes are common lesions. The morphological description of the lesion and whether or not it is painful will help you identify it. Lesions are often related to STDs, but they may also be caused by carcinoma or a systemic problem.

Vaginal Bleeding

Abnormal uterine bleeding includes **menorrhagia** (excessive menstrual flow), **breakthrough bleeding** (vaginal bleeding between periods), or **amenorrhea** (decreased or absent menstrual bleeding). Determine your patient's usual menstrual cycle and flow. Changes may be associated with normal reproductive changes or signal an underlying problem, so careful assessment is needed to determine the cause.

Pain

Pain directly related to menstrual flow is called **dysmenorrhea**. It is the most frequent complaint of young women seeking healthcare. **Primary dysmenorrhea** is menstrual pain or discomfort not usually associated with a physical abnormality or pathology and can be diagnosed with a normal physical examination. **Secondary dysmenorrhea** is menstrual pain or discomfort associated with pathology, such as endometriosis, infection, intrauterine device (IUD) placement, pelvic mass, adhesions, cervical obstruction, or congenital malformations. Other causes of pelvic pain include pelvic inflammatory disease (PID) and renal and gastrointestinal (GI) problems.

Pain in the lower abdomen or back can also be renal, GI, or musculoskeletal in nature. Therefore, it is important to perform a thorough symptom analysis. Perform a review of systems (ROS) to identify any relevant data.

Amenorrhea

Amenorrhea is the absence of menses. **Primary amenorrhea** means that menses never occurred. In **secondary amenorrhea**, menses have ceased for 6 months, if the cycle was normally regular, and for 12 months, if the cycle was normally irregular. Primary amenorrhea may be caused by a genetic or hormonal disorder. Secondary amenorrhea may be caused by a normal process such as pregnancy or menopause or may indicate an underlying problem. A thorough symptom analysis will help you to determine the underlying cause.

Urinary Symptoms

Urinary symptoms are often associated with female reproductive problems caused by the anatomical proximity of the systems. Urinary symptoms often cause discomfort, prompting the patient to seek treatment. Dysuria, burning, frequency, change in urinary patterns, and hematuria are common problems. Because the two systems work so closely together, a problem in one often eventually affects the other. A urinary problem may cause a reproductive problem, but more frequently, a reproductive problem results in a urinary problem. Urinary incontinence is a common complaint of older women, but it should not be considered a normal change associated with aging. (For more information, see Chapter 27, Assessing the Older Adult.)

Past Health History

The various components of the past history are linked to female genitalia pathology and female genitalia-related information. You may be able to identify a direct link from the past health history to your patient's current health status.

Past Health History

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Childhood Illnesses

- Did you have communicable diseases transmitted at birth?
- Were you exposed to DES before birth?

- Diseases passed from mother to infant during birth include hepatitis B, herpes, human immunodeficiency virus (HIV), chlamydia, and HPV.
- Exposure to DES in utero increases risk of cervicovaginal cancer and reproductive problems.

Hospitalizations/Surgeries

- Have you ever had gynecological surgery: hysterectomy, myomectomy, salpingectomy, oophorectomy, dilation and curettage (D & C), dilatation and evacuation, laparoscopy, vulvectomy, tubal ligation, colpotomy, cesarean section, female circumcision?

- Surgical procedures affect reproductive health.

Diagnostic Procedures

- When was your last Pap test?
- Have any biopsies been done?

- Pap test screens for cervical cancer.
- Biopsies detect malignancies.

Serious Injuries

- Have you ever had abdominal or vaginal trauma or injuries?
- Have you ever been raped or sexually abused?

- May explain abnormal physical findings or sexual dysfunction.

Serious/Chronic Illnesses

- Do you have an STD, such as gonorrhea, syphilis, herpes, HIV/AIDS, hepatitis, chlamydia, condyloma (HPV)?
- Do you have any infertility problems?

- STDs can cause infertility.
- Recurrent incidence of STDs may indicate untreated sexual partner or multiple sexual partners.
- Existing infertility problems may help explain physical findings.

- Do you have diabetes mellitus, thyroid disease, incontinence, constipation, UTIs?

- Inability to conceive after 1 year of regular coitus without contraception may indicate fertility problem and warrants referral.

- Diabetes mellitus increases incidence of vaginal yeast infections. There are also specific precautions for pregnant diabetics. (See Chapter 23, Assessing the Mother-to-Be.)

- A history of thyroid disease may explain menstrual irregularities.

- Recurrent UTIs may be a symptom of untreated vaginal infection.

- Voiding pattern change may be a symptom of infection or reproductive structure changes.

- Are you a paraplegic or quadriplegic?
- Do you have a chronic neurological problem?

- Disabled or handicapped patients are at increased obstetric risk depending on level of injury, tone of uterus, and competency of cervix. They are also at risk for urinary complications such as UTIs.

Immunizations

- Were you immunized against hepatitis B?
- Were you immunized against HPV?

- Hepatitis B can be transmitted through sexual activity and to newborns from infected mothers.
- Gardasil vaccine protects against four types of HPV, which cause 70% of cervical cancers and 90% of genital warts.

Allergies

- Do you have any allergies?

- Contact dermatitis caused by condoms, nonoxynol 9 or other spermicides, or latex identifies additional health promotion needs.

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Medications</p> <ul style="list-style-type: none"> Are you taking any prescribed or over-the-counter medications, including antibiotics, oral contraceptives, estrogen replacement therapy, antidepressants, antihypertensives? 	<ul style="list-style-type: none"> Antibiotics increase incidence of <i>Candida</i> vaginosis and may lessen effectiveness of oral contraceptives. Estrogens may cause altered menstrual flow. Antidepressants and antihypertensive medications can alter libido. Antidepressants may also cause menstrual irregularities. (See <i>Drugs That Adversely Affect the Female Reproductive System</i>.)
<p>Recent Travel/Military Service/Exposure to Infectious or Contagious Diseases</p> <ul style="list-style-type: none"> Have you been exposed to chemicals? Do you have a history of STDs? 	<ul style="list-style-type: none"> Increased risk for infertility. Could present an educational opportunity for disease prevention.

Drugs That Adversely Affect the Female Reproductive System

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Androgens	Danazol	Vaginitis with itching, dryness, burning, or bleeding; amenorrhea
	Fluoxymesterone, methyltestosterone, testosterone	Amenorrhea and other menstrual irregularities; virilization, including clitoral enlargement
Antidepressants	Tricyclic antidepressants Selective serotonin reuptake inhibitors	Changed libido, menstrual irregularity Decreased libido, anorgasmia
Antihypertensives	Clonidine, reserpine Methyldopa	Decreased libido Decreased libido, amenorrhea
Antipsychotics	Chlorpromazine, perphenazine, prochlorperazine, thioridazine, trifluoperazine, haloperidol	Inhibition of ovulation (chlorpromazine only), menstrual irregularities, amenorrhea, changed libido
Beta-blockers	Atenolol, labetalol hydrochloride, nadolol, propranolol hydrochloride, metoprolol	Decreased libido
Cardiac glycosides	Digoxin	Changes in cellular layer of vaginal walls in postmenopausal women
Corticosteroids	Dexamethasone, hydrocortisone, prednisone	Amenorrhea and menstrual irregularities
Cytotoxics	Busulfan	Amenorrhea with menopausal symptoms in premenopausal women, ovarian suppression, ovarian fibrosis and atrophy
	Chlorambucil	Amenorrhea
	Cyclophosphamide	Gonadal suppression (possibly irreversible), amenorrhea, ovarian fibrosis
	Methotrexate	Menstrual dysfunction, infertility
	Tamoxifen	Vaginal discharge or bleeding, menstrual irregularities, pruritus vulvae (intense itching of the female external genitalia)
	Thiotepa	Amenorrhea

(continued)

Drugs That Adversely Affect the Female Reproductive System (continued)

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Estrogens	Conjugated estrogens, esterified estrogens, estradiol, estrone, ethinyl estradiol	Altered menstrual flow, dysmenorrhea, amenorrhea, cervical erosion or abnormal secretions, enlargement of uterine fibromas, vaginal candidiasis
	Dienestrol	Vaginal discharge, uterine bleeding with excessive use
Thyroid hormones	Levothyroxine sodium, thyroid USP, and others	Menstrual irregularities with excessive doses
Progestins	Medroxyprogesterone acetate, norethindrone, norgestrel, progesterone	Breakthrough bleeding, dysmenorrhea, amenorrhea, cervical erosion, abnormal secretions
Miscellaneous	Lithium carbonate	Decreased libido
	L-tryptophan	Decreased libido
	Spironolactone	Menstrual irregularities, amenorrhea, postmenopausal bleeding
	Valproic acid	Menstrual irregularities, amenorrhea, possible polycystic ovarian syndrome

Family History

The family history can identify predisposing or causative factors of a health problem. It identifies not only genetic

ally linked disorders but also familial risk factors that may predispose your patient to GU disease.

Family History

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Mother's Reproductive History <ul style="list-style-type: none"> ■ Did your mother take DES while pregnant with you? ■ Did she transfer STDs during pregnancy or delivery, have multiple pregnancies, or have babies with congenital anomalies? ■ Do you have a family history of reproductive cancers? 	<ul style="list-style-type: none"> ■ Exposure to DES in utero can cause cervicovaginal cancer and reproductive problems. ■ STDs can cause infertility. ■ Spontaneous abortion, menstrual irregularities, multiple births, congenital anomalies, and diseases of female genitalia may have familial tendency. ■ Gynecological cancers such as ovarian and uterine have familial/genetic links.

Review of Systems

Changes in reproductive system function have an effect on other body systems. The ROS can identify changes in

other systems and help you identify something you might otherwise have missed.

Review of Systems

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
General Health Survey <ul style="list-style-type: none"> ■ How have you been feeling? ■ Have you had any changes in your energy level? 	<ul style="list-style-type: none"> ■ Fatigue or activity intolerance is often a sign of underlying health problem. For example, fatigue from

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Have you had a fever? ■ Have you experienced any weight changes? 	<p>HIV/AIDS would interfere with activity tolerance. Fatigue influences desire and ability to perform sexual intercourse and may also be associated with anemia.</p>
<p>Integumentary <i>Skin/Hair/Nails</i></p>	<ul style="list-style-type: none"> ■ Rashes, growths, and lumps may indicate pathology. ■ Pruritus in hair or on labia may indicate scabies, kidney or liver problems. ■ Estrogen and progesterone can cause changes in skin, hair, and nails, such as those seen during pregnancy and menopause. ■ Changes in hair texture, moisture, and distribution occur during pregnancy and menopause.
<ul style="list-style-type: none"> ■ Have you had any rashes, lesions, or skin color changes? ■ Any changes in hair growth? 	
<p>HEENT <i>Head and Neck</i></p>	<ul style="list-style-type: none"> ■ Swelling of lymph nodes associated with infection, malignancy. ■ Enlarged thyroid may affect sexual function. ■ Headaches are a common side effect of oral contraceptives.
<ul style="list-style-type: none"> ■ Do you have any swollen glands or nodes? 	
<ul style="list-style-type: none"> ■ Do you have headaches? 	
<p><i>Eyes, Mouth, and Throat</i></p>	<ul style="list-style-type: none"> ■ Oropharynx and eyes are sites for STDs, such as herpes simplex virus 2 (HSV-2) and gonorrhea. ■ Secondary syphilis can result in oral mucous patches. ■ Gonorrheal eye infections occur in newborns (ophthalmia neonatorum) and gonorrheal conjunctivitis in adults. ■ <i>Tearing, photophobia, eyelid edema, conjunctival edema: Chlamydia trachomatis.</i>
<ul style="list-style-type: none"> ■ Do you have a sore throat? 	
<ul style="list-style-type: none"> ■ Watery eyes? Eye drainage? 	
<p>Respiratory</p>	<ul style="list-style-type: none"> ■ Respiratory disease such as chronic obstructive pulmonary disease (COPD) may affect sexual activity.
<ul style="list-style-type: none"> ■ Do you have any breathing problems? 	
<p>Cardiovascular</p>	<ul style="list-style-type: none"> ■ Some heart murmurs are associated with anemia. ■ Cardiovascular symptoms are associated with oral contraceptive use. ■ Chronic cardiovascular disease can affect ability to perform sexually.
<ul style="list-style-type: none"> ■ Do you have a heart murmur? 	
<ul style="list-style-type: none"> ■ Did you ever have vascular problems, thrombophlebitis, cardiovascular disease? 	
<p>Breasts</p>	<ul style="list-style-type: none"> ■ Cyclic hormonal changes are associated with breast tenderness. ■ Female hormones are associated with fibrocystic disease and breast cancer.
<ul style="list-style-type: none"> ■ Do you have any breast tenderness, lumps, or discharge? 	
<ul style="list-style-type: none"> ■ Where are you in your menstrual cycle? 	
<p>Gastrointestinal</p>	<ul style="list-style-type: none"> ■ Liver disease associated with gynecologic malignancy may cause ascites. ■ May indicate underlying eating disorder, malignancy, or AIDS. ■ May indicate pathology of reproductive organs.
<ul style="list-style-type: none"> ■ Do you have a history of liver disease? 	
<ul style="list-style-type: none"> ■ Do you have appetite or weight loss? 	

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Abdominal pain? Reproductive/Menstrual History ■ How old were you when you started to menstruate? ■ When was your LMP? ■ How long are your periods? How long is your cycle? ■ Are your periods regular? ■ Is your flow heavy, moderate, or light? ■ Are your periods painful or extremely heavy? ■ Do you skip periods or have you stopped getting them altogether? ■ Do you have spotting between periods? Premenstrual Syndrome ■ Before your period, do you have breast tenderness, bloating, moodiness; cravings for salt, sugar, or chocolate; fatigue; weight gain; headaches; or joint pain? Obstetric History ■ When was your LMP? ■ Have you ever used fertility drugs? ■ How many pregnancies have you had? ■ How many living children? ■ Have you ever had an abortion, miscarriage, C-section, or other complication with pregnancy? ■ How long were your labors? ■ Did you have any postpartum complications? Contraceptive History ■ Do you use contraceptives? If yes, what types, how often, and have you had any problems with contraception? ■ How do you prevent STDs? ■ Do you or did you smoke? If yes, how much and for how long? Perimenopause ■ Do you have spotting between periods, hot flashes, heart palpitations, numbness, tingling, drenching sweats, mood swings, vaginal dryness, or itching? ■ Are you taking estrogen replacement therapy? ■ How do you feel about menopause? Musculoskeletal ■ Do you have any weakness, limitation, joint pain, swelling? ■ Are you past menopause? ■ Have you ever had a bone density test? ■ Did you ever break a bone? Neurological ■ Have you had any changes in your menstrual cycle? ■ Have you had mood swings? ■ Have you had weakness, paralysis, tremors? Mental or personality changes? 	<ul style="list-style-type: none"> ■ <i>Late onset of menarche (age 16–18)</i>: Can result from inadequate nutrition caused by eating disorders, chronic diseases such as Crohn's disease, environmental stresses, intensive athletic training, hypothyroidism, or opiate or steroid use. ■ Premenstrual syndrome (PMS) signs and symptoms may be alleviated through stress reduction and avoidance of certain foods. ■ Past obstetric health is a predictor of future reproductive health. ■ IUDs increase risk of PID; diaphragms may cause urinary discomfort; oral contraceptives have a variety of side effects; and some people are allergic to spermicides or latex in condoms. ■ Oral contraceptives and smoking increase risk of cardiovascular problems. ■ Take this opportunity to teach woman about various contraception methods. ■ Women may have various discomforting signs and symptoms associated with menopause. ■ Hormonal and physical changes may affect self-concept. ■ Identifying symptoms helps you plan appropriate interventions. ■ The physical act of sexual intercourse requires the musculoskeletal system to be functional. ■ Postmenopausal women are at greater risk for developing osteoporosis. ■ Amenorrhea may result from problems with hypothalamus of diencephalon. ■ Hormonal fluctuations may affect emotional state (e.g., during pregnancy). ■ Neurosyphilis occurs in late syphilis and can cause paresis, tremors, personality changes, and psychosis.

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Endocrine</p> <ul style="list-style-type: none"> ■ Do you have diabetes? ■ Do you have thyroid disease? <p>Lymphatic/Hematologic</p> <ul style="list-style-type: none"> ■ Do you have HIV or AIDS? ■ Do you have a history of cancer? ■ Do you have a heavy menstrual flow? 	<ul style="list-style-type: none"> ■ Diabetes increases risk for candidal vaginal infections. ■ Hypothyroidism causes menorrhagia, decreased libido, and infertility. ■ Hyperthyroidism can cause abnormal menses or amenorrhea. ■ AIDS affects the immune system. ■ Gynecological cancers can metastasize to local lymphatic tissue. ■ Heavy menses (menorrhagia) can lead to anemia.

Psychosocial Profile

The psychosocial profile can provide a bridge between

the health maintenance activities and female genital function.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Health Practices and Beliefs/Self-Care Activities</p> <ul style="list-style-type: none"> ■ Do you get a yearly gynecological exam? ■ When was your last Pap test? What were the results? ■ Have you ever had a dexascan? Mammogram? If yes, when and what were the results? ■ What is your personal hygiene routine? ■ Do you douche frequently? <p>Typical Day</p> <ul style="list-style-type: none"> ■ What is your typical day like? <p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ What did you eat in the last 24 hours? ■ What foods do you usually eat? ■ Have you lost or gained weight recently? <p>Activity and Exercise Patterns</p> <ul style="list-style-type: none"> ■ Do you exercise on a routine basis? ■ If so, how often and what type of exercise? <p>Recreation/Pets/Hobbies</p> <ul style="list-style-type: none"> ■ Do you have any hobbies? ■ What do you do for recreation? 	<ul style="list-style-type: none"> ■ Annual exams help in early detection of problems. ■ Pap smear can detect precancerous and cancerous cell changes in cervix and may also detect HPV. ■ Regular Pap tests indicate patient's participation in self-care. ■ Vaginal smears are done for women who have had a hysterectomy. ■ Postmenopausal women are at greater risk for developing osteoporosis. ■ The risk of breast cancer increases with age and use of HRT. ■ Poor personal hygiene can lead to infections of genitalia or urinary tract. ■ Excessive cleansing or frequent douching can promote rashes or genital infection. ■ Ability to care for self indicates a high level of wellness. ■ May identify health teaching needs. ■ Increased levels of refined sugars, salt, and caffeine enhance PMS symptoms. ■ Extreme dieting can lead to amenorrhea. ■ Exercise may diminish dysmenorrhea and menorrhagia. ■ Excessive exercise may result in amenorrhea. ■ Extended periods of exposure to moisture increases likelihood of <i>Candida</i> vaginosis.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Do you swim or ride horses? ■ Do you have cats? 	<ul style="list-style-type: none"> ■ High incidence of genital trauma from saddle injuries. ■ Pregnant women can develop toxoplasmosis from touching cat litter.
<p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ How many hours of sleep do you average a night? ■ Is your sleep restful or interrupted? 	<ul style="list-style-type: none"> ■ Lack of sleep or extreme fatigue can lead to amenorrhea.
<p>Personal Habits</p> <p><i>Alcohol Use</i></p> <ul style="list-style-type: none"> ■ Do you drink alcohol? If so, what type, how much, and how often? 	<ul style="list-style-type: none"> ■ Significant correlation between alcohol use and date rape in college-age women. ■ Alcohol use during pregnancy can cause fetal alcohol syndrome.
<p><i>Drug Use</i></p> <ul style="list-style-type: none"> ■ Do you use street drugs, such as cocaine, barbiturates, amphetamines, or narcotics? If so, how often and how much? 	<ul style="list-style-type: none"> ■ Impairs judgment, increasing risk of STD exposure. Can cause decreased libido.
<p><i>Tobacco Use</i></p> <ul style="list-style-type: none"> ■ Do you smoke? If so, what type, how much per day, and for how many years? 	<ul style="list-style-type: none"> ■ Smoking is a risk factor for cervical cancer. ■ Increased incidence of strokes in smokers who use oral contraceptives.
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ What type of work do you do? ■ Are you exposed to arsenic, glycol ethers, lead, radiation, vinyl chloride, or PCBs? 	<ul style="list-style-type: none"> ■ Exposure to some chemicals, lead, and radiation has been linked to birth defects and spontaneous abortions. ■ Oncology nurses exposed to antineoplastic drugs may have spontaneous abortions, fetal anomalies, and changes in menstrual cycles.
<p>Environmental Health Patterns</p> <ul style="list-style-type: none"> ■ Do you live in an apartment or single home? Is sanitation adequate? ■ Is space adequate, or are you overcrowded? ■ Are you homeless? 	<ul style="list-style-type: none"> ■ Poor sanitation may lead to numerous forms of vaginitis. ■ Overcrowding can cause mite infestation. ■ Hygiene is a significant problem for homeless people.
<p>Roles/Relationships/Self-Concept</p> <ul style="list-style-type: none"> ■ How do you see yourself? 	<ul style="list-style-type: none"> ■ Poor self-image and lack of self-confidence predispose a patient to sexual dysfunction.
<p>Cultural/Religious Influences</p> <ul style="list-style-type: none"> ■ What is your cultural and religious background? 	<ul style="list-style-type: none"> ■ Cultural and religious practices influence GU health. For example, abortion and contraception are prohibited in Catholic faith, and some cultures/religions have specific beliefs related to menstruation and sexual practices.
<p>Family Roles and Relationships</p> <ul style="list-style-type: none"> ■ What is your present relationship status? 	<ul style="list-style-type: none"> ■ Grief from loss of a relationship can affect sexual desire. ■ Abusive relationships can affect sexual functioning.
<p>Social Supports/Stress and Coping</p> <ul style="list-style-type: none"> ■ Who are your supports? ■ Do you have much stress in your life? If so, how do you deal with it? 	<ul style="list-style-type: none"> ■ Stress can cause amenorrhea, exacerbate genital herpes simplex, and lead to sexual dysfunction. ■ Feelings of emotional or social isolation can increase stress.

Sexual History

Because the GU system encompasses sexuality, a more detailed sexual history may be indicated. Leaving this until the end of your history gives you time to establish rapport with the patient before asking sensitive or embarrassing questions.

Asking about sexual functions and practices is important for several reasons:

- Many women have sex-related questions or problems that they want to discuss.
- Sexual practices may be directly related to the patient's symptoms.
- Sexual dysfunction may be related to medications and may be reversible.
- Sexual practices are related to the patient's risk for cross-contamination to other areas, STDs, hepatitis, cervical dysplasia, and cervical carcinoma. Risk for disease increases as the number of sexual partners increases.
- Changes in sexual function may indicate pain, infection, hormonal changes, disease, changes in mental status, or altered role and relationship patterns.

Depending on the individual patient, ask the following questions to assess sexual function:

- Have you ever been sexually active?
- Are you currently sexually active? That is, have you had sex with anyone in the past few months? (If the answer is yes, ask the next question.)
- Do you have sex with men, women, or both (heterosexual, homosexual, or bisexual)?
- What types of sexual activity do you engage in? Oral, anal, genital?
- How often do you have intercourse?
- Do you have more than one partner? How many partners have you had in the last 6 months? How many partners in your lifetime? Do you trade sexual favors for drugs or money?
- Do you use birth control? What kind? How often?
- Are you worried about the acquired immunodeficiency syndrome (AIDS) virus or other STDs?
- Do you take any precautions to avoid infections? If so, what?
- Are you able to achieve orgasm? Do you have any problems or concerns about your sexual function?
- Have you had surgery on any of your reproductive organs? If so, what kind and when?



Case Study Findings

Reviewing Mrs. James's

Biographical data:

- 29-year-old African-American female.
- Married, mother of a 4-year-old girl.

- American by birth; Protestant religion.
- Attended 1 year of college after high school; homemaker at present; husband is stockbroker for investment banking company.
- Healthcare insurance with husband's company.
- Source of referral: Self, seems reliable.

Current health status:

- Scant midcycle vaginal bleeding for the last 2 months with no other complaints at present.
- Seeking care for annual Pap smear, gynecological examination, and renewed prescription for oral contraceptives.

Past health history:

- Rheumatic fever at age 7.
- Usual childhood illnesses except for measles, mumps, and rubella.
- Denies any previous surgeries or serious injuries.
- Has taken oral contraceptives since birth of child 4 years ago. No other medications except occasional aspirin.
- No history of STDs.
- Gravida 1, para 1, no complications with pregnancy or vaginal delivery.
- Menarche at age 12, last menstrual period (LMP) 21 days ago. Menses last 5 days with moderate to light flow and occurs every 21 to 23 days. Denies dysmenorrhea.
- Denies exposure to diethylstilbestrol (DES).

Family history:

- No gynecological carcinomas or maternal exposure to DES.
- Parents are alive and well.
- Has two healthy female siblings, ages 33 and 35.

Review of systems:

- *General Health Survey*: Usual state of health good, no changes in weight or energy level.
- *Integumentary*: No changes in skin, hair, or nails.
- *Head, Eyes, Ears, Nose, and Throat (HEENT)*: No sore throat, headaches, lumps, or masses in neck.
- *Respiratory*: No history of respiratory disease.
- *Cardiovascular*: No history of cardiovascular disease.
- *Gastrointestinal*: Appetite good, no GI problems.
- *Musculoskeletal*: No weakness, joint pain, or fractures.
- *Neurological*: No history of neurological problems.
- *Endocrine*: No history of diabetes or thyroid disease.
- *Lymphatic/Hematologic*: No lymphatic or hematologic disorders.

(continued on page 628)



Case Study Findings (continued)

Psychosocial profile:

- Last gynecological examination and Pap smear 1 year ago with no abnormalities found. Had one cold in last year without seeking medical attention. Bathes daily and does not douche.
- Typical day includes waking at 6:00 A.M., preparing family breakfast, and performing household chores. Daughter goes to day care 2 days a week while patient volunteers at indigent healthcare facility.
- Eats three meals a day, drinks two cups of coffee. Does not eat at restaurants often, and attempts to make healthy meals.
- Exercise/leisure activities include gardening, reading, and outside activities with family.
- Family pet is a 1-year-old beagle.
- Asleep by 10:30 P.M. (7½ hours a night).
- Does not smoke, drinks one glass of wine occasionally with evening meals, denies recreational drug use.
- Lives in a suburban house of adequate size for a large family.
- No cultural or religious influences on healthcare. Family attends church most Sundays.
- Describes loving relationship with spouse, and states they intend to have three more children. Enjoys being a homemaker.
- Copes with stress by communicating with spouse.

Sexual history findings:

- Sexually active in a monogamous, heterosexual relationship with husband.
- Satisfied with sexual performance.
- Uses oral contraceptives.



Case Study Evaluation

Before you proceed with the physical examination of the female GU system, review the data you have obtained from Mrs. James's history. Document the GU history related to Mrs. James.



CRITICAL THINKING ACTIVITY #1

In view of the subjective data obtained from Mrs. James, what parts of the physical assessment will you be focusing on?



CRITICAL THINKING ACTIVITY #2

Consider the subjective information that you have obtained from Mrs. James. What other historical information should you focus on?

Physical Assessment

Once you have obtained the subjective data, focus on collecting objective data by performing the physical examination.

Approach

Instruct the patient not to use vaginal sprays, douche, or have coitus for 24 to 48 hours before the examination. Doing so might affect the Pap smear and other vaginal cultures. Make sure the examination room is warm, comfortable, and private. Ideally, the foot of the examination table should point away from the door.

Many women feel anxious and fearful about having a genital assessment, so be thoroughly professional and especially sensitive to your patient's needs. This will make the examination easier for both of you. Also listen for any clues that might help you uncover concerns that the woman might be reluctant to express. Having a female attendant present is also advisable. In addition, keep in mind cultural differences. For example, married Muslim women must have their husbands present during the examination. Women of some cultures will not disrobe or allow a man to perform a physical examination on them. Others will not allow a sample of their body fluids to be taken and examined by strangers.

Before you begin, ask the woman to empty her bladder. Have her completely disrobe and put on a hospital gown. She may leave on socks if the stirrups are not covered with padding. Then explain each assessment technique and its purpose and show her the speculum and other equipment. Encourage her to express any anxieties or concerns. Reassure her that the position for the examination and the assessment techniques used may be uncomfortable at times, but they should not be painful.

Assessment of the female reproductive system consists of inspection and palpation of the abdomen, external genitalia, and internal reproductive structures. You will use the speculum to view the internal genitalia and collect specimens for laboratory analysis. Thorough palpation of the internal structures includes a bimanual examination and rectovaginal assessment.



TOOLBOX

Assemble the following equipment for a routine examination: Large hand mirror, gooseneck lamp, disposable gloves, drape, vaginal speculum, cytological materials (Ayre spatula, cytobrush, cotton-tipped applicators, obstetric (OB) swabs, microscope slides), labeled Thayer-Martin culture plate, cytology fixative spray, reagents (normal saline solution or potassium hydroxide [KOH]), Hemoccult slide and developer, acetic acid, warm water, and water-soluble lubricant.

Performing a General Survey

Before you focus on the GU assessment, perform a general survey and head-to-toe assessment, looking for clues that might suggest a GU problem. For example, obtain vital signs. Elevations in temperature may indicate an infection that might be accompanied by a GU infection.

Note the woman's emotional status. This examination can produce anxiety. Ensure privacy and expose only the area being assessed. Note the woman's posture and nonverbal behavior. Does her posture reflect pain or discomfort?

Performing a Head-to-Toe Physical Assessment

The GU system affects many different systems. So examine the patient for specific changes that may affect the GU system.

Positioning the Woman

Unfortunately, the GU assessment requires uncomfortable positioning for the patient. The lithotomy position is preferred, but if a woman is unable to assume this position, it

can be modified or a side-lying Sims' position can be used instead. Before performing the assessment, do the following:

- Drape the woman's torso and thighs while she is sitting.
- Help her into the lithotomy position with her heels in the stirrups. This abducts the legs and flexes the hips.
- Help her move her buttocks down to the end of the examination table until they are flush with the edge of the table.
- Readjust the drape to cover her as much as possible.
- Adjust the stirrups until she feels comfortable.
- Sit on a stool at the foot of the examination table facing the woman's external genitalia.
- Adjust the lighting source.
- Touch the inside of the woman's thigh before you touch her vulva. This helps her get used to your touch.



HELPFUL HINT

Change gloves as needed during the examination to prevent cross-contamination

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

General Health Survey

INSPECT

- Well developed physically.
- No acute distress.
- Awake, alert, oriented. Memory intact.
- Affect appropriate.
- Weight stable, no loss of height.

Integumentary

Skin

INSPECT

- Intact, no lesions.

Hair

- Even distribution of hair.

HEENT

Head and Neck

INSPECT/PALPATE

- No palpable lymph nodes.
- Thyroid not palpable.

ABNORMAL FINDINGS/RATIONALE

- Personality changes and mental deterioration can accompany late-stage syphilis.
- Depression can affect sexual functioning.
- Loss of height may be seen in postmenopausal women with osteoporosis.
- Weight loss resulting from anorexia can cause amenorrhea. Weight gain is associated with pregnancy. Unexplained weight loss may be associated with malignancy.

- Rashes and skin lesions are associated with many STDs.
- *Systemic rash*: Secondary syphilis.
- *Vesicles*: Herpes simplex types 1 and 2.
- *Increased skin pigmentation*: Increased hormones during pregnancy.

- *Alopecia*: Secondary syphilis.
- Abnormal increase in body hair (hirsutism): Decrease in female hormones.

- *Palpable lymph nodes*: May indicate systemic infection or malignancy.
- *Enlarged thyroid (hypo- or hyperthyroidism)*: May affect reproductive and sexual functioning.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<p><i>Eyes, Mouth, and Throat</i></p> <ul style="list-style-type: none"> ■ Conjunctiva pink, no drainage. ■ Oral mucous membranes pink, moist, and intact. <p>Respiratory <i>INSPECT/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Lungs clear; no shortness of breath (SOB). <p>Cardiovascular <i>INSPECT/PALPATE/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Regular heart rate and rhythm, no extra sounds. ■ Positive pedal pulses, negative Homans' sign, no edema. <p>Breasts <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ Nontender, no discharge (unless lactating), no lesions. <p>Abdomen <i>INSPECT/PALPATE/AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Positive bowel sounds; abdomen soft, nontender, no masses. ■ No organomegaly. <p>Musculoskeletal <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ +5 muscle strength, full range of motion, no joint deformities. <p>Neurological <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ No weakness, paralysis. ■ Awake, alert, and oriented x 3, memory intact. ■ Affect appropriate. <p>Lymphatic/Hematologic <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ No palpable lymph nodes. 	<ul style="list-style-type: none"> ■ <i>Conjunctivitis</i>: Can be caused by gonorrhea. ■ <i>Oral lesions</i>: Associated with STDs. ■ Chronic lung disease may impair sexual functioning. ■ Anemia can cause tachycardia and a systolic flow murmur. ■ Oral contraceptives associated with increased risk of thrombus formation. ■ Cyclic hormonal changes may cause breast fullness and tenderness. ■ <i>Palpable abdominal masses</i>: May be a fetus or fibroid tumor. ■ <i>Enlarged liver and ascites</i>: Associated with metastasis of gynecological cancers. ■ Weakness or joint pain may limit sexual functioning. ■ <i>Weakness/paralysis</i>: Neurosyphilis. ■ <i>Charcot joints</i>: Late syphilis. ■ <i>Unexplained fractures or spinal changes (dowager's bump)</i>: Osteoporosis in postmenopausal women. ■ <i>Weakness and paralysis</i>: Neurosyphilis. ■ <i>Changes in mental status/psychosis</i>: Late syphilis. ■ Depression may affect sexual functioning. ■ <i>Palpable lymph nodes</i>: May indicate infection. ■ <i>Inguinal lymph nodes</i>: Associated with metastatic disease.



HELPFUL HINT

Culture any abnormal discharge.

Performing a Genitourinary Physical Assessment

Assessment of the female GU system consists of inspection and palpation, first of the external genitalia and then of the internal genitalia.

Assessing the External Genitalia

Inspection

Take note of hair pattern and distribution. With gloved hand, separate the pubic hair, checking for skin condition and the presence of lesions, discharge, or parasites. If dis-

charge is present, obtain a specimen for culture. Sometimes a complete examination is not needed. In this case, if you are caring for a patient in the home or hospital, examine her external genitalia while you bathe her.

The color and odor of the discharge may suggest a specific organism. For example, frothy, malodorous, watery, green or gray discharge is associated with *Trichomonas* or *Haemophilus* organisms; purulent green-yellow discharge may indicate gonorrhea; heavy gray-white discharge is a sign of *Chlamydia trachomatis*; and a cheesy discharge signals candidiasis.

Palpation

Gently palpate the external genitalia, noting any masses, nodules, discharge, or areas of tenderness. Be alert for vertical and horizontal lymph node enlargement that

INSPECTING THE EXTERNAL GENITALIA

1. External Genitalia

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Skin

- With patient in lithotomy position, note pubic hair distribution, and check skin for lesions, edema, ecchymosis, or pubic pediculosis.



Inspecting the skin

- *Hair distribution shaped like inverse triangle. May be some growth on abdomen and upper inner thighs. A diamond-shaped pattern from the umbilicus may be caused by cultural or familial differences.*
- *No parasites should be present.*
- *Skin over mons pubis is clear except for nevi and normal hair distribution. Labia majora and minora are symmetrical with smooth to moderately wrinkled, slightly pigmented skin without ecchymosis, excoriation, nodules, edema, rash, or lesions*

Clitoris

- *Note position, redness or lesions.*



Touching inside of the thigh

- *Clitoris about 2 cm long and 0.5 cm in diameter.*
- *No redness or lesions*

Urethral Meatus

- Use thumb on dominant hand to separate labia minora to expose urethral meatus, which is very sensitive to touch.



Using thumb and index finger to separate the labia minora

- *Note shape, color, and size.*
- *Note color, position, redness, edema, lesions, or discharge.*
- *Urethral opening slitlike, midline, and free of discharge, swelling, redness, or lesions.*

Vaginal Introitus

- *With labia minora retracted, ask patient to bear down while you observe for patency and bulging.*
- *Note color, discharge, lesions, or bulges.*
- *Introitus mucosa pink and moist. Normal vaginal discharge is clear to white and free of foul odors. Some white clumps may be seen that are mass numbers of epithelial cells. Introitus is patent, and there is no bulging or tenderness.*

Perineum and Anus

- *Note color, lesions, bulges, or hemorrhoids.*



Inspecting the perineum and anus

- *Perineum is smooth and slightly darkened. A well-healed episiotomy scar is normal after vaginal delivery. Anus is dark pink to brown and puckered. Skin tags are common around anal area.*

(continued)

INSPECTING THE EXTERNAL GENITALIA

1. External Genitalia (continued)

ABNORMAL FINDINGS/RATIONALE

- Diamond-shaped pattern not associated with cultural or familial differences is abnormal and may be hirsutism, which indicates endocrine disorder.
- Pubic lice, nits, or flecks of residual blood on skin is abnormal. Pediculosis pubis (pubic lice) is an infestation of hairy regions of body. It usually occurs in the pubic area, but may involve hair on the abdomen, chest, and axillae. Nits are minute white louse eggs that attach close to the pubic hair shaft.



Pubic louse

- *Ecchymosis*: May be caused by blunt trauma.
- *Labial varicosities*: Pregnancy or uterine tumor.
- *Edema*: Hematoma formation, obstruction of lymphatic system, or Bartholin's cyst.
- *Broken areas of skin*: Ulcerations or abrasions caused by infection or trauma.
- Rash over mons pubis and labia is abnormal.
- *Chancere*: Primary syphilis. Painless, reddish, round ulcer with depressed center and indurated edges. Lasts for 4 weeks and then disappears without leaving a scar.
- *Condylomata acuminatum (venereal warts)*: Caused by HPV. White, dry, painless growths with narrow bases.
- *Herpes simplex*: Small, red, painful vesicles that progress to ulcer stage. Pruritus may be present.
- *Hypertrophy of clitoris*: May indicate female pseudohermaphroditism caused by androgen excess.
- *Chancroid*: Painful ulcer with rough floor and purulent yellow exudate heals, leaving a scar.
- **Female circumcision** is removal of all or part of the clitoris, labia minora, and labia majora, usually in early childhood or early adolescence. This practice is widespread in many African countries and among some Muslim groups.
- *Discharge of any color from meatus*: May indicate UTI.

- *Swelling or redness around meatus*: Possible infection of Skene's glands, urethral caruncle, urethral carcinoma, or prolapse of urethral mucosa.
- *Pale color, dryness*: Possible atrophy from topical steroids and aging.
- Foul-smelling discharge that is not clear to slightly pale white is abnormal.
- Gonorrhea, chlamydia, *Candida*, *Trichomonas*, bacterial vaginosis, atrophic vaginitis, or cervicitis are possible infectious processes that produce an abnormal vaginal discharge.
- *External tear*: May indicate trauma from sexual activity or abuse.
- *Fissure*: May indicate congenital malformation or childbirth trauma.
- **Cystocele** is bulging of bladder into anterior vaginal wall.
- **Cystourethrocele** is bulging of anterior vaginal wall, bladder, and urethra into vaginal introitus.
- **Rectocele** is bulging of rectal wall into posterior vaginal wall.
- *Fissure or tear of perineum*: May be caused by trauma, abscess, or unhealed episiotomy.
- *Venous prominences in anal area*: May indicate external hemorrhoids.



Herpes vulvovaginitis

PALPATING THE EXTERNAL GENITALIA

1. External Genitalia

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS



HELPFUL HINT

Maintain standard precautions and wear gloves.

Labia

- Palpate each labium between thumb and index finger of your dominant hand.
- *Labia soft and uniform in structure with no swelling, pain, induration, or purulent discharge.*



Palpating Bartholin's glands

Urethral Meatus and Skene's and Bartholin's Glands

- Insert index finger of your dominant hand into vagina, and apply pressure to anterior aspect of vaginal wall to milk urethra.
- Swab any discharge with cotton-tipped applicator for microscopic exam.
- *Milking urethra no pain or urethral discharge.*
- *Skene's glands surround urethral meatus.*
- *Bartholin's glands deep in perineal structures.*



Palpating Skene's glands and milking the urethra

Vaginal Introitus

- Keep your finger in the vagina, and ask the patient to squeeze the vaginal muscles around it.
- *Vaginal muscle tone tight and strong in nulliparous women and diminished in parous women.*

Perineum

- Partially remove your finger from the introitus until it is posterior to the perineum, with your thumb anterior to the perineum.
- Assess tone and texture.
- *Perineum smooth, firm, and homogenous in nulliparous women and thinner in parous women.*

ABNORMAL FINDINGS/RATIONALE

- *Swelling, redness, induration, or purulent discharge from labial folds with hot, tender areas:* May indicate Bartholin's gland infection caused by gonococci or *Chlamydia trachomatis*.
- *Pain and discharge:* May indicate Skene's gland infection or UTI.
- *Significantly diminished/absent muscle tone:* May result from injury, age, childbirth, or medication.
- *Bulging from vagina:* Cystocele, rectocele, or uterine prolapse.
- *Thin perineum, fissures, tears:* May indicate atrophy, trauma, or an unhealed episiotomy.

2. Bladder and Kidneys (see Chapter 17, Assessing the Abdomen)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Bladder and kidneys not palpable.*

may indicate an infection, malignancy, or other systemic problem.

Assessing the Internal Genitalia

Next, perform the internal examination. It includes the speculum examination and specimen collection and bimanual palpation of the internal structures.

Inspection

Select the appropriate size speculum. You should have several sizes and types of sterilized, metal specula on hand. The two most commonly used types are the Graves and the Penderson. The Graves speculum is available in 3½" to 5½" lengths and ¾" to 1½" widths. It has a spoon-shaped flare at the distal blade ends, which helps you view the cervix and examine multiparous women. The narrow, flat Penderson speculum is used for children, women who have never been sexually active, nulliparous women, and some postmenopausal women.

Different-sized plastic specula are also available. Some of them make a loud clicking noise when the blades are opened, so be sure to warn your patient before the examination. Plastic specula may be more difficult to insert into the vagina. Be very careful with insertion. Surface tension factors and rough edges on some low-cost brands may also tear the vaginal mucous membrane lining.



HELPFUL HINT

Before insertion, lubricate the speculum with warm water. Commercial lubricants may interfere with the accuracy of the cytological samples and cultures.

To perform the examination, hold the speculum in your dominant hand with the blades closed. Rest your index finger at the proximal end of the superior blade. Wrap your other fingers around the handle with your thumb over the thumbscrew. Insert your nondominant index and middle fingers ventral sides down just inside the vagina and apply pressure to the posterior vaginal wall to help the perineal muscles relax. If the woman is tense, instruct her to take deep breaths. When you feel the muscles relax, insert the speculum at an oblique angle. After withdrawing your finger, gently rotate the speculum to a horizontal angle and advance it at a 45-degree downward angle against the posterior vaginal wall. Using your dominant thumb, depress the lever to open the blades and visualize the cervix. If you cannot see the cervix, close the blades and pull the speculum back 2 to 3 cm and reinsert it at a different angle. Once the cervix is visualized, lock the speculum blades into place. Adjust the light source. If any discharge obstructs the cervix, use an OB swab to clear the cervix. Inspect the cervix and os for color, position, size, surface, discharge, and shape.



HELPFUL HINTS

- Avoid pinching the labia or pulling the woman's pubic hair while inserting the speculum. If insertion causes undue pain, stop and evaluate your speculum position.
- If the woman seems tense or anxious, suggest that she take deep breaths during the examination.

SPECULUM EXAMINATION OF THE INTERNAL GENITALIA


1. Cervix

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Lubricate speculum with warm water.
- Hold speculum in dominant hand with thumb on thumbscrew, insert your nondominant index finger and middle finger into vagina, and apply pressure to the posterior vaginal wall to relax the perineal muscles.
- Insert speculum with blades closed at an oblique angle.
- Open blades and rotate speculum to horizontal angle and advance it at a 45-degree downward angle against the posterior wall, open blades and lock screw with thumb once cervix is visualized.
- Inspect color, shape, position, size, surface, lesions, and discharge.



Inserting the speculum



Proper position of the speculum in the vagina

SPECULUM EXAMINATION OF THE INTERNAL GENITALIA

1. Cervix (continued)



Opening the speculum



View through the speculum

- *Cervix glistening pink; pale pink after menopause or blue (Chadwick's sign) during pregnancy.*



Chadwick's sign

- *No lesions. Located midline in vagina with anterior or posterior position relative to vaginal vault. Size is about 2.5 cm. Normal discharge is white. No odor.*
- *Nulliparous woman has small round or oval os. In parous woman, os is a horizontal slit.*



A

B

A. Nulliparous cervical os. B. Parous cervical os

ABNORMAL FINDINGS/RATIONALE

- *Cyanosis without pregnancy:* Venous congestion or systemic hypoxia as in congestive heart failure.
- *Redness or friable appearance:* Infection and inflammation, such as in chlamydia or gonorrhea.
- *Lateral positioning of cervix:* Tumor or adhesions.
- *Projection of cervix into vaginal vault greater than 2.5 cm:* Uterine prolapse.
- *Cervical size greater than 4 cm:* Hypertrophy caused by inflammation or tumor.
- Reddish circle around os may be abnormal.
- *Ectropion or eversion:* Occurs when squamocolumnar junction appears on ectocervix. Can result from lacerations during childbirth or be congenital variation.
- *Nabothian cysts:* Small, round, yellow, benign lesions from obstruction of cervical glands.
- *Polyps:* Bright-red, soft protrusions through cervical os.
- *Hemorrhages over cervical surface (strawberry spots):* Associated with trichomonal infection.
- *Unilateral transverse, bilateral transverse, stellate, or irregular cervical os:* Caused by cervical tears occurring during rapid, second-stage childbirth delivery, forceps delivery, or trauma.
- *Greenish-yellow, mucopurulent discharge that adheres to vaginal walls, pus in os:* Gonococcal infection.
- *White, cottage cheese-like discharge that adheres to vaginal walls, with patches on os:* *Candida* infection.
- *Red spots on cervix; grayish-yellow, purulent, often bubbly discharge that smells fishy and often pools in fornix:* *Trichomonas*.

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> Changes in energy level Weight changes Fevers	General	<i>Measure:</i> Height and weight Vital signs, checking for temperature elevations, HTN <i>Inspect:</i> Signs of discomfort Affect
Integumentary	<i>Ask about:</i> Changes in hair growth Rashes, lesions	Integumentary	<i>Inspect:</i> Skin lesions Hair distribution
HEENT <i>Head and Neck</i>	<i>Ask about:</i> Headaches Swollen glands, nodes	HEENT <i>Head and Neck</i>	<i>Palpate:</i> Lymph node enlargement Thyroid enlargement
<i>Eyes, Mouth, Throat</i>	<i>Ask about:</i> Eye drainage Oral lesions Sore throat	<i>Eyes and Mouth</i>	<i>Inspect:</i> Oral mucosa for redness and lesions Conjunctiva for drainage
Respiratory	<i>Ask about:</i> History of respiratory disease	Respiratory	<i>Inspect:</i> Signs of respiratory distress <i>Auscultate:</i> Abnormal breath sounds
Cardiovascular	<i>Ask about:</i> History of cardiovascular disease, HTN, thrombophlebitis, heart murmurs	Cardiovascular	<i>Inspect:</i> Signs of impaired circulation Skin changes <i>Palpate:</i> Pulses, edema, Homans' sign <i>Auscultate:</i> Extra heart sounds
Gastrointestinal	<i>Ask about:</i> History of liver disease Loss of appetite Abdominal pain	Gastrointestinal	<i>Inspect:</i> Ascites <i>Palpate/percuss:</i> Liver enlargement Masses
Neurological	<i>Ask about:</i> History of neurological problems Paralysis Tremors Personality changes Depression	Neurological	<i>Test:</i> Changes in mental status and affect Sensory deficits
Musculoskeletal	<i>Ask about:</i> Weakness/limitations Joint pain, swelling Unexplained fracture	Musculoskeletal	<i>Test:</i> Muscle strength <i>Inspect:</i> Joint swelling and deformity Spinal deformities
Endocrine	<i>Ask about:</i> History of diabetes and thyroid disease	Endocrine	<i>Palpate:</i> Thyroid
Lymphatic/Hematologic	<i>Ask about:</i> History of malignancies, HIV, AIDS Abnormal menses	Lymphatic/Hematologic	<i>Palpate:</i> Lymph nodes

Assessment of Female GU System's Relationship to Other Systems

Collecting Specimens

Collect specimens (smears and cultures) after inspecting the cervix and cervical os. Collect the Pap smear first, using three different slides or one slide for all three smears. The **Pap smear** is a collection of three specimens taken from the endocervix, the cervix, and the vaginal pool. The purpose is to evaluate cervicovaginal cells for pathology that might indicate cancer. All women over age 21 or 3 years after the onset of sexual activity should have an annual Pap screening.

Endocervical Smear

Using your dominant hand, insert a cotton-tipped swab, cytobrush, or the longer, serrated end of an Ayre spatula through the speculum into the cervical os about 1 cm. Rotate the instrument between your index finger and your thumb, first 360 degrees clockwise and then counterclockwise. Keep the instrument in contact with the cervical tissue. If you use a swab, leave it in the cervical os for 30 seconds to ensure saturation. Remove the instrument and spread the cells in a rolling motion on the slide. If you are putting all three smears on one slide, spread the cells on one-third of the slide only. (Some cells are marked “E” to show you where to spread the cells.) Do not press down hard or wipe the instrument back and forth because this will destroy the cells. Discard the instrument after use.

Cervical Smear

Insert the bifurcated end of the Ayre spatula through the speculum base. Place the longer end of the spatula into the cervical os and put the shorter part snug against the cervix. Rotate the spatula 360 degrees one time only. Remove the spatula and gently spread the specimen on another slide or on the second third of the same slide (may be marked “C”).

Vaginal Pool Smear

Reverse the Ayre spatula and place the rounded end into the posterior vaginal fornix. Gently scrape the area. You may also use a cotton-tipped applicator. Remove and gently spread the specimen on a separate slide or on the last one-third of the slide (may be marked “V”). Dispose of the spatula. Spray the entire slide (or slides) with cytological fixative within 1 minute and send to the laboratory. Slides and fixatives are not needed if you use the thin prep Pap test.

Other Smears

Next, collect other smears, if needed:

- **Gonococcal smear:** Insert a sterile cotton swab 1 cm into the cervical os. Hold the applicator in place 20 to 30 seconds. Remove the swab and roll it on a Thayer-Martin culture plate in a Z pattern. Rotate the swab as you roll it to ensure that all of the specimen is used.
- **Saline mount (wet prep):** Spread a sample of the cervical or vaginal pool specimen onto a microscope slide, add one drop of normal saline solution, and apply a cover slip.
- **KOH prep (rapid test for *Candida*):** Spread a sample of the cervical or vaginal pool specimen onto a microscope slide, add one drop of KOH, and apply coverslip.
- **Acetic acid wash:** After all other specimens have been collected, swab the cervix with a 5 percent acetic acid solution.
- **Anal culture:** Insert a sterile cotton swab 1 cm into the anal canal and hold in place for 20 to 30 seconds. Remove the swab. Discard the applicator if fecal material is collected and restart. Roll the swab in a Z pattern on a Thayer-Martin culture plate. Dispose of the swab.

COLLECTING SPECIMENS FOR CULTURE


1. Pap Smear

TEST/NORMAL VARIATIONS

- **Endocervical smear:** With dominant hand, insert cotton-tipped swab, cytobrush, or Ayre spatula into cervical os 1 cm and rotate 360 degrees clockwise, then counterclockwise.
- **Cervical smear:** Insert long end of spatula into os and shorter end against cervix and rotate 360 degrees.
- **Vaginal pool smears:** Insert rounded end into the posterior vaginal fornix and gently scrape.
- **For all Pap smears:** Apply specimens to slides and apply fixatives. (Slides and fixatives not needed if using thin prep Pap test.)
- **Cytology report says “within normal limits” using Bethesda system.** This denotes a lack of pathogenesis.



Taking a vaginal pool smear



Taking a cervical smear

(continued)

COLLECTING SPECIMENS FOR CULTURE

1. Pap Smear (continued)

ABNORMAL FINDINGS/RATIONALE

- *“Benign cellular changes” on report:* May be caused by fungal, bacterial, protozoan, or viral infections.
- *“Atypical squamous cells” on report:* Causes include inflammatory or infectious processes, a preliminary lesion, or an unknown cause.
- *“Epithelial cell abnormalities” on report:* May indicate squamous intraepithelial lesion, squamous cell carcinoma, or glandular cell abnormalities often seen in postmenopausal women not on HRT.

2. Gonococcal Smear

TEST/NORMAL VARIATIONS

- Insert sterile cotton-tipped swab 1 cm into os and hold in place for 20 to 30 seconds. Remove swab and roll swab on Thayer-Martin culture plate in Z pattern.
- *Cervicovaginal tissues are normally free of Neisseria gonorrhoeae.*

ABNORMAL FINDINGS/RATIONALE

- It is abnormal to have a large number of gram-negative diplococci present.

3. Saline Mount (Wet Prep)

TEST/NORMAL VARIATIONS

- Spread cervical or vaginal pool specimen on microscope slide with 1 drop of saline. Apply cover clip and examine.
- *Fewer than 10 white blood cells (WBCs) per field.*
- *No protozoa or other organisms.*

ABNORMAL FINDINGS/RATIONALE

- *Large number of WBCs (inflammatory response):* *Chlamydia* or a bacterial infection. Protozoa are indicative of trichomoniasis.

4. KOH Prep

TEST/NORMAL VARIATIONS

- Spread cervical or vaginal pool specimen on microscope slide with 1 drop of KOH. Apply cover clip and examine.
- *Cervicovaginal tissues are free of C. albicans.*

ABNORMAL FINDINGS/RATIONALE

- *Presence of yeast or pseudohyphae (chains of budding yeast):* Indicates *Candida* infection.

5. Acetic Acid Wash

TEST/NORMAL VARIATIONS

- Swab cervix with 5 percent acetic acid after all other specimens have been collected.
- *No change in appearance of cervix.*

ABNORMAL FINDINGS/RATIONALE

- *Rapid acetowhitening or blanching:* May indicate presence of HPV, causative agent of genital warts.

COLLECTING SPECIMENS FOR CULTURE

6. Anal Culture

TEST/NORMAL VARIATIONS

- Insert sterile cotton-tipped swab 1 cm into anus, hold for 20 to 30 seconds. If fecal material present, discard and obtain another sample. Roll swab in “Z” pattern on Thayer-Martin plate.
- *Anal tissues are free of N. gonorrhoeae.*

ABNORMAL FINDINGS/RATIONALE

- *Large number of gram-negative diplococci:* Indicates *N. gonorrhoeae*.

Inspecting the Vaginal Wall

Once you have obtained the specimens, inspect the vaginal wall for color, texture, and lesions as you withdraw the speculum. To remove the speculum, disengage the

locking device and slowly withdraw the speculum without closing the blades. Rotate the speculum into an oblique position as you remove it.

INSPECTING THE VAGINAL WALL

1. Vaginal Wall

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Inspect vaginal walls as you withdraw speculum.
- Note color, rugae, lesions.
- *Vaginal wall pink, moist, deeply rugated, and without lesions or redness. In postmenopausal women, walls may be smooth, shiny, and transparent.*

ABNORMAL FINDINGS/RATIONALE

- *White spots on vaginal wall:* May be leukoplakia from *C. albicans*. Repeated occurrences may indicate HIV infection.
- *Pallor of vaginal walls:* Anemia or menopause.
- *Redness of vaginal walls:* Inflammation, hyperemia, or trauma from tampon insertion or removal.
- *Vaginal lesions or masses:* Carcinoma, tumors, and DES exposure.

Palpation

Bimanual technique is used to assess the internal genitalia. Explain the assessment to the woman. As you perform the examination, observe her face for signs of discomfort. Apply a water-soluble lubricant to the first two fingertips of your dominant hand. As you stand between the woman's legs, place your nondominant hand on her abdomen below the umbilicus. Insert your dominant index and middle fingers into the vagina with your palm up. Advance slowly while palpating the vaginal walls. Keep your thumb abducted and away from the urethral meatus and clitoris. Rotate your wrist so you can palpate all surfaces in the vagina.

Next, keeping your dominant hand in the same position, with your palm up, assess the cervix for consistency, position, shape, and tenderness. Move the cervix from side to side to assess mobility. Palpate the fornices with your fingertips, noting nodules or irregularities.

Now, reach under and behind the cervix and lift the uterus toward the abdomen and your external hand as you apply pressure toward the internal fingers. Keep your wrist straight. Try to grasp the uterus between your hands. Evaluate the uterus for size, shape, consistency, mobility, tenderness, masses, and position. (Note: A retroverted, retroflexed uterus can be assessed only rectovaginally.)

After assessing the uterus, palpate the fallopian tubes and ovaries. Move your intravaginal hand to the right lateral fornix, and move your external hand to the right lower quadrant of the abdomen. Push your external hand inward and downward while pushing your intravaginal hand inward and upward. Palpate for size, shape, consistency, and mobility. Repeat on the left side.

Fallopian tubes are rarely palpable. Palpation of the ovaries depends on the woman's age and size. The ovaries may be tender during palpation if the woman is in the luteal phase of the menstrual cycle.

BIMANUAL PALPATION OF THE INTERNAL GENITALIA

1. Vaginal Wall

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Apply water-soluble lubricant to first 2 fingers of dominant hand and insert into the vagina.
- Note consistency of wall.
- *Nontender, with smooth or rugated surface with no lesions, masses, or cysts.*



Inserting the finger into the vagina

ABNORMAL FINDINGS/RATIONALE

- *Lesions, masses, scarring, or cysts:* May be benign, such as inclusion cysts, myomas, or fibromas; or may be malignant. Most common site for vaginal malignancies is upper one-third of posterior wall.

2. Cervix

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- With fingers inserted into the vagina, palpate cervix.
- Note position, mobility, consistency, and tenderness.
- *Mobile, nontender, smooth and firm, symmetrically round, and in midline, movable 1 to 2 cm in each direction without discomfort.*
- *Softening of cervix (Goodell's sign) is seen at 5th to 6th week of pregnancy.*

ABNORMAL FINDINGS/RATIONALE

- *Pain on palpation or to assess mobility (positive Chandleier's sign or cervical motion tenderness):* PID or ectopic pregnancy.
- Irregular surface, immobility, or nodular surface of cervix is abnormal. Causes may include malignancy, nabothian cysts, or polyps.

3. Fornices

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Palpate the vaginal walls around the cervix.
- *Walls are smooth and without nodules.*

ABNORMAL FINDINGS/RATIONALE

- *Nodules or irregularities:* Malignancy, polyps, herniations.

BIMANUAL PALPATION OF THE INTERNAL GENITALIA

4. Uterus

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- With fingers inserted into the vagina and palm upward, place nondominant hand on the abdomen above the symphysis pubis.
- With dominant hand, reach under and behind cervix, and lift uterus up toward your other hand on the abdomen and palpate uterus.
- Note size, shape, consistency, mobility, position, tenderness, masses.
- *Size varies according to parity. Pear shaped in nonpregnant woman and more rounded in parous woman. Smooth, firm, mobile, nontender, and without masses.*



Palpating the uterus

ABNORMAL FINDINGS/RATIONALE

- **Enlargement/changes in shape:** Enlargement may indicate intrauterine pregnancy or tumor.
- **Nodules:** May be myomas, tumors containing muscle tissue.

5. Adnexa

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- With fingers inserted into the vagina and palm upward, move toward the right lateral fornix, press upward while other hand on abdomen presses downward in the right lower quadrant.
- Note size, shape, consistency, mobility, and tenderness.
- Use same technique to palpate left ovary but position hands to left.
- *Ovaries almond-shaped, firm, smooth, and mobile without tenderness.*
- *Nonpalpable in postmenopausal women.*



Palpating the ovaries

ABNORMAL FINDINGS/RATIONALE

- **Enlarged, irregular, nodular, painful, immobile ovaries:** Ectopic pregnancy, ovarian cyst, PID, malignancy.

Rectovaginal Examination

After completing the bimanual internal genitalia examination, remove your dominant hand from the vagina and change gloves. Apply lubricant to the fingertips of your dominant hand. Explain the procedure to the woman. Ask her to strain downward while you assess anal sphincter tone. To do this, insert your dominant

index finger back into the vagina, and insert your middle finger into the rectum. Advance the rectal finger forward as you depress the abdomen with your nondominant hand. Assess the cervix, uterus, and rectovaginal septum for patency. Withdraw your fingers. If there is stool on your glove, you may test it for occult blood at this time.

RECTOVAGINAL EXAMINATION

1. Rectum, Anal Sphincter, Rectovaginal Septum

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Change gloves, and apply lubricant to fingers of dominant hand.
- As patient bears down, insert index finger into the vagina, and middle finger into the rectum, while palpating lower abdomen with nondominant hand.
- Note rectal wall consistency, anal sphincter tone, masses.
- Withdraw fingers; if stool present, test for occult blood.
- *Rectal walls smooth and free of lesions. Rectovaginal septum patent. Rectal pouch rugated and free of masses.*
- *Anal sphincter tone is strong. Cervix and uterus are smooth.*
- *Test for occult blood negative.*



Inserting the fingers for the rectovaginal exam



Performing the rectovaginal exam



Proper positioning of the hands

ABNORMAL FINDINGS/RATIONALE

- *Masses or lesions:* Consider malignancy or internal hemorrhoids.
- *Lax sphincter tone:* Perineal trauma from childbirth, anal intercourse, or neurological disorders.
- *Stool test positive for occult blood:* Warrants further investigation.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. James have concluded. The following are your findings:

■ General Health Survey

- Well-developed 29-year-old African American female in no acute distress.
- Head-to-toe scan revealed no pertinent findings related to the GU system.

■ Vital Signs

- Temperature, 98.8°F
- Pulse, 80 beats per minute (BPM), strong and regular.
- Respirations, 12/min, regular and unlabored.
- Blood pressure (BP), 122/76.
- Height, 5'5".
- Weight, 122 lb.

■ GU

- Appropriate pubic hair distribution for age.
- External genitalia pink and moist with no varicosities, lesions, organisms, edema, or abnormal discharge.
- Small amount of white, odorless discharge.
- No swelling, tenderness, or discharge on palpation of Bartholin's glands and Skene's glands.
- No masses, lesions, or anatomical deviations of vulva and perineum. Skin intact, smooth.
- Cervix smooth, moist, shiny pink, 2 to 3 cm in diameter at midline, firm, nontender, and midplane.
- No lesions, nodules, masses, discharge, or bleeding. Vaginal walls are pink, moist, rugose, and without lesions, swelling, or masses.

- Concentric rugae around vaginal wall, no nodules or tenderness.
- Uterus is appropriate size for age and condition. Firm, pear-shaped, symmetrical, slightly mobile, and anteverted. No masses or tenderness.
- Ovaries and fallopian tubes are nonpalpable, nontender.
- No hemorrhoids, painful areas, masses, or nodules in rectum.



Case Study Analysis and Plan

Now that you have completed your assessment of Mrs. James, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mrs. James, then use this information to identify appropriate nursing diagnoses. Here are some possible ones. Cluster the supporting data.

1. Knowledge deficit related to adverse effects of oral contraceptives
2. Potential disturbance in reproductive health related to abnormal vaginal bleeding

Identify any additional nursing diagnoses.

Identify any collaborative nursing diagnoses.



CRITICAL THINKING ACTIVITY #3

Develop a nursing care plan and a brief teaching plan for Mrs. James, including learning outcomes and teaching strategies, using one of these nursing diagnoses.

RESEARCH TELLS US

Current guidelines by the U.S. Preventative Services Task Force recommend cervical cancer screening every 3 years instead of annually for women who have had three or more previous negative tests. Even with this recommendation, many clinicians still perform annual Pap tests for fear that the risk for cervical cancer is still too high.

To determine the risk of cervical cancer associated with less frequent screenings, Sawaya and colleagues (2003) compared annual screenings for 3 years with

screenings performed every 3 years after a last negative Pap test in women 30 to 64 years of age who had had three or more consecutive negative Pap tests. The risk of cervical cancer for those who extended the screening to 3 years was approximately 3 in 100,000. These findings support the recommendation of extending the screening interval to every 3 years for women with three or more consecutive negative Pap test results. Women should be informed that this is a safe option.

Source: Sawaya, G., McConnell, K., Kulasingam, S., Lawson, H., Kerlikowske, K., Melnikow, J., Lee, N., Gildengorin, G., Meyers, E., and Washington, E. (2003). Risk of cervical cancer associated with extending the interval between cervical-cancer screening. *The New England Journal of Medicine* 349(16):1501–1509.

HEALTH CONCERNS

Risk Factors for Female Genital Cancers

■ Cervical cancer

- Early age at first intercourse.
- Multiple sex partners.
- Previous history of human papillomavirus (HPV).
- Tobacco use.
- Positive family history.
- Gardasil vaccine protects against four types of HPV, which cause 70% of cervical cancer.

■ Endometrial cancer

- History of infertility.
- Failure to ovulate.
- Unopposed estrogen therapy.
- Use of tamoxifen.
- Obesity.
- Positive family history.

■ Ovarian cancer

- Advanced age.
- Nulliparity.
- History of breast cancer.
- Positive family history.

■ Vaginal cancer

- Mother who took DES during pregnancy.

Warning Signs of Reproductive Organ Cancer

- Abnormal vaginal bleeding and vaginal discharge.
- Persistent aching pelvic or lower back pain.
- Postmenopausal bleeding.
- Abdominal pain.
- Abnormal Pap smear.

(health concerns continued on page 644)

HEALTH CONCERNS (continued)

Risk Factors for AIDS

- Sex with partner who engaged in homosexual activity.
- History of multiple sex partners or having a partner with a history of multiple partners, especially without using condoms.
- Intravenous drug use, sharing needles.
- Male partner who had contacts with prostitutes, especially without using condoms.
- Frequent or multiple blood transfusions.

Risk Factors for Major Sexually Transmitted Diseases

- Being young and sexually active (ages 15 to 30).
- History of multiple sexual partners.
- Early onset of sexual activity.
- Previous STD.
- Urban dweller.
- Low income.
- Unmarried.

Common Abnormalities

ABNORMALITY

ASSESSMENT FINDINGS

Bacterial Vaginosis

- Inflammation of the vagina caused by one or more bacteria or bacteria-like organisms.
- Minimal vulvar pruritus.
- Yellowish-white vaginal discharge.
- Occasional foul odor to discharge.
- Wet mount shows WBCs and/or clue cells.

Chlamydia

- A sexually transmitted bacterial infection that can compromise reproductive health.
- Women are often asymptomatic.
- *Genital symptoms*: Increase in clear to white vaginal discharge, dysuria, and postcoital bleeding.
- *Abdominal symptoms*: Increase in menstrual cramping and pelvic pain.
- Dysmenorrhea (menstrual cramping).
- Cramping or aching in lower abdomen midline.
- Aching in lower back.
- Mild tenderness on pelvic and abdominal exam bilaterally.
- No fever.
- Past history of dysmenorrhea.

Endometrial Cancer

- Malignancy of the endometrium.
- Menopausal age.
- History of previous estrogen therapy, nulliparity, obesity, possible diabetes or hypertension.
- Previous curettage, sterility, or poor fertility.
- Red or brown vaginal discharge or abnormal uterine bleeding, spotting for days to months.
- Uterine or adnexal mass, usually nontender, detected on palpation.
- Cervical lesion.
- Pain in later invasive stage.

Endometriosis

- Inflammation of the endometrium.
- Ages 25 to 45.
- Excessive, prolonged uterine bleeding.
- Menstrual pain referred to rectum and lower sacrum.
- Pain on defecation.
- Dysuria.
- Constipation.
- History of menstrual disturbances.

ABNORMALITY	ASSESSMENT FINDINGS
<p>Functional Ovarian Cysts</p> <ul style="list-style-type: none"> ■ May be follicle cysts or corpus luteum cysts 	<ul style="list-style-type: none"> ■ Multiple tender nodules palpable along the uterosacral ligaments or in the rectovaginal septum. ■ Pain on palpation of uterus.
<p>Genital Herpes</p> <ul style="list-style-type: none"> ■ Viral STD. 	<ul style="list-style-type: none"> ■ Ages 20 to 40. ■ Mild pelvic discomfort, low back pain, or dyspareunia. ■ Localized pain and tenderness. ■ Abnormal uterine bleeding. ■ Menstrual irregularities. ■ Delayed menstruation followed by persistent bleeding. ■ Cysts detected on bimanual palpation.
<p>Monilial Vaginitis (Yeast Infection)</p> <ul style="list-style-type: none"> ■ Fungal infection causing vaginal inflammatory process associated with the <i>Candida</i> organism. 	<ul style="list-style-type: none"> ■ Age teens or early 30s. ■ Watery discharge with lesions or sores and blisters on external genitalia. ■ Mild itching and pain. ■ Recent urinary tract or gynecologic exam. ■ Frequent intercourse without male partner using condom. ■ Fever. ■ Enlarged inguinal lymph nodes. ■ Yellow-gray film on cervix.
<p>Ovarian Carcinoma</p> <ul style="list-style-type: none"> ■ Malignancy of ovaries. 	<ul style="list-style-type: none"> ■ Intense vulvar itching and/or burning. ■ Vulvar swelling and excoriation. ■ No vaginal bleeding. ■ Cheesy or curdlike vaginal discharge often present in labial folds. ■ Monilial rash may spread to upper thighs. ■ Wet mount shows yeast or pseudohyphae (chains of budding yeast).
<p>Urinary Tract Infection</p> <ul style="list-style-type: none"> ■ Inflammation of bladder, urethra, ureters, or kidneys. ■ Most common invading organism is <i>Escherichia coli</i>. ■ Sexual intercourse may contaminate urethra with bacteria. ■ Afebrile or low-grade temperature below 101°F. 	<ul style="list-style-type: none"> ■ Early-stage menopause or postmenopause. ■ History of urinary frequency and constipation. ■ Abdominal pain in later stage. ■ Lower abdominal mass. ■ Possible ascites. ■ Irregular postmenopausal bleeding (possible but infrequent). ■ Displaced cervix. ■ Solid ovarian mass on bimanual palpation.
<p>Urinary Tract Infection</p> <ul style="list-style-type: none"> ■ Inflammation of bladder, urethra, ureters, or kidneys. ■ Most common invading organism is <i>Escherichia coli</i>. ■ Sexual intercourse may contaminate urethra with bacteria. ■ Afebrile or low-grade temperature below 101°F. 	<ul style="list-style-type: none"> ■ Suprapubic tenderness on palpation. ■ Low back discomfort occasionally. ■ Possible costovertebral angle tenderness. ■ No vaginal or urethral discharge. ■ <i>Bladder symptoms:</i> Dysuria, hematuria, nocturia, and urinary frequency.

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the female GU system.

Ask the patient the following:

- If premenopausal, when was your LMP?
- Do you have any pain? Location?
- Can you describe your menstrual cycle? What was your age of menarche? What was your age at menopause? Do you have any abnormal vaginal bleeding?
- What are your sexual practices? How many partners have you had? What type of sexual activity: oral, genital, anal? What is your sexual preference: heterosexual, homosexual, or bisexual? Do you practice safe sex? What do you use? Do you use contraceptives? If, yes, what type?
- Obstetric history, Have you ever been pregnant? Number of live deliveries? Number of miscarriages? Abortions? Age of first full-term pregnancy?
- Do you have any vaginal discharge? If so, can you describe it?
- Do you have a history of GU surgery, STDs, or cancer?
- Are you currently taking any medications? HRT? BCP?
- Do you have any urinary symptoms, such as burning, frequency, or dysuria?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam.

Gloves
Drape
Light
Vaginal speculum
Cytological materials for Pap test
Culture plates
Reagents
Hemocult test

Warm water
Water-soluble lubricant

Position

Lithotomy

Technique

Remember to scan your patient from head-to-toe looking for signs related to the female GU assessment.

Inspection, Palpation, Speculum Examination

Inspect

- **External Genitalia**
 - Hair distribution
 - Skin
 - Clitoris
 - Urethral meatus
 - Vaginal introitus
 - Perineum and anus
- **Internal Genitalia**
 - Cervix
 - Obtain specimens
 - Vaginal wall

Palpate

- **External Genitalia**
 - Labia
 - Bartholin's and Skene's glands
 - Vaginal introitus
 - Perineum
- **Internal Genitalia**
 - Vaginal wall
 - Cervix
 - Fornices
 - Uterus
 - Adnexa
 - Rectovaginal exam
 - Obtain specimens
 - Hematest stool if present

DOCUMENT

Document your findings and develop your plan of care.

SUMMARY


- The female reproductive system is an area where you can have a major impact on your patient's health through routine screening and education.
- A thorough assessment relies on health history data, physical assessment findings, and laboratory results. It requires an understanding of normal anatomy and physiology for various age groups, good interviewing

and communication skills, skill in inspecting and palpating, and expertise in the use of equipment such as the vaginal speculum.

- The assessment also requires sensitivity in responding to the woman's need for privacy and in respecting her personal boundaries.

R E V I E W Q U E S T I O N S

1. Mary Johnson, age 25, has come to the emergency room with the chief complaint of lower abdominal pain. Considering her age, which question would be most appropriate to identify a gynecological problem?
 - a. When was your last menstrual period?
 - b. What was your age of menarche?
 - c. Are your period flows heavy?
 - d. When did you last move your bowels?
2. As you obtain a gynecological history, Mary states that she is a marathon runner. What effect might strenuous exercise have on her menstrual cycle?
 - a. Dysmenorrhea
 - b. Menorrhagea
 - c. Amenorrhea
 - d. No effect
3. Ms. Johnson also reports that she has white, cheesy vaginal discharge. This type of vaginal discharge is often associated with:
 - a. Gonorrhea.
 - b. *Trichomonas*.
 - c. HPV.
 - d. Candidiasis.
4. Sharon Stewart, age 30, is scheduled for her annual gynecological exam. Part of the exam includes a Pap test. The Pap test is a screening test for:
 - a. Cervical cancer.
 - b. Gonorrhea.
 - c. Syphilis.
 - d. HPV.
5. On examination of Ms. Stewart's external genitalia, you note a painless, white, wartlike lesion. The lesion described is most likely:
 - a. Gonorrhea
 - b. Syphilis
 - c. Condylomata acuminatum.
 - d. Herpes simplex
6. When performing a gynecological examination including a Pap test, what lubricant should be used for the speculum?
 - a. No lubricant
 - b. Water
 - c. Water-soluble jelly
 - d. Vaseline
7. You are performing a pelvic examination of Joan Kelly, age 33. She is 10 weeks' pregnant. You examine the color of the cervix. Considering that Mrs. Kelly is pregnant, what color change might you see in her cervix?
 - a. Pink
 - b. Pale pink
 - c. Red
 - d. Blue
8. You examine the color of the cervix. What would be the normal color of the cervix of a premenopausal, nonpregnant woman?
 - a. Pink
 - b. Pale pink
 - c. Red
 - d. Blue
9. What is the normal pubic hair distribution of a 25-year-old female patient?
 - a. Triangle shape
 - b. Circular shape
 - c. Inverse triangle shape
 - d. Diamond shape
10. You are obtaining a vaginal pool specimen to perform a KOH prep test. The KOH prep test is best for assessing:
 - a. *Candida albicans*
 - b. HPV
 - c. Gonorrhea
 - d. Chlamydia



Assessing the Male Genitourinary System

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent male genitourinary history questions
- Obtain a male genitourinary history
- Perform a physical assessment of the male genitourinary system
- Document male genitourinary findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

Assessing the genitourinary (GU) system is essential to obtaining a picture of your male patient's overall health status. Assessment includes both the male reproductive system and the urinary system. The male reproductive system is anatomically divided into external and internal genital organs. The penis and scrotum are external organs and are easily inspected and palpated, whereas the internal structures have limited accessibility.

The kidneys, ureters, bladder, and urethra make up the urinary system. Because these structures are distributed among the retroperitoneal space, abdomen, and genitals, examination of the urinary system is also incorporated into other areas of a total physical assessment. For example, you examine the kidneys during your assessment of the posterior thorax, back, and abdomen.

Your assessment should focus on detecting acute health problems, identifying risks for GU problems, and pinpointing health promotion teaching needs. Be alert for sexual and reproductive problems. Sexually transmitted diseases (STDs) are the most common communicable diseases in the United States. Early detection is critical because untreated STDs can have permanent, devastating effects. Likewise, structural or functional problems in the urinary system can be devastating. Because this system is vital in waste removal and acid-base balance, malfunctions can affect homeostasis and even be life-threatening.

As you read this chapter, remember that the GU system affects and is affected by every other body system, having far-reaching effects on the patient's quality of life.

Anatomy and Physiology Review

Before beginning your assessment, you need a clear understanding of the male GU system. Knowing normal functions and structures will enable you to detect those that are abnormal.

Functions of the male GU system include:

- Manufacturing and protecting sperm for fertilization.
- Transporting sperm.
- Regulating hormonal production and secretion of male sex hormones.
- Providing sexual pleasure.
- Excreting urine.

Structures and Functions

The male genitourinary system (Fig. 19.1) includes both internal and external structures. The penis and scrotum are the external genitalia. The internal genitalia include the testes, epididymis, two vas deferens (also called *ductus deferens*), two seminal vesicles, the ejaculatory duct, the prostate gland, and two bulbourethral (Cowper) glands. Because of the anatomical location of the prostate

gland, access to the prostate for assessment is through digital rectal examination. The **urethra** (about 8 inches or 20 cm long) serves as a passageway for both urine and ejaculated semen, extending from the urinary bladder to the urethral meatus at the tip of the penis.

These structures are further classified as accessory ducts and glands. The vas deferens, ejaculatory duct, and urethra are accessory ducts, and the seminal vesicles, prostate gland, and bulbourethral gland are accessory glands. These glands are responsible for producing a secretion that combines with sperm to create semen. (See *Structures and Functions of the Male Genitourinary System*.)

The Process of Reproduction

Mechanical Process

Male sexual function consists of four stages: erection, lubrication, emission, and ejaculation. Erection of the penis is achieved through physical and psychogenic stimulation of sensory nerves in the genital area. Parasympathetic impulses from the sacral area of the spinal cord result in vascular dilation of the arterioles, causing 20 to 50 mL of blood to fill the corpora cavernosa (erectile tissue), thus expanding the penis. The veins from the tissue compress to occlude venous outflow. At the same time, parasympathetic impulses cause the bulbourethral glands to secrete mucus, which provides lubrication during intercourse. Once the sexual stimulus reaches a critical intensity, the reflex centers of the spinal cord send sympathetic impulses to the genital organs, and orgasm occurs. It begins with contraction of the epididymis and the vas deferens, causing expulsion of sperm through the ejaculatory duct and prostate and into the internal urethra. During emission, the bladder neck closes to prevent the flow of semen into the bladder. Ejaculation follows, with contractions of the penile urethra expelling semen from the urethra, producing a pleasurable sensation known as an **orgasm**. Once sexual stimulation ceases and ejaculation occurs, the penile arteries constrict and the penis returns to its flaccid state, called **detumescence**.

Spermatogenesis

Spermatogenesis is the process by which the testes produce sperm. This process begins during puberty, at around age 13, and continues throughout life. Cells in the seminiferous tubules, called **interstitial cells of Leydig**, secrete the male hormone testosterone. **Testosterone** is responsible for the development of secondary sexual characteristics and reproductive capacity. It is also responsible for male sexual sensation and performance as well as muscle development. Sperm is stored in the epididymis and vas deferens for about 1 month. If it is not ejaculated within that time, the sperm degenerates and is reabsorbed. The normal sperm count is 300 to 500 million sperm with each




FIGURE 19.1. Male GU system.

ejaculation. A man with a sperm count below 20 to 30 million per ejaculation is considered infertile, and a man who is unable to produce viable sperm is considered sterile. (See *Understanding Spermatogenesis*.)

Interaction With Other Body Systems

The cardiovascular, urinary, musculoskeletal, neurological, and endocrine systems contribute to maintaining the male reproductive system.

The Cardiovascular System

Nutrients and sex hormones are transported to the reproductive organs through the vascular system. Sexual excitement causes vasodilation of the arterioles in the penis, resulting in erection.

The Urinary System

The male ejaculates through the urethra, which also carries urine from the bladder to the outside of the body.

Structures and Functions of the Male Genitourinary System

STRUCTURE	DESCRIPTION/PRIMARY FUNCTION
Scrotum	<ul style="list-style-type: none"> Loosely hanging pouch behind penis. Has two sacs that contain testes. Protects testes, epididymides, and part of spermatic cord. Also protects sperm production and viability by maintaining an appropriate surface temperature.
Testes	<ul style="list-style-type: none"> Two essential organs located in scrotum, each approximately 5 cm long by 2.5 cm wide. Each contains tightly coiled seminiferous tubules. Testes are attached to spermatic cord, which allows them to hang vertically in scrotum. Tubules are main component of testes, and they produce sperm by spermatogenesis. Testes secrete some testosterone.
Spermatic Cord	<ul style="list-style-type: none"> Fibrous connective tissue that forms a protective sheath around vas deferens, nerves, blood vessels, lymphatic structures, and muscle fibers associated with the scrotum.
Seminal Vesicles	<ul style="list-style-type: none"> Two pouches that are accessory organs, lying posteriorly to and at the base of the bladder. They contribute about 70 percent of the volume of semen. The fluid is alkaline and rich in fructose, water, and vitamin C and helps provide a source of energy for sperm metabolism. Vesicles also produce prostaglandins, which contribute to sperm motility and viability. The alkaline pH acts to protect the sperm by neutralizing vaginal acidity.
Bulbourethral Glands	<ul style="list-style-type: none"> Two accessory organs located just below the prostate. Also called <i>Cowper's glands</i>, they secrete an alkaline substance that protects sperm by neutralizing the acidic environment of the vagina. These glands also provide lubrication within the urethra to facilitate ejaculation and at the end of the penis during sexual intercourse.
Prostate Gland	<ul style="list-style-type: none"> An accessory gland that borders the urethra near the lower part of the bladder. The prostate produces a milky, alkaline solution that flows into the urethra during sexual intercourse. The liquid aids sperm transport and neutralization of the acidic vaginal secretions. The prostate also produces about 20 percent of all semen.
Epididymis	<ul style="list-style-type: none"> Tightly coiled tube located on the top and behind the testis inside the scrotum. The primary function of the epididymis is to store sperm till maturity. Sperm mature and develop as they pass through the epididymis to the vas deferens.
Vas Deferens	<ul style="list-style-type: none"> Muscular, tubular structure that stretches from the end of the epididymis to the ejaculatory (ductus deferens) duct. Mature sperm remain in ampulla, a wide portion of the vas deferens, before ejaculation. Smooth muscle contraction of the vas deferens propels sperm forward into the ejaculatory duct.
Urethra	<ul style="list-style-type: none"> The terminal end of the urethra transports both urine and semen to the outside of the body. About 6 to 8 inches long, the urethra passes through the prostate, the wall of the pelvic floor, and the length of the penis.
Urethral Meatus	<ul style="list-style-type: none"> External meatus of the urethra. Slitlike orifice located on ventral surface just millimeters from the tip of the glans.
Ureters	<ul style="list-style-type: none"> Hollow tubelike structures connecting the renal pelvis of each kidney with the urinary bladder. They lie within the retroperitoneal space. Ureters propel urine from the renal pelvis into the urinary bladder. Each kidney has a single ureter.

STRUCTURE	DESCRIPTION/PRIMARY FUNCTION
Urinary Bladder	<ul style="list-style-type: none"> ■ Muscular sac that temporarily stores urine. ■ Located anterior to rectum, directly behind the pubic symphysis.
Penis	<ul style="list-style-type: none"> ■ Male sex organ that serves as passageway for sperm to exit and be deposited into the vagina during sexual intercourse. ■ Also serves as an exit for urine.
Glans Penis	<ul style="list-style-type: none"> ■ Tip of penis with many nerve endings. Very sensitive and important in sexual arousal.
Inguinal Canal	<ul style="list-style-type: none"> ■ Encases vas deferens as it passes from the scrotum through the abdominal muscles into the pelvis. The external inguinal ring, located above and lateral to pubic tubercle, is the opening as the spermatic cord travels through the inguinal canal and exits from the scrotum. The spermatic cord enters the abdomen through the internal ring, located about 1 cm above the midpoint of the inguinal ligament. The inguinal ligament is located midway between the anterior superior iliac spine and the pubic tubercle of the symphysis pubis. The inguinal canal is located between the two rings. It runs parallel and above the inguinal ligament (Poupart's ligament), which connects the pubic tubercle and the anterior superior iliac spine. The femoral canal runs parallel and below the inguinal ligament. ■ Lymph nodes from the penis, scrotal surface, and anus drain into the inguinal lymph nodes. ■ Lymph nodes from the testes drain into perivenacaval nodes in abdomen.

Hypertrophy of the prostate gland compresses the urethra, making urination difficult and increasing the risk of urinary tract infections (UTIs) and urinary retention.

The Musculoskeletal System

The bones of the pelvis and muscles of the pelvic floor provide protection and support for some reproductive organs. Androgens (male reproductive hormones) increase bone density. Bladder continence is achieved with an intact detrusor muscle, internal sphincter muscle tone, and external sphincter contraction. Structural and functional integrity of the muscles of the bladder, urethra, and pelvic floor are crucial.

The Neurological System

Parasympathetic nervous system impulses regulate events of the male sexual response, including the vasodilation that results in erection of the penis. **Micturition** is a reflex of the parasympathetic control that stimulates detrusor muscle contraction with the simultaneous relaxation of the external sphincter and pelvic floor muscles.

The Endocrine System

Gonadotropin-releasing hormone (GnRH), which is released by the hypothalamus, and luteinizing hormone (LH) and follicle-stimulating hormone (FSH), which are produced

by the anterior pituitary gland, trigger testosterone secretion and spermatogenesis.

Developmental, Cultural, and Ethnic Variations

Infants

Characteristics of male genitalia are used to help determine gestational age at birth. Premature infants may have undescended testes and few rugae. Breech-delivered infants may have scrotal edema and ecchymoses. Male infants should be assessed for **hypospadias** (urethral meatus opening ventrally on the glans) and **epispadias** (urethral meatus opening dorsally on the glans). Hydroceles and hernias are common findings in boys under age 2.

Children


You need to address specific areas when assessing a male child's GU system. Here are some questions to ask the parent or caregiver:

- Have you noticed any redness, swelling, discharge, or foul-smelling odors in the child's genital area? (Helps identify an infection.)
- Have you noticed asymmetry, lumps, or masses in the genitals? If so, where are they? Are they movable? Hard or soft? Painful to the touch? (Suggests hernia or lymphadenopathy, which may reveal an underlying problem.)

Understanding Spermatogenesis

Spermatogenesis—the formation of mature sperm within the seminiferous tubules—occurs in several stages:

- Spermatogonia, the primary germinal epithelial cells, grow and develop into primary spermatocytes. Both spermatogonia and primary spermatocytes contain 46 chromosomes, consisting of 44 autosomes and the 2 sex chromosomes, X and Y.
- Primary spermatocytes divide to form secondary spermatocytes. No new chromosomes are formed in this stage—the pairs only divide. Each secondary spermatocyte contains half the number of autosomes, 22—1 secondary spermatocyte contains an X chromosome; the other, a Y chromosome.
- Each secondary spermatocyte then divides again to form spermatids.
- Finally, the spermatids undergo a series of structural changes that transform them into mature spermatozoa, or sperm. Each spermatozoon is composed of a head, neck, body, and tail. The head contains the nucleus; the tail, a large amount of adenosine triphosphate (ATP), which provides energy for sperm motility.



- Does the child cry and hold his genitalia as if something were hurting? (Pain may be caused by an acute problem such as a strangulated hernia, but it may also suggest sexual abuse and warrants further investigation. See Appendix C for more information on assessing sexual abuse.)
- Have you ever been told that the child has a hydrocele or hernia?
- Has there been trauma to the child's genitalia, either through injury or during play?
- Is the child toilet trained (usually begins by age 2)? Are you having any toilet training problems? Does your child wet the bed (enuresis)? (Helps you assess developmental progress.)
- Does the child have any problem urinating, and is the urine stream straight? (May identify epispadias or hypospadias.)

Adolescents


Many of the questions you ask adolescents are similar to those you ask adults. But when assessing adolescent males, tailor your questions to younger and older boys. Identifying the patient's understanding of sexuality and sexual practices is important because it provides an opportunity to clarify any misinformation and provide education. Above all, be nonjudgmental. A judgmental approach will almost certainly limit the amount of data you collect.

Identify any areas of concern that your patient has regarding the sexual developmental changes occurring in his body. Often, adolescents need to know that these changes are normal. Also determine if he is sexually active, and if so, whether he practices safe sex. Ask if he has any concerns regarding sexual practices and whether he ever had sex education classes in school.

Tanner staging is used to track the sexual maturation of males during puberty by characterizing pubic hair distribution and penile and testicular size. Assessment techniques are similar for patients of all ages, but your observations and the patients' histories are unique.


Older Adults

Normal physiological changes occur in the male reproductive system with aging. Pubic hair grays and becomes thinner; the penis appears somewhat atrophic; and the testicles may appear smaller and feel slightly softer than in a younger man. The prostate may feel larger than in youth. The scrotal sac loses its elasticity, and testosterone levels usually decrease by age 50. Although normal spermatogenesis continues, sperm output gradually decreases with age. Although the physiological response to sexual stimulation does not change, the time required to achieve erection increases and ejaculation may be less intense, with decreased seminal fluid and rapid detumescence.



NEUROLOGICAL

Sympathetic and parasympathetic nervous systems necessary for sexual function.



INTEGUMENTARY

Androgens influence body hair growth.
Sensory receptors on external genitalia provide sexual sensations.


ENDOCRINE

Hypothalamus secretes GnRH.
Anterior pituitary secretes LH and FSH.
Testes secrete testosterone.



RESPIRATORY

Provides oxygen and removes carbon dioxide for reproductive system.
Respiratory system demands are increased during sexual arousal.




DIGESTIVE

Provides nutrients needed for genitourinary system.



CARDIOVASCULAR

Delivers nutrients and hormones to genitourinary system.
Vascular system needed to achieve erection during sexual arousal.




LYMPHATIC

Secretions have bactericidal chemicals and lysozymes that provide nonspecific defense against STDs.
Provides IgA that assists in tissue repair and defense against infection.



MUSCULAR

Testosterone affects muscle growth.
Contractions needed for ejaculation depend on intact musculoskeletal system.



SKELETAL

Androgens stimulate bone growth and density.
Pelvis protects internal reproductive structures.

Relationship of Male GU System to Other Systems

Tanner Staging

STAGE	PUBIC HAIR	PENIS	TESTES AND SCROTUM
1: PREADOLESCENT	No pubic hair except for fine body hair similar to that on abdomen.	Same size and proportions as in childhood.	Same size and proportions as in childhood.
2	Sparse growth of long, slightly pigmented, downy hair, straight or only slightly curled, chiefly at base of penis.	Slight or no enlargement.	Testes larger; scrotum larger, somewhat reddened and altered in texture.
3	Darker, coarser, curlier hair spreading sparsely over pubic symphysis.	Larger, especially in length.	Further enlarged.
4	Coarse and curly hair, as in adult; area covered greater than in stage 3 but not as great as in adult.	Further enlarged in length and breadth, with development of glans.	Further enlarged; scrotal skin darkened.
5	Hair same as adult in quantity and quality, spreading to medial surfaces of thighs but not up over abdomen.	Adult in size and shape.	Adult in size and shape.

Some chronic health problems and a variety of other disease processes may have a direct effect on penile function. For example, renal failure and hypothyroidism or hyperthyroidism can alter hormonal release and metabolism at various levels. Erectile dysfunction (impotence) seems to increase with age, but it is not a normal part of aging. The most common cause of impotence in older men is vascular disease. Other causes include diabetes mellitus, hypogonadal states, medications such as antihypertensives and antidepressants, and psychological problems such as depression.

People of Different Cultures or Ethnic Groups

Cultural and ethnic variations often influence the decision regarding circumcision. The United States has always had a high rate of circumcision, and until recently, it was recommended for its medical benefits. However, The American Academy of Pediatrics now states that the medical benefits are not significant enough to endorse circumcision as a routine procedure. This opinion is also held in Canada, England, and Sweden, where circumcision is considered unnecessary. However, circumcision is a religious practice for Jews and Muslims.

Focused Health History

If you are unable to perform a detailed history, ask key questions that focus on a history of male GU problems, presence of symptoms, identification of risk factors, and health promotion activities. The following questions focus on the warning signs of male GU problems:

- Do you have penile pain? Do you have any lesions? If yes, please describe.
- Could you tell me about your sexual practices? Do you practice oral, genital, or anal sex? How many partners have you had? What is your sexual preference: heterosexual, homosexual, or bisexual? Do you practice safe sex? What do you use?
- Do you have any discharge? Type?
- Did you have a history of GU surgery, STDs, or cancer?
- Are you taking any medications? If yes, what are the medications?
- Do you have any medical problems, such as diabetes mellitus, HTN, or peripheral vascular disease?
- Do you have any problems with urination? Burning, frequency, dysuria, hematuria?

Introducing the Case Study



Sam Richards, age 35, works for the telephone company as a lineman. He is complaining about burning on urination for the past 3 days. He requests a "VD test" and appears anxious. He states that he has not slept for 2 nights because he is worried.

comfortable for you and your patient, but the more professional you are, the easier the examination will be for both of you. A nonthreatening, nonjudgmental approach is also essential. Remember to collect data about the urinary system as well as the reproductive system.

Questions about sexuality may be embarrassing for the patient, or he may feel threatened. Sexual performance is often associated with a man's sense of masculinity and youth. If your patient is having sexual function problems, he may be reluctant to share this information with you. Being sensitive and in tune to your patient's nonverbal clues may help you uncover concerns that he is reluctant to volunteer.

Your assessment should focus on detection of any acute health problems and risks for GU problems, as well as identifying any health promotion teaching needs, especially STD prevention. STDs are the most common communicable diseases in the United States, and if left untreated, they can have devastating, permanent effects. If time is an issue and you are unable to perform a complete health history, perform a focused history on the male GU system.

Performing the Male Genitourinary Assessment

Assessment of the male GU system consists of taking a complete health history and performing a physical examination. As you perform your assessment, be alert for signs and symptoms of actual and potential problems involving the various organs and structures in the GU system.

Health History

The purpose of the history is to identify potential or existing problems and risk factors. The history will provide you with subjective data that will guide your physical examination.

Begin by establishing rapport with your patient. Assessing the male GU system may be somewhat un-

Biographical Data

Review the patient's biographical data for clues that relate to the GU system. Begin by looking at his age—it influences the questions you ask during the health history. Because of changes in sexual development, the

Sexually Transmitted Diseases

Diseases and infections transmitted through sexual activity include chlamydia, cytomegalovirus, hepatitis B, herpes simplex, HIV/AIDS, HPV, molluscum contagiosum virus, gonorrhea, and syphilis. They may cause local or systemic effects and be long lasting or even fatal. Risk factors include:

- Multiple sex partners or having a partner with multiple partners.
- Intravenous drug use and sharing contaminated needles.
- Unprotected sex.
- Contacts with prostitutes, especially without using condoms.
- Frequent, multiple blood transfusions.

focus of the adolescent history will be slightly different than that of the older adult history. The age of your patient may also identify risk factors. For example, testicular cancer, epididymitis, and testicular torsion most frequently affect young men, whereas benign prostatic hypertrophy (BPH) and prostate cancer are more prevalent in older men.

Your patient's occupation may put him at risk for GU problems if he is exposed to chemicals or radiation that may cause infertility or increase the risk for testicular cancer. Even religion has an impact on the GU system because different cultures or ethnic groups have different circumcision practices.

Current Health Status

The current health status focuses on the patient's chief complaint. Common complaints include pain, lesions, swelling, discharge, urinary problems, and erectile dysfunction. Explore these complaints using the PQRST format. Begin by asking questions about GU symptoms; leave questions about sexual performance until the end.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Genital Pain

Pain in the genital area usually has an acute onset and often follows trauma or infection. STDs and UTIs can cause epididymitis. Testicular torsion, a surgical emergency,

causes excruciating pain and is not accompanied by urinary symptoms, fever, or elevated white blood cell (WBC) count. Prostatitis can cause perineal pain and may be bacterial or nonbacterial in origin.

Genital Lesions

Genital lesions such as vesicles, ulcers, warts, or rashes may indicate a variety of problems. The morphological description of the lesion will help with identification. Whether or not the lesion is painful is also important in differentiating lesions.

Lesions are often symptoms of STDs, but they may also be caused by carcinoma or a systemic problem.

Genital Swelling

Genital swelling or edema may result from local trauma or infection, but it may also be associated with systemic problems such as congestive heart failure (CHF). Swelling in the inguinal area may signal the presence of a hernia.

Penile Discharge

The only normal drainage from the penis is seminal fluid and urine. Any other discharge may signal a problem. The patient's description of the discharge may give clues to the underlying problem. Copious amounts of thick, yellow discharge are associated with gonorrhea. Yellow discharge is also seen with chlamydia. Thin, watery discharge is associated with prostatitis. Bloody discharge may occur with trauma, infection, or cancer. Discharge of infectious origin is usually accompanied by pain.

Urinary Problems

Urinary problems are often associated with male reproductive problems because the penis serves as both a reproductive and a urinary structure. Urinary problems often cause discomfort, prompting the patient to seek treatment. Dysuria, burning, frequency, change in urinary patterns, and hematuria are common complaints.

Because the two systems work so closely together, a problem in one system often eventually affects the other. However, reproductive problems more often cause urinary problems than vice versa.

Past Health History

When taking your patient's past health history, think about what information is related to male genitalia pathology. You may be able to identify a direct link between the past health history and your patient's current health status. Remember to save questions about sexual performance until the end.

Past Health History

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Childhood Illnesses</p> <ul style="list-style-type: none"> ■ Did you ever have mumps (parotitis), undescended testicles, or an inguinal hernia as a child? <p>Hospitalizations/Surgeries</p> <ul style="list-style-type: none"> ■ Have you ever had a prostatectomy, transurethral prostatectomy, circumcision, orchiectomy, vasectomy, lesion or nodule removal, hernia repair, epispadias or hypospadias repair? <p>Diagnostic Procedures</p> <ul style="list-style-type: none"> ■ Have you ever had a PSA test? ■ Have you ever had a rectal exam? When? <p>Injuries</p> <ul style="list-style-type: none"> ■ Have you ever had trauma to the genital area? <p>Serious/Chronic Illnesses</p> <ul style="list-style-type: none"> ■ Do you have a history of STD, prostatitis, UTI, kidney stones, genital area trauma, cancer, congenital or acquired genital deformity, or infertility? ■ Do you have a history of diabetes or vascular disease? <p>Immunizations</p> <ul style="list-style-type: none"> ■ Were you immunized for the mumps? <p>Allergies</p> <ul style="list-style-type: none"> ■ Do you have contact dermatitis? ■ Are you allergic to condoms, nonoxynol 9 or other spermicides, or latex? <p>Medications</p> <ul style="list-style-type: none"> ■ Are you on any medications, such as antibiotics, hormone replacements, alpha blockers, 5-alpha-reductase inhibitors or antidepressants? <p>Recent Travel/Military Service/Exposure to Infectious or Contagious Diseases</p> <ul style="list-style-type: none"> ■ Have you been exposed to chemicals? ■ Do you have a history of HIV, HPV, molluscum contagiosum, condyloma acuminata, syphilis, penile lesion, chlamydia, or gonorrhea? 	<ul style="list-style-type: none"> ■ <i>Mumps</i>: May cause infertility as a result of orchitis. ■ <i>Undescended testicles (cryptorchidism)</i>: Risk factor for testicular cancer. ■ <i>Inguinal hernia as a child</i>: Risk factor for testicular cancer. ■ May explain patient complaints or physical findings. ■ PSA is a screening test for prostate cancer. ■ Rectal exams for prostate evaluation should be done yearly for men over age 50. ■ May explain physical findings. ■ History of STDs indicates a lifestyle that predisposes patient to subsequent infections. ■ Other problems may explain physical problems. ■ May also be used as baseline information. ■ Can cause GU problems such as erectile dysfunction. ■ Mumps during adolescence may cause sterility. ■ Further identifies health promotion needs. ■ Certain medications can contribute to sexual dysfunction. See <i>Drugs That Adversely Affect Male Sexual Function</i>. ■ Increased risk for infertility. Could present an educational opportunity for disease prevention.

Drugs that Adversely Affect Male Sexual Function

CLASSIFICATION	MEDICATION	EFFECT
Antiandrogen	Finasteride	Impotence
Antianxiety/sedative	Benzodiazepines, chlordiazepoxide	Changes in libido
Anticholinergic	Atropine	Impotence

(continued)

Drugs that Adversely Affect Male Sexual Function (continued)

CLASSIFICATION	MEDICATION	EFFECT
Antidepressants	Tricyclic antidepressants	Increased/decreased libido and impotence
	Trazodone	Decreased libido, impotence, priapism, retrograde ejaculation
	Selective serotonin reuptake inhibitors	Decreased libido, delayed orgasm, anorgasmia, ejaculatory dysfunction
Antiepileptic	Primidone	Impotence
Antihypertensive	Methyldopa	Ejaculatory failure
	Prazosin	Impotence, priapism
	Clonidine	Impotence, decreased sexual activity, decreased libido
Antipsychotics	All beta blockers	Impotence, decreased libido
	All antipsychotics	Priapism, impotence, ejaculatory inhibition
Diuretics	Chlorothiazide, spironolactone	Impotence
Estrogen	Conjugated estrogen	Impotence, testicular atrophy
Tranquilizers	Diazepam, alprazolam	Changes in libido

Family History

The family history can identify factors that might cause a health problem or predispose the patient to one. It identi-

fies genetically linked disorders as well as familial risk factors that put your patient at risk for genitourinary disease.

Family History

RISK FACTORS/QUESTIONS TO ASK

Genetically Linked Diseases of Male Genitalia

- Do you have a family history of varicocele, testicular cancer, or infertility?

- Did your mother take DES during her pregnancy?

Familial/Genetically Linked Disorders with Genitourinary Symptoms

- Do you have a family history of diabetes or vascular disease?

RATIONALE/SIGNIFICANCE

- Increased risk for cancer and infertility.

- May indicate congenital deformities.

- Increased risk for testicular cancer.

- Can affect sexual function.

Review of Systems

The review of systems (ROS) allows you to assess the interrelationship of the GU system to every other system. You can determine how the GU system affects or

is affected by every other system. During the ROS, you often uncover important facts that your patients failed to mention earlier.

Review of Systems

SYSTEM/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- Have you had any changes in your energy level?
- Have you had any fevers?
- Have you experienced any weight change?

- Fatigue, activity intolerance, fever, and weight loss are often signs of underlying health problems, such as HIV/AIDS or prostatitis.
- Scrotal abnormalities may cause discomfort and affect activity tolerance.
- Fatigue influences desire and ability to perform sexual intercourse.

Integumentary

Skin

- Have you had any rashes or lesions?

- May be associated with STDs, but not restricted to genitalia. Primary syphilis chancre found on lip as well as penis. Bilateral, symmetrical rash of secondary syphilis usually occurs on palms and soles.

Hair

- Have you had any hair loss?

- Male hormones affect body hair growth.
- Secondary syphilis can cause alopecia.

HEENT

Head and Neck

- Do you have swollen glands or nodes?

- *Swelling of lymph nodes:* Associated with infection and malignancy.
- *Enlarged thyroid:* May affect sexual function.

Eyes, Nose, and Throat

- Do you have a sore throat?
- Have you had watery eyes? Drainage from your eyes?

- STDs can be found in the oropharynx and are often not assessed in patient complaining of sore throat. Secondary syphilis can result in oral mucous patches.
- Gonorrheal eye infections occur in newborns (ophthalmia neonatorum) and adults (gonorrheal conjunctivitis).
- *Tearing, photophobia, edema of eyelids, and conjunctival edema:* *Chlamydia trachomatis* infection.

Respiratory

- Do you have any breathing problems?

- Respiratory disease such as COPD may affect sexual activity.

Cardiovascular

- Do you have cardiovascular disease, such as HTN or CHF?

- May compromise sexual activity.
- Vascular problems and HTN medications may cause erectile dysfunction.
- Scrotal edema may accompany CHF.

Gastrointestinal

- Do you have a history of liver disease?
- Have you had a loss of appetite or weight loss?

- Liver disease associated with ascites may cause scrotal edema.
- Weight/appetite loss may be associated with malignancy or AIDS.

Musculoskeletal

- Do you have weakness, limitation, joint pain, or swelling?

- Physical act of sexual intercourse requires functional musculoskeletal system. Late stage syphilis damages joints (Charcot's joint).

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Neurological <ul style="list-style-type: none"> Do you have any neurological problems? Do you have any weakness, paralysis, or tremors? Mental or personality changes or depression? 	<ul style="list-style-type: none"> May affect sexual function. Neurosyphilis occurs in late syphilis and can cause paresis, tremors, personality changes, and psychosis. Depression is a risk factor for erectile dysfunction.
Endocrine <ul style="list-style-type: none"> Do you have diabetes or thyroid disease? 	<ul style="list-style-type: none"> Diabetic neuropathy can cause impotence. Hypothyroidism causes decreased libido and infertility. Hyperthyroidism can cause impotence and gynecomastia.
Immunologic/Hematologic <ul style="list-style-type: none"> Do you have HIV or AIDS? Do you have sickle cell anemia? 	<ul style="list-style-type: none"> AIDS and STDs affect the immune system. Sickle cell anemia can cause priapism (painful, sustained erection without sexual desire).

Psychosocial Profile

The psychosocial profile can reveal patterns in the patient's life that may affect his reproductive health and also assist in determining the cause of a health concern

such as erectile dysfunction. Evaluating lifestyle may also identify specific areas in which health maintenance education is needed.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Health Practices and Beliefs/Self-Care Activities <ul style="list-style-type: none"> Do you do TSEs? Have you had your prostate checked? Do you use condoms and practice "safe sex"? How often do you bathe? 	<ul style="list-style-type: none"> Monthly TSE is recommended as well as an annual exam. Annual prostate exam starting at age 50. Use of condoms minimizes the risk for STDs. Poor hygiene can lead to infections of genitalia or urinary tract.
Typical Day <ul style="list-style-type: none"> What is your typical day like? 	<ul style="list-style-type: none"> Performing activities of daily living (ADLs) indicates level of wellness. Fatigue associated with AIDS.
Nutritional Patterns <ul style="list-style-type: none"> What did you eat in the last 24 hours? What foods do you usually eat? 	<ul style="list-style-type: none"> <i>High-saturated-fat, high-cholesterol diet:</i> Associated with arteriosclerosis, which indirectly contributes to erectile dysfunction. <i>Weight loss:</i> Associated with cancer, AIDS, and anorexia.
Activity and Exercise Patterns <ul style="list-style-type: none"> Do you exercise? How often? What type of exercise? Do you participate in sports or activities that require heavy lifting or that may cause genital injury? Do you wear protective equipment while participating in sports? 	<ul style="list-style-type: none"> May indicate reproductive health and self-esteem. Trauma to testicle may cause a hydrocele. Heavy lifting can result in hernias. Use of protective equipment, such as a jockstrap or cup, decreases risk of genital trauma.
Recreation/Hobbies <ul style="list-style-type: none"> What do you do in your leisure time? 	<ul style="list-style-type: none"> Some hobbies can expose people to carcinogens.

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Sleep/Rest Patterns</p> <ul style="list-style-type: none"> ■ How many hours of sleep do you average a night? ■ Do you wake up to urinate? How often? <p>Personal Habits</p> <p><i>Drug Use</i></p> <ul style="list-style-type: none"> ■ Do you use street drugs, such as cocaine, barbiturates, amphetamines, or narcotics? If so, how often and how much? <p><i>Alcohol Use</i></p> <ul style="list-style-type: none"> ■ Do you drink alcohol? What type, how much, and how often? <p><i>Tobacco Use</i></p> <ul style="list-style-type: none"> ■ Do you use tobacco? What type, how much per day, and for how many years? <p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ Does it expose you to radiation or toxic chemicals? ■ Does your work involve the use of heavy equipment? <p>Environmental Health Patterns</p> <ul style="list-style-type: none"> ■ Do you live in an apartment or a single home? ■ What type of heating do you have? ■ Is the plumbing adequate? <p>Roles/Relationships/Self Concept</p> <ul style="list-style-type: none"> ■ How do you see yourself? ■ Would you say you have a positive self-image? <p>Cultural/Religious Influences</p> <ul style="list-style-type: none"> ■ Do you have any cultural or religious beliefs that affect your sexual health? <p>Family Roles and Relationships</p> <ul style="list-style-type: none"> ■ Have GU problems affected your marriage or family life? <p>Social Supports/Stress and Coping</p> <ul style="list-style-type: none"> ■ Who are your supports? ■ How do you deal with stress? 	<ul style="list-style-type: none"> ■ Fatigue can cause sexual dysfunction. ■ <i>Nocturia</i>: May be underlying symptom of urethritis. ■ May impair judgment, increasing risk of STD exposure. ■ Street drugs may cause priapism, impotence, increased sexual excitability, increased libido, and delayed orgasm. ■ Impairs gonadotropin release and accelerates testosterone metabolism, causing impotence and loss of libido. Large doses can acutely depress sexual reflexes. ■ Chronic alcoholism causes high levels of circulating estrogens, which decrease libido. ■ Alcohol intoxication may impair judgment, decreasing incidence of safe sex practices and increasing risk of exposure to STDs. ■ Cigarette smoking increases risk of atherosclerotic disease, which may decrease penile blood flow. ■ <i>Radiation/toxic chemicals</i>: Linked to cancer of male genitalia and infertility. ■ <i>Work with heavy machinery</i>: Increases risk for hernias and injury to genitalia. ■ Poor sanitation may indicate poor hygiene, which can cause infections of genitalia or urinary tract. ■ Poor self-image and lack of self-confidence can predispose patient to sexual dysfunction. ■ Circumcision is a religious ceremony for Jewish faith. ■ Abortion and contraception are prohibited for members of Catholic faith. ■ Premature ejaculation, impotence, and other sexual problems may affect relationships. ■ Stress within a relationship may cause sexual dysfunction. ■ Stress can cause sexual dysfunction. ■ Emotional or social isolation can increase stress.

Sexual History

Because the GU system encompasses sexuality, a more detailed sexual history may be warranted. Wait until the end of the history before asking sensitive or embarrassing questions. This gives you time to establish rapport

with your patient. Asking about sexual functions and practices is important for several reasons:

- Many patients have sex-related questions or problems that they want to discuss.

- Sexual practices may be directly related to the patient's symptoms.
- Sexual dysfunction may be a consequence of medications and may be reversible.
- Sexual practices are related to the patient's risk for STDs.

Questions that are helpful in assessing sexual function include the following:

- Have you ever been sexually active?
- Are you currently sexually active? That is, have you had sex with anyone in the past few months? How often? If yes, do you have sex with men, women, or both (heterosexual, homosexual, or bisexual)?
- What types of sexual activity do you engage in? Oral, anal, genital?
- Do you have more than one partner? How many partners have you had in the last 6 months?
- Do you use birth control? What kind? How often?
- Are you worried about the acquired immunodeficiency syndrome (AIDS) virus or other STDs?
- Do you take any precautions to avoid infections? If so, what?
- Do you have any problems or concerns about your sexual function?
- Have you had surgery on any of your reproductive organs? If so, what and when?
- Have you been taught to examine your testicles?

Erectile Dysfunction

Erectile dysfunction is the inability to achieve satisfactory sexual performance. It includes the inability to achieve erection, achieve complete erection, maintain erection, or ejaculate. A specific description of the dysfunction is essential to identify the most appropriate therapy. Remember: Take a look at your patient's medical history because diabetes, vascular problems, and certain medications are risk factors for erectile dysfunction. Additional questions that address erectile dysfunction include:

- Are you still interested in sex?
- Are you able to achieve and maintain an erection?
- Do you have morning erections?
- Were there any changes in your relationship with your partner or in your life situation when the problem began?
- How long does intercourse last?
- Do you sometimes feel that you cannot ejaculate?
- Are you satisfied with your sex life as it is now?



HELPFUL HINT

If your patient can achieve a morning erection, the cause of erectile dysfunction is usually not organic.



Case Study Findings

Reviewing Mr. Richards'

Biographical data:

- 35-year-old white male.
- American by birth; Protestant religion.
- Recently divorced, has a 10-year-old daughter.
- Employed as telephone lineman for 10 years; healthcare insurance through work.
- High school education.
- Source of referral: self, seems reliable.

Current health status:

- Onset of dysuria 3 days ago.
- Clear urethral discharge occasionally at end of urine stream.
- Denies low back pain, blood in urine, or recent trauma.
- Denies suprapubic discomfort.
- Had two different sex partners within the last 6 months.
- Uses condoms "sometimes."

Past health history:

- Usual childhood illnesses except for measles, mumps, and rubella.
- Tonsillectomy in 1967, no other surgeries or hospitalizations.
- Had chlamydia approximately 1 year ago and received antibiotics.
- No history of renal calculi.
- Immunizations current.
- Denies taking any prescription drugs; admits using aspirin occasionally for aches and pains.

Family history:

- Father died at age 52 of heart attack.
- Mother, age 62, alive and well.
- Three siblings are healthy.

Review of systems:

- *General Health Survey*: Usual state of health good.
- *Integumentary*: No changes in skin, hair, and nails; no lesions.
- *HEENT*: No swollen glands/nodes; denies sore throat; no eye drainage.
- *Respiratory*: No history of chronic obstructive pulmonary disease (COPD).
- *Cardiovascular*: No history of hypertension (HTN) or cardiovascular disease.
- *Gastrointestinal*: No history of liver disease.
- *Musculoskeletal*: Denies joint pain or swelling.
- *Neurological*: No neurological problems, weakness, or paralysis; denies personality changes or depression.
- *Endocrine*: No history of thyroid disease or diabetes.

- *Immunologic/Hematologic:* Has never been tested for human immunodeficiency virus (HIV); no history of sickle cell anemia.

Psychosocial profile:

- Cannot remember when he last had a physical or went to a doctor for illness.
- Does not perform testicular self-examination (TSE).
- *Typical day:* Awakens at 6:30 A.M., has only coffee for breakfast, gets to work at 7:30 A.M. Work is very physical and includes climbing up telephone poles and onto roofs of houses and crawling under houses. Usually arrives home by 5:30 P.M. Spends evenings watching television or dating. Spends time with child every other weekend. Usually in bed about 10 P.M. on workdays. Skips breakfast, drinks 4 cups of coffee daily, eats out a lot, snacks on junk food, rarely has a nutritious meal.
- Besides work, hikes about every other weekend, plays tennis occasionally, does yard work.
- Sleeps about 7 hours a night on workdays, less on nights he has dates. Naps on weekends to “catch up” on sleep.
- Has smoked a pack of cigarettes daily for the last 15 years. Drinks beer about three times per week and an additional 6 to 10 beers on weekends. Denies recreational drug use.
- Job is physically taxing and potentially dangerous because of climbing.
- Has good relationship with his one child and with his mother, who cleans, irons, and occasionally cooks for him.
- No community involvement, does not attend church. Lives alone since divorce 1 year ago, in single-family home with more space than he needs.
- Copes with stress by ignoring it.

Sexual history:

- Currently sexually active.
- Heterosexual.
- Two sexual partners within last 6 months.
- Inconsistent use of condoms.
- Satisfied with sexual performance.
- No problems achieving erection or ejaculation.
- Does not perform TSE.



Case Study Evaluation

Before you proceed with the physical examination of the male GU system, review and document the data you have obtained from Mr. Richards' history.



CRITICAL THINKING ACTIVITY #1

Based on your history findings, what teaching needs can you identify for Mr. Richards?

Physical Assessment

Once you have obtained the subjective data, focus on collecting the objective data by performing the physical exam.

Approach

Ensure that the exam room is warm and comfortable and that the light in the room is sufficient to adequately observe the patient. Instruct him to empty his bladder and to undress to expose the groin area. Drape him appropriately for privacy. Because reproductive system assessment involves exposing and handling the genitals, the patient may feel anxious and embarrassed. To help minimize discomfort, explain each assessment step before performing it and expose only the necessary areas. Keeping a professional demeanor will help you feel more at ease, too.

During the examination, the patient can lie on the table in the supine position with his legs spread slightly or stand in front of you while you are sitting eye level to the genitalia. A rectal examination is used to assess the prostate gland. Your patient can either assume a side-lying (Sims') position or lean over the examination table.

You will use the techniques of inspection, palpation, and auscultation to assess this patient.



TOOLBOX

Assemble the following equipment for a routine exam: Nonsterile gloves, water-soluble lubricant, penlight, and stethoscope. To obtain a urethral specimen and perform a culture, you will also need cotton swabs or 1½" to 2" gauze wrap, urogenital alginate swabs, and a Thayer-Martin plate (gonorrhea culture).

Performing a General Survey

Before you begin your physical exam, perform a general survey, looking for clues that suggest a GU problem. For example:

- Obtain the patient's vital signs. Elevations in temperature may indicate an infection that might be accompanied by a GU infection. HTN may affect sexual functioning.
- Because the examination can produce anxiety, take note of your patient's emotional status.
- Take note of your patient's posture and nonverbal behavior. Does his posturing reflect pain or discomfort?

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> Changes in energy level, weight Fevers	General	<i>Take:</i> Vital signs, checking for temperature elevations, HTN <i>Note:</i> Signs of discomfort Affect
Integumentary	<i>Ask about:</i> Changes in hair growth Rashes, lesions	Integumentary	<i>Inspect:</i> Skin lesions Hair distribution Areas of alopecia
HEENT <i>Head, Neck, Eyes, Mouth and Throat</i>	<i>Ask about:</i> Swollen glands/nodes Eye drainage Sore throat Oral lesions	HEENT <i>Head, Neck, Eyes, Mouth and Throat</i>	<i>Palpate:</i> Lymph node enlargement Thyroid enlargement <i>Inspect:</i> Conjunctiva for drainage Oral mucosa for redness and lesions
Respiratory	<i>Ask about:</i> History of respiratory disease	Respiratory	<i>Inspect:</i> Signs of respiratory distress <i>Auscultate:</i> Abnormal breath sounds
Cardiovascular	<i>Ask about:</i> History of CVD, HTN, CHF	Cardiovascular	<i>Inspect:</i> Signs of impaired circulation Skin changes <i>Palpate:</i> Pulses, edema <i>Auscultate:</i> Extra heart sounds
Gastrointestinal	<i>Ask about:</i> History of liver disease Loss of appetite	Gastrointestinal	<i>Inspect:</i> Ascites <i>Palpate/Percuss:</i> Liver enlargement
Musculoskeletal	<i>Ask about:</i> Weakness, limitations Joint pain, swelling	Breasts	<i>Inspect:</i> Gynecomastia
Neurological	<i>Ask about:</i> History of neurological problems Paralysis Tremors Personality changes Depression	Musculoskeletal	<i>Test:</i> Muscle strength <i>Inspect:</i> Joint swelling and deformity
Endocrine	<i>Ask about:</i> History of thyroid disease and diabetes	Neurological	<i>Test for:</i> Changes in mental status, affect Sensory deficits
Lymphatic/ Hematologic	<i>Ask about:</i> History of HIV, AIDS, sickle cell anemia		

Performing a Head-to-Toe Physical Assessment

The GU system affects many different systems. Therefore,

examine the patient for specific changes that may affect the GU system.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

General Health Survey

- Well developed physically.
- No acute distress.
- Awake, alert, oriented. Memory intact.
- Affect appropriate.

Integumentary

Skin

- Intact, no lesions.

Hair

- Even distribution of hair.

HEENT

Head and Neck

- No palpable lymph nodes.
- Thyroid not palpable.

Eyes, Mouth, and Throat

- Conjunctiva pink, no drainage.
- Oral mucous membranes pink, moist and intact.

Respiratory

- Lungs clear; no SOB.

Cardiovascular

- Regular heart rate and rhythm, no extra sounds.
- +2 pulses, extremities warm bilaterally, no edema.

Breasts

- No enlargement

Gastrointestinal

- Abdomen flat, positive bowel sounds, no organomegaly.
- No palpable lymph nodes.

Musculoskeletal

- >+5 muscle strength, full range of motion.

- No joint deformities.

Neurological

- No weakness or paralysis.
- Awake, alert, and oriented $\times 3$, memory intact, affect appropriate.

ABNORMAL FINDINGS/RATIONALE

- Personality changes and mental deterioration can accompany late-stage syphilis.
- Depression can affect sexual function.

- Rashes and skin lesions are associated with many STDs.
- *Systemic rash*: Secondary syphilis.
- *Vesicles*: Herpes simplex types 1 and 2.
- *Alopecia*: Associated with secondary syphilis.
- *Decrease in body hair*: Possible decrease in male hormones.

- *Palpable lymph nodes*: May indicate systemic infection or malignancy.
- *Enlarged thyroid (hypo- or hyperthyroidism)*: May affect sexual functioning.

- *Conjunctivitis*: Can be caused by gonorrhea.
- *Oral lesions*: Associated with STDs.

- Chronic lung disease may impair sexual functioning.

- HTN and CHF may impair sexual function.
- CHF may cause scrotal edema.
- Vascular disease is risk factor for erectile dysfunction.

- *Gynecomastia*: Associated with decreased male hormones.

- Liver disease associated with ascites may also cause scrotal edema.
- *Palpable inguinal lymph nodes*: Possible infection or malignancy.

- Weakness or joint pain may limit sexual functioning.
- *Weakness and paralysis*: Associated with neurosyphilis.
- *Charcot joints*: Associated with late syphilis.

- *Weakness and paralysis*: Associated with neurosyphilis.
- *Changes in mental status/psychosis*: Late syphilis.
- *Depression*: Risk factor for psychogenic erectile dysfunction.

Performing a Genitourinary Physical Assessment

Assessment of the male GU system includes inspection, palpation, and auscultation.

Inspection

Begin with inspection of the genitalia. The patient should be lying on the table in a supine position with his legs slightly spread. Or he can stand in front of you and you can sit so that the genital area is at your eye level. Remember to put on gloves. After inspecting the genitalia,

inspect the rectal area. Have the patient move to a side-lying position or stand and bend over the examination table.



HELPFUL HINT

Change gloves as needed during the exam to prevent cross-contamination.

INSPECTING THE GENITALIA

1. Hair Distribution

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note hair distribution pattern.
- *Pubic hair distribution triangular, sparsely distributed on scrotum and inner thigh, and absent on penis. Genital hair coarse. No nits or lice.*



Inspecting hair distribution

ABNORMAL FINDINGS/RATIONALE

- *Sparse or absent hair (alopecia):* May result from genetic factors (developmental defects and hereditary disorders), aging, or local or systemic disease. Diseases include infection, neoplasms, endocrine diseases, and nutritional or metabolic deficiencies. Other causes are physical or chemical agents and destruction or damage to hair follicles.

INSPECTING THE GENITALIA

2. Penis

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note color, lesions, swelling, discharge, or pubic pediculosis on dorsal, lateral, and ventral sides.
- If uncircumcised, retract foreskin. Note ease of retraction of foreskin and inspect glans.
- *No lesions or inflammation. Shaft skin loose and wrinkled without erection. Glans smooth and free of lesions, swelling, or inflammation. No penile discharge. Dorsal vein sometimes visible. If patient is uncircumcised, loose skin on the penis shaft folds to cover glans, forming the foreskin. Foreskin retracts easily.*

ABNORMAL FINDINGS/RATIONALE

- **Chancre:** Signals primary syphilis. Painless, ulcerated, exudative, papular lesion with an erythematous halo, surrounding edema, and a friable base.
- **Chancroid:** Caused by *Haemophilus* through small breaks in epidermal tissue. Pinhead papules to cauliflower-like groupings of painful, filiform, skin-colored, pink, or red lesions.



Chancre



Chancroid

- **Condyloma acuminatum (genital warts):** Caused by HPV infection. Multifocal, wartlike, maculopapular lesions that are tan, brown, pink, violet, or white.



Genital warts on the penis



Genital warts on the scrotum

- **Candida:** Superficial mycotic infection of moist cutaneous sites. Erythematous plaques with scaling, papular lesions with sharp margins and occasionally clear centers and pustules.
- **Herpetic lesion:** Herpes simplex virus 1 and 2 cause painful eruptions of pustules and vesicles that rupture. Fever, headache, dysuria, dyspareunia, and urinary retention may occur.



Genital herpes

(continued)

INSPECTING THE GENITALIA

2. Penis (continued)

- *Tinea cruris*: Fungal infection of the groin often referred to as *jock itch*.



Tinea cruris

- *Phimosis*: Occurs in uncircumcised males. Foreskin is unable to retract and may become swollen.
- *Priapism*: Often associated with leukemia, metastatic carcinoma, or sickling hemoglobinopathies.
- *Chordee*: Ventral or dorsal curvature of penis. Ventral chordee seen mostly with epispadias.

3. Scrotum

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note color, hair distribution, lesions, size, shape, and position.
- *Scrotal skin rugated, thin, and more deeply pigmented than body color. Skin firmer in young men and elongates with flaccidity in older men.*
- *Skin free of lesions, nodules, swelling, inflammation, and erythema. Scrotal size and shape vary greatly. Left scrotal sac lower than right.*



Inspecting the scrotum

ABNORMAL FINDINGS/RATIONALE

- *Masses*: Can arise from benign or malignant conditions.
- *Scrotal swelling*: Seen with inguinal hernia, hydrocele, varicocele, spermatocele, tumor, and edema.
- *Hydrocele*: Nontender accumulation of fluid between two layers of tunica vaginalis in scrotum. May be idiopathic or a result of trauma, inguinal surgery, epididymitis, or tumor. Mass transilluminates.
- *Spermatocele*: Nontender, well-defined cystic mass on superior testis or epididymis caused by blockage of efferent ductules of rete testis.
- *Varicocele*: Varicose veins of spermatic cord that feel like a “bag of worms” and slowly collapse when scrotum is elevated. Caused by dilated veins in pampiniform plexus of spermatic cord. Right-sided may indicate obstruction at vena cava.
- *Sebaceous cyst*: Round, firm, nontender cutaneous cyst confined to scrotal skin. May result from decrease in localized circulation and closure of sebaceous glands or ducts.

INSPECTING THE GENITALIA

4. Urethral Meatus

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note position of urinary meatus.
- *Located centrally on glans. Pink in white males and darker pink in darker-skinned males. No discharge.*


ABNORMAL FINDINGS/RATIONALE

- *Epispadias:* Urethral meatus opens on dorsal (upper) side of penis.
- *Hypospadias:* Urethral meatus opens on ventral (under) side of penis.

5. Inguinal Area

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note condition of skin, enlarged lymph nodes.
- Have patient bear down and note any bulges.
- *No swelling or bulges. Lymph nodes nonpalpable.*



ABNORMAL FINDINGS/RATIONALE

- *Bulge:* May indicate a hernia or enlarged lymph node.

6. Rectal Area

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note skin condition, lesions, rectal prolapse, hemorrhoids, fissures, bleeding, or discharge.
- *Skin intact with slightly darker pigmentation around anus. Anus intact; no lesions, hemorrhoids, fissures, bleeding, or rectal prolapse.*

ABNORMAL FINDINGS/RATIONALE

- STD lesions, warts, hemorrhoids, fissures, bleeding, rectal prolapse.

Palpation

Next, palpate the penis and scrotum for tenderness and abnormalities. Palpate the inguinal and femoral areas for hernias and enlarged lymph nodes. And palpate the rec-

tum for masses or nodules, and note the tone of the anal sphincter. Also check the prostate and bulbourethral glands for tenderness, masses, or swelling. Remember to wear gloves.

PALPATING THE GENITALIA**1. Bladder and Kidneys (See Chapter 15: Assessing the Abdomen)**

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Bladder and kidneys not palpable.*

**HELPFUL HINT**

Maintain standard precautions and wear gloves whenever palpating any part of the male GU system.

2. Penis

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Lightly palpate shaft with thumb and index finger.
- Note consistency, tenderness, masses, or nodules.
- *Pulsations on dorsal sides. Nontender. No masses or firm plaques.*



Palpating the penis

ABNORMAL FINDINGS/RATIONALE

- *Diminished/absent palpable pulse:* Possible vascular insufficiency.
- Normal blood flow may be affected by systemic disease, localized trauma, or localized disease.
- Priapism is associated with spinal cord lesions or sickle cell anemia.
- Penis should not be enlarged in nonerect state.
- Phimosis and paraphimosis (foreskin retracts but does not return) are abnormal.

**ALERT**

Seek immediate assistance if foreskin cannot be retracted. Prolonged constriction of vessels can decrease blood flow and lead to necrosis.

3. Urethral Meatus

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Note discharge.
- *No discharge and drainage.*

PALPATING THE GENITALIA

3. Urethral Meatus (continued)

ABNORMAL FINDINGS/RATIONALE

- *Purulent discharge or mucus shreds*: Bacterial infection of GU tract. Color, consistency, and amount may vary.
- Bacterial infection causes inflammation with leukocytes, shedding tissue cells, and bacteria.

4. Scrotum

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Use thumb and two fingers to gently palpate scrotum, and note size, shape, consistency, mobility, tenderness, masses, or nodules of testes.
- Palpate the epididymis and vas deferens on posterolateral surface and note swelling or nodules.
- *Transillumination*: If a mass, lump, or swelling is present, shine light over area and note transmission of light.



Palpating the scrotum and testicles



Palpating the epididymis



Transilluminating the scrotum



ALERT

Do not squeeze or pinch any mass, lesion, or other structure that is palpated.

- *Scrotum contains a testicle and epididymis on each side.*
- *Testicles firm but not hard, oval, smooth, equal in size, and sensitive to pressure but not tender.*
- *Epididymis comma-shaped, distinguishable from testicle, and insensitive to pressure. Spermatic cord smooth and round.*

ABNORMAL FINDINGS/RATIONALE

- A unilateral mass palpated within or about the testicle is abnormal. Intratesticular masses are nodular and painless and should be considered malignant until proven otherwise. (See section on inspecting the genitalia for description of spermatocele, hydrocele, and varicocele.)
- *Testicle that is enlarged, retracted, in a lateral position, and extremely sensitive*: Testicular torsion.
- Indurated, swollen, tender epididymis is abnormal.
- Undescended testes is abnormal.
- *Cryptorchidism (absence of testes and epididymis in scrotal sac)*: Related to testicular failure, deficient gonadotrophic stimulation, mechanical obstruction, or gubernacular defects. The undescended testes have a histologic change by age 6, so refer patient as early as possible.
- **Orchitis** is acute, painful swelling of testicle and warm scrotal skin. Patient may feel heaviness in scrotum.
- Atrophic testicle and scrotal edema are abnormal.
- *Red glow on transillumination*: Serous fluid within scrotal sac. Occurs in hydrocele and spermatocele.

(continued)

PALPATING THE GENITALIA

5. Inguinal Area

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- **Horizontal lymph nodes:** Palpate inguinal area.
- **Vertical lymph nodes:** Palpate inner aspect of thigh.
- **Hernias:** Place gloved index finger in scrotal sac above testicle and invaginate sac. Follow spermatic cord to Hesselbach's triangle, advance through external inguinal ring, ask patient to bear down, and feel for bulges.



Palpating the horizontal lymph nodes



Palpating the vertical lymph nodes



Palpating hernias



ALERT

Do not have hypertensive patients bear down during exam. Observe for discomfort and light-headedness. Some patients cannot tolerate bearing down for more than a few seconds.



HELPFUL HINTS

- Instruct patient to inform you of any pain or discomfort during palpation. This is a sensitive, ticklish area. Use firm and deliberate movements.
- If you cannot insert your finger into the inguinal canal with gentle pressure, stop. Never force it into the opening.

- *No red glow on transillumination.*
- *Nodes if palpable should be less than 1 cm and freely mobile. No bulges present in inguinal area. No palpable masses in inguinal canal. No portions of the bowel enter the scrotum. No palpable masses at femoral canal.*

PALPATING THE GENITALIA

5. Inguinal Area (continued)

ABNORMAL FINDINGS/RATIONALE

- *Unilateral or bilateral enlargement of inguinal lymph nodes:* May indicate bacterial infection. Nodes may be tender or painless. Lymph node enlargement may also be associated with metastatic disease.
- **Indirect inguinal hernia** comes down inguinal canal and is palpated at inguinal ring or in scrotum.
- **Direct inguinal hernia** is a mass that enlarges with coughing.
- **Femoral hernia** is palpated medial to femoral vessels and inferior to inguinal ligament.



ALERT

A strangulated hernia is a surgical emergency.

6. Rectum/Prostate

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- If indicated, change gloves and apply water-soluble lubricant on index finger.
- Have patient bend over exam table and bear down.
- Gently insert finger into rectum, and note sphincter tone, tenderness, masses, hemorrhoids, or bleeding.
- *Prostate:* Note size, shape, symmetry, consistency, mobility, nodules, tenderness.
- Withdraw fingers; if stool present, test for occult blood.
- *Rectum smooth and nontender with good anal sphincter tone.*
- *Prostate small, smooth, mobile, and nontender.*
- *Medial sulcus palpable.*
- *Test stool for occult blood, negative.*



Palpating the prostate

ABNORMAL FINDINGS/RATIONALE

- *Soft, nontender, enlarged prostate:* BPH. Related to aging and presence of dihydroxy-testosterone.
- *Firm, tender, or fluctuant mass on prostate:* Acute bacterial prostatitis. UTIs commonly occur concurrently. Increased risk for prostatic abscess.
- *Firm, hard, or indurated single or multiple nodules:* Possible prostate cancer.
- *Extremely tender, warm prostate:* May indicate bacterial prostatitis. Do not palpate vigorously. This would be painful and might cause bacteremia.
- Positive occult blood test warrants further evaluation.
- *Palpable mass or nodule:* Polyp or internal hemorrhoid.
- *Hard mass:* Possible rectal cancer.

Obtaining a Urethral Specimen

Gonorrhea and chlamydia are classically associated with a urethral discharge. To obtain a specimen, have the patient either lie supine or stand. Put on gloves and hold the penis with your nondominant hand. Use your dominant hand to clean the urethral meatus with sterile gauze or a cotton swab. Then insert a thin urogenital alginate swab 1 to 2 cm into the urethra and rotate the swab from side to side. Leave it in place for several seconds. A Thayer-Martin medium is used when collecting urethral discharge specimens for gonorrhea. A Gram stain can be done to diagnose gonorrhea, but it should be followed by a culture. A variety of tests for chlamydia are available, but the collection technique remains the same. Some laboratories can perform both tests with one swab, and some will require two specimens. Chlamydia tests can also be done on urine.

Penis

To palpate, stand in front of the patient's genital area. Palpate the entire length of the penis between your thumb and your first two fingers. Note any pulsations, tenderness, masses, or plaques. If your patient is not circumcised, retract the foreskin. As you retract, you may normally find white, cheesy material called **smegma**. Gently squeeze the glans of the penis to expose the meatus between your thumb and your forefinger. If discharge is present or the patient complains of discharge, take a culture.

Scrotum

Next, gently palpate a testicle between your thumb and your first two fingers. Note the size, shape, consistency, and presence of masses. Palpate the spermatic cord from the epididymis to the external ring. Note the consistency and presence of tenderness or masses. Repeat on the opposite side. This is also a good time to teach TSE. If you detect a scrotal mass or enlargement, use transillumination to visualize it. Darken the room, apply a light source to the unaffected side behind the scrotum, and direct it forward. Repeat with the affected side.

Femoral and Inguinal Hernias

Now, palpate the skin overlying the inguinal and femoral areas for lymph nodes, noting size, consistency, tenderness, and mobility. To palpate for inguinal hernias, ask the patient to bear down while you palpate the inguinal area. Place your dominant index finger in the patient's scrotal sac above the testicle and invaginate the scrotal skin. Follow the spermatic cord until you reach a triangular, slitlike opening (Hesselbach's triangle), which is the ex-

ternal inguinal ring. The nail of your examining finger should be facing inward and your finger pad outward. If possible, continue to advance your finger along the inguinal canal and ask the patient to cough or bear down. Note any masses. Repeat the procedure on the opposite side. To assess for femoral hernias, palpate the femoral canal, which is located below the femoral artery, and have the patient bear down or cough (Fig. 19.2).

Rectum, Prostate Gland, and Bulbourethral Gland

To examine the rectum and prostate, have the patient stand and bend over the examination table. Place your lubricated, gloved index finger at the anal orifice and instruct the patient to bear down. Then, gently insert your finger into the anal sphincter, with the tip flexed toward the anterior rectal wall. Palpate the anterior and posterior rectal walls. They should be smooth and nontender, with good anal sphincter tone. Next, palpate the posterior surface of the prostate gland, noting size, shape, consistency, sensitivity, and mobility. Slowly withdraw your finger. If your glove contains fecal matter, perform an occult blood test. If the prostate is enlarged, determine the size and

Teaching Testicular Self-Examination

Teach TSE to your patient during the scrotal examination. Follow these steps:

- First, explain the rationale for the exam. Monthly TSEs permit earlier detection of testicular cancer, which occurs most often in 16- to 35-year-olds. Suggest that he perform the exam on the same day each month.
- Tell him that the scrotum is easier to examine after taking a warm bath or shower.
- Now, show him how to perform the exam: With one hand, lift the penis and check the scrotum for any changes in shape, size, or color. The left side of the scrotum normally hangs slightly lower than the right.
- Next, check the testes for lumps and masses. Locate the epididymis, the crescent-shaped structure at the back of each testis. It should feel soft.
- Use the thumb and first two fingers of your left hand to squeeze the left spermatic cord gently. The cord extends upward from the epididymis, just above the testis. Repeat on the right side using your right hand. Check for lumps and masses.
- Finally, examine each testis. Place your fingers on the underside and your thumb on the top and gently roll the testis between your thumb and your fingers. The testis should be egg-shaped and movable and feel rubbery, smooth, and firm, with no lumps. Both testes should be the same size.
- Tell the patient that if he finds any lumps, masses, or other changes, he should call his doctor.




FIGURE 19.2. Palpating for inguinal and femoral hernias.

grade: Grade I, <1 cm protrusion into rectum; grade II, 1 to 2 cm protrusion into rectum; grade III, 2 to 3 cm protrusion into rectum; grade IV, >3 cm protrusion into rectum.

Next, perform a bidigital examination of the bulbourethral gland by pressing your thumb into the perianal tissue while pressing your index finger toward it. Assess for tenderness, masses, or swelling. Release your thumb and advance your index finger (Fig. 19.3).

Auscultation

If you detect a scrotal mass on inspection and palpation, auscultate the scrotum. Ask the patient to lie in a supine position, then stand at his side next to the genital area. Place your stethoscope over the scrotal mass to listen for bowel sounds. The presence of bowel sounds suggests a hernia. (See *Auscultating the Genitalia*.)




FIGURE 19.3. Palpating the prostate and bulbourethral glands.

AUSCULTATING THE GENITALIA

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- If scrotal mass present, auscultate over mass for bowel sounds.
- No bowel sounds.

ABNORMAL FINDINGS/RATIONALE

- *Bowel sounds in scrotum:* Indirect inguinal hernia.



Case Study Findings

Mr. Richards' general survey and head-to-toe assessment include:

- **General Health Survey**
 - 35-year-old, well-developed white male in no acute distress.
 - Head-to-toe scan revealed no pertinent findings related to the GU system, except positive tenderness in suprapubic area, several palpable inguinal nodes noted and tender.
- **Vital Signs**
 - Temperature, 97.9°F
 - Pulse, 88 beats per minute (BPM), strong and regular.
 - Respirations, 14/min.
 - Blood pressure (BP), 128/76.
 - Height, 5'11".
 - Weight, 182 lb.
- **GU**
 - Triangular distribution of pubic hair, no nits or lice.
 - Erythema at urethral meatus, skin free of lesions and inflammation, rounded glans penis, no penile curvature, circumcised.
 - Pulsations present on dorsal aspect of penis, no masses or firm plaques.
 - Scrotum rugated, no testicular swelling, no erythema, no masses or nodules, left testis slightly lower than right.
 - Testicles palpated bilaterally without masses or tenderness, epididymis not enlarged and nontender, no swelling or tenderness, spermatic cords intact without nodules bilaterally.
 - Urethral meatus centrally located.
 - Clear to white thin discharge expressed from urethral meatus.

- No inguinal bulges, right inguinal lymph nodes slightly elevated (1.5 cm) and tender, no inguinal masses.



Case Study Analysis and Plan

Now that your examination of Mr. Richards is complete, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mr. Richards, then use this information to identify appropriate nursing diagnoses. Here are some possible ones. Cluster the supporting data.

1. Anxiety, related to dysuria
2. Fear, related to infectious process in reproductive system
3. Pain
4. Ineffective Health Maintenance related to insufficient knowledge of safe sex practices
5. Ineffective Health Maintenance related to insufficient knowledge of self-care

Identify any additional nursing diagnoses and any collaborative nursing diagnoses.



CRITICAL THINKING ACTIVITY #2

Develop a nursing care plan and a brief teaching plan for Mr. Richards using one of the nursing diagnoses from above. Include learning outcomes and teaching strategies.

RESEARCH TELLS US

Prostate cancer is the most common form of cancer among men. The use of prostate-specific antigen (PSA) screening has resulted in early detection of prostatic tumors even before they become evident on clinical examination. However, treatment of early stage prostate cancer is controversial, thus questioning the benefits of PSA screening for prostate cancer.

A study of 977 patients with nonpalpable prostate cancer from 1991 through 2002 were classified according to pretreatment PSA, Gleason scores (most commonly used prostate cancer grading system), percent positive biopsy, and age to determine the aggressiveness of the disease. Of the 977 patients, 601 (61.5%) were found to have significant tumor (Smith et al., 2004). Considering that the number of patients with prostate cancer diagnosed by PSA had significant disease, these findings support the continued use of the PSA screening.

Another study of prostate screening addresses the racial disparity of prostate screening in the elderly. African American men have a 60 percent greater incidence of prostate cancer than do white men with twice the mortality rate as white men. The low screening rates among African American men may account for the higher mortality rate. A case-control study of racial differences in the use of PSA screening among beneficiaries in New Jersey revealed that elderly African Americans were less likely to have PSA screening than were elderly whites (Gilligan et al., 2004). These findings reinforce the racial disparity in healthcare services in the United States.

It's important to remember that PSA screening has proven effective in early detection and treatment of prostate cancer. As health promoters, nurses are in a key position to educate the public and increase public awareness of prostate cancer and preventative screening.

Source: Smith, R., Malkoicz, S., Whittington, R., Van Arsdalen, K., Tochner, Z., and Wein, A. (2004). Identification of clinically significant prostate cancer by prostate-specific antigen screening. *Archives of Internal Medicine* 164(11) 1227–1230; and Gilligan, T., Wang, P., Levin, R., Kantoff, P., and Avorn, J. (2004). Racial differences in screening for prostate cancer in the elderly. *Archives of Internal Medicine* 164(17) 1858–1864.

HEALTH CONCERNS

Risk Factors for Testicular Cancer

- Cryptorchidism
- Age 20–40
- Family history of testicular cancer
- Mumps orchitis
- Inguinal hernia during childhood
- Maternal exposure to diethylstilbestrol (DES)
- Testicular cancer in other testicle

Signs of Testicular Cancer

- Hard, fixed, nontender mass or nodules on testicle that are not visible with transillumination
- Scrotal swelling
- Scrotal heaviness

Risk Factors for Prostate Cancer

- Family history of prostatic cancer
- Advanced age
- African American ethnicity

Signs of Prostate Cancer

- Urinary symptoms such as dysuria, frequency, nocturia, hesitancy, dribbling, hematuria, and retention
- Enlarged, hard, fixed prostate
- Back pain (may indicate metastasis)

Screening Tests

In addition to monthly TSE, the American Cancer Society recommends an annual digital rectal exam (DRE) for all men over age 50. Serum PSA is a glycoprotein exclusive to the prostate epithelium that is elevated in prostate cancer because of the accelerated metabolic rate and catabolism of the tumor. A serum PSA along with DRE increases the chance of early detection. Transrectal ultrasound may also be used to visualize the outer lobes of the prostate and localize possible cancerous sites.

Common Abnormalities

ABNORMALITY	ASSESSMENT FINDINGS	
<p>Benign Prostatic Hypertrophy</p> <ul style="list-style-type: none"> Benign enlargement of prostate. 	<ul style="list-style-type: none"> Age over 50 Changes in urination pattern such as hesitancy, incontinence with dribbling, reduced caliber and force of urine stream, and possibly retention. Enlarged, firm, slightly elastic, smooth, possibly tender prostate. Burning on urination if accompanied by UTI. 	
<p>Chancroid</p> <ul style="list-style-type: none"> Caused by <i>Haemophilus</i> through small breaks in epidermal tissue. 	<ul style="list-style-type: none"> Pinhead papules to cauliflower-like groupings of filiform, skin-colored, pink or red painful lesions on genitalia heal, leaving a scar. 	
<p>Chlamydia</p> <ul style="list-style-type: none"> Caused by <i>C. trachomatis</i>. 	<ul style="list-style-type: none"> Dysuria and clear or white penile discharge. 	
<p>Condyloma Acuminatum</p> <ul style="list-style-type: none"> Genital warts caused by HPV. 	<ul style="list-style-type: none"> Soft, pink papillomatous or raised warty lesions on genitalia. 	
<p>Cryptorchidism</p> <ul style="list-style-type: none"> Undescended testicles. 	<ul style="list-style-type: none"> Blanching of areas on penis with 5 percent acetic acid wash. Undeveloped scrotum (undescended testicle). Increases risk of testicular cancer. Only one testis or epididymis palpable in scrotal sac. Testicular atrophy. 	
<p>Epididymitis</p> <ul style="list-style-type: none"> Acutely inflamed epididymis. 	<ul style="list-style-type: none"> Occurs chiefly in adults and may coexist with a UTI or prostatitis. Causes scrotal pain, reddened scrotum, inflamed vas deferens, and UTI symptoms. 	

ABNORMALITY	ASSESSMENT FINDINGS
Epispadias	<ul style="list-style-type: none"> ■ Urinary meatus on dorsal side of penis.
Genital Herpes <ul style="list-style-type: none"> ■ Viral STD that is treatable but not curable. 	<ul style="list-style-type: none"> ■ Fever, malaise, arthralgia, inguinal lymphadenopathy. ■ Multiple small vesicles on penis followed by ulcers on red bases.
Gonorrhea <ul style="list-style-type: none"> ■ Caused by <i>Neisseria gonorrhoeae</i>. 	<ul style="list-style-type: none"> ■ Pain. ■ Possible dysuria, pruritus, urethral discharge. ■ Recurrent infection.
Hydrocele <ul style="list-style-type: none"> ■ Lymphatic fluid-filled tunica vaginalis around testis. 	<ul style="list-style-type: none"> ■ Urinary symptoms. ■ Urethral discharge.
Hypospadias	<ul style="list-style-type: none"> ■ Causes enlarged, nontender scrotum. ■ Can be transilluminated. ■ Urinary meatus on ventral side of the penis.
Indirect Inguinal Hernia <ul style="list-style-type: none"> ■ Protrusion of hernial sac containing intestine at inguinal area. 	<ul style="list-style-type: none"> ■ Sac protrudes through internal inguinal ring into inguinal canal and may descend into scrotum. ■ Causes bulging in right or left inguinal area with heavy lifting or coughing. ■ Examination reveals mass in inguinal canal or scrotum. ■ May or may not cause tenderness or pain.
Orchitis <ul style="list-style-type: none"> ■ Inflammation of the testis. 	<ul style="list-style-type: none"> ■ Scrotal or inguinal mass. ■ Unilateral or bilateral scrotal swelling with pain or tenderness. ■ Fever. ■ Possible nausea and vomiting. ■ Recent infection, especially epididymitis or mumps. ■ Impotence. ■ Possible infertility. ■ Inflammation.



(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Paraphimosis</p> <ul style="list-style-type: none"> ■ Inability to return foreskin over penis because of edema. 	<ul style="list-style-type: none"> ■ Yeast or bacterial infection under foreskin.
<p>Phimosis</p> <ul style="list-style-type: none"> ■ Inflammation of foreskin around penis. 	<ul style="list-style-type: none"> ■ Causes constriction.
<p>Priapism</p> <ul style="list-style-type: none"> ■ Painful, sustained erection of penis without sexual desire. 	<ul style="list-style-type: none"> ■ Erect penis. ■ History of sickle cell anemia. ■ History of spinal cord lesion.
<p>Prostate Cancer</p> <ul style="list-style-type: none"> ■ Malignancy of prostate. 	<ul style="list-style-type: none"> ■ Symptoms same as BPH. ■ Prostate enlarged, hard, fixed, irregular on rectal exam. ■ Lumbosacral pain radiating to hips or legs (may be seen with metastasis).
<p>Prostatitis</p> <ul style="list-style-type: none"> ■ Inflammation of prostate gland. 	<ul style="list-style-type: none"> ■ Changes in urination pattern such as frequency, urgency, hesitancy, and nocturia. ■ Dysuria and lower back pain. ■ Penile discharge and thin, watery, possibly blood-tinged semen. ■ Enlarged, tender, boggy prostate (firm in chronic cases). ■ Fever and chills. ■ Diminished libido, impotence. ■ Recent UTI. ■ Possible infertility. ■ Nontender palpable cyst in scrotum.
<p>Spermatocele</p> <ul style="list-style-type: none"> ■ Sperm-filled cyst of epididymis. 	<ul style="list-style-type: none"> ■ Can be transilluminated.
<p>Syphilis</p> <ul style="list-style-type: none"> ■ Caused by <i>Treponema pallidum</i>. ■ Primary, secondary, and tertiary forms. 	<ul style="list-style-type: none"> ■ Primary: Genital chancre, heals without leaving a scar.
	<ul style="list-style-type: none"> ■ Secondary: Alopecia, rash, malaise, headache, arthralgia. ■ Tertiary: Gummas, aortic valve, thoracic aneurysm, personality changes, ataxia, paresthesia, Charcot's joints.



ABNORMALITY

ASSESSMENT FINDINGS

Testicular Torsion

- Twisting of spermatic cord.
- Surgical emergency.

Varicocele

- Varicose veins of spermatic cord.

- Pain and swelling in testes.
- Nausea and vomiting.

- Associated with infertility.
- Scrotum is purplish and soft.
- Scrotum feels like a “bag of worms” and collapses when it is elevated in supine position.
- Dull ache along spermatic cord.
- Slight dragging sensation in scrotum.

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the GU system. Ask the patient the following:

- Do you have penile pain? Do you have any lesions? If yes, please describe.
- Could you tell me about your sexual practices? Do you practice oral, genital, or anal sex? How many partners have you had? What is your sexual preference: heterosexual, homosexual, or bisexual? Do you practice safe sex? What do you use?
- Do you have any discharge? Type?
- Did you have a history of GU surgery, STDs, or cancer?
- Are you taking any medications? If, yes, what are the medications?
- Do you have any medical problems, such as diabetes mellitus, HTN, or peripheral vascular disease?
- Do you have any problems with urination, such as burning, frequency, dysuria, or hematuria?

PHYSICAL ASSESSMENT

Equipment

Gather equipment needed for the exam.

Gloves

Penlight

Stethoscope

Cotton swab or gauze

Urogenital alginate swabs and Thayer-Martin plate

Hemoccult test

Water-soluble lubricant

Position

Supine or standing for genitalia assessment; side-lying or leaning over exam table for rectal exam.

Technique

Remember to scan your patient from head-to-toe looking for signs related to male GU assessment.

Inspection, Palpation, Auscultation

Inspect

Genitalia

- Hair distribution
- Penis
- Scrotum
- Urethral meatus
- Inguinal area

Palpate

- Penis
- Scrotum and testes
- Horizontal and vertical lymph nodes
- Hernias (inguinal, femoral)
- Rectum
- Prostate

Auscultate

- Scrotum, if hernia present.

Specimens

- Obtain specimens for diagnostic tests.
- Hematest stool if present.

DOCUMENT

Document your findings and develop your plan of care.


S U M M A R Y

- Assessing the male reproductive system requires knowledge of the anatomy and physiology of the male reproductive organs. These include the scrotum, testes, spermatic cord, duct system, accessory glands, lymphatic system, penis, and prostate.
- The physical assessment of the male reproductive system requires the examiner to be sensitive in responding to the patient's need for privacy.

R E V I E W Q U E S T I O N S

1. Mathew Wall, age 25, discovered a lump in his left testicle while performing a TSE. Which finding describes the normal consistency of the testes?
 - a. Soft
 - b. Rubbery
 - c. Nodular
 - d. Hard
2. The health history is helpful in identifying risk factors for testicular cancer. Which childhood medical problem increases the risk for testicular cancer?
 - a. Mumps
 - b. Undescended testicles
 - c. Umbilical hernia
 - d. Chickenpox
3. Which family history factor increases the risk for testicular cancer?
 - a. Maternal use of birth control pills (BCPs)
 - b. Maternal history of smoking
 - c. Maternal history of alcohol use
 - d. Maternal use of DES
4. The PSA test is a screening test for:
 - a. BPH.
 - b. Prostatitis.
 - c. Prostatic cancer.
 - d. Epididymitis.
5. Bill Brown, age 20, comes to the student health center complaining of thick, yellow penile discharge. This type of penile discharge is often associated with:
 - a. Gonorrhea.
 - b. Syphilis.
 - c. Human papillomavirus (HPV).
 - d. Candidiasis.
6. On examination of the external genitalia, you note a painless, white, wartlike lesion. The lesion described is most likely:
 - a. Gonorrhea.
 - b. Syphilis.
 - c. Condylomata acuminatum.
 - d. Herpes simplex.
7. While obtaining a health history of Jack Madison, age 27, you note that he has a positive history of sickle cell anemia. Sickle cell anemia increases his risk for developing:
 - a. Priapism.
 - b. Phimosis.
 - c. Hydrocele.
 - d. Chordee.
8. While performing a newborn assessment on Baby Boy Kelly, you note that the urethral meatus opens on the dorsal side of the penis. This condition is referred to as:
 - a. Epispadias.
 - b. Hypospadias.
 - c. Varicocele.
 - d. Hydrocele.
9. While performing a newborn assessment on Baby Boy Kelly, you note that there is scrotal swelling. What is the most likely cause of this condition?
 - a. Epispadias
 - b. Hypospadias
 - c. Varicocele
 - d. Hydrocele
10. You are working at a health fair, performing cancer risk assessments and promoting cancer screenings. Joe Devlin, age 50, has never been screened for prostate cancer. What is the recommended age for prostate screening for this patient?
 - a. 20
 - b. 30
 - c. 40
 - d. 50

CHAPTER 20



Assessing the Motor-Musculoskeletal System

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent motor-musculoskeletal history questions
- Obtain a motor-musculoskeletal history
- Perform a physical assessment of the motor-musculoskeletal system
- Document motor-musculoskeletal findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

The musculoskeletal system provides shape and support to the body, allows movement, protects the internal organs, produces red blood cells in the bone marrow (hematopoiesis), and stores calcium and phosphorus in the bones. Although examining this system is usually only a small part of the overall physical assessment, everything we do depends on an intact musculoskeletal system. How extensive an assessment you perform depends largely on each patient's problems and needs.

Perform a comprehensive musculoskeletal assessment if you detect a musculoskeletal abnormality or uncover a symptom that suggests musculoskeletal involvement. Musculoskeletal problems are common in all age groups. Primary problems may result from congenital, developmental, infectious, neoplastic, traumatic, or degenerative disorders of the system itself. Secondary problems can result from disorders of other body systems.

The goal of a complete musculoskeletal assessment is to detect risk factors, potential problems, or musculoskeletal dysfunction early and then to plan appropriate interventions, including teaching health promotion and disease prevention and implementing treatment measures. By doing so, you can play a significant role in preventing pain and dysfunction in your patients.

Anatomy and Physiology Review

Before beginning your assessment, you need to understand how the musculoskeletal system works. It consists of three major components: bones, muscles, and joints. Tendons, ligaments, cartilage, and bursae serve as connecting structures and complete the system. (See *Structures and Functions of the Musculoskeletal System*.) Figure 20.1 illustrates the musculoskeletal system, anterior and posterior views.

Bones

Composed of osseous tissue, bones are divided into two types: **compact bone**, which is hard and dense and makes up the shaft and outer layers, and **spongy bone**, which contains numerous spaces and makes up the ends and centers of the bones. **Osteoblasts** and **osteoclasts** are the cells responsible for the continuous process of creating and destroying bone. Osteoblasts form new bone tissue, and osteoclasts break down bone tissue. Bones also contain **red marrow**, which produces blood cells, and **yellow marrow**, which is composed mostly of fat. The outer covering of bone, called the **periosteum**, contains osteoblasts and blood vessels that promote nourishment and formation of new bone tissue.

Bones vary in shape and include long bones (Fig. 20.2), such as the humerus and femur; short, flat

bones, such as the sternum and ribs; and bones with irregular shapes, such as the hips and vertebrae.

Muscles

The body is composed of skeletal, smooth, and cardiac muscle. Made up of fasciculi (long muscle fibers) that are arranged in bundles and joined by connective tissue, skeletal muscles attach to bones by way of strong, fibrous cords called **tendons**. **Ligaments** are dense, flexible, strong bands of fibrous connective tissue that tie bones to other bones.

Cartilage is dense connective tissue consisting of fibers embedded in a strong, gel-like substance. Cartilage lacks nerve innervation, blood vessels, and lymph vessels, so it is insensitive to pain and regenerates slowly and minimally after injury. Regeneration occurs primarily at sites where the articular cartilage meets the synovial membrane. Cartilage may be fibrous, hyaline, or elastic. **Fibrous cartilage** forms the symphysis pubis and the intervertebral discs. **Hyaline cartilage** covers the articular bone surfaces (where bones meet at a joint), connects the ribs to the sternum, and is found in the trachea, bronchi, and nasal septum. **Elastic cartilage** is located in the auditory canal, the external ear, and the epiglottis.

Joints

The **joint** or *articulation* is the place where two or more bones meet. Joints provide range of motion (ROM) for the body parts and are classified three ways: by the degree of movement they permit, by the connecting tissues that hold them together, and by the type of motion the structure permits. Figure 20.3 illustrates the fibrous and cartilaginous joints and Figure 20.4 the synovial joints. (See *Types of Synovial Joints*.)

Bursae

Bursae are small, disc-shaped synovial fluid sacs located at points of friction around joints. They act as cushions, thereby reducing the stress to adjacent structures, and facilitate movement. Two examples of bursae are the **prepatellar bursa** (in the knee) and the **subacromial bursa** (in the shoulder).

Interaction with Other Body Systems

The neurological and respiratory systems contribute to maintaining musculoskeletal functioning. A problem in any of these systems may affect the functioning of the musculoskeletal system.

Neurological System

The neurological system is responsible for coordinating the functions of the skeleton and muscles. If your patient




FIGURE 20.1. Cross-section of musculoskeletal system.





FIGURE 20.2. Cross-section of long bone.

FIBROUS JOINTS (synarthroses)

Syndesmosis

Suture



CARTILAGINOUS JOINTS (amphiarthroses)

Symphysis

Synchondrosis



FIGURE 20.3. Fibrous and cartilaginous joints.

has neurological complaints, combine the musculoskeletal and neurological assessments because the spinal cord and nerves originate from the spine and innervate the musculoskeletal structures of the back and the extremities. A dysfunction in the neurological system is often reflected as pain, abnormal movement, or paresthesias in the extremities and/or back. The patient's gait may provide information on muscular weakness or neurological disease.

Back pain is a major source of disability in the United States. A large proportion of the population complains of back pain at one time or another, with the most common complaint being low back pain radiating into the hip and down the leg. This pain is usually of neurological origin and emanates from the sciatic nerve. Pain may also be caused by arthritic disease of the spine or hip or muscle spasm of the lower back. Understanding the anatomy of the back and spinal nerve tracks will help you determine the pain's origin.

Respiratory System

The respiratory system depends on the thorax, bony structures, and muscles of the chest to protect the lungs and assist with breathing. The accessory muscles, which include the sternocleidomastoid, anterior serrati, scalene, trapezius, intercostal, and rhomboid muscles, come into play when a person is involved in aerobic activities or when the body has intrapulmonary resistance to air movement (e.g., chronic pulmonary lung disease). These accessory muscles enhance ventilation by increasing chest expansion and lung size during inspiration. Intercostal muscles also coordinate rib movement; external intercostal muscles pull the ribs up and out, and internal intercostals pull the ribs down and inward. Contraction of these muscles facilitates air movement into the lungs by decreasing intrathoracic pressure. As these muscles relax, exhalation occurs as the lung recoils. Abdominal muscles can also assist with deep breathing, tachypnea (rapid breathing), exercise, coughing, and sneezing. An intact thoracic cage and normal accessory and abdominal muscles are necessary for respiratory function. A musculoskeletal injury or problem in these areas can result in altered respiratory functioning.

Developmental, Cultural, and Ethnic Considerations

Infants and Children

Before birth, a skeleton forms in the fetus; it is first composed of cartilage and then later ossifies into true growing bone. After birth, bone growth continues rapidly during infancy and then steadily during the childhood years. Another growth spurt occurs for both boys and girls during adolescence.

Long bones increase in diameter by depositing new bone tissue around the shafts. Lengthening occurs at the




FIGURE 20.4. Types of synovial joints.

epiphyses, which are specialized growth centers (growth plates) located at the ends of long bones. Any injury or infection at these growth plates puts the growing child at risk for bone deformity. Longitudinal growth of the bones continues until closure of the epiphyses, which occurs at age 20.

Skeletal contour changes are also apparent in infants and children. At birth, the spine has a single C-shaped curve. At 3 to 4 months of age, an infant is able to raise his or her head from the prone position, allowing the

development of the anterior curve in the cervical neck region. As development progresses and the infant is able to stand and walk, the anterior curve develops in the lumbar region. This occurs between 12 and 18 months of age. A toddler stands with feet wide apart to provide balance as he or she learns to walk.

The school-age child usually stands with the normal adult curvature, which should continue until old age. Throughout childhood, the skeleton continues to grow linearly; muscles and fat are responsible for significant

Structures and Functions of the Musculoskeletal System

STRUCTURE

FUNCTION

Bones

- 206 bones make up the axial skeleton (head and trunk) and appendicular skeleton (extremities, shoulders, and hips).

Muscles

- 650 skeletal (voluntary) muscles under conscious control.

Joints (Articulations)

- The place where two adjacent bones or cartilage meet.

- Provide structure and protection, serve as levers, produce blood cells, and store calcium.

- Allow the body to move, stand erect, and produce body heat.

- Provide ROM for body parts.

Classification Based on Movement They Permit

- *Synarthrosis*: Immovable (e.g., sutures of skull).
- *Amphiarthrosis*: Limited movement (e.g., symphysis pubis).
- *Diarthrosis*: Freely movable (e.g., hip).

Classification Based on Connecting Tissues That Hold Them Together

- *Fibrous joints*: No joint cavity; fibrous connective tissue joins bones; usually allow no movement. Types include:
 - *Sutures*: Bones fused together (e.g., skull).
 - *Syndesmoses*: Bones very close together; held together by ligament that gives strength and support to joint and also limits movement (e.g., tibiofibular joint).
 - *Gomphoses*: Peg and socket (e.g., root of tooth).
- *Cartilaginous joints*: Bones joined by hyaline cartilage or fibrocartilaginous disc; allows slight movement. Types include:
 - *Synchondrosis (primary cartilaginous joint)*: Allows for growth, but not movement (e.g., epiphyseal plate joins diaphysis and epiphysis of long bones and allows growth). Once growth is complete, joint becomes synostotic (sealed).
 - *Symphyses (secondary cartilaginous joint)*: Articulating bones covered with hyaline cartilage; fibrocartilaginous discs that act as shock absorbers; allows for some movement (e.g., symphysis pubis moves during pregnancy to allow for fetal growth).
- *Synovial*: Most movable and complex. Has cavity filled with lubricating (synovial) fluid to help ends of bones slide. Enclosed by fibrous capsule of connective tissue and connected to periosteum of bone. Contain free-floating synovial cells and various leukocytes that phagocytose joint debris and microorganisms. Some contain bursae. Synovial joints may be:
 - *Uniaxial*: Movement limited to one axis (e.g., elbow).
 - *Biaxial*: Movement on two axes (e.g., hand).
 - *Multiaxial or triaxial*: Movement on three axes (e.g., shoulder).

Connecting Structures

- Tendons
- Ligaments
- Cartilage
- Bursae

- Connect muscles to bone. They enable bones to move when skeletal muscles contract.
- Strong, dense bands of tissue that connect the joint (articular) ends of bones, either facilitating movement or providing stability or limiting movement.
- Supports and shapes various structures (e.g., auditory canal), and cushions and absorbs shock that otherwise would be transmitted directly to bone (e.g., intervertebral discs).
- Small, disc-shaped, flattened sacs filled with synovial fluid located at points of friction around joints. Act as cushions, reducing stress to adjacent structures and facilitating movement. Examples are prepatellar bursa in knee and subacromial bursa in shoulder.

Types of Synovial Joints

JOINT	DESCRIPTION	EXAMPLE/MOVEMENT
Pivot	<ul style="list-style-type: none"> Permits rotation in one axis. Axis is longitudinal, with bone moving around a central axis without any displacement from that axis. 	<ul style="list-style-type: none"> <i>Example:</i> Proximal radioulnar joint. <i>Movement:</i> Supinates, pronates, and rotates.
Hinge	<ul style="list-style-type: none"> Allows movement in only one axis, namely flexion or extension, with axis situated transversely. 	<ul style="list-style-type: none"> <i>Example:</i> Elbow, knee. <i>Movement:</i> Flexion and extension.
Condylloid	<ul style="list-style-type: none"> Permits movement in two axes. Described as an “egg-in-spoon joint,” with long diameter of oval serving as one axis, and short diameter of oval serving as other axis. 	<ul style="list-style-type: none"> <i>Example:</i> Wrist. <i>Movement:</i> Flexion, extension, abduction, adduction, and circumduction.
Saddle	<ul style="list-style-type: none"> Has two axes like condylloid joint. Articular surfaces are saddle shaped and move in similar fashion to condylloid joint. 	<ul style="list-style-type: none"> <i>Example:</i> Thumb. <i>Movement:</i> Abduction, adduction, opposition, and reposition.
Ball and Socket	<ul style="list-style-type: none"> Moves across many possible axes. Articular surfaces are reciprocal segments of a sphere. 	<ul style="list-style-type: none"> <i>Example:</i> Shoulder and hip. <i>Movement:</i> Flexion, extension, internal rotation, external rotation, abduction, adduction, and circumduction.
Plane/Gliding	<ul style="list-style-type: none"> Moves across many axes. Articular surfaces are flat, and one bone rides over the other in many directions. 	<ul style="list-style-type: none"> <i>Example:</i> Patellofemoral and acromioclavicular joints, some carpal and tarsal bones, and articular vertebral processes. <i>Movement:</i> Limited in many directions.

weight increases. Individual muscle fibers continue to grow as the child grows, with a marked growth period noted during adolescence. At this time, muscles are responding to increased growth hormone, adrenal androgens, and testosterone in boys. Muscles vary in growth rate, size, and strength, depending on genetic factors, nutritional status, and amount of exercise.

Common knee deviations in children include **genu varum** (bowlegs) and **genu valgum** (knock knees). In a child with genu varum, the knees are approximately 5 cm apart and the medial malleoli touch when the child stands. This variation is common during the first years when the child is beginning to walk, but usually does not persist beyond 2 to 3 years. When genu valgum is present in a child, the knees touch and the medial malleoli are 7.5 cm or more apart when the child is standing. These deviations are considered normal for a child aged 2 to 3½ years and may persist until age 7.

Toddlers often have “potbellies” and **lordosis** (accentuated lumbar curve). This posture is normal and helps the child adjust to the change in the center of gravity. It should disappear as the child grows. Spinal deformities in children may be structural, but more commonly are caused by poor posture. **Scoliosis** (lateral curvature)


may become apparent during adolescence, with girls at a higher risk than boys. The spine does not grow straight, and the shoulders and iliac crests are not the same height. Assessing for scoliosis is an important component when working with adolescents.

Additional history questions for parents of infants and children include:

- Was there a normal labor and delivery without trauma to the infant? (Trauma increases the risk for fractures of the humerus and clavicle.)
- Did the infant need to be resuscitated? (Anoxia may result in hypotonia of muscles.)
- Did the infant develop as expected?
- Has the child ever had a broken bone or dislocation? If so, how were these injuries treated?
- Have you noticed any bone deformity, spinal curvature, or unusual shape of the legs, toes, or feet?

Additional history questions for adolescents include:

- Are you involved in any sports? How often do you play or practice?
- Do you use protective or special equipment?
- Do you warm up before participating in an athletic activity?



Relationship of Musculoskeletal System to Other Systems

These questions help assess the safety of the sport for the height and weight of the adolescent and gauge the risk of injuries.

Pregnant Women

Progressive lordosis is the most characteristic change in posture in pregnant women. It compensates for the enlarging fetus by shifting the center of gravity and moving the weight of the enlarging fetus back on the lower extremities. This shift in balance may create strain on the low back muscles, which may be felt as low back pain during late pregnancy. Anterior cervical flexion during the third trimester, kyphosis (humpback), and slumped shoulders are other postural changes that compensate for lordosis. Upper back changes may put pressure on the ulnar and median nerves, creating aching, numbness, and weakness in the upper extremities of some pregnant women. Increased mobility in the sacroiliac, sacrococcygeal, and symphysis pubis joints in the pelvis occurs in preparation for delivery, which also contributes to the characteristic waddling gait.

Older Adults

With aging, bone density decreases because loss of bone tissue occurs more rapidly than formation of new bone tissue. This results in osteoporosis, with women at a higher risk than men and whites affected more often than African Americans.

Kyphosis with a backward head tilt to compensate and slight flexion of the hips and knees are postural changes that occur with aging. Decreased height is the most noticeable postural change. It occurs as a result of shortening of the vertebral column secondary to the loss of water content and thinning of the intervertebral discs. Both men and women can expect a decrease in height beginning in their 40s and continuing until age 60. After this age, height decreases occur secondary to osteoporotic changes in the height of individual vertebrae. This collapse of vertebrae results in a shortening of the trunk and comparatively long extremities. This change in the spine, kyphosis and loss of height due to collapse and wedging of the vertebral column, is often referred to as *dowager's hump*.

Contour changes also occur as a result of the distribution of subcutaneous fat. Most men and women gain weight in their fourth and fifth decades, losing fat in the face and depositing it in the abdomen and hips. This distribution pattern continues into later years, but fat continues to decrease in the periphery, especially the forearms. The loss of subcutaneous fat in certain parts of the body contributes to marked bony prominences. The tips of the vertebrae, ribs, and iliac crests may become very noticeable and the body hollows (e.g., cheeks and axillae) become deeper. There is loss of muscle mass, a decrease in muscle size, and some muscle atrophy, which contributes to generalized weakness. These muscle changes can cause decreased agility, an abnormal gait with uneven rhythm and short steps, and a wide base of support. Other compli-

cating factors include fear of falling, osteoporosis, painful arthritic joints, poor vision, and peripheral neuropathy.

Lifestyle can affect the musculoskeletal changes that occur with aging. A sedentary lifestyle leads to decreased muscle mass and strength and increased muscle atrophy. Decreased speed and strength, resistance to fatigue, reaction time, and coordination are changes that can be prevented by physical exercise. Physical activity also prevents or delays bone loss in aging adults.

Additional history questions for the aging adult should elicit information about loss of function, self-care deficits, or safety risks that may occur as a result of aging, injury, or illness. Ask the following questions:

- Have you broken any bones recently? If so, how?
- Have you noticed any weakness over the past months?
- Have you had any increase in stumbling or falls?
- Do you use mobility aids like a cane or walker to help you get around?

People of Different Cultures/Ethnic Groups

When assessing the musculoskeletal system, you need to consider your patient's ethnicity because it can affect musculoskeletal anthropomorphic and physical characteristics. (See *Culturally Bound Musculoskeletal Health Problems*.)

Introducing the Case Study



Maria O'Malley, age 68, is concerned with osteoporosis because she is postmenopausal and feels as though she is "shrinking." Before her routine physical examination, Mrs. O'Malley begins to ask questions about how osteoporosis occurs, what can be done to treat it, and what complications can occur from having it.

Performing the Musculoskeletal Assessment

Health History

Obtaining a thorough and accurate health history is crucial to your assessment of the musculoskeletal system. As the health history interview progresses, clarify exactly what the patient means by certain subjective complaints, including the location, character, and onset of any symptoms. Only the patient can give you data regarding pain, stiffness, ability to move, and how daily activities have been affected. The history provides the subjective data that will direct the physical examination. If time is an issue and you are unable to perform a complete health history, perform a focused history on the motor-musculoskeletal system.

Culturally Bound Musculoskeletal Health Problems

CULTURAL GROUP	HEALTH PROBLEM
African Americans	<ul style="list-style-type: none"> ■ Tendency toward hyperplasia (overgrowth) of connective tissue accounts for increased incidence of SLE. ■ Greater bone density than Europeans, Asians, and Hispanics accounts for decreased incidence of osteoporosis.
Amish	<ul style="list-style-type: none"> ■ Dwarfism syndrome found in nearly all Amish communities.
Chinese Americans	<ul style="list-style-type: none"> ■ Generally shorter than Westerners. Bone structure also differs from that of Westerners: <ul style="list-style-type: none"> ■ Ulna longer than radius. ■ Hip measurements smaller (females 4.14 cm smaller; males 7.6 cm smaller). ■ Bone length shorter. ■ Bone density less.
Egyptian Americans	<ul style="list-style-type: none"> ■ Relatively short in stature. Average adult height for males 5'10"; for females 5'4".
Filipino Americans	<ul style="list-style-type: none"> ■ Short in stature. Average adult height ranges from 5' to average European American size.
Irish Americans	<ul style="list-style-type: none"> ■ Taller and broader than average European American and Asian. Hip width greater. ■ Bone density less than that of African Americans.
Navajo Native Americans	<ul style="list-style-type: none"> ■ Taller and thinner than other Native American peoples.
Vietnamese Americans	<ul style="list-style-type: none"> ■ Small in stature. Average adult height 5' for women, with men being a few inches taller.

Source: Adapted from Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.

Biographical Data

A review of the biographical data may identify patients who are at risk for musculoskeletal problems. Consider the patient's age, sex, marital status, race, ethnic origin, and occupation as possible risk factors. For example, women are at a higher risk than men for osteoporosis and rheumatoid arthritis (RA); an occupation that requires heavy lifting increases the risk for injury to muscles, joints, and supporting structures; and the risk for degenerative joint disease (DJD; osteoarthritis) increases with age. Even a patient's address may have significance. For

example, Lyme disease, which has musculoskeletal repercussions, is endemic on the northeast coast from Massachusetts to Maryland, in the midwest states of Wisconsin and Minnesota, and on the west coast from northern California to Oregon.

Current Health Status

Begin your assessment with questions about the patient's current health status, because she or he is most interested in this. The major symptoms to watch for, in order of importance, are pain, weakness, and stiffness.

Focused Musculoskeletal History

If your patient's condition or time restraints prohibit a detailed health history, take a focused musculoskeletal history instead. The patient's response to the questions below will help direct your assessment.

- Do you have a history of musculoskeletal problems, pain, or disease? If yes, are you taking any medications or undergoing any treatments?
- Do you have any other medical problems?
- Are you taking any medications? Prescribed, OTC?
- Have any accidents or trauma ever affected your bones or joints?
- Do your joint, muscle, or bone problems limit your usual activities?
- Do you have any occupational hazards that could affect your muscles and joints?
- Have you been immunized for tetanus and polio?
- Do you smoke or consume alcohol or caffeine? If yes, how much and how often?

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Pain

Pain can result from bone, muscle, or joint problems. Bone pain is not usually associated with movement unless there is a fracture, but muscle pain is. Current or recent illness can cause muscle aches. Bone pain is deep, dull, and throbbing; muscle pain takes the form of cramping or soreness. In RA, joint pain and stiffness are worse in the morning. In osteoarthritis, the joints are stiff after rest and pain is worse at the end of the day.

Weakness

Muscle weakness associated with certain diseases migrates from muscle to muscle or to groups of muscles. Knowing the symptom patterns and when they occur can help identify the disease process responsible for weakness. Identifying the weakness related to proximal or distal muscle groups is also helpful. Proximal weakness is usually a myopathy; distal weakness is usually a neuropathy.

Be sure to ask the patient if the weakness interferes with his or her ability to perform everyday activities. For example, proximal weakness of the upper extremities often presents itself as difficulty lifting objects or combing hair. Proximal weakness of the lower extremities presents as difficulty with walking and crossing the knees. Patients with polymyalgia rheumatica have proximal muscle weakness. A distal weakness of the upper extremities is manifested by difficulty in dressing or turning a doorknob. Patients with myasthenia gravis have difficulty with diplopia, swallowing, and chewing, along with generalized muscle weakness.

Stiffness

Stiffness is a common musculoskeletal complaint. It is important to ascertain if the stiffness is worse at any particular time of the day. For example, patients with RA are stiff on arising because of the period of joint rest that occurs during sleep.

Balance and Coordination Problems

Problems with balance and coordination often indicate a neurological problem. These problems often manifest as gait problems or difficulty in performing activities of daily living (ADLs). The patient may complain of falling or losing balance or may have ataxia—irregular and uncoordinated voluntary movements. Gait problems are associated with cerebellar disorders, Parkinson's disease, multiple sclerosis (MS), herniated disc, stroke, Huntington's chorea, brain tumor, inner ear problems, medications, and exposure to chemical toxins.

Other Related Symptoms

Certain musculoskeletal diseases may produce multiple symptoms. For example, acute rheumatic fever, gout, and autoimmune inflammatory diseases may cause fever and joint pain. A woman with back pain and vaginal discharge may have a gynecological disease. Lower back pain and weight loss suggest tuberculosis (TB) of the spine. Bowel and bladder dysfunction suggest a herniated disc. Therefore, also ask the patient: "Do you have a sore throat, fever, joint pain, rash, weight loss, diarrhea, numbness, tingling, or swelling?"

Past Health History

After taking the past health history, compare it with the patient's present musculoskeletal status or uncover risk factors that might predispose the patient to musculoskeletal disorders.

(Text continued on page 697)

Past Health History

RISK FACTORS/QUESTIONS TO ASK

Childhood Illnesses

- Do you have a history of juvenile RA (JRA)?
- Do you have cerebral palsy? Muscular dystrophy?

Surgery

- Have you ever had surgery or other treatment involving bones, muscles, joints, or other supporting structures?

Diagnostic Procedures

- *Postmenopausal women:* Have you ever had a bone densitometry (dexascan)?

RATIONALE/SIGNIFICANCE

- Thirty percent of children with JRA have symptoms as adults. Residual deformities from JRA may be seen in adults.
- History of childhood musculoskeletal problems may account for current physical findings.
- Provides baseline information for physical assessment and explains physical findings.
- Perimenopausal and postmenopausal women should be screened for osteoporosis.

(continued)

Past Health History (continued)

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Serious Injuries</p> <ul style="list-style-type: none"> Have you had any past problems or injuries to your joints, muscles, or bones? If so, what? What treatment was given? Do you have any residual effects from the injury or problem? <p>Serious/Chronic Illnesses</p> <ul style="list-style-type: none"> Do you have any other medical problems? Have you had a recent infection? 	<ul style="list-style-type: none"> Data on past injuries or problems provide baseline data for physical assessment. Past trauma may affect ROM. SLE can cause arthritis, synovitis, and myositis. HIV increases risk for infectious arthritis, osteomyelitis, and polymyositis. RA and SLE may improve with HIV. Lyme disease can cause migratory joint and muscle pain; if untreated, can lead to arthritis. TB can affect bones and joints. Neuropathies associated with diabetes can lead to muscle atrophy and neuropathic arthropathy (Charcot's joint). Tertiary syphilis can also cause Charcot's joints. Recent infections such as Epstein-Barr virus may trigger RA.
<p>Immunizations</p> <ul style="list-style-type: none"> What immunizations have you had and when? Have you been immunized against tetanus and polio? 	<ul style="list-style-type: none"> Tetanus and polio have symptoms that mimic other musculoskeletal disorders. Immunization information can aid in differential diagnoses.
<p>Allergies</p> <ul style="list-style-type: none"> Are you allergic to dairy products? Do you have lactose intolerance? 	<ul style="list-style-type: none"> Allergies to dairy products or lactose intolerance may lead to inadequate calcium intake. Patients who are immobile or who have a reduced intake of calcium and vitamin D are especially prone to osteoporosis.
<p>Medications</p> <ul style="list-style-type: none"> Are you taking any prescribed or OTC medications? <i>Middle-aged women:</i> Are you menopausal? Are you receiving estrogen replacement therapy? 	<ul style="list-style-type: none"> Certain medications can affect the musculoskeletal system. See <i>Drugs That Adversely Affect the Musculoskeletal System</i>. Women who begin menopause early are at greater risk for osteoporosis because of decreased estrogen levels, which tend to decrease bone mass density.
<p>Recent Travel/Military Service/Exposure to Infectious or Contagious Diseases</p> <ul style="list-style-type: none"> Have you been hiking or camping recently? Were you ever in the military? 	<ul style="list-style-type: none"> Lyme disease is endemic in certain areas of the country. Exposure to chemicals during military service may manifest in musculoskeletal symptoms.

Drugs That Adversely Affect the Musculoskeletal System

DRUG NAME	DRUG	POSSIBLE ADVERSE REACTIONS
Adrenocorticotrophic hormone (ACTH)	Corticotropin	Muscle weakness, muscle wasting, osteoporosis, vertebral compression fractures, aseptic necrosis of humeral or femoral heads
Anticonvulsants	Phenytoin	Ataxia
Antidepressants	Trazadone	Musculoskeletal aches and pains

DRUG NAME	DRUG	POSSIBLE ADVERSE REACTIONS
Antigout agents Antilipemic agents	Colchicine 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitors ("statins"), fibric acid derivatives, niacin	Myopathy with prolonged administration Myopathy, rhabdomyolysis, myalgias
Benzodiazepines Central nervous system stimulants	All benzodiazepines Amphetamine sulfate	Ataxia Increased motor activity
Corticosteroids	Prednisone, dexamethaxone, hydrocortisone	Muscle weakness, muscle wasting, osteoporosis, vertebral compression fractures, aseptic necrosis of humeral or femoral heads
Diuretics Phenothiazines	Thiazides and loops All phenothiazines	Muscle cramps Extrapyramidal symptoms (dystonic reactions, motor restlessness, and Parkinsonian signs and symptoms)
Miscellaneous skin agents	Isotretinoin	Bone or joint pain, general muscle aches

Family History

The purpose of the family history is to identify predisposing or causative factors for musculoskeletal prob-

lems. After assessing the current and past musculoskeletal health status, investigate possible familial tendencies toward problems.

Family History

RISK FACTORS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Familial Musculoskeletal Problems</p> <ul style="list-style-type: none"> Do you have a family history of gout, arthritis, or osteoporosis? 	<ul style="list-style-type: none"> Positive family history for these diseases increases risk to patient.

Review of Systems

The review of systems (ROS) allows you to assess the interrelationship of the musculoskeletal system to every

other system. Often, you may uncover important facts that your patient failed to mention earlier.

Review of Systems

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>General Health Survey</p> <ul style="list-style-type: none"> How have you been feeling? Have you had any changes in weight or height? Have you had a fever? 	<ul style="list-style-type: none"> <i>Fatigue/activity intolerance:</i> RA, SLE, and Lyme disease. Obesity is risk factor for DJD, osteoarthritis, and low back pain. <i>Anorexia/weight loss:</i> RA. <i>Loss of height:</i> Osteoporosis. RA, Lyme disease.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Integumentary</p> <ul style="list-style-type: none"> Have you had any rashes or lesions? Have you had a change in hair growth or hair loss? 	<ul style="list-style-type: none"> <i>Bull's-eye rash</i>: Lyme disease. <i>Ulcers and subcutaneous rheumatoid nodules on forearm and elbow</i>: RA. <i>Butterfly facial rash</i>: SLE. <i>Patchy alopecia and short broken hair above forehead</i>: SLE.
<p>HEENT <i>Head and Neck</i></p> <ul style="list-style-type: none"> Do you have any swollen glands or nodes? 	<ul style="list-style-type: none"> <i>Enlarged lymph nodes</i>: Systemic infection. <i>Lymphadenopathy</i>: RA. Enlarged thyroid as a result of thyroid disease can affect musculoskeletal system.
<p><i>Eyes, Ears, Nose, and Throat</i></p> <ul style="list-style-type: none"> Do you have dry eyes or mouth? Do you have red eyes? 	<ul style="list-style-type: none"> Sjögren's syndrome causes dry mouth and eyes and is associated with RA. <i>Episcleritis and keratoconjunctivitis</i>: RA.
<p>Respiratory</p> <ul style="list-style-type: none"> Do you have any breathing problems? 	<ul style="list-style-type: none"> <i>Interstitial fibrosis and pleuritis</i>: RA. <i>Pneumonitis and pleural effusion</i>: SLE.
<p>Cardiovascular</p> <ul style="list-style-type: none"> Do you have any cardiovascular problems? Have you had changes in the color of your hands in response to cold? 	<ul style="list-style-type: none"> Rheumatoid heart disease associated with RA. <i>Pericarditis, myocarditis, Raynaud's disease</i>: RA and SLE
<p>Gastrointestinal</p> <ul style="list-style-type: none"> Do you have abdominal pain, diarrhea, nausea, vomiting, or difficulty swallowing? Do you have any bowel problems or incontinence? 	<ul style="list-style-type: none"> <i>Gastrointestinal complaints</i>: SLE. <i>Bowel problems, incontinence, loss of bowel function</i>: Cauda equina syndrome caused by compression of sacral nerve, as can occur with a herniated disc or spinal stenosis.
<p>Genitourinary</p> <ul style="list-style-type: none"> Do you have blood in your urine? 	<ul style="list-style-type: none"> <i>Hematuria</i>: SLE.
<p>Reproductive</p> <ul style="list-style-type: none"> Do you have a history of sexually transmitted diseases (STDs)? <i>Women</i>: Are you menopausal? If yes, are you on hormone replacement therapy (HRT)? Do you have menstrual irregularities? Do you have any gynecological disorders or vaginal discharge? 	<ul style="list-style-type: none"> Tertiary syphilis can cause ataxia and Charcot's joints. Postmenopausal women at greater risk for osteoporosis. Menstrual irregularities are associated with SLE. Gynecological disorders such as pelvic inflammatory disease (PID) or ovarian cancer may cause back pain.
<p>Neurological</p> <ul style="list-style-type: none"> Do you have any loss of sensation, numbness, or tingling? Are you experiencing paralysis? 	<ul style="list-style-type: none"> <i>Loss of sensation</i>: Herniated disc. <i>Neuropathies</i>: RA, SLE. Paralysis can lead to muscle atrophy from disuse. Tertiary syphilis can cause ataxia and Charcot's joints.
<p>Endocrine</p> <ul style="list-style-type: none"> Do you have diabetes mellitus? Do you have thyroid disease? 	<ul style="list-style-type: none"> Diabetes can cause neuropathies and muscle wasting. Hypothyroidism can cause weakness, muscle aches, pain, and arthralgia. Hyperthyroidism can cause osteoporosis.
<p>Immunologic/Hematologic</p> <ul style="list-style-type: none"> Do you have sickle cell anemia? Do you bruise easily? 	<ul style="list-style-type: none"> Can cause joint pain. SLE can cause thrombocytopenia.

Psychosocial Profile

The psychosocial profile can reveal patterns in the

patient's life that may affect the musculoskeletal system and put her or him at risk for musculoskeletal disorders.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Health Practices and Beliefs/Self-Care Activities

- What do you do to promote the health of your musculoskeletal system?
- Do you get an annual physical exam?
- Do you wear protective equipment when playing sports?
- Do you understand proper body mechanics?

Typical Day

- How would you describe your typical day?

Nutritional Patterns

- What did you eat in the last 24 hours?
- Do you drink milk?
- Have you lost or gained weight recently? If so, how much?

Activity/Exercise Patterns

- Do you exercise? How much and what type?

Recreation/Hobbies

- Do you participate in sports or activities that require heavy lifting?
- Do you wear protective equipment?
- Do you participate in outdoor activities, such as hiking or camping?

Sleep/Rest Patterns

- How many hours of sleep do you get per night?

Personal Habits

- Do you smoke?
- Do you drink alcohol or caffeinated beverages?
- Do you use drugs?

- Helps identify health promotion and preventive activities.

- Musculoskeletal problems, such as arthritis, may affect ability to perform ADLs.
- Sedentary lifestyle increases risk of osteoporosis. Prolonged immobility leads to muscle wasting.
- Impairment of musculoskeletal system may impair ability to perform normal activities. Assess for correct use of assistive devices.

- Assess diet for adequate vitamin/mineral intake:
 - Calcium deficiency increases risk of osteoporosis.
 - Adequate protein promotes healthy muscle tone and bone growth.
 - Diet high in purines (organ meats and sardines) can trigger gouty arthritis.
- Obesity is risk factor for osteoarthritis and low back problems.
- Weight loss is associated with RA.

- Exercise is important to a healthy musculoskeletal system. Moderate exercise builds up and maintains bone mass.
- Strenuous exercise increases risk for injury.
- Regular exercise promotes strength and flexibility and can help slow musculoskeletal changes associated with aging.

- Recreational activities may increase risk for injury.
- Use of protective equipment decreases risk of trauma.
- Lyme disease is caused by deer ticks.

- Musculoskeletal pain may interfere with sleep.
- Patients with herniated discs may have trouble finding comfortable sleeping position.

- Cigarette smoking increases risk of osteoporosis.
- Alcohol and caffeine are risk factors for osteoporosis and gout.
- Drug use increases risk of HIV, which is associated with inflammatory arthritis.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Occupational Health Patterns <ul style="list-style-type: none"> ■ What type of work do you do? Does it involve lifting or repetitive movements? ■ How would you describe your usual posture? ■ What kind of shoes do you wear to work? ■ Have you ever lost work time because of a musculoskeletal problem? ■ Are you exposed to toxic chemicals at work? 	<ul style="list-style-type: none"> ■ Occupations that require physical labor increase risk for musculoskeletal problems. ■ Repetitive motion increases risk for carpal tunnel syndrome. ■ Poor body posture and body mechanics can lead to back problems. ■ High-heeled shoes can lead to contracture of the Achilles tendon. ■ Musculoskeletal problems may prevent patient from returning to previous work. ■ Arsenic and mercury can cause ataxia.
Environmental Health Patterns <ul style="list-style-type: none"> ■ Where do you live? ■ How many stairs do you have to climb? 	<ul style="list-style-type: none"> ■ Helps determine discharge planning needs.
Roles/Relationships/Self-Concept <ul style="list-style-type: none"> ■ How did you view yourself before you had this problem, and how do you view yourself now? ■ Have musculoskeletal problems interfered with your ability to interact with others? 	<ul style="list-style-type: none"> ■ Musculoskeletal deformities or disabling or crippling problems may affect patient's self-image and cause low self-esteem. ■ Chronic musculoskeletal problems can disable the patient and impair socialization and independence.
Cultural Influences <ul style="list-style-type: none"> ■ What is your ethnic background? 	<ul style="list-style-type: none"> ■ Osteoporosis has a higher incidence among Caucasians. ■ Asians and African Americans have 10 percent more bone density than whites. ■ Back problems, joint pain, and stiffness may interfere with sexual activities.
Sexuality Patterns <ul style="list-style-type: none"> ■ Have musculoskeletal problems interfered with your usual sexual activity? 	<ul style="list-style-type: none"> ■ Identifying supports is helpful in planning care.
Social Supports <ul style="list-style-type: none"> ■ Who are your supports? 	<ul style="list-style-type: none"> ■ Stress can exacerbate SLE and precipitate RA.
Stress and Coping <ul style="list-style-type: none"> ■ How do you deal with stress? 	



Case Study Findings

Reviewing Mrs. O'Malley's

Biographical data:

- 68-year-old Caucasian female, widow.
- Works as housekeeper for parish priest.
- Born and raised in the United States, Polish descent, Catholic religion.
- *Current health insurance:* Medicare parts A and B, supplemented with Medicaid.
- Source: Self, seems reliable.

Current health status:

- "Several-year" history of aching joints in the evening after working all day.
- Discomfort first occurred in back and neck.
- Takes over-the-counter (OTC) analgesic for pain with some relief.

Past health history:

- History of hypertension (HTN).
- Wrist fracture at age 65, caused by a fall on the ice. No pinning was performed.
- Cholecystectomy at age 52, appendectomy at age 8, hysterectomy at age 35.
- Vaginal deliveries of two children with no complications.
- Denies allergies to food, drugs, or environmental factors.
- Medications include Zestril 10 mg once daily for HTN and Alleve once daily for arthritic pain.

Family history:

- Positive history of osteoporosis and HTN.
- Mother had osteoporosis and HTN; died of cardiac problems at age 82.



Case Study Findings (continued)

- Father had HTN and died of heart attack at age 65.
- Sister, age 72, positive for osteoporosis and HTN.
- One daughter and one son, alive and well.
- All other known relatives alive and well.

Review of systems:

- *General Health Survey:* Describes usual state of health as “fair.”
- *Integumentary:* Reports no changes in skin, hair, or nails.
- *Head, Eyes, Ears, Nose, and Throat (HEENT):* Complains of pain and stiffness in neck; no swollen glands.
- *Respiratory:* No respiratory problems.
- *Cardiovascular:* Positive history of HTN, controlled with medication.
- *Gastrointestinal:* No abdominal problems.
- *Genitourinary:* No problems.
- *Reproductive:* Postmenopausal since age 35.
- *Neurological:* No changes in sensations.
- *Endocrine:* No history of diabetes or thyroid disease.
- *Immunologic/Hematologic:* No infections or bleeding.

Psychosocial profile:

- Has a yearly exam because of HTN. Has not seen a gynecologist since her hysterectomy. Visits family physician every 3 months for HTN check.
- Arises at 6 A.M. Does chores around own home and dresses for work. Workday is 9 A.M. to 5 P.M. Typical day consists of daily chores around employer’s home, including cooking, cleaning, and running errands. After work, she relaxes and watches the news and some television shows before going to bed at 9 P.M.
- Admits she is a “tea and toast lady.” Lives alone, so if she cooks, it is something quick, unless she goes out to eat with friends or family. Drinks two to four cups of caffeinated coffee a day. Has never been overweight.
- Activities include housecleaning and running errands uptown. No formal exercise program.
- Has no pets. Knits for a hobby while watching TV. Occasionally has dinner out or sees a movie with friends or family.
- Sleeps 4 to 5 hours uninterrupted, but often gets up at night to void.
- Denies smoking, alcohol use, or illicit drug use.
- Job requires some lifting (e.g., carrying vacuum cleaner upstairs) as well as bending.

- Says her self-esteem is “pretty good.” Says she feels proud she can still earn a living. Admits she worries about living alone and developing health problems as she gets older, especially those that limit her mobility.
- Lives alone in a ranch home in suburbs; no steps. Is not afraid to be alone because of good neighbors.
- Has lunch or dinner once a week with daughter or son. Talks on phone to sister several times a week. Has friends in neighborhood, including younger neighbors who check on her.
- Copes with stress by talking with children or priest.



Case Study Evaluation

Before you move on to the physical examination, consider what you have learned from your interview with Mrs. O’Malley and document her musculoskeletal history.



CRITICAL THINKING ACTIVITY # 1

Based on her health history, what musculoskeletal problem do you think Mrs. O’Malley is at risk for? What are the risk factors?

Physical Assessment

Now that you have collected the subjective data, begin collecting the objective data by means of the physical examination. Keep all pertinent history findings in mind as you proceed. These findings, along with those from the physical examination, will complete the assessment picture. Then you can analyze the data, formulate nursing diagnoses, and develop a plan of a care. Because a musculoskeletal problem might be present, you will need to know normal musculoskeletal function before you can determine abnormal findings. The anatomy will direct your assessment.

Approach

Physical assessment of the musculoskeletal system provides data on the patient’s posture, gait, cerebellar function and bone structure, muscle strength, joint mobility, and ability to perform ADLs. You will use inspection, palpation, and percussion to assess the musculoskeletal system. Be systematic in your approach, working from head to toe and comparing one side with the other. Inspect and palpate each joint and muscle; then assess range of motion (ROM) and test muscle strength.

Perform inspection and palpation simultaneously during this assessment.



TOOLBOX

You will use a tape measure and a goniometer. Use the tape measure to measure limb lengths and circumferences. Use the goniometer to measure joint ROM (Fig. 20.5). To obtain a goniometer reading, match the angle of the joint being measured to the arms of the goniometer; then describe the limited motion of the joint in degrees.



FIGURE 20.5. Using a goniometer.

Performing a General Survey

Before examining specific areas, scan the patient from head to toe, looking for general appearance and signs of musculoskeletal problems. For example:

- Obtain the patient's vital signs. Elevations in temperature are associated with Lyme disease, RA, and infections such as osteomyelitis.
- Be alert for signs of pain or discomfort as your patient performs ROM. If signs or symptoms of pain are present with movement, never force a joint.

Performing a Head-to-Toe Physical Assessment

Now look for changes in every system that might signal a musculoskeletal problem.

Performing a Musculoskeletal Physical Assessment

Now that you have completed a head-to-toe scan, focus on the specifics of the examination. Proceed in this order:

- Assess posture, gait, and cerebellar function.
- Measure limbs.
- Assess joints and test joint movement.
- Assess muscle strength and ROM.
- Perform additional tests to assess for wrist, spine, hip, and knee problems, if necessary.

Assessing Posture, Gait, and Cerebellar Function

Posture

Begin by assessing your patient's **posture**, or his or her position in relation to the external environment. Posture should be erect, with the head midline. The spine is best assessed with the patient standing. Inspect from the front, back, and side, looking for alignment and noting symmetry of shoulders, scapula, and iliac crests. Adults have four normal curvatures. The cervical and lumbar curves are concave, and the thoracic and sacral curves are convex. When assessing for spinal curvature deformities, it is important to determine if the deformity is structural or functional (postural) in nature.

To assess for kyphosis and scoliosis, have the patient bend forward from the waist with her or his arms relaxed and dangling. Then stand behind the patient to check for curvatures. The curvatures disappear in functional kyphosis and scoliosis but remain in the structural form. To assess for lordosis, have the patient stand against a wall and flatten her or his back against it. If you can freely slide your hand through the lumbar curve, the patient has structural lordosis. An alternative method is to have the patient lie supine with the knees slightly flexed and attempt to flatten her or his lumbar curve. If she or he can, the lordosis is functional.

Next, have your patient stand erect with the feet together and note the position of the knees. Are the knees midline? Draw an imaginary line from the anterior superior iliac crest to the feet. Normally, this line should transect the patella. Note if the patella deviates away from the midline. Normally, the knees should be less than 2.5 cm (1") apart and the medial malleoli (ankle bones) less than 3 cm (1½") apart.

Gait

Next, assess your patient's gait. Pay particular attention to the base of support (distance between the feet) and stride length (distance between each step). Is there a wide base of support? The average base of support for an adult is about 2 to 4 inches. The average stride length for an adult is about 12 to 14 inches. Stride length depends in part on leg length, so the longer the legs, the greater the stride length.



HELPFUL HINT

A wide base of support and shortened stride length often reflect a balance problem.

Also pay attention to the phases of the gait—stance and swing. The stance phase has four components: heel strike, foot flat, midstance, and push-off. During midstance, the weight shifts to bearing the full body weight. The swing phase has three components: accelerate, swing-

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

General Health Survey

INSPECT

- Well-developed; no acute distress.

Integumentary

INSPECT

- No rashes or lesions; hair evenly distributed; no alopecia.

HEENT

Head, Face, Neck

INSPECT/PALPATE

- Facial expression appropriate; thyroid and nodes not palpable.

Eyes

- Eyes clear and bright.

Respiratory

PALPATE/AUSCULTATE

- Respiratory rate 18 breaths/min. Lungs clear.

Cardiovascular

AUSCULTATE

- Regular heart rate and rhythm.

Gastrointestinal

INSPECT/PALPATE/AUSCULTATE

- Positive bowel sounds; abdomen soft, nontender.

Genitourinary/Reproductive

INSPECT

- No lesions.

Neurologic

INSPECT

- Sensory intact; no numbness, paralysis, or parasthesia.

ABNORMAL FINDINGS/RATIONALE

- Obesity associated with osteoarthritis.

- *Bull's-eye rash*: Lyme disease.
- *Butterfly rash, patchy alopecia*: SLE.

- *Masklike facial expression*: Parkinson's disease.
- *Enlarged lymph nodes*: Infection, RA.
- *Weakness, pain with weight bearing, osteoporosis, decreased muscle tone*: Hyperthyroidism.
- *Muscle weakness and cramps, difficulty walking*: Hypothyroidism.

- *Red eye (episcleritis)*: RA.

- *Interstitial fibrosis*: RA.
- *Pleural effusion, pneumonitis*: SLE.

- *Pericarditis, myocarditis, Raynaud's disease*: RA and SLE.

- *Gastrointestinal complaints*: SLE.

- STDs, such as syphilis, can cause tabes dorsalis, ataxia, areflexia, and damaged joints (Charcot's joint).

- Decreased sensations and numbness associated with neuropathies.
- Paralysis can result in muscle atrophy.

through, and decelerate. Figure 20.6 illustrates the two phases and their components. If you do not detect a gait problem, chart “phases conform.” If you do detect a problem, try to identify the specific portion of the gait that is abnormal. For example, someone with left-sided weakness from a stroke may have problems with the acceleration portion of the swing phase and the midstance portion of the stance phase. Continue to observe your patient's gait, taking note of posture and cadence as he or she walks. Ask yourself: Do the arms swing in opposition? Is there toeing in or out? Are movements smooth and coordinated?



HELPFUL HINT

- Inspect the soles of your patient's shoes. How they wear is a good indicator of gait problems.
- When performing the Romberg test, stand nearby in case the patient loses her or his balance.

Cerebellar Function

Now, assess cerebellar function, including balance, coordination, and accuracy of movements. (See *Assessing Posture, Gait, and Cerebellar Function*.)

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> General health Weight gain Height loss Fever	General	<i>Measure:</i> Vital signs Height and weight
Integumentary	<i>Ask about:</i> Rashes Hair loss	Integumentary	<i>Inspect:</i> Rashes, lesions, alopecia
HEENT	<i>Ask about:</i> Swollen glands Dry, red eyes	HEENT	<i>Inspect:</i> Eyes for redness <i>Palpate:</i> Lymph nodes, thyroid
Respiratory	<i>Ask about:</i> Breathing difficulty	Respiratory	<i>Auscultate:</i> Lungs
Cardiovascular	<i>Ask about:</i> History of CV problems	Cardiovascular	<i>Auscultate:</i> Heart sounds
Gastrointestinal	<i>Ask about:</i> Nausea, vomiting, diarrhea	Gastrointestinal	<i>Auscultate:</i> Bowel sounds
Genitourinary/ Reproductive	<i>Ask about:</i> History of STDs <i>Menopausal women:</i> HRT	Genitourinary/ Reproductive	<i>Inspect:</i> Lesions
Neurological	<i>Ask about:</i> Changes in sensation Numbness, tingling	Neurological	<i>Test:</i> Sensory deficits, paralysis
Endocrine	<i>Ask about:</i> History of diabetes mellitus, thyroid disease		
Lymphatic/ Hematologic	<i>Ask about:</i> Bruising		

Assessment of Musculoskeletal System's Relationship to Other Systems




FIGURE 20.6. *Stance phase: A. Heel strike. B. Foot flat. C. Midstance. D. Push-off. Swing phase: E. Accelerate. F. Swing-through. G. Decelerate.*

Balance

To assess balance, look at your patient's gait. If he or she has a gait problem, you will be unable to proceed further. If there are no gait problems, have the patient tandem (heel-to-toe) walk, heel-and-toe walk, hop in place, do a deep knee bend, and perform the Romberg test. To perform the Romberg test, have the patient stand with feet together and eyes open; then have him or her close the eyes. If cerebellar function is intact, he or she will be able to maintain balance with minimal swaying with eyes closed (negative Romberg test). Keep in mind that balance problems may also occur with inner ear disorders.

Coordination

Next, assess coordination. Note the patient's dominant side. Usually this side is more coordinated. To test upper extremity coordination, have the patient perform rapid alternating movements (RAMs) by patting her or his thigh with one hand, alternating between the supinate and the pronate hand position. Have her or him perform finger-thumb opposition to further test hand coordination. To test lower extremity coordination, have the patient perform rhythmic toe tapping and then run the heel of one foot down the shin of the opposite leg. Remember to test each side separately and compare findings.

Accuracy of Movements

Point-to-point localization is used to assess accuracy of movements. Have the patient touch his or her finger to

the nose with the eyes open, then closed. Or have him or her touch your finger at various positions.

Taking Limb Measurements

Limb measurements include both length and circumference. Measure arm lengths from the acromion process to the tip of the middle finger. Measure leg lengths from the anterior superior iliac crest crossing over the knee to the medial malleolus. This is referred to as **true leg length**. An **apparent leg length** measurement is measured from a nonfixed point, the umbilicus, to the medial malleolus. A true leg length discrepancy results from bone length inequality, whereas an apparent leg length discrepancy may result from a pelvic tilt or flexion deformity of the hip. Discrepancies between right and left should not exceed more than 1 cm. Obviously, a leg length discrepancy usually results in clinical problems such as gait disorders or hip or back pain.

Limb circumference reflects actual muscle size or muscle mass. Measure circumference on forearms, upper arms, thighs, and calves. Take note of the patient's dominant side, which may normally be up to 1 cm larger than her or his nondominant side. To ensure accurate circumference measurement, determine the midpoint of the extremity being assessed. For example, if you are measuring upper arm circumference, measure the distance from the acromion process to the olecranon process, and use the midpoint to determine circumference; then do the same for the opposite arm. (See *Taking Limb Measurements*.)

ASSESSING POSTURE, GAIT, AND CEREBELLAR FUNCTION

1. Posture

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Inspect position, posture.
- Inspect normal curves of the spine with the patient standing.



Assessing for normal curves

- If you suspect a spinal deformity, test for kyphosis and scoliosis by having patient bend at the waist and assume a “dive” position. Stand behind patient and inspect and palpate spinal curvature.



Assessing for kyphosis and scoliosis

- **Test for lordosis:** Have the patient stand against a wall and flatten her or his back against it while you attempt to slide your hand behind patient's back.
Alternate way to test for lordosis: With patient supine and knees slightly flexed, instruct patient to flatten back against mattress while you attempt to slide hand under back.

- **Assessing for knee deviation:** Note position of knees, and draw an imaginary line from the anterior superior iliac crest through the knees to the feet. Imaginary line should transect patella.



Assessing for lordosis



Assessing for deviation of the knees

ASSESSING POSTURE, GAIT, AND CEREBELLAR FUNCTION

1. Posture (continued)

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- *Posture erect, head midline, four normal spinal curves.*
- *In pregnancy, accentuated lumbar curve (lordosis) with a compensatory cervical flexion is normal to compensate for change in center of gravity.*
- *In newborns, C-shaped spine without cervical curve is normal. Cervical curve develops once infant begins to hold head upright.*
- *In toddlers, lordosis is normal as they learn to walk.*
- *Senile kyphosis (dowager's hump) may occur in older patients.*



Pregnancy lordosis



Senile kyphosis

- *Some African Americans have accentuated lumbar curves.*
- *Knees midline.*
- *Genu valgum and genu varum often seen in toddlers and preschoolers, but should not persist beyond age 7.*

(continued)

ASSESSING POSTURE, GAIT, AND CEREBELLAR FUNCTION

1. Posture (continued)

ABNORMAL FINDINGS/RATIONALE

- *Spinal deformities include:*
 - *Kyphosis:* Accentuated thoracic curve.
 - *Scoliosis:* Lateral “S” spinal deviation.



Scoliosis



HELPFUL HINT

To differentiate structural from functional scoliosis, have patient bend at waist. In true structural scoliosis, deviation is apparent. In functional scoliosis, deviation disappears.

- *Lordosis:* Accentuated lumbar curve.



Lordosis

- *Genu valgum:* Knees touch and medial malleoli are 7.5 cm (3 inches) or more apart.
- *Genu varum:* Knees are greater than 5 cm (2 inches) apart and medial malleoli touch.

2. Gait

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Inspect the patient as he or she walks.
- Note posture, head position, conformity of phases, toe position, arm swing, cadence, coordination of movements, stride length, and base of support.



Observing gait



HELPFUL HINT

To inspect gait, observe patient as she or he walks into the examination room. (See *Abnormal Gaits*.)

- *Posture erect, head midline, phases conform, weight evenly distributed, both feet point straight ahead, no toeing in or out, all movements coordinated and rhythmic, arms swing in opposition, stride length appropriate.*
- *In pregnancy, with older patients, and in toddlers learning to walk, wider base of support gives greater stability and is normal.*

ASSESSING POSTURE, GAIT, AND CEREBELLAR FUNCTION

2. Gait (continued)

ABNORMAL FINDINGS/RATIONALE

- *Uneven weight bearing*: Associated with joint pain.
- *Gait disorders such as wide base of support*: Associated with cerebellar dysfunction.
- *Ataxia (impaired coordinated movement with erratic muscular activity), spasticity (increased tone or contractions of muscles causing stiff, awkward movements), and tremors*: Parkinson's disease, MS, cerebral palsy.
- *Scissors gait (legs cross over)*: Disorders of motor cortex or corticospinal tracts, such as bilateral spastic paresis.
- *Spastic movements*: Upper motor neuron disorders.
- *Flaccidity (defective or absent muscle tone)*: Lower motor neuron disorders.
- *Flaccidity and foot drop (affected leg drags and is unable to clear during swing phase)*: Peripheral nerve disorders.

3. Cerebellar Function

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

Balance

- Inspect gait, and note balance.
- Have patient tandem walk, heel to toe.
- Have patient heel-and-toe walk.



Tandem walking

Heel-and-toe walking.
A. Walking on heels.

B. Walking on toes

- Have patient do deep knee bend.



Deep knee bend

- Have patient hop in place.



Hopping in place

(continued)

ASSESSING POSTURE, GAIT, AND CEREBELLAR FUNCTION

3. Cerebellar Function (continued)

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- **Perform the Romberg test:** Have patient stand with feet together, arms at side, eyes open and then with eyes closed. Stand close by in case the patient loses balance.
- **Balance intact.** Patient can tandem walk, heel-and-toe walk, perform deep knee bend, and hop in place. *Negative Romberg test.*
- **Cerebellar function cannot be tested on infants under 3 months.** Instead, observe sucking, swallowing, and kicking movements to estimate cerebellar function.
- **Cerebellar function cannot be tested on toddlers because of their immature neuromuscular system.**



Romberg test. A. Eyes open.



B. Eyes closed.

Coordination

- **Determine patient's dominant side.**
- **Test upper extremities:** Have patient perform RAMS by patting thigh with one hand, alternating with supination and pronation.



RAMs. A. Pronate position



B. Supinate position

- **Have patient perform finger-thumb opposition.**



Finger-thumb opposition

- **Test lower extremities:** Have patient perform rhythmic toe tapping or run heel of foot down shin of opposite leg.



Toe tapping



Running heel down shin

ASSESSING POSTURE, GAIT, AND CEREBELLAR FUNCTION

3. Cerebellar Function (continued)

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- *Upper and lower extremity movements coordinated. Right side dominant and slightly more coordinated. Positive RAM.*

- *Positive finger-thumb opposition, toe tapping, and heel down shin.*

Accuracy of Movement

- **Have patient perform point-to-point localization:** Finger to nose, or while you stand in front of patient, hold your finger about 12 inches in front and instruct patient to touch your finger, repeat at various positions.



Finger to finger



Finger to nose

- *Movements accurate.*

ABNORMAL FINDINGS/RATIONALE

- *Balance problems:* Cerebellar disorder.

- *Positive Romberg test:* Cerebellar disorder if patient has difficulty maintaining balance with eyes open or closed. If the patient loses balance only when eyes are closed, there may be damage to the dorsal column.

- *Slowness and awkwardness in performing movements:* Cerebellar disorder or motor weakness associated with extrapyramidal disease.

- *Inaccurate movements:* Cerebellar disorder.

Abnormal Gaits

TYPE OF GAIT

DESCRIPTION/CAUSE

Propulsive gait



- Rigid, stooped posture with head leaning forward and arms, knees, and hips stiffly flexed.

- Rapid, short, shuffling steps.

- *Causes:* Parkinson's disease (classic gait).

(continued)

Abnormal Gaits (continued)

TYPE OF GAIT

DESCRIPTION/CAUSE

Scissors gait




- Bilateral spastic paresis of legs; arms not involved.
- Legs flexed at hip and knees.
- Knees adduct and meet or cross like scissors.
- Short steps, foot plantarflexed, walks on toes.
- *Causes:* Cerebral palsy, MS, spinal cord tumors.

Spastic gait (hemiplegic)



- Unilaterally stiff, dragging leg from leg, muscle hypertonicity.
- *Causes:* Stroke, MS, brain tumor.

Waddling gait




- Ducklike walk with wide base of support, chest thrown back, exaggerated lumbar curve (lordosis), and protruding abdomen.
- Normal in toddlers and late stages of pregnancy.
- Weak pelvic girdle muscles (gluteus medius, hip flexors, and extensors).
- *Causes:* MS, hip dislocation.

TYPE OF GAIT

DESCRIPTION/CAUSE

Steppage gait (equine, prancing, parietic, or weak)



- Foot drop with external rotation of hip and hip and knee flexion. Foot slaps when it hits ground.
- *Causes:* MS, herniated lumbar disc, Guillain-Barré syndrome, perineal muscle atrophy, or nerve damage.

TAKING LIMB MEASUREMENTS

1. Length

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Arms

- Measure from the acromion process to the tip of the middle finger.

Legs

- Measure from the anterior superior iliac crest, crossing over knee to the medial malleolus.



Measuring arm length



Measuring leg length. A. Apparent



B. True

(continued)

TAKING LIMB MEASUREMENTS

1. Length (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Arm and leg lengths ≤ 1 cm difference between right and left.*

ABNORMAL FINDINGS/RATIONALE

- Leg length discrepancies can cause back and hip pain, gait problems, and pseudoscoliosis, or apparent scoliosis.
- Equal true leg lengths but unequal apparent leg lengths are seen with hip and pelvic area abnormalities.

2. Circumference

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Determine patient's dominant side.
- Measure midpoint of extremity and measure circumference.



Measuring midpoint of arm



Measuring arm circumference



Measuring thigh circumference

- *Arm and leg circumferences ≤ 1 cm difference between right and left.*

ABNORMAL FINDINGS/RATIONALE


- Circumference differences > 1 cm may reflect muscular atrophy or hypertrophy.

Assessing Joints and Muscles


Inspect the size, shape, color, and symmetry of each joint, noting masses, deformities, and muscle atrophy. Compare muscle and joint findings bilaterally and palpate for edema, heat, tenderness, pain, nodules, crepitus, and

stability. Test active ROM (in which the patient performs the movement) and passive ROM (in which you put the joint through the ROM) of the joint (Fig. 20.7). Changes in articular cartilage, scarring of the joint capsule, and muscle contractures all limit motion. It is important to


Depression: Lowering a body part
Elevation: Raising a body part



Retraction: moving backward
Protraction: moving forward




Circumduction: moving in a circular fashion




Opposition

Resposition



Extension: straightening, increasing the joint angle



Flexion: bending, decreasing the joint angle

Inversion: turning inward

Eversion: turning outward



Pronation: turning downward



Supination: turning upward




External rotation: turning away from midline

Internal rotation: turning toward midline



Abduction: moving away from midline



Adduction: moving toward midline

FIGURE 20.7. Body movement.

determine the types of motion the patient can no longer perform, especially the ADLs.



ALERT

Never force a joint.

Joint clicking or crepitus may occur normally sometimes or may be associated with dislocations of the humerus, temporomandibular joint (TMJ) problems, displacement of the biceps tendon from its groove, damaged knees, or DJD.

Joint deformity may result from a congenital malformation or a chronic condition. In any patient with a deformity, ask the following questions:

- When did you first notice the deformity?
- Did the deformity occur suddenly?
- Did it occur as the result of trauma?
- Has the deformity changed over time?
- Has it affected your range of motion?

Test muscle strength by asking the patient to put each joint through ROM while you apply resistance to the part being moved. If the patient is unable to move the part against resistance, then ask him or her to perform the ROM against gravity. If this cannot be performed, attempt to passively put the muscle through ROM. If this is not possible (because of a contracture), palpate the muscle while the patient attempts to move it. Document muscle strength as shown in the box entitled Rating Scale for Muscle Strength.

Begin your assessment by inspecting and palpating the muscles of the upper and lower extremities. Assess muscle tone and mass in both relaxed and contracted state, and then assess each joint and test muscle strength as the patient performs ROM.

Additional Tests

You can perform the following additional tests to assess for wrist, spinal, hip, and knee problems.

Tests for Wrist Problems

Phalen's Test. Have the patient flex the hands back to back at a 90-degree angle and hold this position for about

Joint Movements

MOVEMENT	DESCRIPTION
Extension	Straightening or increasing the angle of a joint
Flexion	Bending or decreasing the angle of a joint
Hyperextension	Straightening beyond the normal angle of a joint
Abduction	Moving away from the midline
Adduction	Moving toward the midline
Circumduction	Moving in a circular fashion
Internal rotation	Turning inward toward the midline
External rotation	Turning outward away from the midline
Pronation	Turning down
Supination	Turning up
Inversion	Turning inward (medially)
Eversion	Turning outward (laterally)
Retraction	Moving backward
Protraction	Moving forward
Opposition	Movement of thumb toward fingerpad
Reposition	Movement of thumb back to anatomic position
Depression	Movement by lowering body part
Elevation	Movement by raising body part
Ulnar deviation	Movement of hand toward ulnar side
Radial deviation	Movement of hand toward radial side

1 minute (Fig. 20.8). If the patient complains of numbness or tingling anywhere from the thumb to the ring finger, the test is positive for carpal tunnel syndrome.

Tinel's Test. Percuss lightly over the median nerve, located on the inner aspect of the wrist (Fig. 20.9). If numbness and tingling occur on the palmar aspect of the wrist and extend from the thumb to the second finger, the test is positive for carpal tunnel syndrome.

(Text continued on page 728)

Rating Scale for Muscle Strength

RATING SCALE	EXPLANATION	CLASSIFICATION
5	Active motion against full resistance	Normal
4	Active motion against some resistance	Slight weakness
3	Active motion against gravity	Average weakness
2	Passive ROM (gravity removed and assisted by examiner)	Poor ROM
1	Slight flicker of contraction	Severe weakness
0	No muscular contraction	Paralysis



FIGURE 20.8. Phalen's test.



FIGURE 20.9. Tinel's test.

ASSESSING JOINTS AND MUSCLES

1. Muscles

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- **Tone:** Palpate muscles of upper and lower extremities in relaxed and contracted state.
- **Strength:** Screen muscle strength of upper and lower extremities.
- **Upper extremities:** Test hand grip by crossing your index and middle fingers and asking patient to squeeze.
- **Lower extremities:** Have patient raise leg against your hand as you apply resistance.
- **Comprehensive muscle strength:** As the patient performs ROM, apply resistance to part being moved, then grade strength 0 to 5.



Palpating muscle tone in lower extremities

Upper and Lower Extremities

- Inspect and palpate for muscle mass and tone in relaxed and contracted state.
- *Muscles soft, pliable, and nontender in relaxed state; firm and nontender in contracted state. No abnormal movements.*

ABNORMAL FINDINGS/RATIONALE

- Atrophy (muscle wasting), unexplained hypertrophy (excessive muscle size), flaccidity (atony), weakness (hypotonicity), fasciculations (involuntary twitching of muscle fibers), or tremors (involuntary contraction of muscles).

(continued)

ASSESSING JOINTS AND MUSCLES

2. Joints

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Test ROM of joints, and note condition of skin, erythema, edema, heat, deformity, crepitus, tenderness, and stability of all joints.
- Demonstrate movement of each joint for patient, then have patient return movement.

Temporomandibular Joint (Hinge Joint and Gliding Joint)

- Inspect, palpate, and test ROM and muscle strength.
- Assess flexion (depression), extension (elevation), side-to-side movement, protraction (pushing out), and retraction (pulling in) of jaw.
- Palpate strength of masseter and temporalis muscles as patient clenches teeth.
- Ask patient to laterally move the jaw left and right and then open the mouth against resistance. Feel for contraction of temporal and masseter muscles to test integrity of cranial nerve V (trigeminal).



Assessing the TMJ

■ **Normal ROM:**

- *Normal opening 1–2 cm with ease.*
- *Lateral movement 1–2 cm. Snapping and clicking are normal.*
- *+5 muscle strength, no pain or spasms on contraction.*

■ Sternoclavicular Joint (Gliding Joint)

- Inspect and palpate for location to midline, color, swelling, pain, and masses with patient sitting.
- *Joint midline with no visible bony overgrowth, swelling, or redness; nontender.*

Cervical, Thoracic, and Lumbar Spine (Gliding Joint for Vertebrae, Pivotal Joint for Atlantoaxial Joint)



- **Inspect and palpate spinous processes and paravertebral muscles for tenderness or pain.**
- **Test ROM.**

Cervical Spine

- **Test ROM by having patient flex (touch chin to chest), extend (put head in neutral position), hyperextend (look up), bend laterally, put ear to shoulder and rotate, and turn head side to side. To test muscle strength, repeat against resistance.**



Testing ROM and muscle strength of neck



A–B. Normal cervical spine ROM

ASSESSING JOINTS AND MUSCLES


2. Joints (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Joints midline with no visible bony overgrowth, swelling, or redness; nontender.
- Nontender spinous processes; paravertebral muscles well developed, smooth, firm, and nontender.
- **Normal ROM:**
 - 55 degrees of hyperextension.
 - 45 degrees of flexion.
 - 40 degrees of right and left lateral bending.
 - 70 degrees of left and right rotation.
 - +5 muscle strength.

Thoracic and Lumbar Spine

- Test ROM by having patient flex (bend at waist), extend (stand upright), hyperextend (bend backward), bend laterally (side to side), and rotate (turn upper body side to side). To test muscle strength, repeat against resistance.




C. Normal cervical spine ROM



HELPFUL HINT

Stand behind patient and stabilize his or her pelvis with your hands as he or she performs ROM of thoracic and lumbar spine.



A–C. Normal thoracic and lumbar spine ROM

(continued)

ASSESSING JOINTS AND MUSCLES

2. Joints (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

■ **Normal ROM:**

- 75 to 90 degrees of flexion, smooth movement.
- 30 degrees of hyperextension.
- 35 degrees of left and right lateral bending.
- 30 degrees of left and right rotation.

Shoulders (Ball and Socket Joint)

- Inspect for symmetry, swelling, color, and masses.
- Palpate for tenderness.
- Test ROM by having patient flex (move arms forward), extend (move arms backward with elbows straight), abduct (bring both hands together over head with elbows straight), and adduct (bring both hands in front of body past midline with elbows straight). Also have patient perform external rotation (with elbows flexed, bring hands behind head) and internal rotation (with elbows flexed, bring arms behind back).
- **Test muscle strength of shoulder by performing ROM against resistance.**
- *Symmetrically round; no swelling, redness, or deformity. Muscles fully developed. Clavicles and scapulae even and symmetrical. No tenderness, swelling, or heat. Flexion, extension, abduction, adduction, rotation, and shrug of shoulders should be performed without difficulty.*

■ **Normal ROM:**


- 180 degrees of forward flexion.
- 50 to 60 degrees of backward extension.
- 180 degrees of abduction.
- 45 to 50 degrees of adduction.
- 90 degrees of external rotation.
- 90 degrees of internal rotation.
- + 5 muscle strength.



Testing ROM of the shoulder



Testing muscle strength of the shoulder



ASSESSING JOINTS AND MUSCLES

2. Joints (continued)


ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Upper Arm and Elbow (Hinge Joint)

- Inspect for symmetry, swelling, color, and masses.
- Palpate for tenderness.
- Test ROM by having patient flex (bend arm at elbow), extend (straighten arm at elbow), supinate (with elbow flexed, turn arm with palm up), pronate (with elbow flexed, turn arm with palm down). To test muscle strength of upper arms and elbow, repeat against resistance.
- *Symmetrical; no redness or swelling; nontender without nodules.*
- **Normal ROM:**
 - 150 degrees of flexion.
 - 80 degrees of extension.
 - 90 degrees of supination and pronation.
 - + 5 muscle strength.



Testing ROM and muscle strength of the upper arm and elbow



Wrist (Condyloid Joint)

- Inspect and palpate for tenderness and nodules.

(continued)

ASSESSING JOINTS AND MUSCLES


2. Joints (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS


- Test ROM by having patient flex (bend wrist down), extend (straighten wrist), hyperextend (bend wrist back), perform ulnar deviation (move hand toward ulnar side), and perform radial deviation (move hand toward radial side).
- Test muscle strength by repeating against resistance.
- *Symmetrical; no redness or swelling; nontender without nodules.*
- **Normal ROM:**
 - 90 degrees of flexion.
 - 70 degrees of hyperextension.
 - 55 degrees of ulnar deviation.
 - 20 degrees of radial deviation.
 - Full ROM against resistance.
 - +5 muscle strength.



Testing ROM and muscle strength of the wrist



A



B

A and B. Normal wrist ROM

Hands and Fingers (Hinge, Saddle, and Condylloid Joints)

- Inspect for size, shape, symmetry, swelling, and color.
- Palpate for nodules and tenderness.
- Test ROM by having patient abduct (spread fingers apart), adduct (bring fingers together), flex (make a fist), extend (open fist), hyperextend (bend fingers back), perform palmar adduction (bring thumb to index finger), perform palmar abduction (move thumb away from index finger), and opposition and reposition.



Testing ROM of the hands and fingers



Palpating the finger joints

ASSESSING JOINTS AND MUSCLES

2. Joints (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Test muscle strength by repeating movements against resistance and hand grip.



HELPFUL HINT

When testing hand grip, cross your index and middle fingers and then have patient squeeze your crossed fingers.

- *Symmetrical fingers and hands; fingers lie straight; no swelling or deformities; nontender and without nodules.*

■ **Normal ROM:**

- 20 degrees of abduction.
- Adduction of fingers.
- 90 degrees of flexion.
- 30 degrees of hyperextension.
- Easily moves away from other fingers, palmar abduction.
- 50 degrees of flexion, palmar adduction.
- Full ROM against resistance.
- + 5 muscle strength.

Hips (Ball and Socket Joint)


- Inspect for symmetry and shape with patient standing.
- Palpate for stability, tenderness, tenderness, and crepitus.
- Test ROM by having patient extend (patient supine or standing, leg straight), flex (raise extended leg and flex knee to chest while keeping other leg extended), abduct (move extended leg away from midline of body as far as possible), adduct (move extended leg toward midline of body as far as possible), perform internal rotation (bend knee and turn leg inward) and external rotation (bend knee and turn leg outward), hyperextend (lie prone and lift extended leg off table), and stand and swing extended leg backward.
- Test muscle strength by repeating ROM against resistance.



Testing muscle strength of the hip



Testing muscle strength of the hands and fingers



Normal hand and finger ROM



Testing ROM of the hips

(continued)

ASSESSING JOINTS AND MUSCLES

2. Joints (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS



ALERT


Do not adduct, internally rotate, or flex more than 90 degrees on patients with hip replacements. Doing so may cause hip displacement.

■ *Buttocks equal in size; iliac crests symmetrical in height; nontender, stable, and without crepitus.*

■ **Normal ROM:**

- 0 degrees of extension.
- 90 degrees of flexion.

- 120 degrees of flexion with other leg remaining straight.
- 45 to 50 degrees of abduction.
- 20 to 30 degrees of adduction.
- 40 degrees of internal hip rotation.
- 45 degrees of external hip rotation.
- 15 degrees of hip hyperextension.
- Full ROM against resistance.
- + 5 muscle strength.



Normal hip ROM

ASSESSING JOINTS AND MUSCLES

2. Joints (continued)


ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Knees (Hinge Joint)

- Inspect knees for size, shape, symmetry, deformities, and swelling in both supine and sitting positions with knees dangling.
- Palpate for tenderness, warmth, consistency, and nodules, beginning approximately 10 cm above knee and moving downward over patella.
- Test ROM by having patient flex (bend each knee up to chest) and extend (straighten knee). Test muscle strength by repeating ROM against resistance.
- *Symmetrical; hollow in appearance on both sides of patella; no swelling or deformities; lower leg in alignment with upper leg; nontender and cool; muscles firm and no nodules apparent.*
- **Normal ROM:**
 - 120 to 130 degrees of flexion.
 - ROM not restricted against resistance.
 - 0 degrees of extension.
 - +5 muscle strength.



Testing ROM and muscle strength of the knee



Normal knee ROM

Ankles and Feet (Hinge, Gliding, and Condyloid)

- Inspect position, alignment, shape, and skin while patient is sitting, standing, and walking.
- Palpate for heat, swelling, tenderness, or nodules.
- Test ROM by having patient dorsiflex (point toes upward), plantar flex (point toes downward), evert (turn soles outward), invert (turn soles inward), abduct (rotate foot outward), and adduct (rotate foot inward).



Testing ROM of the ankles and feet



Testing muscle strength of the ankles and feet

- Test muscle strength by repeating ROM against resistance.

(continued)

ASSESSING JOINTS AND MUSCLES


2. Joints (continued)

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

■ *Toes and feet in alignment with lower leg; toes pointing forward and lying flat (may point in [pes varus] or out [pes valgus]); skin intact, smooth, and free of corns or calluses; longitudinal arch; weight bearing on foot midline; no heat, pain, nodules, or swelling.*

■ **Normal ROM:**

- 20 degrees of dorsiflexion.
- 45 degrees of plantarflexion.
- 20 degrees of eversion.
- 30 degrees of inversion.
- 10 degrees of abduction.
- 20 degrees of adduction.
- Full ROM against resistance.
- +5 muscle strength.



ABNORMAL FINDINGS/RATIONALE

- *Decreased ROM, tenderness, swelling, crepitus:* Arthritis.
- *Pain, swelling, popping, clicking, or grating sounds:* TMJ dysfunction.



ALERT

TMJ dysfunction may present as ear pain and headache.



Bursitis of the elbow

- *Decreased muscle strength:* Muscle and joint disease.
- *Pain and spasms:* Myofascial pain syndrome.
- *Less-than-full contraction:* Lesion of cranial nerve V.
- *Swelling, redness, enlargement, tenderness:* Inflammation.
- A neck that is not straight and erect is abnormal.
- *Inability to perform ROM because of pain:* Cervical disc degenerative disease, spinal cord tumor. Pain may radiate to back, shoulder, or arms.
- *Neck pain associated with weakness/loss of sensation in legs:* Cervical spinal cord compression.
- *Inability to perform ROM against resistance:* Muscle and joint disease.
- *Weakness and limited ROM:* Torn rotator cuff.
- *Redness, swelling, and tenderness at elbow (olecranon process):* Bursitis.
- **Tennis elbow** (lateral epicondylitis) is inflammation of forearm extensors or supinator muscles of fingers and wrist, and/or tendon attachment to lateral epicondyle or lateral collateral ligament caused by repetitive supination of forearm against resistance.
- **Golf elbow** (medial epicondylitis) is same as tennis elbow, except flexor and pronator muscles and tendons are affected.
- *Swelling, tenderness, nodules, ulnar deviation, limited ROM:* RA.
- *Nontender, round, enlarged, swollen, fluid-filled cysts on wrists:* Ganglion cyst.
- *Pain with movement:* Tendinitis
- *Pain on extension of wrist against resistance:* Epicondylitis.
- *Flexion of wrist against resistance:* Medial epicondylitis.
- *Decreased muscle strength:* Muscle and joint disease.

ASSESSING JOINTS AND MUSCLES

2. Joints (continued)

ABNORMAL FINDINGS/RATIONALE

- *Swollen, stiff, tender finger joints:* Acute RA.



Rheumatoid arthritis

- *Boutonnière deformity* (flexion of proximal interphalangeal joint and hyperextension of distal interphalangeal joint) and *swan-neck deformity* (hyperextension of proximal interphalangeal joint with flexion of distal interphalangeal joint) can both be seen in long-term RA.
- *Atrophy of thenar prominence:* Carpal tunnel syndrome.
- Heberden's nodes are hard, painless nodules over the distal interphalangeal joints.
- *Pain on extension of a finger:* Tenosynovitis (infection of the tendon sheath).
- *Inability to extend ring finger:* Dupuytren's contracture.
- *Decreased muscle strength:* Muscle and joint disease may present.
- *Unequal gluteal folds:* Dislocated hip.
- *Inability to abduct hip:* Common sign of hip disease.
- *Decrease in internal hip rotation:* Early sign of hip disease.
- *Decreased muscle strength against resistance:* Muscle and joint disease.
- *Tenderness, warmth, boggy consistency:* Synovitis.
- *Crepitation:* Osteoarthritis.
- *Decreased ROM:* Synovial thickening.
- *Inability to extend knee fully:* Flexion contracture of knee.
- *Decreased muscle strength against resistance:* Muscle and joint disease.



Heberden's nodes

- Bouchard's nodes are hard painless nodules over proximal interphalangeal joints.
- Both types of nodes are seen in osteoarthritis and RA.
- Gouty arthritis can cause deformities and nodules of the hands.



Gouty arthritis



DJD of the knees

- *Hallux valgus*, often on medial side, may present with laterally deviated great toe with overlapping of second toe. A *bunion* is an enlarged, painful, inflamed bursa that often occurs with hammer toe, which is hyperextension of metatarsophalangeal joint and flexion of proximal interphalangeal joint.



Hallux valgus, bunion, and hammer toe

(continued)

ASSESSING JOINTS AND MUSCLES

2. Joints (continued)

ABNORMAL FINDINGS/RATIONALE

- Flat feet (pes planus) have no arches or high arches (pes cavus).
- Corns are painfully thickened skin over bony prominences and pressure points.



Corns and bunions.

- A callus is painless, thickened skin that covers pressure points.



Callus

- Plantar warts (verruca vulgaris) are painful warts that occur under a callus.
- *Tender, painful, reddened, hot, swollen metatarsophalangeal joint of great toe:* Gouty arthritis.
- *Nodules on posterior ankle:* RA.
- *Pain and tenderness of metatarsophalangeal joints:* DJD, RA, joint inflammation.



HELPFUL HINT

Many foot problems are caused by poorly fitting shoes.

Test for Arm Problems

Pronator Drift. Perform this test if you detect muscle weakness of the arms. Have the patient stand with arms extended, hands supinated, and eyes open and then closed for at least 20 to 30 seconds (Fig. 20.10). Check for downward drift and pronation of the arms and hands. Pronation and drift of one arm is called *pronator drift* and may indicate a mild hemiparesis. Flexion of the fingers and elbow may accompany pronator drift. A lateral



FIGURE 20.10. Assessing for pronator drift.

and upward drift may also occur in patients with loss of position sense. If your patient is able to hold her or his arms extended without drift, gently tap downward on the arms. If she or he has normal muscle strength, coordination, and position sense, the arms will return to the horizontal position. A weak arm is easily displaced and does not return to the horizontal position.

You can also assess drifting and weakness by having the patient hold his or her arms over the head for 20 to 30 seconds. Then, try to force the arms down to the sides as the patient resists. Drifting or weakness may indicate a hemiparesis.

Test for Spinal Problems

Straight Leg Raising (Lasègue's Test). Perform this test when the patient complains of low back pain that radiates down the leg (sciatica). This test checks for a herniated nucleus pulposus. Ask the patient to lie flat and raise the affected leg to the point of pain (Fig. 20.11). Pain and sciatica that intensify with dorsiflexion of the foot are a positive sign for a herniated disc.



FIGURE 20.11. Straight leg raising.

Tests for Hip Problems

Thomas Test. This test assesses for hip flexure contractures hidden by excessive lumbar lordosis. Have the patient lie supine with both legs extended and then flex one leg to her or his chest (Fig. 20.12). The test is positive if the opposite leg raises off the table. Repeat the same maneuver on the opposite side.

Trendelenburg Test. This test is used to assess for a dislocated hip and gluteus medius muscle strength. Have patient stand erect and check the iliac crest—it should be level. Then, have the patient stand on one foot and check again. If the iliac crest remains level or drops on the side opposite the weight-bearing leg, the gluteus medius muscle is weak or the joint is not stable and there may be a hip dislocation on the weight-bearing side.

Tests for Knee Problems

Perform one of the following two tests if you noted swelling secondary to fluid accumulation or soft tissue damage.

Bulge Test. Perform this test if you suspect small amounts of fluid. With the patient supine, stroke the medial side of the knees upward several times to displace the fluid. Then, press the lateral side of the knee, and inspect for the appearance of a bulge on the medial side (Fig. 20.13).

Patellar Ballottement. Perform this test if you suspect large amounts of fluid. With the patient supine, press firmly with your left thumb and index finger on each side of the patella (Fig. 20.14). This displaces fluid into the



FIGURE 20.12. Thomas test.



FIGURE 20.13. Bulge test.

suprapatellar bursa between the femur and the patella. Then, gently tap on the kneecap. If fluid is present, the patella will bounce back to your finger (floating kneecap).



HELPFUL HINT

Check for hollows on either side of the patella. If hollows are absent, fluid may be present.

Lachman Test. If your patient complains that his or her knee gives way or buckles, test anterior, posterior, medial, and lateral stability. To test medial and lateral stability, have the patient extend the knee and attempt to abduct and adduct it. Normally, no movement should occur if the knee is stable. To assess the anterior and posterior plane, have the patient flex the knee at least 30 degrees. Stabilize and grasp the leg below the patella, and attempt to move it forward and back. If the joint is stable, no movement should occur (Fig. 20.15). Another similar test for stability of the anterior cruciate ligament (ACL) and posterior cruciate ligament (PCL) is the drawer test. With the patient's knee flexed at a 90-degree angle, apply anterior and posterior pressure against the tibia, and feel for movement. Movement of the tibia is a positive draw sign, indicating an ACL or PCL tear.

To test stability of the collateral ligament, have the patient lie supine with the knee slightly flexed. Place your hand at the head of the fibula and apply pressure medially; then reverse and apply pressure laterally. If the joint is



FIGURE 20.14. Patellar ballottement.



FIGURE 20.15. Lachman test.



FIGURE 20.17. Apley's test.

unstable, movement will occur and create a palpable medial or lateral gap at the joint.

Perform one of the following tests if your patient complains of clicks or knee-locking and you suspect a torn meniscus.

McMurray's Test. To perform McMurray's test, position the patient supine with her or his knee fully flexed (Fig. 20.16). Place one hand on the heel and the other on the knee and gently internally and externally rotate the foot as you bring the leg to full extension. The test is positive if audible or palpable clicks occur or the knee locks.

Apley's Test. Position the patient supine with his or her knee flexed at 90 degrees (Fig. 20.17). Place one hand on the heel and the other hand on the knee. Apply pressure with both hands and gently rotate the foot. The test is positive if audible or palpable clicks occur.



FIGURE 20.16. McMurray's test.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. O'Malley have concluded. The following are your findings:

■ General Health Survey

- Well-developed 68-year-old white female in no acute distress.

- Head-to-toe scan revealed no pertinent findings related to the musculoskeletal system.

■ Vital Signs

- Temperature, 98.7°F
- Pulse, 78 beats per minute (BPM), strong and regular.
- Respirations, 18/min., unlabored.
- Blood pressure (BP), 148/68.
- Height, 5'5" (1½" height loss since onset of menopause).
- Weight, 125 lb.

■ Musculoskeletal

- Gait steady, no support needed for ambulation.
- Mild muscle atrophy noted in quadriceps, all other muscles without atrophy. No involuntary movements.
- Skin intact over joints without rashes, deformities, erythema, ecchymosis, swelling, masses, or nodules.
- Shape and tone of muscles within normal limits.
- Contour smooth over joints; no tenderness, heat, redness, or pain palpated in any joint.
- Full ROM of all joints within normal limits.
- Muscle strength 5/5.
- Leg length 80.5 cm bilaterally, thigh circumference 40 cm bilaterally.
- Mild kyphosis when standing.



Case Study Analysis and Plan

Now that you have completed a thorough physical assessment of Mrs. O'Malley, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mrs. O'Malley and use this information to identify a list of nursing diagnoses. Some possible ones are provided as follows. Cluster the supporting data.

1. Pain
2. Impaired physical mobility
3. Risk for injury
4. Health-Seeking Behaviors

Identify any additional nursing diagnoses.



CRITICAL THINKING ACTIVITY #2

List all of the strengths and areas of concern that have a bearing on Mrs. O'Malley's health status.



CRITICAL THINKING ACTIVITY #3

Now that you have identified some nursing diagnoses for Mrs. O'Malley, select one and develop a nursing care plan and a brief teaching plan for her, including learning objectives and teaching strategies.

RESEARCH TELLS US

The risk for osteoporosis is increased in postmenopausal women. However, osteoporosis may not be diagnosed until the patient becomes symptomatic, which often occurs as the result of a fracture. The National Osteoporosis Risk Assessment (NORA), a longitudinal observational study of more than 200,000 postmenopausal women, studied the relationship between osteoporosis risk factors and low bone mineral density (BMD), and between BMD, other risk factors and short-term (1-year) fracture incidence.

NORA included peripheral bone densitometry, self-report risk assessment questionnaires, and 1-year follow-up for skeletal fractures. Siris and colleagues (2001) report an unexpectedly high prevalence of unsuspected osteopenia and osteoporosis in healthy postmenopausal women. Forty percent of the women tested had BMD measurements denoting osteopenia; 7 percent of the women had BMD findings suggestive of osteoporosis.

The study confirmed risk factors for osteoporosis, including:

- Age.
- History of previous fracture.
- Smoking.
- Glucocorticoid use.
- Asian and Hispanic ethnicity.

Although the study was limited in that women previously diagnosed or treated for osteoporosis were excluded, use of self-report, inability to assess asymptomatic spinal fractures, and lack of control of BMD devices, the large sample of geographically and ethnically diverse women in primary care was a definite asset. These findings suggest that a significant number of postmenopausal women in primary care settings are underidentified as at risk for osteoporosis; therefore, they're also undertreated when it comes to preventing osteoporosis.

Another study by Kemmler and associates (2002) examined the effects of exercise in fitness and BMD in early postmenopausal women. Fifty-nine women were enrolled in an intensive exercise training program. Forty-one women served as the control group without exercise. Significant differences were seen between the groups with changes in bone density ($p < .001$), isometric strength ($p < .001$), and quality of life parameters as reflected by report of low back pain. The findings support the effects of intensive exercise training in improving strength, endurance, quality of life, and BMD in early postmenopausal women.

Health promotion activities should include screening for osteoporosis by identifying risk factors, bone densitometry, and encouraging exercise for those at risk.

Sources: Siris, E.S., Miller, P.D., Barrett-Connor, E., Abbot, T., Sherwood, L., and Berge, M. (2001). Identification and fracture outcomes of undiagnosed low bone mineral density in postmenopausal women: Results from the National Osteoporosis Risk Assessment. *Journal of the American Medical Association* 286:2815-2822; Chestnut, C. (2001). Osteoporosis, an underdiagnosed disease. *Journal of the American Medical Association* 286(22):2865-2866; and Kemmler, W., Engelke, K., Lauber, D., Weineck, J. Hensen, J., and Kalender, W. (2002). Exercise effects on fitness and bone mineral density in early postmenopausal women: 1-year EFOPS results. *Medicine & Science in Sports & Exercise* 34(12):2115-2123.

HEALTH CONCERNS

The Arthritis Foundation reports that in 2005, arthritis affected nearly 66 million Americans (1 in 3 adults) with 42.7 million being diagnosed as having arthritis, and 23.2 not yet diagnosed but living with chronic joint symptoms. Arthritis is the leading cause of work disability for adults in the United States.

Types of Arthritis

- **Osteoarthritis** (degenerative joint disease) is deterioration of the bone cartilage in the joint, leading to pain and limited movement. DJD is the most common type of arthritis and is some times called “wear-and-tear” arthritis. 21 million people in the United States have osteoarthritis (CDC, 2005).
- **Rheumatoid arthritis** is an autoimmune disease causing inflammation of the joint. RA is the most serious and disabling form of arthritis mainly affecting women. Three million adults in the United States have rheumatoid arthritis (CDC, 2005).
- **Gout** is a hereditary metabolic disorder with a build up of uric acid (hyperuricemia) in the joints, resulting in inflammation and pain. Affecting men more than women, gout is easily controlled with medication and diet. In the United States, 5.1 million adults have this disorder (CDC, 2005).
- **Ankylosing spondylitis** is arthritis that specifically affects the spine, causing the spine to grow together.

- **Juvenile arthritis** affects children; it may be RA type or the result of another autoimmune disorder such as lupus.
- **Systemic lupus erythematosus (SLE)** is an autoimmune disorder that can result in inflammation and destruction of the joints and connective tissue.
- **Scleroderma** is affects connective tissue causing thickening and hardening of the tissue.
- **Fibromyalgia** is systemic muscle and joint pain, most common in women. An estimated 3.7 million adults in the United States have fibromyalgia (CDC, 2005).

Risk Factors for Arthritis

- **Nonmodifiable risk factors**
 - *Age:* Risk increases with age.
 - *Gender:* Generally more common in women, with gout being more common in men.
 - *Genetic:* RA and SLE associated with higher genetic risk.
- **Modifiable risk factors**
 - *Overweight and obesity:* Can contribute to onset and progression.
 - *Joint injuries:* May contribute to osteoarthritis of that joint.
 - *Infection:* Can potentially cause development of various forms of arthritis.
 - *Occupation:* Repetitive knee bending associated with osteoarthritis of the knee.

<http://www.arthritis.org>;

http://www.cdc.gov/arthritis/arthritis/risk_factors.htm; and Ignatavicius, D., and Workman, M. (2002).

Medical-Surgical Nursing: Critical Thinking for Collaborative Care, ed. 4. Philadelphia: Saunders.

Common Abnormalities

ABNORMALITY	ASSESSMENT FINDINGS
<p>Ankylosing Spondylitis (Marie-Strümpell Disease) Chronic, progressive disease involving joints. Bilateral sclerosis of the sacroiliac joints is a diagnostic sign.</p>	<ul style="list-style-type: none"> ■ Males ages 10 to 30. ■ Familial disorder with genetic predisposition. ■ Low backache. ■ Flattening of lumbar curvature. ■ Decreased joint ROM. ■ Muscle stiffness. ■ Joint swelling and redness. ■ Deformity and immobility. ■ Progressively limited back movement and chest expansion.

ABNORMALITY	ASSESSMENT FINDINGS
<p>Carpal Tunnel Syndrome Compression of the medial nerve of the hand.</p>	<ul style="list-style-type: none"> ■ Decreased spinal ROM with abnormal vertebral alignment, flattened lumbar curve, and dorsal kyphosis. ■ Atrophy of trunk muscles. ■ Fever. ■ Fatigue. ■ Weight loss. ■ Fusion of entire spine in severe cases. ■ Cardiac conduction disturbances in severe and long-standing disease. <ul style="list-style-type: none"> ■ History of predisposing factors such as wrist trauma or injury, RA, gout, myxedema, diabetes mellitus, leukemia, acromegaly, edema of the hand associated with pregnancy. ■ Pain that worsens after manual activity, especially that involving wrist rotation, or at night. ■ Radiates up the arm. ■ May be intermittent or constant. ■ Causes numbness, burning, or tingling of the thumb and the index and middle fingers. ■ Atrophy of thenar eminence (mass of tissue on the lateral side of the palm). ■ Positive Tinel's sign (tingling during wrist percussion). ■ Muscle weakness. ■ Dry skin over thumb and first two fingers. ■ Inability to oppose thumb and little finger.
<p>Lumbosacral Disc Herniation Prolapse of the nucleus pulposus of a ruptured intervertebral disc into the spinal canal.</p>	<ul style="list-style-type: none"> ■ Pain on coughing, sneezing, straining, bending, or lifting. ■ Pain increases with sitting. ■ Accompanied by mild to severe low back, buttock, or leg pain. ■ May be associated with spasms. ■ Decreased sensation, paresthesias. ■ Absent or diminished deep tendon reflexes (DTRs) of affected dermatomes. ■ Voiding (particularly urinary retention) or defecating difficulties. ■ History of predisposing factors such as recent spinal trauma, heavy or awkward lifting or occupational stress on back, lack of exercise, obesity, degenerative changes. ■ Decreased spinal ROM. ■ Unequal limb circumference. ■ Abnormal posture, scoliosis, flattening of lumbar curvature. ■ Diminished DTRs of knee and ankle. ■ Positive straight leg raising test.

(continued)

ASSESSMENT CHECKLIST (continued)

- Coordination (RAM, finger thumb opposition, toe tapping, heel down shin)
- Accuracy of movements (finger to nose, point to point localization)

Measurements

- Arm lengths from acromion process to tip of middle finger
- Arm circumferences
- Leg lengths from anterior superior iliac crest to medial malleolus
- Leg circumference

Muscles

- Tone (relaxed and contracted state) of upper and lower extremities
- Muscle strength

Joints

- ROM, edema, redness, tenderness, crepitus, deformities, stability

Additional Tests

- Carpal tunnel syndrome*: Phalen test or Tinel test
- Arm weakness*: Pronator drift
- Spinal problems*: Straight leg raise
- Hip problems*: Thomas test, Trendelenburg test
- Knee problems*: Bulge sign, Lachman test, draw sign, McMurray's test, Apley's test

Percussion

- Knee for fluid (patellar tap)

DOCUMENT

Document your findings and develop your plan of care.

SUMMARY


- This chapter taught you how to do a thorough musculoskeletal assessment, including a health history and physical examination.
- Before you begin the assessment, you need an understanding of the anatomy of the musculoskeletal system including ROMs of the joints.
- Since the musculoskeletal system has a major impact on a person's well-being and ability to function, begin by taking a detailed health history.
- The physical exam of the musculoskeletal system uses inspection, palpation, and percussion. The exam includes assessment of gait, posture, cerebellar function, limb measurement, joints, and muscles.
- Once you have completed your assessment, analyze your findings, identify actual and potential health problems, and write nursing diagnoses and a plan of care.

REVIEW QUESTIONS

1. What musculoskeletal changes would you find in a woman who is 7 months pregnant when performing a spinal assessment?
 - a. Scoliosis
 - b. Lordosis
 - c. Kyphosis
 - d. No change
2. You are assessing your patient's gait. Which finding should alert you that you patient may have a balance problem?
 - a. Wide base of support
 - b. Increased stride length
 - c. Toeing out
 - d. Rhythmic cadence
3. As a school nurse, you are performing scoliosis screenings. What patient position is best for assessing scoliosis?
 - a. Supine
 - b. Sitting
 - c. Standing
 - d. Bending from waist
4. Mary Belamy, age 63, has HTN and asthma. Which medication would increase her risk for osteoporosis?
 - a. Steroids
 - b. Lasix
 - c. Aspirin
 - d. Beta blocker
5. Your patient, age 69, has had a total right hip replacement. The physical therapist uses a goniometer to assess his progress. The goniometer is useful in assessing:
 - a. ROM of the joint.
 - b. Muscle strength.
 - c. Muscle tone.
 - d. Joint stability.
6. You are assessing your patient's cerebellar function. You have your patient perform the Romberg test by instructing him to:
 - a. Hop on one foot.
 - b. Do a deep knee bend.
 - c. Tandem walk.
 - d. Stand with feet together, eyes open then closed.

7. You are assessing your patient's cerebellar function. Circle all of the tests you would use to assess his balance.
 - a. Tandem walk
 - b. Rapid alternating movements
 - c. Heel and toe walking
 - d. Point-to-point localization
 - e. Romberg test
8. Which is the correct technique for measuring arm lengths?
 - a. Measure from the acromion process to the wrist.
 - b. Measure from the acromion process to the tip of the middle finger.
 - c. Measure from the head of the humerus to the distal interphalangeal joint.
 - d. Measure from the head of the humerus to the proximal interphalangeal joint.
9. Joint ROM is part of a musculoskeletal assessment. ROM of the knee includes (circle all that apply):
 - a. Flexion.
 - b. Extension.
 - c. Eversion.
 - d. Inversion.
10. Joint ROM is part of a musculoskeletal assessment. ROM of the temporomandibular joint includes (circle all that apply):
 - a. Retraction.
 - b. Adduction.
 - c. Elevation.
 - d. Protraction.

CHAPTER 21



Assessing the Sensory-Neurologic System

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent sensory-neurologic system history questions
- Obtain a sensory-neurologic system history
- Perform a physical assessment of the sensory-neurologic system
- Document sensory-neurologic system findings
- Identify actual/potential health problems stated as nursing diagnoses
- Differentiate between normal and abnormal findings

The human nervous system is a unique system that allows the body to interact with the environment as well as to maintain the activities of internal organs. It is composed of structures that transmit electrical and chemical signals between the body's systems and the brain. The multiple functions of the nervous system are so automatic that most people are unaware of their magnitude until a problem occurs.

An impaired nervous system can manifest in many ways, from subtle weakness to drastic loss of mobility. The nervous system acts as the main “circuit board” for every body system. Because the nervous system works so closely with every other system, a problem within another system or within the nervous system itself can cause the nervous system to “short-circuit.”

A major goal of nursing is early detection to prevent or slow the progression of disease. So it is important for nurses to accurately perform a thorough neurologic assessment and to understand the implications of subtle changes in assessment findings. By doing so, you can initiate timely interventions that can save lives.

Because of the complexity of this system, assessment can be challenging, especially for beginning nurses. All body organs and most tissues are innervated; therefore, neurologic activity affects functioning within all body systems. For example, the nervous system responds to stress by increasing the heart rate, resulting in increased blood pressure (BP), whereas it responds to the sight or smell of food by increasing the production of digestive juices. Furthermore, a seemingly straightforward neurologic symptom such as headache, insomnia, or forgetfulness can have a wide range of causes, contributing factors, and consequences.

Another assessment challenge may be interacting with the patient. Neurologic impairment can reduce the ability to communicate, cause confusion, alter personality, and otherwise confound your attempts to obtain an accurate health history. Finally, factors such as drug and alcohol use, medication interactions, diet, amount of sleep, and other self-care factors can influence neurologic health. Therefore, careful sleuthing is required when taking the health history and gathering and evaluating the physical findings.

The previous chapter discussed neuromuscular function, just one aspect of the highly complex neurologic system. This chapter presents the sensory-neurologic assessment, including information on assessing cerebral function, cranial nerves (CNs), sensation, and reflexes. Throughout the assessment, you will identify risk factors for neurologic impairment and health promotion teaching needs. Remember to document all findings, including normal data, precisely and thoroughly, to provide a baseline against which future neurologic function can be measured. These baseline data are critically important for evaluating degeneration in patients with progressive neurologic disease, as well as for evaluating

patients' progress toward rehabilitation after neurologic trauma and other acute problems.

Anatomy and Physiology Review

Before beginning the neurologic assessment, you need to understand the anatomy and physiology of the system, including its anatomic and functional divisions and the type and extent of its interaction with other body systems. In addition to the motor aspects discussed in Chapter 20, general functions of the neurologic system include:

- Cognition, emotion, and memory.
- Sensation, perception, and the integration of sensory-perceptual experience.
- Regulation of homeostasis, consciousness, temperature, BP, and other bodily processes.

Structures and Functions of the Sensory-Neurologic System

There are two types of nerve cells: neuroglia and neurons. **Neuroglia** act as supportive tissue, nourishing and protecting the neurons. They also maintain homeostasis in the interstitial fluid around the neurons and account for about 50 percent of the central nervous system (CNS) volume. Neuroglia have the ability to regenerate and respond to injury by filling spaces left by damaged neurons.

Neurons have the ability to produce action potentials or impulses (excitability or irritability) and to transmit impulses (conductivity). They are composed of a cell body, dendrites, and axons. The **cell body** contains the nucleus. **Dendrites** are short, branchlike structures that receive and carry impulses to the cell body. **Axons** are long fibers that carry the electrical impulses generated by the cell bodies. Some axons are covered with a myelin sheath that allows rapid impulse transmission.

For neurons to communicate with one another, the axon of one neuron has to convey its impulse to the dendrite of the next neuron across a space called a **synapse**. It does this with the help of **neurotransmitters**, chemicals released by the axon that carry the impulse to the receiving dendrite. The impulses conveyed by neurons include sensory data about what the body is seeing, hearing, feeling, and so on, as well as motor data telling the muscles how to respond.

Sensory impulses are transmitted to the brain through **afferent**, or ascending, pathways. Motor impulses are transmitted from the brain to muscles through the **efferent**, or descending, pathways (Fig. 21.1).

Neurons band together into peripheral nerves, spinal nerves, the spinal cord, and the tissues of the brain. These structures make up the neurologic system, which is divided into the CNS and the peripheral nervous system (PNS).




FIGURE 21.1. Neurotransmission and neural pathways.

The Central Nervous System

The CNS consists of the brain and spinal cord.

The Brain

The brain is composed of gray matter, made up of neuronal cell bodies, and white matter, composed of axons and dendrites. The brain consists of four major structures: the cerebrum, diencephalon, cerebellum, and brainstem. These and other components of the brain are shown in Figure 21.2 and described in *Structures and Functions of the Brain*.

The Spinal Cord

The **spinal cord** descends through the foramen magnum (large aperture) of the occipital bone of the skull, through the first cervical vertebra (C1), and through the remainder of the vertebral column to the first or second lumbar vertebra. At this point, the cord itself terminates

and its roots branch off into the cauda equina (“horse’s tail”).

A cross-section of the spinal cord reveals gray matter composed of neuronal cell bodies clustered into an “H” shape, with two anterior and two posterior “horns.” This gray matter is surrounded by white matter composed of myelinated axons and dendrites. Cord fibers associate into **ascending tracts**, which carry sensory data to the brain, and **descending tracts**, which carry motor impulses from the brain.

Sensory Pathways

Sensory pathways, either ascending or afferent, allow sensory data, such as the feeling of a burned hand, to become conscious perceptions. The pathways by which a variety of somatic sensations travel to the cerebral cortex are illustrated in Figure 21.3. The two major sensory pathways are the lateral and anterior spinothalamic tracts and the posterior column. The lateral and anterior spinothalamic tracts




FIGURE 21.2. Lobes of the brain.

Structures and Functions of the Brain

STRUCTURE	FUNCTION
Cerebrum	<ul style="list-style-type: none"> ■ Largest part of the brain. ■ Consists of two hemispheres connected by a band of neurons called the <i>corpus callosum</i>, the great commissure of the brain. ■ Each hemisphere has four lobes.
■ Cerebral cortex	<ul style="list-style-type: none"> ■ Outermost layer of cerebrum. ■ Consists of a ridge of gray matter formed from clustered cell bodies of neurons. ■ Folds of the cerebral cortex are called convolutions or gyri, and the grooves between them are called fissures or sulci.
■ Frontal lobes	<ul style="list-style-type: none"> ■ Controls most conscious processes. ■ Regulate emotional expression, behavior, intellect; influence personality; control voluntary movement. ■ Contain Broca's area (left lobe in most people), which controls expressive language function.
■ Temporal lobes	<ul style="list-style-type: none"> ■ Influences hearing, smell, taste, and memory. ■ Left temporal lobe (in most people) contains Wernicke's area, which allows language comprehension.
■ Limbic system	<ul style="list-style-type: none"> ■ Contains limbic system. ■ Lies deep in frontal and temporal lobes. ■ Influences emotional and sexual arousal, behavioral expression, recent memory, smell.

STRUCTURE	FUNCTION
<ul style="list-style-type: none"> ■ Parietal lobes <ul style="list-style-type: none"> ■ Insula (central lobe) ■ Occipital lobes 	<ul style="list-style-type: none"> ■ Perceive and interpret sensory input, such as pain, temperature, touch, texture, and proprioception. ■ Located beneath the parietal, frontal, and temporal lobes. ■ Involved with GI and visceral activity.
Reticular Activating System	<ul style="list-style-type: none"> ■ Perceive and interpret visual stimuli, including spatial relationships. ■ Diffuse network of hyperexcitable neurons in brainstem and cerebral cortex that screens and channels incoming sensory input. ■ Helps maintain consciousness or wakefulness.
Diencephalon	<ul style="list-style-type: none"> ■ Connects cerebrum to brainstem. ■ Contains thalamus and hypothalamus.
<ul style="list-style-type: none"> ■ Thalamus 	<ul style="list-style-type: none"> ■ Clusters multiple sensory stimuli (e.g., the warmth, smell, and taste of tea) into a coherent whole before sending it to the cerebral cortex for perception. ■ Tunes out continuing unimportant stimuli (e.g., sound of distant radio when you are reading).
<ul style="list-style-type: none"> ■ Hypothalamus 	<ul style="list-style-type: none"> ■ Controls and integrates autonomic nervous system and pituitary gland activities. ■ Regulates production of several hormones; stimulates visceral responses, such as heart rate, in response to emotions; regulates temperature by prompting shivering and sweating. ■ Helps regulate how much we eat, producing feelings of hunger or satiation in response to changes in cellular levels of nutrients. However, the conscious mind can overrule this feeling. ■ Also helps establish and maintain wake/sleep patterns.
Cerebellum	<ul style="list-style-type: none"> ■ Located at posterior base of the brain. ■ Regulates involuntary aspects of movement, such as coordination, muscle tone, kinesthetics, posture, equilibrium.
Brainstem	<ul style="list-style-type: none"> ■ Anterior to cerebellum and superior to spinal cord. ■ Includes midbrain, medulla, and pons. ■ Responsible for involuntary survival behaviors, such as breathing, heart rate, coughing, vomiting. Also transmits impulses from spinal cord to brain. ■ Attachment site for 10 of the 12 cranial nerves.
<ul style="list-style-type: none"> ■ Midbrain 	<ul style="list-style-type: none"> ■ Regulates visual, auditory, and other reflexes and controls eye movements, focusing, and pupil dilation.
<ul style="list-style-type: none"> ■ Medulla 	<ul style="list-style-type: none"> ■ Regulates heart and respiratory rate, BP, and protective reflexes such as swallowing, vomiting, sneezing, and coughing.
<ul style="list-style-type: none"> ■ Pons 	<ul style="list-style-type: none"> ■ Helps control respiratory function, facial movement and sensation, and eye movement.
Meninges	<ul style="list-style-type: none"> ■ Three connective tissue membranes that cover and protect brain and spinal cord: <ul style="list-style-type: none"> ■ <i>Pia mater</i>: Inner layer, contains blood vessels. ■ <i>Arachnoid</i>: Middle layer, provides space for cerebrospinal fluid. ■ <i>Dura mater</i>: Outer layer that adds support and protection.
Ventricles	<ul style="list-style-type: none"> ■ Four cavities within the brain whose capillaries continuously produce and reabsorb cerebrospinal fluid (CSF) from blood plasma. CSF cushions the brain, delivers nutrients, and removes wastes. ■ CSF circulates in the subarachnoid space between the arachnoid mater and the pia mater.

transmit nerve impulses for pain, temperature, itching, tickling, pressure, and crude touch, whereas the posterior column transmits nerve impulses for proprioception, discriminative sensations, and vibrations. Some visceral sensory neurons also enter the spinal cord as shown in the figure; however, other visceral stimuli are carried by the vagus nerve or another of the cranial nerves and enter the brainstem directly, bypassing the spinal cord.

All somatic sensory stimuli—for example, sensations of pain, heat, or pressure—are conveyed from the body periphery into the spinal cord via the posterior root of the appropriate spinal nerve. At that point, sensations of fine touch, proprioception, and vibration continue into the posterior column of the spinal cord and into the brainstem. Here they cross to the opposite side of the brainstem before they continue into the




FIGURE 21.3. Sensory pathways.

thalamus and from there to the sensory areas of the cerebral cortex.

In contrast to this, sensations of pain, temperature, crude touch, and pressure enter the posterior horn and cross the spinal cord into the opposite side of the cord before they begin their ascent. Pain and temperature are carried in the **lateral spinothalamic tract**, whereas crude touch and pressure are carried in the **anterior spinothalamic tract**. These tracts continue to ascend straight upward, entering the thalamus and then the cerebral cortex.

As you might imagine, a patient with a particular type of sensory loss may have a neurologic impairment affecting a particular column or tract. For example, someone who loses the ability to perceive vibration and changes in positioning may have a lesion in the posterior column.

Major Sensory Tracts

TRACT	SENSATION
Lateral Spinothalamic	■ Pain and temperature
Anterior Spinothalamic	■ Light (crude) touch
Posterior Column	■ Touch-pressure, two-point discrimination, vibratory sensation, position sense (proprioception), and some light (crude) touch




FIGURE 21.4. Motor pathways.

Motor Pathways

Motor pathways (descending or efferent) transmit impulses from the brain to the muscles. The three major motor pathways of the CNS are the corticospinal (pyramidal or direct), including the corticobulbar; the extrapyramidal or indirect; and the cerebellum. The descending pathways can be direct or indirect. **Direct (pyramidal) pathways** carry impulses from the cerebral cortex to lower motor neurons that innervate the skeletal muscles, resulting in voluntary movement. The direct pathways include the anterior and lateral corticospinal and corticobulbar tracts. The corticospinal tracts control voluntary skilled movement of the extremities and fine movement of the fingers. The corticobulbar tract connects the motor cortex to lower motor neurons of the cranial nerves. The **indirect (extrapyramidal)**

pathways carry impulses from the brainstem and other parts of the brain, resulting in automatic movement, coordination of movement, and maintenance of skeletal muscle tone and posture. The indirect pathways consist of the reticulospinal, rubrospinal, tectospinal, and vestibulospinal tracts. The possible pathways of the corticospinal system are shown in Figure 21.4. The cerebellar system, which is involved primarily in coordination, equilibrium, and posture, is not shown. Also note that this discussion pertains to somatic, not visceral, motor neurons.

From the motor cortex, all somatic motor impulses descend via one of three tracts through the thalamus and into the brainstem, at which point the configuration of the three tracts resembles a pyramid. The anterior corticospinal and corticobulbar (uncrossed pyramidal tract)




FIGURE 21.5. Reflex arc.

and the reticulospinal and vestibulospinal (uncrossed extrapyramidal tract) continue straight down into the spinal cord. There, the impulse must cross over to the opposite anterior horn before it can exit the spinal cord via the anterior root of the spinal nerve. The lateral corticospinal (crossed pyramidal tract) and the rubrospinal (crossed extrapyramidal tract) both cross to the opposite side of the brainstem before descending directly into the anterior horn and exiting into the anterior root of the spinal nerve.

Motor impulses from the motor cortex to the periphery are controlled by upper and lower motor neurons. The upper motor neurons are located in the cerebral cortex and brainstem and regulate the responses of the lower motor neurons. The lower motor neuron cell bodies lie in the anterior horn of the spinal cord, then exit the cord through the nerve roots. The lower motor neurons are the final link between the CNS and the skeletal muscles. An upper motor neuron lesion causes increased muscle tone (spasticity) and hyperreflexia. A lower motor neuron lesion causes decreased tone, flaccidity, and absent reflexes or hyporeflexia.

Spinal Reflexes

Spinal reflexes do not depend on conscious perception and interpretation of stimuli, nor on deliberate action; in other words, they do not involve the brain. They occur involuntarily, with lightning speed, and are identical in all healthy children and adults, although they are less developed in infants. When we experience a spinal reflex, we are not aware of the reflexive activity itself, only its result.

The simplest spinal reflexes are known as **deep tendon reflexes** (DTRs). They are **monosynaptic**, involving just one sensory neuron communicating across a single synapse to one responding motor neuron. A reflex arc is shown in Figure 21.5.

The classic example of a DTR is the **patellar reflex**, commonly called the *knee-jerk reflex*. The patellar tendon is tapped with a reflex hammer, triggering a stretch response in the sensory fibers of the quadriceps femoris muscle. This stimulus is received by a sensory neuron, which carries it to the anterior horn of the spinal cord, where it is conveyed across a synapse to a motor neuron that then causes the muscle to contract. Because

Major Motor Tracts

TRACT

MOTOR EFFECT

Lateral Corticospinal (Crossed Pyramidal)

- Major tract controlling movement of extremities, fine movements of fingers

Anterior Corticospinal (Uncrossed Pyramidal)

- Voluntary movement of axial muscles

Corticobulbar (Uncrossed Pyramidal)

- Cranial nerves

Rubrospinal (Crossed Extrapyramidal)

- Control and coordination of skilled muscle movement

Reticulospinal (Uncrossed Extrapyramidal)

- Minor influence on movement

Vestibulospinal (Uncrossed Extrapyramidal)

- Integration of axial body movements

- Maintenance of body posture

- Integration of body and limb movements

- Minor influence on movement

- Maintenance of body posture

- Integration of body and limb movements

Spinal Reflexes

REFLEX

SPINAL CORD SEGMENT INVOLVED

Deep Tendon Reflexes

- Achilles reflex (plantar flexion at ankle)
- Patellar reflex (knee extension)
- Biceps reflex (elbow flexion)
- Triceps reflex (elbow extension)
- Brachioradialis reflex (forearm flexion and supination)

- S1 and S2
- L2, L3, and L4
- C5 and C6
- C6, C7, and C8
- C5 and C6

Superficial Reflexes

- Upper abdominal reflex
- Lower abdominal reflex
- Plantar reflex
- Cremasteric reflex
- Bulbocavernosus reflex
- Anal reflex

- T8, T9, and T10
- T10 and T11
- L5, S1, and S2
- L1, L2
- S3, S4
- S3, S4, and S5

of the simplicity of the reflex arc, this contraction occurs within less than 1 second after the hammer tap.

Slow or absent DTRs could indicate a health problem affecting the peripheral nerves, such as a degeneration of their myelin sheaths. Spinal cord trauma or lesions also commonly affect reflexes; therefore, it is helpful to know the specific spinal cord segment associated with each specific reflex. The box entitled Spinal Reflexes identifies the five DTRs, as well as superficial reflexes involving stimulation of the skin, along with the segmental level at which they occur. Assessment of these reflexes is discussed later in the chapter.

In addition to the monosynaptic reflexes, the body exhibits protective **flexor (or withdrawal) reflexes**, which help minimize trauma from harmful stimuli. For example, these reflexes allow speedy, involuntary withdrawal from a heat source. Flexor reflexes require a sensory neuron to interact with an intermediate neuron or interneuron in the spinal cord. This interneuron, in turn, sends the impulse across a synapse to the receiving motor neuron.

Peripheral Nervous System

The **peripheral nervous system** consists of the cranial and spinal nerves and the peripheral autonomic nervous system. The PNS can also be divided into the somatic and the visceral nervous systems. The somatic system has both afferent and efferent divisions. The afferent division receives, processes, and transmits sensory information from the skin and the musculoskeletal system, eyes, tongue, nose, and ears. The efferent division has lower motor neurons and regulates voluntary muscle contraction.

The visceral system also has afferent and efferent divisions. The afferent division is responsible for processing information from the visceral organs. The efferent division is responsible for the motor responses of the smooth muscle, cardiac muscle, skin glands, and viscera. The efferent system is known as the **autonomic nervous system**.

Cranial Nerves

The 12 pairs of cranial nerves originate from the brain and are called the *peripheral nerves of the brain*. As shown in Figure 21.6, these nerves originate from the cerebrum, diencephalon, and brainstem. The function of each of the cranial nerves is indicated in *Function of Cranial Nerves*.

Spinal and Peripheral Nerves

Branching from the spinal cord are 31 pairs of spinal nerves: 8 cervical, 12 thoracic, 5 lumbar, 5 sacral, and 1 coccygeal (Fig. 21.7). The spinal nerves contain both ascending and descending fibers, and although there is some overlap, each is responsible for innervation of a particular area of the body. As shown in Figure 21.8, **dermatomes** are regions of the body innervated by the cutaneous branch of a single spinal nerve. For example, nerve C6 innervates a portion of the lateral (thumb) side of the arm and hand. Although there is some individual variation, it is useful to know which spinal nerves typically innervate which segments of the body so that neurologic deficits can be more easily associated with lesions of particular nerves.

Each spinal nerve emerges from the spinal cord from two distinct roots. The **posterior (dorsal) root** contains afferent fibers that receive sensory information

Function of Cranial Nerves

NUMBER	NAME	TYPE AND FUNCTION
CN I	Olfactory	■ <i>Sensory</i> : Smell
CN II	Optic	■ <i>Sensory</i> : Sight
CN III	Oculomotor	■ <i>Motor</i> : Opening eyelid; moving eye superiorly, medially, and diagonally; constricting pupils
CN IV	Trochlear	■ <i>Motor</i> : Moving eye down and laterally
CN V	Trigeminal	■ <i>Motor</i> : Chewing and jaw opening and clenching
		■ <i>Sensory</i> : Conveying sensory data from eyes (cornea), nose, mouth, teeth, jaw, forehead, scalp, and facial skin
CN VI	Abducens	■ <i>Motor</i> : Moving eye laterally
CN VII	Facial	■ <i>Motor</i> : Closing eyes, closing mouth, moving lips and other muscles of facial expression, salivation and lacrimation (secreting saliva and tears)
		■ <i>Sensory</i> : Tasting on anterior tongue
CN VIII	Acoustic	■ <i>Sensory</i> : Hearing, equilibrium
CN IX	Glossopharyngeal	■ <i>Motor</i> : Swallowing, gag sensation, secretion of saliva
		■ <i>Sensory</i> : Tasting on posterior tongue
CN X	Vagus	■ <i>Motor</i> : Palate, pharynx, larynx (speaking and swallowing)
		■ <i>Sensory</i> : Sensations in pharynx and larynx
		■ <i>Sensorimotor</i> : Cardiovascular, respiratory, and digestive systems
CN XI	Accessory	■ <i>Motor</i> : Contracting muscles of neck and shoulders
CN XII	Hypoglossal	■ <i>Motor</i> : Tongue movement, articulating with tongue, swallowing

from the body periphery and convey it to the CNS. The **anterior (ventral) root** contains efferent fibers that convey motor impulses from the CNS to the muscles of the body, directing and refining movement. These anterior and posterior roots then merge just distal to the cord to form one mixed spinal nerve capable of both sensory and motor functioning. Spinal nerves branch further to form the **peripheral nerves** of the body, most of which also carry both sensory and motor fibers.

The autonomic nervous system is divided into the sympathetic and parasympathetic, with efferent fibers to muscle, organs, or glands. Usually, the two systems work opposite each other. The sympathetic allows the body to respond to stressful situations. The parasympathetic functions when all is normal. The sympathetic nerves exit the spinal cord between the level of the first thoracic and the second lumbar vertebrae. The preganglionic nerves descend the cord and exit, and then enter a relay station known as the **sympathetic chain**. The impulse is then transmitted to a postganglionic neuron that goes to the target organ to stimulate a response.

Sympathetic responses include vasoconstriction; increased BP, heart rate, and contractility; increased respiratory rate; smooth muscle relaxation of bronchioles, gas-

traintestinal (GI) tract, and urinary tract; sphincter contraction; pupillary dilation; increased sweat, and decreased pancreatic secretion.

The parasympathetic fibers leave the CNS by way of the cranial nerves from the midbrain and medulla, and between the second and the fourth sacral vertebrae. A long preganglion fiber exits to an area near the target organ, and then synapses to form a postganglionic nerve, which in turn leads to a response. Parasympathetic responses include decreased heart rate, contractility, and velocity; smooth muscle constriction of bronchioles; increased GI tract tone and peristalsis with sphincter relaxation; urinary sphincter relaxation and increased bladder tone; vasodilation of external genitalia causing male erection; pupillary constriction, and increased pancreatic, lacrimal, and salivary secretions. There is little effect on mental and metabolic activity. Figure 21.9 illustrates the autonomic nervous system.

Interaction With Other Body Systems

Because the nervous system is the main circuit board for the body, all body systems interact with it in some way.




FIGURE 21.6. Origin of cranial nerves.




FIGURE 21.7. Spinal nerves.

The Cardiovascular System

Although the conduction system of the heart is completely independent of extrinsic neural control, cardiac centers in the medulla influence heart rate. Also, the vagus nerve can exert significant influence over both heart rate and vessel constriction, thereby influencing BP.

The Respiratory System

Respiratory centers in the medulla of the brainstem regulate inspiration and expiration. Centers in the pons regulate respiratory drive. The vagus nerve relays data on blood gas levels to the medulla. Voluntary regulation of breathing can be achieved via motor neurons, which can stimulate the respiratory muscles.




FIGURE 21.8. Dermatomes.

The Endocrine System

The secretion of some endocrine glands, such as the adrenal glands, is regulated by nerves. Some neurons in the hypothalamus actually act as endocrine cells, secreting antidiuretic hormone, which stimulates the kidneys to reabsorb water and oxytocin, a reproductive hormone. Some other hormones, notably parathyroid hormone, affect metabolism of electrolytes, which in turn affects neurologic function.

The Musculoskeletal System

Somatic and visceral motor neurons receive impulses from the cerebral cortex to carry out both voluntary and reflexive movements.




FIGURE 21.9. Autonomic nervous system.

Developmental, Cultural, and Ethnic Variations

Newborns and Infants

Although neurologic development is not complete until toddlerhood, neurologic screening to detect the presence or extent of obvious pathologies such as spina bifida and fetal alcohol syndrome is performed on most newborns within the first 24 hours after birth. This screening includes assessment of appearance, alertness, motor and sensory function, and pain. In addition, newborns are assessed for a set of primitive reflexes that disappear later in infancy, including the palmar grasp, stepping reflex, and rooting reflex (for more information, see Chapter 24, *Assessing the Newborn and Infant*).

Injury to the facial nerve during vaginal birth can cause a transient palsy. Down's syndrome is a genetic disorder that causes mild to severe mental retardation. Maternal use of drugs and alcohol during pregnancy, as well as dietary deficiencies, antepartal infections, systemic diseases, and birth trauma can result in neurologic disorders such as mental retardation, blindness, deafness, seizures, and neuromuscular impairments.

Children and Adolescents

You can usually examine children past toddlerhood in a manner similar to that used for adults, with obvious adap-

tations for developmental stage. In most cases, any chronic neurologic pathologies will have been discovered during infancy. However, childhood injuries and later-onset disorders, such as obsessive-compulsive disorder and hyperactivity, may only become apparent during the preschool years. Changes in behavior may also signal abuse.

If the parent reports seizure activity, assess for whether it is associated with signs of infection, such as a runny nose, high temperature, or recent immunizations. Fevers in toddlers and preschoolers occasionally cause seizures. Repeated episodes of seizure activity may indicate epilepsy.


If the parent reports that the child complains of headache or eye pain, assess for signs of sinusitis, meningitis, or concussion. If parents express concern over attention deficit disorder, hyperactivity, or other learning disorders, ask whether the child consumes caffeine or sugar daily, because some children are extremely sensitive to these substances. If the family lives in an older home, also assess for signs of lead poisoning, which can cause severe neurologic problems.

Finally, observe for behavioral indicators that may signal child abuse. Even in the absence of bruises or other signs of trauma, perform a special assessment for child abuse if a child is inappropriately uncommunicative for his or her developmental level, extremely apprehensive or clingy, unable to show emotion, makes little or


**CARDIOVASCULAR**

CV system supplies nutrients and removes wastes.

Nervous system can affect heart rate, blood pressure, and contractility.

**LYMPHATIC**

Protects against infections and helps with tissue repair.

**REPRODUCTIVE**

Sex hormones can influence behavior.

Nervous system controls sexual function and behavior.


**ENDOCRINE**

Hormones released by endocrine system affect CNS metabolism.

Endocrine system secretes antidiuretic hormone (ADH) and oxytocin.

Reproductive hormones can affect CNS development and behavior.


Nervous system controls pituitary gland and many other endocrine glands.

**GENITOURINARY**

GU system removes wastes and helps regulate pH and electrolytes.

Nervous system also helps regulate pH and can adjust renal blood flow.

Nerves of bladder sense fullness and control emptying.

**RESPIRATORY**

Respiratory system supplies oxygen and removes carbon dioxide.

Nervous system can change the rate and depth of respirations.


**MUSCULOSKELETAL**

MS system protects brain and spinal cord.

MS system provides calcium if needed for neural impulse transmission.

MS system allows for verbal/nonverbal communication.

Nervous system controls muscle contraction and proprioception.


**INTEGUMENTARY**

Skin has receptors for sensations of touch, pain, temperature, texture, and pressure.

Skin and hair protect and insulate skull, brain, and peripheral nerves.

Nervous system controls contraction of erector pili muscles and sweat gland secretion.

Neural responses regulate blushing, sweat production, etc.

**GASTROINTESTINAL**

GI system provides nutrients to nervous system.

Nervous system controls peristalsis and GI secretions.

Somatic motor nerves allow control of defecation.

no eye contact, or responds inappropriately or not at all to painful procedures.

Pregnant Women

Ask pregnant patients if they are taking folic acid supplements, and if so, at what point in the pregnancy they began doing so. Folic acid deficiency, especially in the first trimester, is closely linked to neural tube defects such as spina bifida. Other nutrients are also important to support the fetus's neurologic development.

During pregnancy, the woman may experience transient episodes of neurologic pain, including carpal tunnel syndrome, foot and leg cramps, numbness or tingling in the thighs, and frequent headaches. These usually resolve spontaneously after childbirth. Hyperactive reflexes during the later stages of pregnancy may signal preeclampsia, a hypertensive disorder that can be accompanied by seizures.

Older Adults

Neurons are continually lost during our lives, and with advanced age, neural impulses slow down. However, the common belief that intelligence, memory, and discrimination normally decrease with advanced age is simply not true. Research has shown that creative, critical, and abstract thinking, as well as problem-solving ability, are more typically increased in many older adults. For this reason, you should carefully evaluate neurologic deficits in the older adult for the presence of underlying factors.

Neurologic deficits in older adults are commonly caused by:

- Adverse effects of medications or medication interactions.
- Nutritional deficiencies.
- Dehydration.
- Cardiovascular disease affecting cerebral perfusion.
- Diabetes and other endocrine disorders.
- Neurologic trauma.
- Brain tumors or metastatic disease.
- Stroke (cerebrovascular accident).
- Degenerative neurologic diseases, such as Alzheimer's disease and Parkinson's disease.
- Alcohol or drug abuse.
- Isolation.
- Psychiatric disorders.
- Abuse or neglect.

Also bear in mind that the number and sensitivity of sensory neurons decrease with age, leading to a diminished sense of touch. Reflexes also diminish; for example, the Achilles tendon reflex may be symmetrically diminished or absent in older adults. Abdominal reflexes also may be decreased or absent, and the plantar reflex may be decreased because of musculoskeletal changes in the feet.

People of Different Cultures and Ethnic Groups

Take care not to mistake language barriers or cultural variations in communication styles for evidence of neurologic impairment, especially during the mental status examination. Also, make sure your questions are not culturally specific. For example, do not ask a first-generation Mexican American patient to recite the names of the last four U.S. presidents.

Some cultures have a tendency for specific neurologic diseases; for example, African Americans have a higher incidence of hypertension (HTN) and subsequently a higher incidence of stroke. The Irish have a high incidence of neural tube defects and PKU (phenylketonuria). Navajo Native Americans have a unique neuropathy that causes complete absence of myelinated fiber, weakness, hypotonia, areflexia, loss of sensation in extremities, corneal ulcers, and painless fractures. It usually causes death by age 24.

Introducing the Case Study



Leon Webster is a 21-year-old African American college senior who plays football. Friday night, he and several friends went to a local club to celebrate the team's winning season. While driving home, Leon lost control of his car and hit a tree. He was brought to the emergency room (ER) by ambulance. He was alert and oriented but complaining of a headache and neck pain. He had a 3-inch laceration on his forehead where his head hit the rear-view mirror. He was not wearing a seat belt.

Performing the Sensory-Neurologic and Cranial Nerve Assessment

A complete assessment of the patient's neurologic system requires you to take a detailed health history, perform a mental status examination, assess the cranial nerves, assess the sensory system, and test deep tendon and superficial reflexes.

Health History

Before you begin the health history, keep in mind that the patient may be experiencing confusion, impaired verbal communication, memory loss, personality changes, or other deficits. This may affect her or his ability to provide reliable information, so verifying subjective data with a family member or friend may be wise.

Focused Sensory-Neurologic History

Your patient's condition or time restraints may prohibit taking a detailed neurologic history. If so, ask questions that focus on a history of neurologic problems, the presence of neurologic symptoms, identification of risk factors, and health promotion activities. The patient's response to these questions will help direct your assessment. Key questions include:

- Do you have any neurologic problems?
- Do you have any other medical problems?
- Are you taking any medications?
- Do you have a history of head trauma, LOC, dizziness, headaches, or seizures?
- Do you have memory problems or changes in your senses?
- Do you have weakness, numbness, or paralysis?
- Do you have problems walking or performing activities of daily living (ADLs)?
- Do you have mood problems or depression?
- Do you use drugs or alcohol?
- Do you have allergies?
- Have you ever been treated for a neurologic or psychiatric problem?
- When did your symptoms start?

Rather than asking a long series of questions, it may be best to encourage the patient to tell her or his story without interruption. You can learn a great deal from the content of the story and also by listening to the way it is told. If time is an issue and you are unable to perform a complete health history, perform a focused history on the sensory-neurologic system.

Biographical Data

As always, review the patient's biographical data for clues that relate to the neurologic system. Obviously, the patient's age and educational level will influence the questions you ask and the type and extent of teaching you provide. Keep in mind that certain neurologic disorders are age related. For example, the incidence of stroke increases with age, and neurologic diseases such as myasthenia gravis (MG) and multiple sclerosis (MS) usually attack young women with a peak age between 20 and 30 years. Spinal cord injuries occur more frequently in young people because of the higher rate of accidents. Also, some neurologic disorders are gender related. For example, women have a higher incidence of hemorrhagic stroke, whereas men have more thrombotic strokes; before age 40, MG occurs in women two to three times more often than in men; and the incidence of MS is higher in women than in men. Even geographical locale influ-

ences some types of neurologic diseases. For instance, MS occurs most often in colder climates such as the northeastern, Great Lakes, and Pacific Northwestern states. Also, the incidence of stroke is higher in the "stroke belt" states of Alabama, Arkansas, Georgia, Indiana, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, and Virginia.

Ask adolescents and adults about their job history. Could they have been exposed to neurotoxins? Have they had a head or back injury? Marital status, such as a recent divorce or death of a spouse, can certainly influence neurologic findings. A patient's spiritual beliefs may also influence how he or she perceives illness (e.g., as a punishment) and how he or she deals with illness (e.g., the Christian Science belief in healing through mental and spiritual means).

Current Health Status

The current health status focuses on the patient's chief complaint. If she or he has a neurologic problem, begin with the chief complaint. Major neurologic symptoms to watch for are headache, memory loss, confusion, dizziness, loss of consciousness, numbness, sensory loss, and problems with any of the five senses. Explore these complaints using the PQRST format.



ALERT

Establishing time of onset of symptoms is critical if you suspect a stroke. If the patient is a candidate for tissue plasminogen activator (tPA), there is a 3-hour window in which tPA must be administered.

Symptom Analysis

Symptom analysis tables for all the symptoms described in the following paragraphs are available for viewing and printing on the compact disc that came with the book.

Headache

Headache is the most common neurologic symptom. The causes are many. The pain may be mild or severe, acute or chronic, localized or generalized. Ninety percent of all headaches are benign in nature, caused by muscle contraction (tension) and/or vascular (migraine and cluster); the other 10 percent have underlying pathology. Because a headache may be a symptom of a serious medical problem, a careful, thorough symptom analysis is needed to determine the cause.

Mental Status Changes

Mental status changes are an early indication of a change in neurologic status. The changes may be very subtle and difficult to detect. They may begin slowly as

forgetfulness, memory loss, or inability to concentrate, or rapidly proceed to unconsciousness. Causes include neurologic problems, fluid and electrolyte imbalance, hypoxia, low perfusion states, nutritional deficiencies, infections, renal and liver disease, hyper- or hypothermia, trauma, medications, and drug and alcohol abuse. If your patient's mental status is severely impaired, ask family members to describe the changes that have occurred.

Dizziness, Vertigo, and Syncope

Dizziness, vertigo, and syncope are common neurologic signs and symptoms that warrant further investigation. **Dizziness** is a “fainting” sensation, whereas **vertigo** is a sensation that the surroundings are spinning around (objective vertigo) or that the person is spinning around (subjective vertigo). Vertigo is often accompanied by nausea and vomiting, nystagmus, and tinnitus. Dizziness can lead to **syncope**, which is a temporary loss of consciousness. The patient may say that he or she “blacked out” or “had a spell.” Although the underlying cause of these signs and symptoms may be benign, they may also indicate a serious problem, such as an impending stroke, and need to be investigated thoroughly.

Numbness or Loss of Sensation

Numbness or tingling is referred to as **paresthesia**. Possible causes include diabetes and neurologic, metabolic, cardiovascular, renal, and inflammatory diseases. Determine the area affected and the onset and progression of symptoms.

Deficits in the Five Senses

Assess changes in any of the five senses. Intact cranial nerves are essential for many of the senses. CN I (olfactory) is responsible for the sense of smell; CN II (optic), III (oculomotor), IV (trochlear), and VI (abducens) are responsible for visual acuity, pupillary constriction, and extraocular movement (EOM); CN VII (facial) and IX (glossopharyngeal) control taste; CN VIII (acoustic) controls hearing; and CN V (trigeminal) and dermatomes control somatic sensations.

Visual problems are a frequent symptom associated with neurologic disorders and should be further assessed. Visual changes can result from ocular, neurologic, or systemic problems, eye or head trauma, or adverse effects from drugs. The anatomic position of the cranial nerves that control the eye makes the nerves vulnerable to increases in intracranial pressure (ICP). These visual changes can be total loss of vision, visual field cuts, blurred vision, diplopia (double vision), photosensitivity, and amaurosis fugax (unilateral vision loss, as if a shade were being pulled down, resulting from insufficient blood supply to the retina and lasting up to 10 minutes). The visual deficit may have an acute or gradual onset and be permanent or temporary.

Past Health History

The purpose of the past medical history is to compare it with the patient's present neurologic status or uncover risk factors that might predispose the patient to neurologic disorders. The following questions will guide you in exploring specific areas related to the sensory-neurologic system.

(Text continued on page 756)

Past Health History

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Childhood Illnesses

- Do you have a history of head injury or seizures?

- Neurologic problems in childhood may have permanent effects.

Surgery

- Have you ever had surgery or other treatment involving the nervous system?

- Provides baseline information for physical assessment and explains physical findings.

Hospitalizations/Diagnostic Procedures

- Have you ever been treated for a neurologic problem?

- Helps identify potential or recurring neurologic problem.

Serious Injuries

- Have you ever had a serious injury? If so, can you describe it? What treatment was given, and do you have any residual effects from the injury?

- Information on past injuries provides baseline data for physical assessment.
- Postconcussion syndrome can present 1 to 2 weeks postinjury and persist up to a year.

(continued)

Past Health History (continued)

RISK FACTORS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

Serious/Chronic Illnesses

- Do you have any other medical problems?

- Do you have any other medical problems?

- Minor head injury in older adults can cause a subdural hematoma that may be asymptomatic for weeks to months.
- Spine and back injuries may explain peripheral neurologic problems.

- One-third of people who experience TIAs will have a stroke.
- History of stroke increases risk for another stroke 10 times. Determine residual deficits.
- HTN increases risk for stroke 6 times.
- Diabetes increases stroke risk 2 to 4 times.
- Atrial fibrillation increases stroke risk 17 times.
- Diabetic neuropathies can lead to sensory losses.
- Tertiary syphilis can cause neurosyphilis.
- Recent upper respiratory infections (URIs) may precede meningitis.
- History of seizures may explain current neurologic findings.
- *New onset of seizures between ages 20 and 30*: Trauma, tumor, vascular disease.
- *New onset of seizures after age 50*: Stroke or metastatic cancer.
- Three-fourths of all epilepsy is idiopathic (exact cause unknown).

Immunizations

- What immunizations have you had and when?

- Diphtheria, pertussis, and tetanus (DPT) vaccine contraindicated in patients who have experienced encephalopathy within 7 days of DPT administration.
- Administer DPT cautiously in patients who had seizures within 3 days after previous administration. Immunization information can aid in differential diagnoses.
- *Haemophilus influenzae* type B (HiB) vaccine prevents bacterial meningitis.
- Meningococcal (menomune) vaccine is also available and is highly recommended for college-bound adolescents.

Allergies

- Do you have any allergies, and if so, what are they? Reaction?

- Iodine allergy may prohibit use of invasive studies that require contrast dyes.

Medications

- Are you taking any prescribed or over-the-counter (OTC) medications? Herbal products? If yes, what are they? When last taken?

- Anticoagulants or antiplatelet medications?
- Birth control pills (BCP) or hormone replacement therapy (HRT)?

- Certain medications can affect the neurologic system (see *Drugs That Adversely Affect the Sensory-Neurologic System*). Anticoagulants or products that affect bleeding may increase risk for intracranial bleeding. They may also prevent treatment of stroke patients with tPA.
- BCP and HRT may increase risk for stroke.

Recent Military Service

- Were you exposed to toxic chemicals during military service?

- Veterans returning from Operation Desert Storm had many neurological complaints.

Drugs That Adversely Affect the Sensory-Neurologic System

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Adrenergics	Albuterol sulfate, epinephrine, isoproterenol hydrochloride, terbutaline sulfate	Nervousness, tremors, dizziness, restlessness, insomnia
Adrenergic blockers	Ergotamine tartrate	Lightheadedness, vertigo, insomnia, euphoria, confusion, hallucinations, numbness and tingling of fingers and toes
Antianginals	Diltiazem hydrochloride	Headache, fatigue
	Isosorbide dinitrate, nitroglycerin	Headache, dizziness, weakness, orthostatic hypotension
	Nifedipine	Headache, dizziness, lightheadedness, flushing
	Verapamil hydrochloride	Headache, dizziness
Antiarrhythmics	Lidocaine	Lightheadedness, dizziness, paresthesia, tremors, restlessness, confusion, hallucinations, headache
Anticonvulsants	Carbamazepine	Dizziness, drowsiness, ataxia, confusion, speech disturbances, involuntary movements
	Phenytoin	Dose-related headache, confusion, ataxia, slurred speech, lethargy, drowsiness, nervousness, insomnia, blurred vision, diplopia, nystagmus
Antidepressants	Tricyclic antidepressants	Drowsiness, weakness, lethargy, fatigue, agitation, nightmares, restlessness; confusion, disorientation (especially in elderly patients)
	Monoamine oxidase inhibitors	Restlessness, insomnia, drowsiness, headache, orthostatic hypotension, HTN
Antihypertensives	Clonidine	Drowsiness, sedation, dizziness, headache, nightmares, depression, hallucinations
	Hydralazine	Headache
	Methyldopa	Drowsiness, sedation, decreased mental acuity, vertigo, headache, psychic disturbances, nightmares, depression
Anti-infectives	All beta-blockers	Fatigue, lethargy, vivid dreams, hallucinations, depression
	Aminoglycosides	Neuromuscular blockade; ototoxicity causing vertigo, hearing impairment, or both
	Acyclovir	Myoclonus, seizures
	Isoniazid	Peripheral neuropathy
Antineoplastic agents	Nitrofurantoin	Peripheral neuropathy
	Fludarabine	Dysarthrias, paresthesias, weakness, seizures, paralysis
	Procarbazine	Paresthesia, neuropathy, confusion
Antiparkinsonian agents	Vinblastine sulfate	Paresthesia
	Vincristine sulfate	Peripheral neuropathy, loss of deep tendon reflexes, jaw pain
	Amantadine	Psychic disturbances, nervousness, irritability, fatigue, depression, insomnia, confusion, hallucinations, difficulty concentrating
	Levodopa, pramipexole, ropinirole	Psychic disturbances, decreased attention span, memory loss, nervousness, vivid dreams, involuntary muscle movements
Antipsychotics	Haloperidol, phenothiazines	Extrapyramidal reactions, tardive dyskinesia, headache, lethargy, confusion, agitation, hallucinations
Cholinergic blockers	Atropine sulfate, benztropine mesylate, glycopyrrolate	Blurred vision, headache, nervousness, drowsiness, weakness, dizziness, insomnia, disorientation

(continued)

Drugs That Adversely Affect the Sensory-Neurologic System (continued)

DRUG CLASS	DRUG	POSSIBLE ADVERSE REACTIONS
Corticosteroids	Dexamethasone, hydrocortisone, methylprednisolone, prednisone	Mood swings, euphoria, insomnia, headache, vertigo, behavior
Gastrointestinal agents	Cimetidine	Confusion (especially in elderly patients), depression
	Metoclopramide	Restlessness, anxiety drowsiness, lassitude, extrapyramidal reactions, tardive dyskinesia
Narcotic analgesics	Morphine sulfate, hydromorphone, meperidine, methadone, oxycodone	Sedation, dizziness, visual disturbances, clouded sensorium
	Butorphanol tartrate, nalbuphine, pentazocine	Sedation, headache, dizziness, vertigo, light-headedness, euphoria
Nonsteroidal anti-inflammatory agents	Ibuprofen, indomethacin	Drowsiness, dizziness
Sedatives and hypnotics	Barbiturates	Drowsiness, lethargy, vertigo, headache, depression, "hangover," paradoxical excitement in elderly patients, hyperactivity in children
	Benzodiazepines	Drowsiness, dizziness, ataxia, daytime sedation, headache, confusion
Skeletal muscle relaxants	Baclofen	Drowsiness
	Chlorzoxazone	Drowsiness, dizziness
Miscellaneous agents	Cyclobenzaprine	Drowsiness, dizziness, headache, nervousness, confusion
	Lithium	Lethargy, tremors, headache, mental confusion, dizziness, seizures, difficulty concentrating

Family History

The family history identifies any predisposing or causative factors for neurologic problems.

Family History

RISK FACTORS/ QUESTIONS TO ASK

Familial/Genetically Linked Neurologic Problems

- Do you have a family history of HTN, stroke, MS, seizures, amyotrophic lateral sclerosis (ALS), Huntington's chorea, Alzheimer's disease, or cancer?
- Do you have a family history of substance abuse or psychiatric problems?

RATIONALE/SIGNIFICANCE

- Positive family history for these diseases increases risk to patient.

Review of Systems

The review of systems (ROS) allows you to assess how the neurologic system affects or is affected by every

other system. Often, you may uncover an important fact that your patient failed to mention earlier.

Review of Systems

SYSTEM/QUESTIONS TO ASK

General Health Survey

- How have you been feeling?

- Have you had a fever?

Integumentary

- Have you had any changes in or loss of sensation? Rashes?

HEENT

Head and Neck

- Do you have a change in any of your senses?
- Headaches?
- URIs? Enlarged glands or lumps in your neck?

- Do you have thyroid disease?
- Speech changes or difficulty speaking?

Eyes, Ears, Nose, Throat

- Do you have difficulty swallowing, vision changes, hearing changes, or ringing in your ears (tinnitus)?

Respiratory

- Do you have any breathing problems?

Cardiovascular

- Do you have any cardiovascular problems?
- Have you ever been told that you have an irregular heartbeat?
- Do your hands become numb, tingle, or change color in response to cold?

Gastrointestinal

- Do you have nausea, vomiting, or trouble swallowing?

RATIONALE/SIGNIFICANCE

- Look for affective changes such as depression.
- Lethargy, emotional lability seen with stroke.
- Associated with meningitis.

- *Loss of sensation*: TIA, stroke, neuropathies.
- *Skin rash, petechiae*: Meningococcal meningitis.

- *Headaches*: HTN, intracranial bleeding.
- *URI, enlarged lymph nodes*: Systemic infection that may precede meningitis.
- Hypo- or hyperthyroidism can affect patient's mood.
- *Dysarthria*: Lack of articulation of speech.



HELPFUL HINT

Patients with dysarthria often have dysphagia (difficulty swallowing) as well.

- *Aphasia*: Total loss of comprehension and use of language. Can be receptive (inability to comprehend what is being said, also called Wernicke's or sensory aphasia) or expressive (inability to conceptualize what words one wants to say, also called Broca's or motor aphasia).
- *Dysphasia*: Impairment of speech. Speech can be nonfluent, slow, deliberate, with few words, or fluent, with intact speech that lacks meaning.

- Associated with TIA and stroke.

- Mental status changes can occur with chronic obstructive pulmonary disease (COPD) as a result of hypoxia or hypercapnea.

- Positive history of cardiovascular disease increases risk for stroke.
- Atrial fibrillation, a common dysrhythmia in older adults, increases risk for stroke.
- Raynaud's and Buerger's diseases can cause paresthesia.

- Nausea is often seen with neurologic problems such as head trauma and increased ICP.
- Dysphagia may occur with stroke, MS, MG, ALS.



ALERT

Patients who have difficulty swallowing are at higher risk for aspiration.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> Do you have any bowel or bladder problems or loss of sensation? <p>Reproductive</p> <ul style="list-style-type: none"> Have you had any sexually transmitted diseases (STDs)? <p><i>Males</i></p> <ul style="list-style-type: none"> Do you have any problems with sexual performance? <p><i>Females</i></p> <ul style="list-style-type: none"> When was your last menstrual period? Have you had miscarriages? Are you taking BCPs? <p>Musculoskeletal</p> <ul style="list-style-type: none"> Do you have any loss of sensation, numbness, or weakness? Have you experienced paralysis? <p>Endocrine</p> <ul style="list-style-type: none"> Do you have diabetes mellitus or thyroid disease? <p>Lymphatic/Hematological</p> <ul style="list-style-type: none"> Do you have any blood disorders? Do you bruise easily? 	<ul style="list-style-type: none"> <i>Bowel problems, incontinence, loss of bowel function:</i> Cauda equina syndrome caused by compression of sacral nerve, can occur with herniated disc or spinal stenosis. <i>Transient loss of bowel and bladder function:</i> Stroke or spinal cord dysfunction. <i>Incontinence:</i> Tonic-clonic seizures. Tertiary syphilis can cause neurosyphilis. Loss of sensation and inability to achieve erection may be neurologic in origin. Problems with erection can occur with neuropathies and CNS disease such as Parkinson's disease. Hypercoagulable states increase the risk for stroke. Associated with many neurologic disorders. Neuropathies may be associated with diabetes mellitus. Paralysis can lead to muscle atrophy from disuse. Diabetes can cause neuropathies and muscle wasting. Hypothyroidism can cause weakness, lethargy, flat affect, and labile affect. Bleeding disorders may increase risk for stroke. Patients with platelet count <100,000 and International Normalized Ratio (INR) >1.7 are not candidates for tPA.

Psychosocial Profile

The psychosocial profile reveals patterns in the patient's life that may affect the neurologic system and put her or

him at risk for neurologic disorders. It may also identify teaching needs.

Psychosocial Profile

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Health Practices and Beliefs/Self-Care Activities</p> <ul style="list-style-type: none"> What activities do you engage in to promote the health of your neurologic system? Do you get an annual physical exam? Do you wear seat belts? Do you own a gun? Do you wear protective equipment when playing sports? Do you understand proper body mechanics? <p>Typical Day</p> <ul style="list-style-type: none"> How would you describe your typical day? Are you able to perform self-care activities? Do you have difficulty performing ADLs? 	<ul style="list-style-type: none"> Helps identify health promotion and preventative activities. Motor vehicle accidents, falls, gunshot wounds, and sports injuries are the most common causes of spinal cord trauma. Back injuries can cause herniated discs and subsequent neurologic deficits. Neurologic system impairment may hinder patient's ability to perform ADLs. Assess correct use of assistive devices.

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Nutritional Patterns</p> <ul style="list-style-type: none"> ■ What did you eat in the last 24 hours? ■ Are you on any special diets? 	<ul style="list-style-type: none"> ■ <i>Vitamin B deficiency</i>: Paresthesia in feet and hands, gait disturbances. ■ <i>Vitamin B₁₂, niacin, folic acid deficiencies</i>: Peripheral neuropathy. ■ Foods that can trigger headaches include those with amines (cheese and chocolate) and nitrites (hot dogs), vinegar, onions, fermented or marinated foods, monosodium glutamate, caffeine, and ice cream. ■ High-cholesterol diet is risk factor for vascular disease.
<p>Activity and Exercise Patterns</p> <ul style="list-style-type: none"> ■ Do you exercise? If so, what type and how much? <p>Recreation/Hobbies</p> <ul style="list-style-type: none"> ■ Do you participate in contact sports or activities that require heavy lifting? If so, do you wear protective equipment? ■ What are your hobbies? 	<ul style="list-style-type: none"> ■ Strenuous activity increases risk for injury. ■ Recreational activities may increase risk for injury. ■ Use of protective equipment decreases risk. ■ Certain hobbies increase risk for exposure to toxic substances (e.g., lead exposure when working with stained glass).
<p>Sleep and Rest Patterns</p> <ul style="list-style-type: none"> ■ How many hours of sleep do you get per night? 	<ul style="list-style-type: none"> ■ Sleep deprivation can affect cerebral function. ■ Increased somnolence may indicate an increase in ICP. ■ Neurologic problems may interfere with sleep. ■ Patients with pain associated with herniated discs may have difficulty finding comfortable sleep position.
<p>Personal Habits</p> <ul style="list-style-type: none"> ■ Do you smoke, drink alcohol or caffeinated beverages, or use drugs? 	<ul style="list-style-type: none"> ■ Cigarette smoking is risk factor for cerebrovascular disease. Alcohol, caffeine, and drug use can alter cerebral function.
<p>Occupational Health Patterns</p> <ul style="list-style-type: none"> ■ What type of work do you do? Does it involve lifting, repetitive movements, or exposure to chemicals or toxins? 	<ul style="list-style-type: none"> ■ Occupations that increase risk for head or back injury may result in neurologic problems. ■ Neurologic disorders may prevent patient from returning to previous work. ■ Exposure to toxic substances can cause neurologic problems. ■ Radiation and carcinogens increase risk for brain tumors. ■ Arsenic and mercury can cause ataxia.
<p>Environmental Health Patterns</p> <ul style="list-style-type: none"> ■ Where do you live? ■ How many floors are there in your house? <p>Roles/Relationships/Self-Concept</p> <ul style="list-style-type: none"> ■ How do you see yourself? ■ How did you see yourself before you had this problem, and how do you see yourself now? ■ What is your role in the family, and has it changed? 	<ul style="list-style-type: none"> ■ Helps determine discharge planning needs. ■ Disabling/crippling problems or deformities associated with some neurologic disorders may cause body image disturbances and affect patient's self-image and self-esteem. ■ Neurologic problems such as MS may alter patient's role in family.

(continued)

Psychosocial Profile (continued)

CATEGORY/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Religious/Cultural Influences ■ What are your religious practices and beliefs or cultural influences in regard to health care?	■ May affect patient's healthcare practices.
Sexuality Patterns ■ Have physical problems interfered with your usual sexual activity?	■ Neurologic problems may interfere with sexual activity.
Social Supports ■ Who are your supports? ■ Have physical problems interfered with your ability to interact with others?	■ Helpful in planning care. ■ Chronic neurologic problems can disable a patient, impair socialization, and affect independence.
Stress and Coping ■ How do you deal with stress?	■ Stress can cause headaches and trigger an exacerbation of MS.



Case Study Findings

Reviewing Mr. Webster's

Biographical data:

- 21 years old.
- African American.
- Baptist religion.
- Senior in college; business major.
- Insured under family's medical plan while in college.
- Lives on campus during school year; home is 1 hour from school.
- Contact person: Janet Webster, mother.
- Source: Self, seems reliable.

Current health status:

- Complains of headache and neck pain from hitting head in motor vehicle accident.
- Had a few beers earlier in night.
- Felt a little confused after accident and is not sure what happened. Does not know whether he lost consciousness.

Past health history:

- No hospitalizations or surgeries.
- Concussion in high school while playing football, otherwise negative for neurologic problems.
- Fractured fibula freshman year while playing football; injury casted and healed without problem.
- Thinks immunizations are up to date but unsure of last tetanus shot.
- No known food, drug, or environmental allergies.
- No medications or other medical problems.

Family history:

- Father, age 55, has HTN.
- Paternal uncle, age 60, has HTN.
- Paternal grandfather died at 80 from stroke.
- Maternal aunt, age 50, has HTN.
- Maternal grandfather died at 75 from stroke.

Review of systems:

- *General Health Survey*: Usual state of health good until accident.
- *Integumentary*: No change.
- *HEENT*: Headache and neck pain; a "little dizzy."
- *Respiratory*: No problems.
- *Cardiovascular*: No problems.
- *Gastrointestinal*: Appetite usually good, but felt "a little sick after accident." No vomiting.
- *Genitourinary*: No problems.
- *Reproductive*: Does not wish to discuss.
- *Endocrine*: No problems.
- *Lymphatic/Hematologic*: No problems.

Psychosocial profile:

- Gets yearly physical exam for sports.
- Typical day consists of awakening at 9 A.M., grabbing some breakfast, and heading to class. Goes to football practice from 3 P.M. to 6 P.M. 7 days a week, has dinner, relaxes, and does school work until about midnight. Usually goes to bed by 1 A.M.
- Usually eats well; diet high in carbohydrates and protein.
- Practices football with team 3 hours a day, 7 days a week. Wears protective gear.
- Usually gets at least 8 hours of sleep a night.

- Denies use of drugs or nicotine. Admits to drinking alcohol on weekends; amount varies.
- Does well in school (grade point average 3.0).
- Does not wish to discuss sexuality patterns.
- Friends and family are supports.
- Copes with stress by exercising, partying with friends on weekends.
- Attends church occasionally, no religious or cultural influence on health.



CRITICAL THINKING ACTIVITY # 1

What other questions would you ask Mr. Webster to further assess his chief complaints?



Case Study Evaluation

Before you move on to the physical examination, consider what you have learned from your interview with Mr. Webster. Document his sensory-neurologic history.



CRITICAL THINKING ACTIVITY # 2

From the subjective information that you have obtained from Leon's history, what neurologic problem is he at risk for? What factors contribute to this situation?

Physical Assessment

Once you have obtained the subjective data, focus on collecting the objective data by performing the physical examination. The components of the neurologic examination include tests of the patient's:

- Mental status.
- Cranial nerve function.
- Sensory function.
- Reflex function.

As a nurse, you will perform an abbreviated screening neurologic examination more often than the comprehensive neurologic examination described here. Although it is briefer, the screening examination still addresses each area of the neurologic assessment and includes:

- Evaluation of loss of consciousness (LOC) with brief mental status examination and evaluation of verbal responsiveness.
- Testing of selected cranial nerves (usually CNs II, III, IV, and VI).
- Motor screening, including strength, movement, and gait (see Chapter 20, Assessing the Musculoskeletal System).
- Sensory screening, including tactile and pain sensations on upper and lower extremities.
- Reflexes.

If the patient's condition warrants it, continue to assess neurologic status by performing rapid, repeated checks to evaluate LOC, pupil size and reaction, responsiveness, extremity strength and movement, and vital signs.



ALERT

Remember, one of the earliest indicators of increased ICP is a subtle change in LOC, a loss of detail, forgetfulness, restlessness, or sudden quietness. Be alert for these changes. Early detection with prompt intervention may not only prevent major neurologic damage but also save your patient's life.

Approach

The neurologic physical assessment sequence is different from that for other body systems, but you still use inspection, palpation, and auscultation. The main difference is that this examination consists of a series of tests. Even though the physical assessment establishes the objective database, many of the findings are still somewhat subjective, based on the patient's perception.

Because the exam is lengthy, make sure that the patient is seated comfortably, preferably in a chair with back support rather than on an examination table. Check from time to time to be sure that he or she is not becoming fatigued.

The patient with a neurologic disorder may find it difficult to understand instructions and participate fully in the exam. In these cases, limit tests that require a lengthy explanation and active cooperation. If the patient is lethargic or exhibits any other possible alteration in LOC, then it is critical to perform an assessment for LOC.

When assessing older adults, keep in mind that their responses to questions and directions may be slower and that you may need to adjust the pace of the examination. Because of the normal changes of aging and the possibility of paresis, stay close to patients to prevent them from falling, and help with position changes.



TOOLBOX

Assemble the following equipment for a routine neurologic examination: stethoscope and BP cuff; penlight; nonsterile gloves; wisp of cotton; sharp object such as toothpick or sterile needle; objects to touch, such as a coin, button, key, or paper clip; something fragrant, such as rubbing alcohol or coffee; something to taste, such as lemon juice (sour substance), sugar, salt, and quinine (bitter substance); tongue blade; two test tubes or other vials; a reflex hammer; and an ophthalmoscope.

Performing a General Survey

Before you begin the specific neurologic assessment, perform a general survey, including scanning your patient from head to toe and taking vital signs. Look at every system as it relates to the neurologic system. If you detect any changes, investigate further as you perform the specific neurologic examination.

The general survey can provide immediate and important information about the patient's neurologic-neurologic status. Affect, hygiene, grooming, speech, posture, and body language can provide clues to your patient's general level of functioning, as well as any pain or impairment. Ask yourself:

- Does the patient appear her or his stated age?
- Is her or his affect or mood appropriate for the situation?
- Are her or his responses appropriate?
- Is she or he following the interview?
- Is she or he well groomed and neatly dressed?
- Are her or his clothes stained with food?
- Is a female patient's make-up appropriately applied?

Scan for symmetry, especially in the face. Ask your patient to smile, and look for symmetry of facial features. Are there any abnormal movements? Shake your patient's hand and note muscle strength. Look at facial expression and eye contact. Is his or her speech clear or slurred? Are responses slow and deliberate or one-word?

Poor grooming, food on clothing, or inappropriate make-up can reflect visual deficits such as visual field cuts; motor deficits, such as weakness or paralysis; impaired cerebral function, such as confusion; and affective problems such as depression. Poor posture or facial asymmetry (ptosis or droopy smile) may be a sign of weakness or hemiparesis. Abnormal movements or balance and coordination problems may indicate a cerebellar dysfunction. Weak hand grip may reflect weakness of a neurologic nature. Inappropriate responses, short attention span, or poor eye contact may indicate an underlying cerebral function disorder or a psychiatric problem.



HELPFUL HINT

Two places to assess for symmetry of facial features are the nasolabial fold (smile crease) and the palpebral fissures (distance between the eyelids).



ALERT

Cushing's triad—widening pulse pressure, bradycardia, and irregular breathing patterns—is a late sign of increasing ICP and impending brain herniation.

Vital Signs

Because the neurologic system plays a vital role in the regulation of vital signs, a change in vital signs can reflect a change in the neurologic system. For example:

- *BP*: HTN is a risk factor for stroke. A widened pulse pressure is a Cushing's sign of increased ICP.
- *Pulse*: Bradycardia is a Cushing's sign indicating increased ICP; atrial fibrillation increases the risk for stroke.
- *Respirations*: Irregular breathing patterns with periods of apnea is a Cushing's sign of increased ICP.
- *Temperature*: A temperature elevation may be associated with infection, meningitis, or brain abscess. A slight temperature elevation may occur after a stroke as a result of the inflammatory response, and a high temperature can occur as a result of a brainstem stroke.

Performing a Head-to-Toe Physical Assessment

The neurologic system affects the functioning of all other body systems. Therefore, it is essential to examine the patient from head to toe to note any unusual findings. Although not every abnormality you note will be related to neurologic dysfunction, you should consider a variety of possibilities. For example, a bruise on the forehead may be caused by an injury sustained when a patient with a preexisting neurologic impairment experienced an LOC. On the other hand, the patient may have been healthy before the injury, and the injury may be causing the current symptoms of pain and disorientation.



HELPFUL HINTS

- If your patient has a spatial perception problem, be aware that she or he may have **neglect**, a spatial perception problem in which she or he does not see the affected side as part of her or his body.
- Remember: Problems with the right side of the brain will be clinically manifested on the left side, and vice versa.
- Before allowing a patient with suspected stroke to eat, test gag, swallow, and cough reflexes to avoid aspiration.



ALERTS

- Safety is an issue for patients with right-side brain injury because they tend to be impulsive and not know their own limits.
- When assessing a possible stroke patient, remember that if your patient has dysarthria, he or she may also have dysphagia.

Stroke Assessment Scales

Stroke is the third leading cause of death in the United States. For those who survive a stroke, it is a leading cause of disability. Prompt recognition of signs and symptoms of stroke along with immediate treatment can prevent permanent neurologic deficits and disability.

Cincinnati Prehospital Stroke Scale

The Cincinnati Prehospital Stroke Scale can be used to quickly assess for signs of stroke. Presence of even one sign, either facial drop, motor weakness, or slurred speech, warrants immediate evaluation.

- *Look for facial droop:* Have patient smile or show teeth.
- *Test for motor weakness:* Have patient perform pronator drift (see Chapter 20, Assessing the Motor-Musculoskeletal System).
- Listen for speech problems: Have patient say, "You can't teach an old dog new tricks."

National Institute of Health Stroke Scale

The National Institute of Health Stroke Scale quantifies severity of stroke. This scale is used for the initial evaluation, and then to monitor the patient's progress and response to treatment. It has 12 categories with a total range of score from 0 (normal) to 42 (most severely impaired). Categories include:

- LOC.
- LOC questions.
- LOC commands.
- Gaze abnormalities.
- Visual loss.
- Facial weakness.
- Motor weakness in arms and legs.
- Limb ataxia.
- Sensory loss.
- Language.
- Dysarthria.
- Extinction and inattention.

Performing a Head-to-Toe Physical Assessment

SYSTEM/NORMAL FINDINGS

ABNORMAL FINDINGS/RATIONALE

General Health Survey

INSPECT

- No acute distress, AAO × 3, responses clear and appropriate, affect pleasant, well groomed.

- *Changes in LOC:* Early sign of developing CNS disorder.
- *Dysarthria:* Associated with many neurological disorders.
- Inability to care for self as reflected by poor grooming may be caused by a neurological problem, such as hemiparesis or a psychiatric problem, such as depression.

Integumentary

INSPECT

- Superficial sensations intact.
- No rashes or lesions.

- Peripheral neuropathies, spinal cord injuries, and stroke: Can cause sensory deficits.
- Skin rash and petechiae: Meningitis.

HEENT

Head and Neck

INSPECT

- Facial expression appropriate and symmetrical, lymph nodes not palpable.
- Cranial nerves intact.

- *Asymmetrical facial features:* CVA.
- *Masklike facial expression:* Parkinson's disease.
- *Enlarged lymph nodes:* Infection.

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

SYSTEM/NORMAL FINDINGS	ABNORMAL FINDINGS/RATIONALE
<ul style="list-style-type: none"> ■ Gag reflex intact (CNS, IX and X intact). <p><i>Eyes</i> <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ Eyes clear and bright; pupils equal, round, and reactive to light accommodation (PERRLA); pupils direct and consensual; EOM intact; peripheral vision intact; no papilledema (CNII, III, IV, and VI). <p><i>Respiratory</i> <i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Respiratory rate 18/pc min. Lungs clear. <p><i>Cardiovascular</i> <i>AUSCULTATE</i></p> <ul style="list-style-type: none"> ■ Regular heart rate and rhythm. ■ No carotid bruits or thrills. <p><i>Gastrointestinal</i> <i>INSPECT/AUSCULTATE/PALPATE</i></p> <ul style="list-style-type: none"> ■ Swallowing intact. ■ Positive bowel sounds. ■ Abdomen soft, nontender. <p><i>Genitourinary/Reproductive</i> <i>INSPECT</i></p> <ul style="list-style-type: none"> ■ Bladder function intact. <p><i>Musculoskeletal</i> <i>INSPECT/PALPATE</i></p> <ul style="list-style-type: none"> ■ Muscle strength +5. ■ Movements smooth and coordinated. ■ Gait balanced and fluid, with conformity of phases. ■ Negative Romberg test. 	<ul style="list-style-type: none"> ■ <i>Abnormal cranial nerves</i>: Neurological problem. ■ <i>Visual changes</i>: Increased ICP. ■ <i>Peripheral field deficits</i>: Neurological problems. ■ <i>Papilledema</i>: Increased ICP. ■ <i>Irregular breathing pattern</i>: Increased ICP. ■ Hypoxia can cause confusion and increased ICP. ■ Atrial fibrillation increases risk for embolic stroke. ■ Bradycardia is a sign of increased ICP. ■ Carotid bruits or thrills reflect carotid stenosis, increasing risk for thrombotic stroke. ■ Dysphagia can occur with neurological disease. ■ Decreased bowel sounds and constipation are most common bowel problem in stroke patients. ■ STDs such as syphilis can cause neurosyphilis. ■ Poor bladder control or incontinence frequently occur during acute phase of CVA. Can also occur with spinal cord injuries and other neurological disorders. ■ <i>Decreased sensations and numbness</i>: Neuropathies. ■ Paralysis can result in muscle atrophy. ■ <i>Balance problems</i>: Cerebellar dysfunctions. ■ Spatial perception problems can result from a stroke.

Performing a Neurologic Assessment

Once you have completed the head-to-toe scan, zero in on the specifics of the sensory-neurologic examination. Begin by assessing cerebral function, and then assess cranial nerve function, sensory function, and reflexes. The motor-musculoskeletal system was assessed in the previous chapter.

Cerebral Function

Assessment of cerebral function includes LOC, mental status and cognitive functioning, and communication. (See *Assessing Cerebral Function*.)

Level of Consciousness

Evaluating LOC involves assessing arousal (wakefulness) and orientation (ability to receive and accurately interpret sensory stimuli).

Assessing Arousal

Determine the arousal state first, using minimal stimuli and increasing intensity as needed. Start with auditory stimuli, move to tactile stimuli, and then use painful stimuli as a last resort.

Auditory and Tactile Stimuli. To assess auditory stimuli, determine whether the patient is sleeping or awake. If the patient is awake, what is she or he doing? If the

(Text continued on page 769)

AREA/SYSTEM	SUBJECTIVE DATA	AREA/SYSTEM	OBJECTIVE DATA
General	<i>Ask about:</i> General health Fever	General	<i>Measure:</i> Vital signs <i>Observe:</i> Mental status, grooming, affect, behavior, symmetry
Integumentary	<i>Ask about:</i> Rashes Changes in sensations	Integumentary	<i>Inspect:</i> Rashes or petechiae <i>Test:</i> Superficial sensations
HEENT	<i>Ask about:</i> Changes in sense of smell, taste, sight, hearing, touch Headaches Changes in speech or swallowing	HEENT	<i>Inspect:</i> Symmetry of facial features <i>Palpate:</i> Lymph nodes, thyroid <i>Examine:</i> Optic disk <i>Test:</i> Cranial nerves
Respiratory	<i>Ask about:</i> Breathing difficulty	Respiratory	<i>Auscultate:</i> Lungs
Cardiovascular	<i>Ask about:</i> History of CV problems	Cardiovascular	<i>Auscultate:</i> Heart sounds, noting rhythm Carotids for bruits and thrills
Gastrointestinal	<i>Ask about:</i> Nausea, vomiting Changes in bowel function Difficulty swallowing	Gastrointestinal	<i>Auscultate:</i> Bowel sounds
Genitourinary/ Reproductive	<i>Ask about:</i> History of STDs Changes in sexual activity, desire, ability Changes in bladder function	Genitourinary/ Reproductive	<i>Inspect:</i> Lesions <i>Palpate:</i> Bladder distention
Musculoskeletal	<i>Ask about:</i> Muscle weakness Paralysis Problems walking Balance problems	Musculoskeletal	<i>Inspect:</i> Gait <i>Test:</i> Muscle strength ROM Cerebellar function
Endocrine	<i>Ask about:</i> History of diabetes mellitus and thyroid disease		
Lymphatic/ Hematologic	<i>Ask about:</i> Fever Bruising		

Assessment of the Sensory-Neurologic System's Relationship to Other Systems

Neurologic Problems

- **Agnosia:** Inability to recognize object by sight (visual agnosia), touch (tactile agnosia), or hearing (auditory agnosia).
- **Akinesia:** Complete or partial loss of voluntary muscle movement.
- **Aphasia:** Absence or impairment of ability to communicate through speech, writing, or signs.
- **Expressive (motor) aphasia:** Inability to express language even though person knows what he or she wants to say (also called *Broca's* or *motor aphasia*). Frontal lobe affected.
- **Fluent aphasia:** Words can be spoken but are used incorrectly.
- **Nonfluent aphasia:** Slow, deliberate speech, few words.
- **Receptive (sensory) aphasia:** Inability to comprehend spoken or written words (also called *Wernicke's* or *sensory aphasia*). Temporal lobe affected in auditory-receptive; parieto-occipital nerve affected in visual-receptive.
- **Apraxia:** Inability to carry out learned sequential movements or commands.
- **Circumlocution:** Inability to name object verbally, so patient talks around object or uses gesture to define it.
- **Dysarthria:** Defective speech; inability to articulate words; impairment of tongue and other muscles needed for speech.
- **Dysphasia:** Impaired or difficult speech.
- **Dysphonia:** Difficulty with quality of voice; hoarseness.
- **Neologisms:** Made-up, nonsense, meaningless words.
- **Paraphrasia:** Loss of ability to use words correctly and coherently; words are jumbled or misused.
- **Tremors:** Involuntary movement of part of body.
- **Intension tremor:** Involuntary movement when attempting coordinated movements.
- **Fasciculation:** Involuntary contraction or twitching of muscle fibers.

Assessing Cerebral Function

COMPONENT/TECHNIQUE/ NORMAL VARIATIONS

Behavior

- Watch patient's facial expression as she or he responds to questions.
- Note his or her posture, grooming, and affect.
- *Well-groomed, erect posture, pleasant facial expression, appropriate affect.*



HELPFUL HINT

Normal findings vary depending on situation (e.g., if patient is in pain, then her or his behavior should reflect this).

Loss of Consciousness

- Test orientation to time, place, and person.
- *Awake, alert, and oriented to time, place, and person (AAO × 3).*

Mental Status and Cognitive Function

Memory

- **Test immediate recall:** Ask patient to repeat three numbers, such as "4, 9, 1." If patient can do so, ask her or him to repeat a series of five digits.

ABNORMAL FINDINGS/RATIONALE

- **Lack of facial expression/inappropriate expression for speech content:** Possible psychological disorder (e.g., depression or schizophrenia) or neurologic impairment affecting cranial nerves.
- **Masklike expression:** Parkinson's disease.
- **Poor grooming/slumped posture:** Depression if psychological in origin; or stroke with hemiparesis if physiological in origin.
- Disorientation may be physical in origin, such as exhaustion, anxiety, hypoxia, fluid and electrolyte imbalance, drugs, or a neurologic problem.
- Disorientation can also be psychiatric in origin such as with schizophrenia.
- Memory problems can be benign or signal a more serious neurologic problem such as Alzheimer's disease. Forgetfulness—especially for immediate and recent events—is often seen in older adults. With benign

**COMPONENT/TECHNIQUE/
NORMAL VARIATIONS**

- *Test recent memory:* Ask what patient had for breakfast.
- *Test long-term memory:* Ask patient to state his or her birthplace, recite his or her Social Security number, or identify a culturally specific person or event, such as the name of the previous president of the United States or the location of a natural disaster.
- *Immediate, recent, and remote memory intact.*


HELPFUL HINTS

- Avoid using number sequences with a meaningful arrangement, such as 2, 4, 6, 8.
- If you ask your patient for her or his birthday or dates of other events, be sure that you can validate them.

Mathematical and Calculative Ability

- Ask patient to perform a simple calculation, such as adding $4 + 4$. If successful, proceed to more difficult calculation, such as 11×9 .
- *Mathematical/calculative ability intact and appropriate for patient's age, educational level, and language facility.*

General Knowledge and Vocabulary

- Ask how many days in a week and months in a year.

Thought Process

- Ask patient to define familiar words such as "apple," "earthquake," and "chastise." Begin with easy words and proceed to more difficult ones. Remember to consider the patient's age, educational level, and cultural background.


HELPFUL HINTS

- Be sure to take note of your patient's responses throughout the assessment.
- Thought processes are difficult to assess in children younger than age 7 or 8 without advanced training.

- *Thought process intact.*

Abstract Thinking

- Ask patient to interpret a culturally appropriate proverb.
- *Able to generalize from specific example and apply statement to human behavior. Children should be able to distinguish like from unlike as appropriate for their age and language facility.*

ABNORMAL FINDINGS/RATIONALE

forgetfulness, person can retrace or use memory aids to help with recall. Pathological memory loss, as in Alzheimer's disease, is subtle and progressive until ability to function is impaired.

- Temporary memory loss may occur after head trauma.
- Retrograde amnesia is memory loss for events just preceding illness or injury.
- Postconcussion syndrome can occur 2 weeks to 2 months after injury and may cause short-term memory deficits.

- Inability to calculate at level appropriate to age, education, and language ability requires evaluation for neurologic impairment.

- Incoherent speech, illogical or unrealistic ideas, repetition of words and phrases, repeatedly straying from topic, and suddenly losing train of thought are examples of altered thought processes that indicate need for further evaluation.

- Inability to define familiar words requires further evaluation.
- Alteration in thought process can be caused by a physical disorder such as dementia, a psychiatric disorder such as psychosis, or drugs and alcohol.

- *Impaired ability to think abstractly:* Dementia, delirium, mental retardation, psychoses.

(continued)

Assessing Cerebral Function (continued)

COMPONENT/TECHNIQUE/ NORMAL VARIATIONS

ABNORMAL FINDINGS/RATIONALE

Judgment

- Observe patient's response to current situation.
- Ask patient to respond to a situation or hypothetical situation.
- *Judgment appropriate and intact.*

Communication

Speech and Language

- Listen to patient's rate and ease of speech, including enunciation.
- *Speech flows easily; patient enunciates clearly. Sophistication of speech matches age, education, and fluency.*

Spontaneous Speech

- Show patient a picture and have him or her describe what he or she sees.
- *Spontaneous speech intact.*

Motor Speech

- Have patient repeat, "do, ray, me, fa, so, la, ti, do."
- *Motor speech intact.*

Automatic Speech

- Have patient say something that is committed to memory, such as days of week or months of year.
- *Automatic speech intact.*

Sound Recognition

- Have patient close eyes and identify familiar sound such as clapping hands.
- *Sound recognition intact.*

Auditory-Verbal Comprehension

- Ask patient to follow simple directions or explain meaning of a series of words.
- *Auditory-verbal comprehension intact.*

Visual Recognition

- Have patient identify familiar object by sight (e.g., cup, pencil, pen).
- *Visual recognition intact.*

Visual-Verbal Comprehension

- Have patient read sentence from newspaper and explain meaning.
- *Visual-verbal comprehension intact.*

Writing

- Have patient write name, address, simple sentence, one word with eyes open and then closed, name of an object.
- *Writing ability intact.*

Copying Figures

- Show patient several figures and ask her or him to copy them, increasing in complexity. (e.g., circle, X, square, triangle, star).
- *Able to copy figures.*

- Impaired judgment can be associated with dementia, psychosis, or drug and alcohol abuse.

- *Hesitancy, stuttering, stammering, unclear speech:* Lack of familiarity with language, deference or shyness, anxiety, neurologic disorder.
- *Dysphasia/aphasia:* Neurologic problems such as stroke.
- Drugs and alcohol can also cause slurred speech.

- *Impaired spontaneous speech:* Cognitive impairment.

- *Impaired motor speech (dysarthria):* Problem with CN XII.

- *Impaired automatic speech:* Cognitive impairment or memory problem.

- *Impaired sound recognition (auditory agnosia):* Temporal lobe affected.

- *Impaired auditory-verbal comprehension:* Temporal lobe affected.

- *Expressive aphasia:* Frontal lobe affected.
- *Auditory-receptive aphasia:* Temporal lobe affected.

- *Impaired visual recognition (visual agnosia):* Parieto-occipital lobe affected.

- *Impaired visual-verbal comprehension (visual-receptive aphasia):* Cognitive impairment.

- Impaired writing ability.

- Impaired figure copying ability.

patient is sleeping, call her or him by name in a normal tone of voice. If she or he does not respond, speak louder. If auditory stimuli fail, try tactile. Gently touch the patient's hand. If she or he does not respond, gently shake her or his shoulder.

Painful Stimuli. If your patient does not respond to tactile stimuli, you will have to resort to painful stimuli. There are acceptable and unacceptable ways to elicit a response to pain. Never perform a nipple twist. Avoid using a pin or needle, because if the skin breaks, you risk infection. Also remember to rotate sites—repeated stimulation at the same site may cause bruising.

Painful stimuli may be central or peripheral. Central painful stimuli include the trapezius squeeze, the sternal rub, supraorbital pressure, and mandibular pressure. Peripheral painful stimuli include nail pressure and the Achilles tendon squeeze. Apply the stimulus for 15 to 30 seconds. A responsive patient will experience pain and move in response to these stimuli:

- **Trapezius squeeze:** Pinch 1 to 2 inches of the trapezius muscle and twist.
- **Sternal rub:** With the knuckles or the palm of your dominant hand, apply pressure in a grinding motion to the sternum. Do not use this site repeatedly because it will cause bruising.
- **Supraorbital pressure:** Apply firm pressure with your thumbs at the notch at the center of the orbital rim below the eyebrows. Because a nerve runs in the notch, pressure to this area will cause sinus pain. Use this stimulus carefully to avoid damage to the eyes.
- **Mandibular pressure:** With your index and middle finger, apply inward and upward pressure at the angle of the jaw.
- **Nail pressure:** Apply pressure over the moon of the nail with a pen or pencil.
- **Achilles tendon squeeze:** Squeeze the Achilles tendon between your thumb and your index finger.

Documenting Arousal

When documenting your findings, record how your patient responds rather than simply giving the response a label. For example, charting that a patient responds slowly to verbal stimuli but drifts back to sleep is much more descriptive than simply writing “lethargic.” Because a change in LOC is an early sign of a neurologic problem, you need to be able to detect subtle changes in your patient. Describing the response to the stimuli is more objective and allows better comparisons during follow-up assessments.

The Glasgow Coma Scale (GCS) provides a more objective way to assess the patient's LOC. It evaluates best eye response, best motor response, and best verbal response on a scale of 3 to 15. Fifteen (highest score) indicates that the patient is awake, alert, oriented, and able to follow simple commands. Three (lowest score)

Terms Used to Describe Level of Consciousness

- **Alert:** Follows commands in a timely fashion.
- **Lethargic:** Appears drowsy, may drift off to sleep during examination.
- **Stuporous:** Requires vigorous stimulation (shaking, shouting) for a response.
- **Comatose:** Does not respond appropriately to either verbal or painful stimuli.

indicates that the patient does not respond to any stimulus and has no motor or eye response, reflecting a very serious neurologic state with poor prognosis. (See *Glasgow Coma Scale*.)

Assessing Orientation

Next, test orientation to time, place, and person. Purpose is also sometimes included as a fourth area of orientation. Avoid asking questions that require only a “yes” or “no” response. Start with specifics; then be more general, if necessary. For example, if your patient does not know the specific date, ask him or her what month it is, and if he or she does not know the month, ask what season it is. If he or she was once oriented to place and is not the next time you ask, this may be an early sign of a deteriorating neurologic status.

Time. Ask the patient to state the date, including the year and day of the week. Hospitalized patients—especially older adults—can easily become disoriented to time, but they usually reorient easily. Be sure your documentation reflects this.



HELPFUL HINT

When an older patient is disoriented to time, she or he usually thinks it is an earlier date, not a later one.



ALERT

If the patient's responses are bizarre—such as thinking it is the year 2050—consider a psychiatric problem.

Place. Ask your patient to state where he or she is. Can he or she identify environmental cues (e.g., bed, equipment, sound of bells or buzzers) to determine location? A person who is usually oriented but becomes confused when hospitalized often temporarily mistakes the hospital room for

Glasgow Coma Scale

OBSERVATION	RESPONSE ELICITED	SCORE
Eye response	■ Opens spontaneously	4
	■ Opens to verbal command	3
	■ Opens to pain	2
	■ No response	1
Motor response	■ Reacts to verbal command	6
	■ Reacts to painful stimuli	
	■ Identifies localized pain	5
	■ Flexes and withdraws	4
	■ Assumes flexor posture	3
	■ Assumes extensor posture	2
Verbal response	■ No response	1
	■ Is oriented and converses	5
	■ Is disoriented, but converses	4
	■ Uses inappropriate words	3
	■ Makes incomprehensible sounds	2
	■ No response	1

Eye and verbal responses are self-explanatory. Under motor responses, “Identifies localized pain” denotes that the patient is not fully conscious but is aware enough to respond to an annoying stimulus. “Flexes and withdraws” is a lower-level motor response and indicates that the patient pulls away from painful stimuli. Be sure to apply the stimulus long enough to elicit more than just a reflex response.

Posturing is a still lower level of response and indicates abnormal positions of flexion or extension of the arms with legs extended and internally rotated and feet plantarflexed. It is an ominous sign that usually has a poor prognosis. In **flexion (decorticate) posturing**, the arms are flexed to the chest and the hands are clenched and internally rotated as a result of a lesion at or above the brainstem in the cerebral cortex. In **extension (decerebrate) posturing**, the arms are extended and the hands are clenched and hyperpronated as a result of a midbrain (brainstem) lesion. This is the more serious of the two.

home. You will need to address safety concerns with patients who are disoriented both at home and in the hospital; for example, those with Alzheimer’s disease, who tend to wander.

Person. Ask the patient to state her or his name. Self-identity usually remains intact the longest, making disorientation to person an ominous sign.

Mental Status and Cognitive Function

Once you have determined that your patient is arousable and oriented, assess level of awareness. Level of awareness reflects mental status and cognitive function. It is the functional state of the mind as judged by a person’s behavior, appearance, response to stimuli, speech, memory, and judgment. It reflects the person’s connection with his or her environment. These areas are at a higher level of functioning than LOC and reflect the cerebral cortex’s ability to process and respond. Be sure to explain to your patient that you need to ask questions that have obvious answers to accurately assess his or her neurologic status.

Like the general survey, a mental status screening is often integrated into the health history interview. A typical screening consists of 10 questions that address each area of the detailed mental status examination. (See Mental Status Screening Questions, page 785). Another instrument that can be used to assess cognitive function is the Mini-Mental State Examination (MMSE) (Folstein, 1975). It is used to screen for and monitor dementia. You will need to perform a rigorous mental status examination in the following situations:

- If data from patient or patient behavior during the health history interview suggests an abnormality.
- If family members or caregivers report changes in the patient’s personality or behavior.
- If the patient has a history of head injury, stroke, dysphasia or aphasia, or mental illness.

A mental status and cognitive function assessment includes: memory, general knowledge and vocabulary, mathematical and calculative skills, and thought process/abstract reasoning/judgment.

Mental Status Screening Questions

QUESTION	FUNCTION SCREENED
What is your name?	Orientation to person
What is today's date?	Orientation to time
What year is it?	Orientation to time
Where are you now?	Orientation to place
How old are you?	Memory
Where were you born?	Remote memory
What did you have for breakfast?	Recent memory
Who is the U.S. president?	General knowledge
Can you count backward from 20 to 1?	Attention and calculation skills
Why are you here?	Judgment

Memory

Assess immediate, recent, and remote memory. Test immediate memory by asking your patient to repeat a series of numbers. Test recent memory by asking what the patient had for breakfast or by asking her or him to name three objects—for example, a pen, a tree, and a ball—and then asking her or him to recall them later. To test remote memory, ask birth dates or anniversary dates if someone can validate the information; if not, ask dates of major historical events.

General Knowledge and Vocabulary

Before you assess general knowledge and vocabulary, you need to consider the developmental level, educational level, and cultural background of your patient so that you can phrase and direct your questions accordingly. To test general knowledge, ask about current events, the name of the president of the United States, or common knowledge questions, such as the number of months in a year or days in a week. To test vocabulary, ask the patient to define words. Begin with easy, familiar words, such as “orange,” and proceed to more difficult or abstract words, such as “dictatorial.”

Mathematical and Calculative Skills

To test mathematical and calculative skills, have your patient solve a simple math problem. Counting backward from 100 by 7s (serial 7s) is frequently used, but this is difficult for many people, especially those dependent on calculators. As an alternative, have the person count backward by 3s or 4s, or ask him or her to solve a simple problem. An example is: “If you purchased a magazine for \$1.50, and you have \$2.00, how much change would you get back?” Whichever method you use, make sure the problem is appropriate for the patient’s educational level.

Thought Process/Abstract Reasoning/Judgment

To assess thought process, examine the appropriateness, organization, and content of your patient’s responses throughout the entire assessment. Be alert for any sensory-perceptual experiences, feelings, or false beliefs that are not based on reality and may indicate illusions, hallucinations, or delusions.

Assess abstract reasoning by asking your patient to explain a simple proverb, such as “People in glass houses shouldn’t throw stones.” If the person’s age or cultural background makes using proverbs inappropriate, use a phrase such as “It’s raining cats and dogs!” Note the degree of concreteness or abstractness of her or his interpretation. You can further assess abstract ability by asking her or him to group similar objects; for example, ask “What do an apple, an orange, and a pear have in common?” Then determine whether the responses are appropriate.

Sound judgment involves considering options and choosing appropriate actions. Assess your patient’s judgment by observing his or her response to the current situation or by giving him or her a hypothetical situation. For example, say: “If you were walking down the street and witnessed a car accident, what would you do?” Then decide whether the response is appropriate and reasonable.

Assessing Communication

When assessing speech, evaluate not only the ability to speak but also the content, appropriateness, speed, and quality of speech. Identify the patient’s primary language, and solicit an interpreter if needed. Various speech problems are associated with neurologic disorders, and the type of problem depends on which area of the brain is affected. Increasing language difficulties may reflect a turn for the worse in your patient’s neurologic status that warrants further medical evaluation. The inability to communicate can be frustrating for both of you, so be patient, allow your patient time to respond, and make referrals to speech therapy as needed.

To pinpoint the exact location of the neurologic problem, assess further for spontaneous speech, sound recognition, auditory-verbal comprehension, visual recognition, visual-verbal comprehension, motor speech, automatic speech, naming, vocabulary, writing, and copying figures. This assessment can also be used to develop an effective speech therapy plan for your patient.

Cranial Nerve Function

Evaluation of the cranial nerves is an essential part of the neurologic examination. However, like other portions of the exam, it depends upon an alert patient who is emotionally, cognitively, and physically able to participate. Assessment techniques are described in the following section. See Chapter 12, Assessing the Eye and the Ear, for assessment techniques for CNs II, III, IV, VI, and VIII. (See *Assessing the Cranial Nerves*.)

(Text continued on page 777)

ASSESSING THE CRANIAL NERVES

1. CN I—Olfactory Nerve

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Before testing nerve function, ensure patency of each nostril by occluding in turn and asking patient to sniff. Once patency is established, ask patient to close eyes. Occlude one nostril and hold aromatic substance such as coffee beneath nose. Ask patient to identify substance. Repeat with other nostril.
- *Patient is able to identify substance. Bear in mind that some substances may be unfamiliar, especially to children.*



Testing the olfactory nerve

ABNORMAL FINDINGS/RATIONALE

- **Anosmia** is loss of sense of smell. It may be inherited and nonpathological, or may result from chronic rhinitis, sinusitis, heavy smoking, zinc deficiency, or cocaine use. It may also indicate cranial nerve damage from facial fractures or head injuries, disorders of base of frontal lobe such as a tumor, or arteriosclerotic changes. Persons with anosmia usually also have taste problems.

2. CNs II, III, IV, and VI—Optic, Oculomotor, Trochlear, and Abducens Nerves

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Test visual acuity, peripheral vision, eye movements, and pupil accommodation as discussed in Chapter 12.



Testing visual acuity



Testing eye movements



Testing pupil accommodation

- Test oculocephalic reflex (“doll’s eyes”) in unresponsive patient. Rotate head quickly from side to side.



A



B

A and B. Testing oculocephalic reflex

ASSESSING THE CRANIAL NERVES

2. CNs II, III, IV, and VI—Optic, Oculomotor, Trochlear, and Abducens Nerves (continued)



ALERT

Never perform this test on a patient with suspected neck injury.

- *Visual acuity intact 20/20, both eyes; PERRLA direct and consensual; EOM intact.*
- **Hippus phenomenon:** *Brisk constriction of pupils in reaction to light, followed by dilation and constriction; may be normal or sign of early CN III compression.*
- **Normal doll's eyes:** *Eyes deviate to side opposite from way head is turned.*

ABNORMAL FINDINGS/RATIONALE

- CN II deficits can occur with stroke or brain tumor.
- Changes in pupillary reactions can signal CN III deficits. Increased ICP causes changes in pupillary reaction. As pressure increases, response becomes more sluggish until pupils finally become fixed and dilated.
- *Abnormal doll's eyes (eyes fixed):* Damage to oculomotor nerves (CN III, IV, VI) or brainstem.

3. CN V—Trigeminal Nerve

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- **Testing motor function:** Ask patient to move jaw from side to side against resistance and then clench jaw as you palpate contraction of temporal and masseter muscles, or to bite down on a tongue blade.
- **Testing sensory function:** Ask patient to close eyes and tell you when he or she feels sensation on the face. Touch jaw, cheeks, and forehead with cotton wisp. Touch same areas with toothpick. Compare both bilaterally.
- **Testing corneal reflex:** Gently touch cornea with cotton wisp.



Testing CN V motor function



Testing CN V sensory function



Testing corneal reflex



HELPFUL HINT

Touching cornea can cause abrasions. Alternative approach is to puff air across cornea with a needleless syringe, or gently touch eyelash and look for blink reflex.

- *Full range of motion (ROM) in jaw and 15 strength.*
- *Patient perceives light touch and superficial pain bilaterally.*

(continued)

ASSESSING THE CRANIAL NERVES

3. CN V—Trigeminal Nerve (continued)

ABNORMAL FINDINGS/RATIONALE

- *Weak or absent contraction unilaterally:* Lesion of nerve, cervical spine, or brainstem.



ALERT

In stroke patients, food may collect in cheek on patient's affected side, increasing risk for aspiration.

- Inability to perceive light touch and superficial pain may indicate peripheral nerve damage.
- *Tic douloureux:* Neuralgic pain of CN V caused by the pressure of degeneration of a nerve.
- Corneal reflex test used in patients with decreased LOC to evaluate integrity of brainstem.

4. CN VII—Facial Nerve

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- *Testing motor function:* Ask patient to perform these movements: smile, frown, raise eyebrows, show upper teeth, show lower teeth, puff out cheeks, purse lips, close eyes tightly while nurse tries to open them.
- *Testing sensory function:* Test taste on anterior two-thirds of tongue for sweet, sour, salty.



Testing CN VII motor function



Testing taste sensation



HELPFUL HINT

Sweet: Tip of the tongue
Sour: Sides of back half of tongue
Salty: Anterior sides and tip of tongue
Bitter: Back of tongue

- *Facial nerve intact; able to make faces. Taste sensation on anterior tongue intact.*
- *Taste decreased in older adults.*

ABNORMAL FINDINGS/RATIONALE

- Asymmetrical or impaired movement: Nerve damage, such as that caused by Bell's palsy or stroke.
- Impaired taste/loss of taste: Damage to facial nerve, chemotherapy or radiation therapy to head and neck.

ASSESSING THE CRANIAL NERVES

5. CN VIII—Acoustic Nerve

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Perform Weber and Rinne tests for hearing (see Chapter 12, Assessing the Eye and the Ear).
- Perform watch-tick test by holding watch close to patient's ear.
- Perform Romberg test for balance (see Chapter 20, Assessing the Motor-Musculoskeletal System).
- *If patient is unresponsive, perform cold caloric test for oculovestibular reflex (tests CNs III, VI, and vestibular portion of VIII):* Irrigate external ear canal with ice-cold water.



Watch-tick test



Cold caloric test

**ALERT**

Do not irrigate an ear unless you are sure the tympanic membrane is intact.

- Hearing intact.
- Negative Romberg test.
- Normal cold caloric test. (Normal response is nystagmus [involuntary rhythmic eye movement] toward irrigated ear.)

ABNORMAL FINDINGS/RATIONALE

- *Hearing loss, nystagmus, balance disturbance, dizziness/vertigo:* Acoustic nerve damage.
- *Nystagmus:* CN VIII, brainstem, or cerebellum problem or phenytoin (Dilantin) toxicity.
- *Abnormal cold caloric test (no eye movement):* Damage to CNs III, VI, and VIII.

6. CNs IX and X—Glossopharyngeal and Vagus Nerves

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Observe ability to cough, swallow, and talk.
- *Test motor function:* Ask patient to open mouth and say “ah” while you depress the tongue with a tongue blade. Observe soft palate and uvula. Soft palate and uvula should rise medially.



Testing CN IX and X motor function

(continued)

ASSESSING THE CRANIAL NERVES

6. CNs IX and X—Glossopharyngeal and Vagus Nerves (continued)

- Test sensory function of CN IX and motor function of CN X by stimulating gag reflex. Tell patient that you are going to touch interior throat, then lightly touch tip of tongue blade to posterior pharyngeal wall. Observe movement.
- Test taste on posterior portion of tongue.
- *Swallow and cough reflex intact. Speech clear.*
- *Elevation and constriction of pharyngeal musculature and tongue retraction indicate positive gag reflex.*
- *Taste on posterior tongue intact.*

ABNORMAL FINDINGS/RATIONALE

- *Unilateral movement:* Contralateral nerve damage.
- Damage to CNs IX and X also impairs swallowing.
- *Changes in voice quality (e.g., hoarseness):* CN X damage.
- CN X damage may also affect vital functions, causing arrhythmias because vagus nerve innervates most of viscera through parasympathetic system.
- *Diminished/absent gag reflex:* Nerve damage. Evaluate further because patient is at increased risk for aspiration.
- *Impaired taste on posterior portion of tongue:* Problem with CN IX.

7. CN XI—Accessory Nerve

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

- Test motor function of shoulder and neck muscles: Ask patient to shrug shoulders upward against your resistance. Then ask her or him to turn head from side to side against your resistance. Observe for symmetry of contraction and muscle strength.
- *Movement symmetrical, with patient moving against resistance without pain.*
- *Full ROM of neck with +5/5 strength.*



Testing CN XI motor function

ABNORMAL FINDINGS/RATIONALE

- *Asymmetrical/diminished/absent movement/pain/unilateral or bilateral weakness:* Peripheral nerve CN XI damage.

8. CN XII—Hypoglossal Nerve

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS



HELPFUL HINT

Precise articulation indicates proper functioning, so further assessment is indicated only when speech is impaired.

(continued)

ASSESSING THE CRANIAL NERVES

8. Hypoglossal Nerve (continued)

- Have patient say “d, l, n, t” or a phrase containing these letters, such as the phrase in the Cincinnati Prehospital Stroke Scale. The ability to say these letters requires use of the tongue.
- Ask the patient to protrude the tongue. Observe any deviation from midline, tumors, lesions, or atrophy. Now ask the patient to move the tongue from side to side.
- *Can protrude tongue medially. No atrophy, tumors, or lesions.*



Testing CN XII motor function

ABNORMAL FINDINGS/RATIONALE

- *Asymmetrical/diminished/absent movement/deviation from midline/protruded tongue:* Peripheral nerve CN XII damage.
- Tongue paralysis results in dysarthria.

Sensory System Function

Assess the patient's sensory function using tests of light (superficial) touch, pain, temperature, vibration, position sense, **stereognosis** (ability to recognize the form of solid objects by touch), **graphesthesia** (ability to recognize outlines, numbers, words, or symbols written on skin), two-point discrimination, point localization, and extinction. Light touch, pain, and temperature are superficial sensations and travel the spinothalamic tracts of the anterolateral system. Vibration, position sense, stereognosis, graphesthesia, two-point discrimination, point localization, and extinction are highly localized and travel the dorsal column–medial lemniscal pathway of the posterior column–medial lemniscal pathway.

Before beginning this portion of the examination, inform the patient that you will apply various stimuli, and that she or he should close the eyes during the entire exam. Instruct her or him to say “now” whenever she or he perceives a sensation. Avoid asking leading questions, such as, “Can you feel anything here?” Also avoid using any recognizable pattern, but do test symmetrical regions. If you notice an area of altered sensation, stimulate areas close to it until you have delineated its borders. Record your findings with a diagram of the area.

The extent of the sensory assessment depends on your findings. For screening purposes, include the upper and lower extremities and the trunk. If you detect a deficit, perform a more thorough assessment.

ASSESSING SENSORY FUNCTION

1. Light Touch

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Brush a light stimulus such as a cotton wisp over patient's skin in several locations, including torso and extremities.



Testing superficial light touch

(continued)

ASSESSING SENSORY FUNCTION

1. Light Touch (continued)



HELPFUL HINT

If light touch sensation is intact distally, do not assume that it is intact proximally.

- Identifies areas stimulated by light touch.

ABNORMAL FINDINGS/RATIONALE

- Diminished/absent cutaneous perception:** Peripheral nerve damage or damage to posterior column of spinal cord. Peripheral neuropathies can also cause sensory deficits.
- Hypesthesia:** Increased sensitivity.
- Paresthesia:** Numbness and tingling.
- Anesthesia:** Loss of sensation.

2. Pain

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Stimulate skin lightly with sharp and dull ends of toothpick.
- Apply stimuli randomly and ask patient to identify whether sensation is sharp or dull.
- Identifies areas stimulated and type of stimulation.



A. Assessing pain in the upper extremity



B. Assessing pain in the lower extremity

- Touch patient's skin with test tubes filled with hot or cold water.
- Apply stimuli randomly, and ask patient to identify whether sensation is hot or cold.
- Identifies areas stimulated and type of stimulation.



A. Testing temperature in the upper extremity



B. Testing temperature in the lower extremity

ASSESSING SENSORY FUNCTION

2. Pain (continued)



HELPFUL HINT

Test temperature only if patient's perception of pain is abnormal. If pain sensation is intact, there is no need to test temperature, because temperature and pain run along the same tract.

ABNORMAL FINDINGS/RATIONALE

- *Diminished or absent pain perception:* Peripheral nerve damage or damage to lateral spinothalamic tract.
- *Hyperalgia:* Increased pain sensation.
- *Hypoalgia:* Decreased pain sensation.
- *Analgesia:* No pain sensation.
- *Diminished/absent temperature perception:* Peripheral nerve damage or damage to lateral spinothalamic tract.

3. Vibration

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Place a vibrating tuning fork over a finger joint, and then over a toe joint.
- Ask patient to tell you when vibration is felt and when it stops.
- If patient is unable to detect vibration, test proximal areas as well.



HELPFUL HINT

If vibratory sensation is intact distally, it is intact proximally.



A. Vibrating the upper extremity.



B. Vibrating the lower extremity.

- *Vibratory sensation intact bilaterally in upper and lower extremities.*

ABNORMAL FINDINGS/RATIONALE

- *Diminished/absent vibration sense:* Peripheral nerve damage caused by alcoholism, diabetes, or damage to posterior column of spinal cord.

4. Kinesthetics (Position Sense)

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Determine patient's ability to perceive passive movement of extremities. Hold fingers on sides and move up and down, and have patient identify direction of movement.
- Flex and extend patient's big toe, and ask patient to describe movement as up or down.

(continued)

ASSESSING SENSORY FUNCTION

4. Kinesthetics (Position Sense) (continued)



A. Testing position sense in the finger



B. Testing position sense in the toe



HELPFUL HINTS

- Avoid moving the patient's finger by placing your finger on top of the patient's because the patient may sense the pressure of your finger rather than a true position change.
- If position sensation is intact distally, it is intact proximally.

- *Position sensation intact bilaterally in upper and lower extremities.*

ABNORMAL FINDINGS/RATIONALE

- *Diminished or absent position sense:* Peripheral nerve damage or damage to posterior column of spinal cord.

5. Stereognosis

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- With patient's eyes closed, place a familiar object, such as a coin or a button, in patient's hand, and ask patient to identify it.
- Test both hands using different objects.
- *Stereognosis intact bilaterally.*



Testing stereognosis

ABNORMAL FINDINGS/RATIONALE

- Abnormal findings suggest a lesion or other disorder involving sensory cortex or a disorder affecting posterior column.

ASSESSING SENSORY FUNCTION

6. Graphesthesia

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- With patient's eyes closed, use point of a closed pen to trace a number on patient's hand, and ask patient to identify the number.
 - *Graphesthesia intact bilaterally.*



Testing graphesthesia

ABNORMAL FINDINGS/RATIONALE

- Abnormal findings suggest lesion or other disorder involving sensory cortex or disorder affecting posterior column.

7. Two-Point Discrimination

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Ability to differentiate between two points of simultaneous stimulation.
- Using ends of two toothpicks, stimulate two points on fingertips simultaneously.
- Gradually move toothpicks together, and assess smallest distance at which patient can still discriminate two points (minimal perceptible distance).
- Document distance and location.



Testing two-point discrimination



HELPFUL HINT

The normal discriminatory distance depends on the area tested, with the fingertips being the most discriminating.

- *Discriminates between two points on fingertips no more than 0.5 cm apart and on hands no more than 2 cm apart.*

ABNORMAL FINDINGS/RATIONALE

- Abnormal findings suggest lesion or other disorder involving sensory cortex or disorder affecting posterior column.

(continued)

ASSESSING SENSORY FUNCTION

8. Point Localization

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Ability to sense and locate area being stimulated.
- With patient's eyes closed, touch an area; then have patient point to where he or she was touched.
- Test both sides and upper and lower extremities.
- *Point localization intact.*



Testing point localization

ABNORMAL FINDINGS/RATIONALE

- Abnormal findings suggest lesion or other disorder involving sensory cortex or disorder affecting posterior column.

9. Sensory Extinction

ASSESSMENT TECHNIQUE/NORMAL VARIATIONS

- Simultaneously touch both sides of patient's body at same point.
- Ask patient to point to where she or he was touched.
- *Extinction intact.*



Testing extinction

ABNORMAL FINDINGS/RATIONALE

- Identification of stimulus on only one side suggests lesion or other disorder involving sensory cortical region in opposite hemisphere.

Reflexes

Reflex assessment evaluates the intactness of the spinal cord. Intact sensory and motor systems are required for a normal reflex response. Normal reflexes include DTRs and superficial reflexes. There are also protective reflexes that are responses of intact cranial nerves. The protective reflexes include the gag, cough and swallow, blink and corneal. CNs IX and X are responsible for the gag, cough and swallow, and CNs V and VII are responsible for the blink and corneal reflexes. Reflexes that normally occur in newborns are termed *pathological* or *primitive reflexes* when they occur in adults.

Deep Tendon and Superficial Reflexes

DTRs include the biceps (C5, C6), triceps (C7, C8), brachioradialis (C5, C6), patellar (L2, L3, L4) and Achilles (S1, S2). Superficial reflexes include the plantar (L4 to S2), abdominal (T8, T9, T10), anal (S3, S4, S5), cremasteric (L1, L2), and bulbocavernosus (S3, S4).

To test DTRs, use a rubber percussion hammer to swiftly tap a slightly stretched tendon to elicit contraction of an associated skeletal muscle. Keep your wrist loose so that immediately upon impact, it swings back spontaneously. As with other neurologic assessments, test reflexes bilaterally. Then grade DTRs on a scale from 0 to 4. Stick figures are often used to chart DTR findings (Fig. 21.10).

Documenting Reflex Findings

Use these grading scales to rate the strength of each reflex in a deep tendon and superficial reflex assessment.

Deep tendon reflex grades


- 0 absent
- + present but diminished
- ++ normal
- +++ increased but not necessarily pathologic
- ++++ hyperactive or clonic (involuntary contraction and relaxation of skeletal muscle)

Superficial reflex grades

- 0 absent
- + present

Use the patient's reflex ratings on a drawing of a stick figure. The figures here show documentation of normal and abnormal reflex responses.

Normal



Abnormal




FIGURE 21.10. Documenting reflex findings.

Because the normal response for a DTR is muscle contraction, eliciting a response may be difficult if your patient is tense and the muscles are contracted. Through **cognitive inhibition**, the patient can override and suppress the peripheral reflex response. Tell your patient to relax and let you support the limb being tested. If you are still having difficulty eliciting a DTR response, use a reinforcement technique to enhance the response. One technique, **distraction maneuvers**, allows the patient to concentrate on something other than the reflex being tested, thereby relaxing the area and eliminating unintentional cognitive inhibition. The other technique, **isometric maneuvers**, overrides the inhibitory message from the brain and increases reflex response. Two isometric maneuvers are having your patient clench his or her teeth (enhances upper extremity reflexes) and having the patient interlock his or her hands and push (enhances lower extremity reflexes). These techniques are shown in Figure 21.11. If you use a reinforcement technique, be sure to document it.

When assessing superficial reflexes, use a tongue blade, the base of the reflex hammer, or the back of your thumbnail and stroke the body briskly, without pressing into the underlying organs. Superficial reflexes are graded as positive or negative. DTRs are graded as follows:

- 0: No response detected.
- 1: Response present but diminished (hypoactive).
- 2: Response normal.
- 3: Response somewhat stronger than normal.
- 4: Response hyperactive with clonus.

Grades 1 and 3 are usually considered normal. Even grade 0 may occur symmetrically in some patients in the absence of any underlying neurologic disorder. For example, a patellar reflex may be difficult to elicit on a patient who has had knee surgery. Grade 4 usually indicates pathology. **Clonus** is the presence of rhythmic involuntary contractions, most often at the foot and ankle. Sustained clonus confirms CNS involvement.

Pathological or Primitive Reflexes

Pathological or primitive reflexes usually indicate a severe underlying neurologic problem. They reflect cerebral degeneration or late-stage dementia and are referred to as **primitive reflexes** because they are normally seen in a newborn with an immature neurologic system. Family members may be misled by these reflexes, which may seem like a sign of improvement. Therefore, be sure to explain that these are reflexive responses and provide support for the family.

Primitive reflexes include the following:

- **Grasp:** Place your fingers in palm of patient's hand. Patient will close her or his fingers and grasp yours.



FIGURE 21.11. Isometric maneuvers. A. Clenching teeth. B. Interlocking hands.

- *Sucking*: Gently stimulate patient's lips with a mouth swab. Patient will start sucking.
- *Snout*: Gently tap oral area with finger. Patient's lips will pucker.
- *Rooting*: Gently stroke side of patient's face. Patient will turn toward stimulated side.
- *Glabellar*: Gently tap on patient's forehead. Patient will blink.



FIGURE 21.12. Kernig's sign.

- *Babinski*: Stroke lateral aspect of sole of foot. Dorsi-flexion of great toe and fanning of toes will occur.

Meningeal Signs

Classic signs of meningitis include nuchal rigidity (extension of neck with resistance to flexion), fever, photosensitivity, headache, nausea, and vomiting. If you suspect that your patient has meningitis, assess for Kernig's and Brudzinski's signs.

To assess for Kernig's sign (Fig. 21.12), have the patient lie supine with one leg flexed. Tell him or her to try to extend the leg while you apply pressure to the knee. Contraction and pain of the hamstring muscles and resistance to extension are positive signs of meningitis.

To assess for Brudzinski's sign (Fig. 21.13), have the patient lie supine with her or his head flexed to her or his chest. Flexion of the hips is a positive sign of meningitis.



FIGURE 21.13. Brudzinski's sign.

ASSESSING REFLEXES

1. Deep Tendon Reflexes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Biceps Reflex

- Rest patient's elbow in your nondominant hand, with your thumb over biceps tendon.
- Strike your thumbnail.
- *Contraction of biceps with flexion of forearm.*
- +2



Testing biceps reflex

ASSESSING REFLEXES

1. Deep Tendon Reflexes (continued)

Triceps Reflex

- Abduct patient's arm and flex it at the elbow.
- Support the arm with your nondominant hand.
- Strike triceps tendon about 1 to 2 inches above olecranon process, approaching it from directly behind.



Testing triceps reflex

- *Contraction of triceps with extension at elbow.*
- +2

Brachioradialis Reflex

- With your nondominant hand, support patient's hand and palpate brachioradialis tendon 3 to 5 cm above wrist.
- Simultaneously strike styloid process of radius.



Testing brachioradialis reflex

- *Flexion at elbow and supination of forearm. You may also notice flexion of fingers.*
- +2

Patellar Reflex

- Have patient sit with legs dangling.
- Strike tendon directly below patella.



Testing patellar reflex

- *Contraction of quadriceps with extension of knee.*
- +2

Achilles Reflex

- Have patient lie supine or sit with one knee flexed.
- Holding patient's foot slightly dorsiflexed, strike Achilles tendon.



Testing Achilles reflex

- *Plantarflexion of foot.*
- +2

Test for Ankle Clonus

- If you get +4 reflexes while supporting leg and foot, quickly dorsiflex foot.



Testing for clonus

- *No contraction*

(continued)

ASSESSING REFLEXES

1. Deep Tendon Reflexes (continued)

ABNORMAL FINDINGS/RATIONALE

- **Absent/diminished DTRs:** Degenerative disease; damage to peripheral nerve such as peripheral neuropathy; lower motor neuron disorder, such as ALS and Guillain-Barré syndrome.
- **Hyperactive reflexes with clonus:** Spinal cord injuries, upper motor neuron disease such as MS.
- Rhythmic contraction of leg muscles and foot is positive sign of clonus and indicates upper motor neuron disorder.

2. Superficial Reflexes

ASSESSMENT TECHNIQUES/NORMAL VARIATIONS

Abdominal Reflex

- Stroke patient's abdomen diagonally from upper and lower quadrants toward umbilicus.
- Contraction of rectus abdominis. Umbilicus moves toward stimulus.



HELPFUL HINT

Obesity and pregnancy can mask this reflex.



Testing abdominal reflex

Anal Reflex

- Gently stroke skin around anus with gloved finger.
- *Anus puckers.*

Cremasteric Reflex

- Gently stroke inner aspect of a male's thigh.
- *Testes rise.*

Bulbocavernosus Reflex

- Gently apply pressure over bulbocavernosus muscle on dorsal side of penis.
- *Bulbocavernosus muscle contracts.*

Plantar Reflex (Babinski's Response)

- Stroke sole of patient's foot in an arc from lateral heel to medial ball.
- *Flexion of all toes.*



Testing plantar reflex

ABNORMAL FINDINGS/RATIONALE

- Absence of superficial reflexes may indicate pyramidal tract lesions.
- **Babinski's response:** Dorsiflexion of big toe, with or without fanning of other toes. In absence of drug or alcohol intoxication, it suggests pathology involving upper motor neurons.



Case Study Findings

Your general health survey and head-to-toe assessment of Mr. Webster have concluded. The following are your findings:

- **General Health Survey**
 - Well groomed, posture erect but guarding neck.
 - Speech clear, affect appropriate.
 - Appears stated age.
 - 2" laceration of forehead.
 - Alcohol on breath.
 - In no acute distress, but grimacing in pain.
 - Head-to-toe scan pertinent findings related to the Neurologic system include, pupils 3 mm, PERRLA (pupils equal, round, reactive to light and accommodation [directly and consensually], negative fundoscopic, 2" sutured laceration on forehead secondary to head injury, equal motor strength, no weakness, gait steady, balance intact.
- **Vital Signs**
 - Temperature, 98°F
 - Pulse, 88 beats per minute (BPM) and regular.
 - Respirations, 20/min.
 - BP, 130/80.
 - Pulse oximetry, 98 percent room air.
- **Sensory-Neurologic**
 - *LOC*: AAO × 3; GCS 15.
 - *Communication*: Intact.
 - *Memory*: Immediate and remote intact, but doesn't remember events of accident.

- *Cognitive functions*: Intact.
- CNs I –XII intact.
- DTR +2; positive plantar.



Case Study Analysis and Plan

Now that you have completed a thorough assessment of Mr. Webster, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mr. Webster and use this information to identify a list of nursing diagnoses. Some possible ones are provided as follows. Cluster the supporting data for each.

1. Pain, related to injuries
2. Risk for altered cerebral tissue perfusion related to head injury
3. High risk for injury related to risky behavior

Identify any additional nursing diagnoses.



CRITICAL THINKING ACTIVITY

Now that you have identified some nursing diagnoses for Mr. Webster, select one and develop a nursing care plan and a brief teaching plan for him, including learning objectives and teaching strategies.

RESEARCH TELLS US

In the emergency department, time is critical. As part of a neurologic assessment, the MMSE is used to assess the mental status of a patient. Although the MMSE is a well-established tool, the amount of time needed to administer the MMSE may prohibit its use in an ED setting. The Quick Confusion Scale (QCS) is an alternate instrument used to measure the patient's mental status. The QCS, a 6-item, 15-point instrument that can be administered quickly, is ideal in the ED setting. Its benefits include:

- Less time needed to administer.
- No forms or writing materials needed
- Familiar scoring, similar to Glasgow Coma Scale (GCS)
- Easily calculated and interpreted scores.
- Easy to remember after several uses.

To assess the validity of the QCS, Huffand colleagues (2001) performed a prospective comparative study of the MMSE and the QCS. A convenience

RESEARCH TELLS US (continued)

sample of ED patients 55 and older ($n = 205$) were administered the MMSE and the QCS. Patients excluded from the study included those with head trauma, multisystem trauma, GCS score less than 15, acute illness, or incomplete data, as well as those who were non-English speaking, had an educational level less than 8 years, or required contact or droplet isolation. Interrater reliability was established for rater's collecting data ($r = .90$). Not only were scores collected, but

also the time needed to administer instruments was recorded. A significant correlation was found between the QCS scores, mean score 11.89 ($SD = 3.5$, range 0–15), and the MMSE scores, mean score 24.8 ($SD = 4.7$; range 2–30) ($r = .783$). The MMSE took significantly more time to administer than the QCS ($p < .01$). Further study by Irons and associates (2002) support the use of the QCS as a viable alternative for assessing mental status in the ED setting.

Source: Huff, J., Farace, E., Brady, W., Kheir, J., and Shaver, G. (2001) The Quick Confusion Scale in the ED: Comparison with the Mini-Mental State Examination. *The American Journal of Emergency Medicine* 19(6):461–464; and Iron, M.J., Farace, E., Brady, W.J., and Huff, J.S. (2002). Mental status screening of emergency department patients: Normative study of the Quick Confusion Scale. *Academic Emergency Medicine*. 9(10):989–994.

The Quick Confusion Scale

Item	No. Correct	x Weight = Total
What year is it now?	0 or 1 (score 1 if correct, 0 if incorrect)	x 2 = _____
What month is it?	0 or 1	x 2 = _____
Present memory phrase: "Repeat this phrase after me and remember it: 'John Brown, 42 Market St., New York.'"		
About what time is it? (Answer correct if within 1 hour)	0 or 1	x 2 = _____
Count backwards from 20 to 1.	0, 1, or 2	x 1 = _____
Say the months in reverse.	0, 1, or 2	x 1 = _____
Repeat memory phrase (each underlined portion correct is worth 1 point).	0,1,2,3,4. or 5	x 1 = _____
Final score is the sum of the totals.		= _____

Explanation of Scoring Using the Quick Confusion Scale

The higher the Quick Confusion Scale scores the better the mental state. Best total score is 15.

- Item 1. "What year is it now?" Score 1 if answered correctly, 0 if incorrect.
- Item 2. "What month is it?" Score 1 if answered correctly, 0 if incorrect.
- Item 3. "About what time it is?" Answer considered correct if within 1 hour; score 1 if correct, 0 if incorrect.
- Item 4. "Count backwards from 20 to 1." Score 2 if correctly performed; score 1 if one error, score 0 if 2 or more errors.
- Item 5. "Say the months in reverse." Score 2 if correctly performed; score 1 if one error, score 0 if 2 or more errors.
- Item 6. Repeat memory phrase, "John Brown, 42 Market St., New York." Each portion correctly recalled is worth 1 point in scoring; score 5 if correctly performed; each error drops score by one.

The final score is achieved by summing up the weighted totals; items 1, 2, and 3 are multiplied by 2 and summed with the other item scores to yield the final score. Best score is 15.

HEALTH CONCERNS

Although the mortality rate for stroke is alarming, the physical, emotional, and financial costs for those who survive are just as alarming. You cannot put a price on the physical and emotional toll, but the direct and indirect financial costs exceed \$41 billion annually. According to the National Institutes of Health Neurologic Institute, stroke risk reduction and prompt treatment could decrease stroke by 80 percent by the end of the decade. Identifying risk factors is the first step in preventing this disease.

Risk Factors for Stroke

- **Nonmodifiable:** Age, gender, race, previous stroke or transient ischemic attack (TIA), asymptomatic carotid stenosis, heredity.
- **Modifiable:** HTN; diabetes mellitus; cardiovascular disease; abnormal serum lipid, triglyceride, cholesterol, low-density lipoprotein (LDL), and high-density lipoprotein (HDL) levels; cigarette smoking; excessive alcohol; drug abuse.
- **Other factors:** Obesity, migraines, oral contraceptive use, hypercoagulable states.

Prompt recognition of signs and symptoms of stroke is the second step in prevention.

Common Abnormalities

ABNORMALITY

ASSESSMENT FINDINGS

Alzheimer's Disease

- A chronic, progressive disorder that accounts for 50 percent of all dementias

- Most common in people over 65
 - Pathological changes in the brain including plaques and neuronal tangles
 - Begins with mild memory loss
 - Progresses to deterioration of intellectual function, personality changes, and speech and language problems
 - Repeated head trauma
 - Exposure to aluminum
 - Previous CNS infection
 - Family history
 - Emotional lability
 - Weight loss, anorexia, malnutrition
 - Incontinence
 - Poor hygiene
 - Gait instability, weakness
 - Inability to perform ADLs
 - Awakening frequently during night with daytime naps
 - Forgetfulness
 - Inability to solve problems
 - Depression, suicidal ideation
 - Disheveled appearance
 - Agitation
 - Memory loss (early sign) that is progressive
 - Disorientation to time
 - Flat affect
 - Impaired cognitive functioning
 - Nocturnal wandering
 - Repetitive behaviors
 - Stubbornness, paranoia
- More Advanced Disease**
- Aphasia, agnosia, apraxia

(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Brain Tumor</p> <ul style="list-style-type: none"> ■ An intracranial mass that can be neoplastic, cystic, inflammatory (abscess), or syphilitic 	<ul style="list-style-type: none"> ■ Seizures ■ Limb rigidity, flexor posturing ■ Early morning headache ■ Subtle personality changes or dysphasia ■ Papilledema, possibly leading to visual loss ■ Changes in pupil size and response ■ Disorders of EOM ■ Focal deficits ■ New onset of seizures
<p>Stroke</p> <ul style="list-style-type: none"> ■ Stroke, caused by either ischemia or hemorrhagic lesions 	<ul style="list-style-type: none"> ■ Hemiparesis or hemiparalysis (loss of tactile sensation on affected side of body) ■ HTN; atherosclerosis; family history of cardiovascular or cerebrovascular disease ■ Sudden onset of symptoms with or without warning signs ■ Intact or altered mental activity; emotional lability ■ Hemiparesis or hemiparalysis, usually affecting the arm more than the leg ■ Facial sagging on affected side ■ Impaired swallowing ■ Homonymous hemianopia (vision defect in the right halves or left halves of the visual fields in both eyes) ■ Language disturbance and aphasia with right-sided weakness ■ Perceptual disturbance and altered visual-spatial perceptions with left-sided weakness ■ Loss of sensation on affected side ■ Disturbed stereognosis, body scheme, and visual-spatial skills
<p>Chronic Subdural Hematoma</p> <ul style="list-style-type: none"> ■ A swelling or mass of blood beneath the dura, usually caused by head injury 	<ul style="list-style-type: none"> ■ Personality changes ■ Gradual progressive decrease in level of consciousness; headache ■ History of trivial head injury weeks or months earlier ■ Progressive deterioration in mental activity and level of arousal ■ Disorientation ■ Focal deficit
<p>Guillain-Barré Syndrome</p> <ul style="list-style-type: none"> ■ Acute autoimmune inflammatory destruction of the myelin sheath leading to rapid, progressive symmetrical loss of motor function with no sensory loss, usually triggered by a viral infection 	<ul style="list-style-type: none"> ■ Muscle weakness that affects the lower extremities ■ First flaccid paralysis, little or no sensory loss ■ History of recent surgery, cancer, pregnancy, childbirth, infection, or vaccination ■ Respiratory insufficiency ■ Labile blood pressure ■ Tachycardia ■ Vasomotor flushes ■ Hyperpyrexia and increased sweating ■ Tracheobronchial secretions ■ Paralytic ileus ■ Blurred vision or diplopia

ABNORMALITY	ASSESSMENT FINDINGS
<p>Herniated Intervertebral Disc</p> <ul style="list-style-type: none"> ■ Prolapse of the nucleus pulposus of a ruptured intravertebral disc into the spinal canal 	<ul style="list-style-type: none"> ■ Facial weakness; impaired swallowing ■ Symmetrical flaccid paralysis, usually beginning in legs and progressing upward ■ Absent deep tendon reflexes ■ Sharp, severe pain in the back, which may radiate down extremity (in a pattern that reflects the nerve involved) and may be intensified by such actions as coughing, sneezing, straining, and moving ■ History of recurring symptoms ■ Decreased muscle tone in affected area ■ Mild motor weakness ■ Paresthesia in affected area ■ Intact, absent, or diminished brachioradialis, biceps, and triceps reflexes, depending on the level of lesion ■ Diminished or absent patellar and Achilles reflexes
<p>Meningitis</p> <ul style="list-style-type: none"> ■ Inflammation of the membranes of the brain or spinal cord 	<ul style="list-style-type: none"> ■ Headache, nuchal rigidity ■ Elevated temperature (up to 105°F) ■ Irritability, restlessness, photophobia, confusion ■ History of adjacent infection, neurosurgery, head trauma, systemic sepsis, or immunosuppression ■ Restlessness, disorientation, lethargy, stupor, coma ■ Decreased visual acuity or loss ■ Brudzinski's sign, Kernig's sign ■ Generalized seizures ■ Opisthotonos (abnormal posture characterized by back arching and wrist flexion) as a late sign ■ Hyperalgesia, photophobia, increased reflexes ■ Elevated temperature ■ Altered respiratory pattern; weak, rapid pulse ■ Vomiting
<p>Migraine, Classic</p> <ul style="list-style-type: none"> ■ An often familial symptom complex of periodic attacks of vascular headache, usually temporal and unilateral in onset 	<ul style="list-style-type: none"> ■ Throbbing headache lasting 2 to 12 hours, often preceded by a visual disturbance or scotomata lasting 15 to 20 minutes ■ Anorexia; nausea; vomiting ■ Diarrhea ■ Photophobia ■ Dizziness ■ Syncope ■ Scalp tenderness ■ History of stress, fatigue, hormonal changes, menstruation, change in amount of sleep, or other predisposing factors ■ Ingestion of certain foods (such as aged cheese, red wine, or chocolate) or oral contraceptives that seem to trigger symptoms ■ Perfectionist, compulsive, intelligent, rigid personality type; family emphasis on achievement ■ Transient mood and personality changes

(continued)

Common Abnormalities (continued)

ABNORMALITY	ASSESSMENT FINDINGS
<p>Parkinson's Disease</p> <ul style="list-style-type: none"> ■ Chronic nervous system disease characterized by a fine, slowly spreading tremor, muscular weakness and rigidity, and a peculiar gait 	<ul style="list-style-type: none"> ■ Prodromal transient neurologic deficits such as hemiparesis, aphasia, ophthalmoplegia, and photophobia ■ Akinesia ■ Cogwheel rigidity ■ Resting tremor ■ Gait disturbance ■ Impaired swallowing ■ Flat affect ■ Monotonous speech ■ Decreased blinking ■ Increased sweating and salivation ■ History of cerebral arteriosclerosis, cerebral hypoxia, trauma, toxin ingestion (carbon monoxide, manganese, or mercury), or use of illegal drugs ■ Slow onset and gradual progression of signs and symptoms ■ Emotional lability, depression, or paranoia ■ Diminished facial movements, decreased blinking, impaired swallowing, and abnormal muscle tone ■ Bradykinesia with difficulty initiating movement, intermittent "freezing," and pill-rolling tremor of hands ■ Small, jerky, cramped handwriting (micrographia) ■ Abnormal gait that is slow to start, includes short and shuffling steps, gradually accelerates, and is difficult to stop ■ Low, monotonous speech; involuntary repetition of words (echolalia) or sentences (palilalia) spoken by others
<p>Spinal Cord Tumor</p> <ul style="list-style-type: none"> ■ Swelling or mass of the spinal cord that can be neoplastic, cystic, or inflammatory 	<ul style="list-style-type: none"> ■ Pain and local tenderness throughout the sensory nerve root ■ Motor weakness below level or lesion ■ Slow progression of sensory and motor dysfunction: initially unilateral, eventually bilateral ■ Dysfunction of CNs VIII through XII (with lesion at C4 or above) ■ Spastic weakness or paralysis below the lesion ■ Flaccid paralysis or weakness at level of lesion ■ Respiratory failure or difficulty, occipital headache, nystagmus, and stiff neck with lesion at C4 or above ■ Sensory loss below the level of the lesion ■ Loss of pain and temperature sensation below and contralateral to lesion when only one side of cord is affected ■ Loss of tactile, position, and vibration sensations ipsilateral to lesion ■ Band of hyperesthesia just above level of lesion ■ Increased DTRS below level of lesion ■ Absent reflexes at level of lesion; intact reflexes above lesion; positive Babinski reflex

ASSESSMENT CHECKLIST

HISTORY

Remember to look at each history component as it relates to the sensory-neurologic system.

Ask the patient the following:

- Do you have:
 - Headache?
 - Mental status change?
 - Dizziness, vertigo, and syncope?
 - Numbness or loss of sensation?
 - Deficits in the five senses?
- Do you have any neurologic problems?
- Do you have any other medical problems?
- Are you taking any medications?
- Do you have a history of head trauma, loss of consciousness, dizziness, headaches, or seizures?
- Do you have memory problems or changes in your senses?
- Do you have weakness, numbness, or paralysis?
- Do you have problems walking or performing ADLs?
- Do you have mood problems or depression?
- Do you use drugs or alcohol?
- Do you have allergies?
- Have you ever been treated for a neurologic or psychiatric problem?
- When did your symptoms start?

PHYSICAL ASSESSMENT

Equipment

Stethoscope	Toothpick
BP cuff	Coin
Penlight	Substances to test smell and taste
Gloves	Tongue blade
Cotton	Reflex hammer

Position

Sitting

Technique

Remember to scan your patient from head-to-toe looking for signs related to sensory-neurologic system assessment

ASSESSING CEREBRAL FUNCTION

- Behavior
 - LOC, orientation to time, place and person
 - Mental status and cognitive function
 - Memory (immediate, recent, remote)
 - Mathematical and calculative ability
 - General knowledge and vocabulary
 - Thought process
 - Abstract thinking
 - Judgment
- Communication

ASSESSING CRANIAL NERVES I THROUGH XII

- Sensory*
 - Superficial sensations:* Light touch, pain, temperature
 - Deep sensations:* Vibration and kinesthetics
 - Discriminatory sensations:* Stereognosis, graphesthesia, two-point discrimination, extinction, point-to-point localization
- Reflexes*
 - DTRs:* Biceps, triceps, brachioradialis, patellar, Achilles
 - Superficial reflexes:* Abdominal, plantar, cremasteric, anal, bulbocavernous
 - Primitive
- Meningeal signs

DOCUMENT

Document your findings and develop your plan of care.


SUMMARY

- This chapter taught you how to perform a thorough sensory-neurologic assessment, including a health history and physical examination. Before you begin the assessment, you an understanding of the anatomy of the neurologic system is essential. Because the neurologic system is the main circuit board for the entire body, a problem in the neurologic system can affect a person's well being and ability to function.
- Begin by taking a detailed health history looking for clues related to the neurologic system.
- The physical examination of the neurologic system differs from other assessments. The exam includes assessment of cerebral function, cranial nerve function, sensory function, and reflexes.
- After you complete your assessment, analyze your findings to identify actual and potential health problems, as well as write nursing diagnoses and plan of care.

R E V I E W Q U E S T I O N S

1. Nick Dombrowski, age 22, was involved in an auto accident and is seen in the ER for head trauma. You are performing a neurologic assessment. Which test is best for assessing level of consciousness?
 - a. Romberg test
 - b. GCS
 - c. NIH scale
 - d. Trendelenburg test
2. You are assessing your patient for signs of increased ICP. What would be an early sign of increased ICP?
 - a. Change in mental status
 - b. Dilated pupils
 - c. Tachycardia
 - d. Hypotension
3. You are assessing Nick's level of orientation. Which level of disorientation would be of greatest concern?
 - a. Time
 - b. Place
 - c. Purpose
 - d. Person
4. You are testing the cranial nerves. You have your patient bite down on a tongue blade and note his or her jaw strength. Which cranial nerve does this test?
 - a. CN V
 - b. CN VI
 - c. CN VII
 - d. CN VIII
5. You are testing the cranial nerves. Which test is best for assessing CN XII?
 - a. Testing taste on the anterior portion of the tongue.
 - b. Having patient say, "You can't teach an old dog new tricks."
 - c. Having patient shrug shoulders against resistance.
 - d. Testing gag reflex.
6. You are testing your patient's sensations. You ask your patient to identify a number written in the palm of her or his hand. The test described is:
 - a. Stereognosis.
 - b. Graphesthesia.
 - c. Extinction.
 - d. Point localization.
7. How would you test for extinction?
 - a. Tell patient to point to where you touched him or her.
 - b. Simultaneously touch patient on opposite sides of body and have him or her identify sites.
 - c. Draw number in palm of patient's hand and have him or her identify the number drawn.
 - d. Place coin in palm of patient's hand and him or her identify object in hand.
8. You are testing your patient's DTRs. Which grade is considered normal?
 - a. 0
 - b. 2
 - c. 4
9. Which of the following reflexes would be considered abnormal in an adult?
 - a. Biceps reflex
 - b. Plantar reflex
 - c. Anal reflex
 - d. Babinski's reflex
10. Mary McMurray, age 20, is admitted to the intensive care unit, with the diagnosis of possible meningitis. Which test would be best for assessing for meningitis?
 - a. Babinski
 - b. Brudzinski
 - c. Glabellar
 - d. Trendelenburg sign

CHAPTER 22



Putting It All Together

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify pertinent history data included in a complete history
- Obtain a complete history
- Differentiate a regional from a systems approach in performing a complete physical assessment
- Perform a complete physical assessment
- Document complete assessment findings
- Identify actual/potential health problems stated as nursing diagnoses with supporting data

Now that you have learned what questions to ask and how to examine each body system, how do you combine the steps into a format for performing a complete assessment of your patient? A complete assessment is a very individualized process that depends on many factors, including your patient's age, developmental level, presenting symptoms, cultural background, and health needs. Your aim is to identify abnormalities, symptoms, health maintenance issues, teaching and learning needs, and your patient's strengths, so you can use them in planning patient care. Once health needs are identified, you can help your patient find appropriate resources to meet them. As a professional nurse, you will also be looking for health problems to address in a care plan, using the nursing process and nursing diagnoses.

In previous chapters, you learned about the many developmental, cultural, and ethnic variations to consider when assessing a patient. To illustrate how to “put it all together,” this chapter focuses on assessing a female patient in the developmental stage of menopause.

Introducing the Case Study



Marcia Malone is a 40-year-old woman who comes to you for a health screening. She is seeking information about routine health screenings recommended for her age group. She is concerned about her risk for osteoporosis and has several questions about menopause.

Developmental Considerations

The focus of this assessment is an adult female patient who is approaching menopause.

Developmental Stage Issues

At age 40, Marcia Malone is at the generativity versus stagnation stage of development, with a focus on being a productive member of both her family and society. Tasks for this developmental stage include raising a family and contributing to the community, usually through work or volunteer efforts. Because this stage marks the halfway point in life, people often reflect back on what they have done and look forward to what they hope to accomplish in relationships, self-development, and career. Toward the end of this stage, a person who has successfully raised a family may be confronted with the “empty nest syndrome.” Some people have a difficult time and search for meaning in life, whereas others see an opportunity to further fulfill personal and career goals. Midlife is also a

time when role reversals occur with parents. With life expectancy increasing, caring for aging parents may become a major concern.

Perimenopause and Menopause

Menopause is a developmental change that occurs in adult women. It is defined as the permanent cessation of menstruation as a result of loss of ovarian follicular activity. It usually occurs at about age 51. Although most women have minimal discomfort at this time, many still have concerns about the increased risk of cancer, cardiovascular disease, stroke, and osteoporosis, as well as the physical and neurological effects of menopause. They may also feel anxious because so many myths surround menopause. You can help your patients by debunking these myths, discussing the real concerns of menopause, and helping them to be active participants in maintaining their health into old age.

Perimenopause is a period of up to 10 years before and after menopause, during which signs and symptoms of ovarian involution are present. Symptoms in perimenopausal women are variable but include:

- Vasomotor
 - Hot flashes
 - Night sweats
 - Dizziness
 - Nausea
 - Lightheadedness
 - Palpitations
 - Nervousness
- Genitourinary
 - Vaginal dryness and itching
 - Urinary frequency
 - Urinary urgency and stress incontinence
 - Decreased libido and sexual response
 - Irregular or missed periods
 - Decreased menstrual flow
- Emotional and cognitive
 - Decreased concentration
 - Forgetfulness
 - Depression
 - Irritability
 - Emotional lability
 - Tension
 - Heightened stress response
- Physical
 - Headache
 - Insomnia
 - Sleep disturbances
 - Fatigue (related to sleep disorders)
 - Breast tenderness
 - Musculoskeletal aching
 - Appetite changes
 - Constipation

During perimenopause, women need preventive health screenings and education. Components of a comprehensive midlife healthcare program include:

- Complete history and physical examination.
- Blood work that includes a lipid panel, blood chemistries, and complete blood count (CBC).
- Thyroid function test.
- Pap smear.
- Hemocult test.
- Mammography.
- Psychosocial screening.
- Flexibility and exercise screening.
- Bone density screening.
- Dietary profile.

All women in midlife benefit from educational programs focused on menopause, osteoporosis, mammography, and breast self-examination. Selected patients also benefit from individualized health education for incontinence, heart disease, smoking cessation, skin cancer, and fibrocystic breast disease.

Performing the Assessment

A complete assessment begins with a comprehensive health history. Findings from the health history provide clues to actual or potential health problems to watch for as you perform the physical examination. During the physical examination, keep the key history findings in mind. If a problem area is revealed, perform a more thorough assessment of that area. The health history and physical examination are based on the patient's concerns and needs. Together, they provide you with a complete picture of her health status.

Health History

A complete health history includes obtaining biographical data, current health status, past health history, family history, a review of systems, and a psychosocial profile. By the end of the history, you should have a good idea of appropriate nursing diagnoses for the patient. You may also uncover concerns by asking age-appropriate or systems-review questions.

Biographical Data

Your patient's age, occupation, family or living situation, religion, ethnic background, and hobbies provide clues about his or her current health. For example, some diseases are age related, and some changes occur normally with aging. Religion, ethnic background, and family or living situation affect how a patient deals with illness and healthcare. Certain occupations and hobbies put a patient at risk for health problems.

As you establish rapport with your patient, also remember to take note of her or his clothing, hygiene, and mental status.

As you talk with Mrs. Malone, you note that she is dressed appropriately and that her hygiene is good. She is oriented to person, place, and time; has appropriate affect; has well-developed language; and presents her ideas and concerns in a relaxed, coherent, and logical manner. She appears well nourished and healthy. Her biographical data, although unremarkable, give a few important clues to her current health status.



Case Study Findings

Reviewing Mrs. Malone's biographical data:

- 40-year-old woman.
- Northern European extraction.
- Protestant religion.
- Married, with three children, ages 12, 15, and 18.
- Has a full-time, sedentary job as an accountant.
- Source: Self; seems reliable.

Current Health Status

Your patient's current health status focuses on her chief complaint. She may seek healthcare because of a diagnosed illness or a set of worrisome symptoms. Or she may feel well and need information about maintaining her current good health. If she identifies an illness, obtain a chronological account of its appearance and development. Be sure to include a symptom analysis (PQRST) of any problems identified.

Mrs. Malone denies any current health problems. Rather, her main concern is staying healthy. She states, "I'm concerned about 'the change' because my mother had so many problems at that time. I want to know what I can do at my age to stay healthy and avoid osteoporosis."



Case Study Findings

Mrs. Malone's current health status includes:

- No current health problems.
- Concerned about physiological changes of approaching menopause and healthcare recommended for her age.
- Occasionally takes acetaminophen for tension headaches. No other medications.

Past Health History

A past health history tells you about any diseases, injuries, or medications in the patient's past that might affect her current health. Mrs. Malone has been healthy all her life, so her past health history is uncomplicated.



Case Study Findings

Mrs. Malone's past health history reveals:

- Had chickenpox, measles, mumps, and rubella as a child. Also had frequent bouts of strep throat as a child.
- Had an appendectomy at age 15. Was hospitalized for birth of her three children. Had a tubal ligation at age 35 for fertility control. Has never had a serious injury requiring hospitalization.
- No history of medical problems.
- No drug, environmental, or food allergies. Has taken penicillin without a reaction.
- Had all childhood immunizations. Last diphtheria-tetanus shot 10 years ago. Negative skin test 5 years ago.
- Has not traveled in a foreign country within last year. Has never served in the armed services.

Family History

Familial health problems are an important indicator of your patient's current and future health. Mrs. Malone is well versed in her family's health history and provides you with detailed information.



Case Study Findings

Mrs. Malone's family history reveals:

- Patient, age 40, alive and well.
- Husband, age 45, alive and well.
- Son, age 18, alive and well.
- Daughter, age 15, alive and well.
- Daughter, age 12, alive and well.
- Mother, age 76, had breast cancer at age 54, osteoporosis and fractured hip at age 63.
- Father, age 78, has history of hypertension (HTN) and had a myocardial infarction (MI) at age 68.
- Younger brother died in car accident at age 35.
- Two sisters, ages 42 and 50, alive and well.
- Maternal grandmother died of breast cancer at age 78.
- Maternal grandfather died of MI at age 81.
- Maternal aunt, age 78, has history of breast cancer and osteoporosis.
- Maternal uncle, age 80, has history of HTN.
- Paternal grandmother died of stroke at age 82.
- Paternal grandfather died of MI at age 86.
- Paternal aunt, age 65, alive and well.
- Paternal aunt, age 69, history of breast cancer.
- Paternal uncle, age 71, history of HTN.

Mrs. Malone's genogram is shown in Figure 22.1.

Review of Systems

The review of systems takes the patient's chief complaint and relates it to every other body system to determine its impact on the patient's total health. Asking your patient questions about every body system may also trigger her memory so that she asks more questions and voices more concerns.

Mrs. Malone reports no serious problems during your review of systems. By all indications, she is healthy and is experiencing normal changes for a woman in midlife.



Case Study Findings

Mrs. Malone's review of systems reveals:

- **General Health Status:** Patient feels that her overall health is good. No fever, chills, fatigue, depression, or anxiety. She has occasional colds and has gained 10 lb in the last year without a change in diet.
- **Integumentary:** No rashes, lesions, mole changes, or bruising. Skin is dry and hair is graying slightly.
- **HEENT:** No headaches; frequent sore throats; hoarseness; or problems with vision, hearing, or sinuses. Last eye exam 2 years ago, doesn't recall last hearing exam. No problems with mouth or teeth, 28 permanent teeth, wisdom teeth removed. Last dental exam 6 months ago.
- **Respiratory:** No shortness of breath, wheezing, or productive or nonproductive cough.
- **Cardiovascular:** No history of HTN, heart problems or murmurs, deep vein thrombosis (DVT) or phlebitis, chest pain, palpitations, edema, orthopnea, or claudication.
- **Breasts:** Performs breast self-examination (BSE) occasionally, when she remembers. Never had mammogram. No breast pain, tenderness, masses, nipple discharge, or change in nipples. Has breast pain 7 days before menstrual cycle. Breastfed all three children.
- **Gastrointestinal:** No nausea/vomiting, abdominal pain, dysphasia, heartburn, jaundice, hemorrhoids, or blood in stool. Has daily, formed bowel movement, brown in color.
- **Genitourinary:** No dysuria, frequency, or urgency, although she reports occasional stress incontinence.
- **Reproductive:** Last menstrual period 21 days ago. Menarche at age 14. Gravida 3, para 3. Menses are irregular, ranging from 23 to 45 days (cycle used to be 28 to 30 days). Menses last 5 days with moderate-to-heavy flow. Contraception method is tubal ligation. Satisfied with sex life except for increasing vaginal dryness.

Genogram for Mrs. Malone




FIGURE 22.1. Genogram for Mrs. Malone.



Case Study Findings (continued)

- **Musculoskeletal:** No joint pains, swelling, or muscle weakness.
- **Neurological:** No dizziness, vertigo, syncope, head injuries, seizures, numbness, or tingling of extremities. Reports forgetfulness and difficulty concentrating. Has never been treated for mental disorders, depression, or anxiety, although she did visit a counselor about a family problem 3 years ago.
- **Endocrine:** No diabetes, thyroid disease, polyuria/polydipsia, or heat or cold intolerance. Weight gain of 10 lb last year.
- **Immunologic/Hematologic:** No current infections, allergies, cancer, bleeding, anemia, or blood transfusions. Blood type A+.

Psychosocial Profile

The psychosocial profile reveals patterns in the patient's lifestyle that affect her health, increase her risk for health problems, or influence her adaptation to them. Mrs. Malone's psychosocial profile pinpoints some behaviors that increase her risk for illness.



Case Study Findings

Mrs. Malone's psychosocial profile includes:

- Does not usually seek routine healthcare or screenings. Feels that she is healthy and does not need healthcare unless she is sick. Does not know her cholesterol level, has never had a mammogram, has not had a Pap smear or physical examination in 3 years.

(continued on page 800)



Case Study Findings (continued)

- Typical day consists of arising at 6:00 A.M., getting her children off to school, and then going to work. She works from 8:00 A.M. to 5:00 P.M. 5 days a week. After work, she makes dinner for her family and reads or watches television. She goes to bed at 11:00 P.M. each night.
- Reports that she eats a balanced diet with foods from each of the major food groups each day, including 3 vegetables and 2 fruit servings. Admits to eating sweets every day. Evaluation of daily calcium intake reveals that she usually ingests 500 mg/day. Has had an unexplained 10-lb weight gain in last year.
- 24-hour recall shows the following: Breakfast was two fried eggs, two slices of bacon, two slices of white toast with butter, one 4-oz glass of orange juice, and a cinnamon bun with butter. Lunch was a chicken salad sandwich with mayonnaise on two slices of white toast, potato chips, two chocolate chip cookies, an 8-oz cola, and a chocolate candy bar. Snack was one apple. Dinner was steak, baked potato with butter, string beans, tossed salad with Russian dressing, apple pie with ice cream, and 8 oz of 2 percent low-fat milk.
- Admits that her only exercise is walking from her car to her office twice a day and walking up and down two flights of stairs several times a day. She walked on a daily basis in the past but stopped this year because of lack of time and inability to fit it into her schedule. For recreation she reads, does housework, and gardens on weekends.
- Hobbies include reading and gardening. Has a house cat.
- States that she usually sleeps well for 7 to 8 hours a night. Has occasional night sweats that awaken her.
- Has never smoked and is not exposed to smoke at home. Has smoke and carbon monoxide detectors at home. Drinks a glass of wine with dinner two to three times a week. Denies ever using illicit drugs.
- Identifies no occupational risks. Is not exposed to smoke at work. Drives 40 miles round trip to work each day. Wears seatbelt when driving.
- Attends church weekly with family. No religious or cultural influences that affect health practices.
- States that her marriage is supportive and comfortable. Describes good relationships with children, siblings, and parents.
- Uses prayer, friendships, and family support to deal with stress.



CRITICAL THINKING ACTIVITY # 1

What problems can you identify for Mrs. Malone after completing her health history?



CRITICAL THINKING ACTIVITY # 2

Based on information provided in Mrs. Malone's health history, what will you include in your physical assessment?



Case Study Evaluation

Before you move on to the physical examination, review and summarize the key information from Mrs. Malone's health history.

Physical Assessment

Once you have completed the subjective data collection, begin to collect objective data by means of the physical examination. Keep the key history findings in mind. These findings, along with those from the physical examination to come, will complete the assessment picture. Then you can analyze the data, formulate nursing diagnoses, and develop a plan of care for Mrs. Malone.

The purpose of the physical assessment is to find signs of disease or abnormalities that the patient may or may not have reported in the history or review of systems. If time is an issue and you are unable to perform a complete health history, perform a focused history.

Approach

Two methods are used for completing a total physical assessment. These methods are a systems approach and a head-to-toe approach. Each approach is systematic and complete, and obtains the same data. The advantages of each assessment method relate to the preferences of the individual nurse, the patient population, and the setting for the assessment. Some nurses find it easier to conceptualize and perform the assessment according to systems, and others are comfortable with a head-to-toe (or regional) approach.

A systems approach allows a thorough assessment of each system, doing all assessments related to one system before moving on to the next. This approach works well when you are performing a focused assessment. A head-to-toe assessment includes the same examinations as a systems assessment, but you assess each region of the body before moving on to the next. With this method, the patient does not need to change positions as often, and

Performing a Focused Physical Assessment

When your patient's condition or time restraints do not permit a comprehensive assessment, perform a focused assessment. This assessment focuses only on systems that relate to your patient's problem. Because it is only a partial assessment, less data are collected during the examination.

A focused physical assessment should include:

- A general survey with vital signs and weight.
- Assessment of level of consciousness.
- Assessment of skin color, temperature, and texture.
- Testing of gross motor balance and coordination.
- Testing of extraocular movements (EOMs).
- Testing of pupillary reaction.
- Testing of gross vision and hearing.
- Inspection of oral mucosa as patient says "ah."
- Auscultation of anterior and posterior breath sounds.
- Palpation of apical impulse, point of maximal impact (PMI).
- Auscultation of heart sounds.
- Auscultation of bowel sounds.
- Percussion of abdomen.
- Palpation of abdomen.
- Palpation of peripheral pulses.
- Testing sensation to touch on extremities.
- Palpation of muscle strength of upper and lower extremities.

you can also compare regions of the body for symmetry. This approach is best suited for a complete physical examination. No matter which approach you use, be systematic and consistent.

For the purposes of this text, a convenient head-to-toe approach is demonstrated. All four assessment techniques—inspection, percussion, palpation, and auscultation—are used to perform a complete examination.

- Inspect for abnormalities and normal variations of visible body parts.
- Palpate to identify surface characteristics, areas of pain or tenderness, organs, and abnormalities, including masses and fremitus.
- Percuss to determine the density of underlying tissues and to detect abnormalities in underlying organs.
- Auscultate for sounds made by body organs, including the heart, lungs, intestines, and vascular structures.

Assessment data are usually charted by systems (e.g., respiratory or neurological) and by regions to a limited extent (e.g., head/neck). Your documentation can focus on only positive findings or on both positive and negative findings. No matter which format you use, always be brief and to the point and avoid generalizations.



TOOLBOX

Because you are performing a complete head-to-toe examination, you will use all four techniques and most of the assessment tools previously described in this text. The tools you use depend on the patient's problems and age and the extent of the examination. You will definitely need a stethoscope, sphygmomanometer, tongue blades, tuning fork, reflex hammer, gloves, supplies for a rectal examination, supplies for a vaginal examination if your patient is a woman, an oto-ophthalmoscope, Snellen charts or visual acuity tests as available, and an audiometer for a hearing acuity test. Be sure to wear gloves when assessing the mucous membranes, genitalia, rectal area, and any area that might expose you to body fluids (such as lesions), infections, or infestation.

Performing a General Survey

The first part of the physical assessment is the general survey. At this time, you will take the patient's vital signs, height, and weight and note his or her overall appearance. Observe nutritional status, emotional status, body habitus, and general appearance, including hygiene and clothing.

Performing a Head-to-Toe Physical Assessment

Now, keeping the patient's chief complaint in mind, examine her or him for specific signs of diseases affecting other organ systems. Have the patient change into a gown and ensure privacy by closing the door or pulling the curtain. Drape the patient, exposing only the area being assessed. As you perform the assessment, tell her or him what you are planning to do and why.



HELPFUL HINTS

- Wash your hands before you begin.
- Listen to your patient.
- Provide a warm environment.
- If your patient has a problem, start at that point.
- Work from head to toe.
- Compare side to side.
- Let your patient know your findings.
- Use your time not only to assess but also to teach your patient.
- Leave sensitive or painful areas until the end of the examination.



CRITICAL THINKING ACTIVITY # 3

What are Mrs. Malone's strengths? Are there any areas of concern?

Below is a head-to-toe physical assessment by region for Mrs. Malone.

Performing a Head-to-Toe Physical Assessment	
AREA/REGION EXAMINED AND TECHNIQUE USED	FINDINGS FOR MRS. MALONE
<p>General Health Survey <i>INSPECT</i> Age: Appearance vs. stated. General health appearance, including nutritional status, hygiene, odors, clothing. Level of consciousness, speech, affect.</p> <p>Signs of distress. Motor activity, gait, posture, gross deformities.</p> <p>Measure Anthropometric data and vital signs.</p> <p>Integumentary <i>INSPECT/PALPATE</i> Inspect visible skin for color and lesions throughout examination. Palpate skin temperature, texture, moisture, turgor. Inspect/palpate hair for texture, color, distribution. Inspect/palpate nails for texture, color, refill, and angle of attachment.</p> <p>HEENT <i>Head/Face</i> <i>INSPECT/PALPATE</i> Inspect size, shape, position. Palpate scalp mobility, tenderness, lesions, masses. Inspect nasolabial folds and palpebral fissures for facial symmetry. Note facial expression and ability to frown, smile (cranial nerve [CN] VII). Palpate temporomandibular joint (TMJ); test range of motion. Palpate temporal arteries. Test facial sensation (CN V).</p> <p><i>Eyes</i> <i>INSPECT/PALPATE/FUNDOSCOPIC</i> Test visual acuity with Snellen or pocket vision screener (CN II). Test color vision with color bars on Snellen chart. Test near vision. Test peripheral vision by confrontation. Test EOMs by six cardinal fields of gaze. Test corneal light reflex, corneal blink reflex. Perform cover-uncover test. Note general appearance of eyes, eyelids, sclera, conjunctiva.</p>	<p>Appears younger than stated age (40 years). Well-developed, well-nourished woman, neatly dressed and well groomed. Oriented to person, place, and time; appropriate affect. Speech clear and responds appropriately. No acute distress at this time. Moves all extremities well, gait balanced and coordinated.</p> <p>Height 5'5"; weight 145 lb. <i>Vital signs:</i> Temperature, 98.8°F; pulse, 74; respirations, 18; BP, 128/80.</p> <p>Skin even in color, no suspicious lesions. 3-cm white scar from appendectomy on right lower quadrant (RLQ). Skin warm, smooth and dry, + turgor Hair clean, coarse, slightly graying, evenly distributed. Nails pink, brisk capillary refill, no clubbing.</p> <p>Normocephalic, erect, midline. Scalp mobile, no lesions, tenderness, or masses. Facial features symmetrical.</p> <p>No TMJ tenderness, locking, or clicking.</p> <p>+2 temporal arteries. CN V intact.</p> <p><i>Snellen:</i> R: 20/20; L: 20/20; both: 20/20.</p> <p>Color vision intact. Difficulty noted with near vision. Visual fields normal by confrontation. EOMs intact; no nystagmus (CN III, IV, VI). Corneal light reflex symmetrical bilaterally. No drifting. Eyes clear and bright; positive blink reflex; no lid lag, ectropion, or entropion, lesions on lids; sclera white; conjunctivae clear and glossy.</p>

AREA/REGION EXAMINED AND TECHNIQUE USED	FINDINGS FOR MRS. MALONE
<p>Inspect cornea, anterior chamber, iris, lens with oblique lighting. Palpate lacrimal glands and ducts. Test pupillary reaction: direct, consensual, convergence.</p> <p>Perform fundoscopic examination and check for red reflex (darken room).</p>	<p>Cornea, iris intact; anterior chamber.</p> <p>Lacrimal glands and ducts nontender. Pupils 3 mm, PERRLA (see Chapter 12) direct and consensual; positive constriction and convergence. Red reflex present bilaterally, discs flat with sharp margins, vessels present without crossing defects, retina even in color without hemorrhages or exudates, macula even in color.</p>
<p><i>Ears</i> INSPECT/PALPATE/OTOSCOPIIC Inspect/palpate external ear.</p> <p>Check angle of attachment.</p> <p>Perform Weber, Rinne, whisper tests (CN VIII).</p> <p>Perform otoscopic examination of canals and tympanic membrane.</p>	<p>Skin of external ear intact with no masses, lesions, or discharge. Angle of attachment <10 degrees. Tragus, mastoid, helix nontender. Positive whisper test. <i>Weber test</i>: No lateralization. <i>Rinne test</i>: ac > bc bilaterally. External ear canals clear without redness, swelling, lesions, or discharge. Tympanic membranes intact, pearly gray, with light reflex and landmarks visible.</p>
<p><i>Nose/Sinuses</i> INSPECT/PALPATE Palpate sinuses for tenderness. Palpate nasal patency. Test sense of smell (CN I). Inspect nasal mucus membranes, septum, turbinates with speculum.</p>	<p>Frontal, maxillary sinuses nontender. Nares patent. Patient recognizes familiar odor. Nasal mucosa pink, septum intact with no deviation.</p>
<p><i>Mouth</i> INSPECT/PALPATE Inspect/palpate lips, oral mucosa. Inspect teeth, gingiva.</p> <p>Inspect pharynx, tonsils, palate. Inspect tongue, test taste on anterior/posterior tongue (CN VII, IX). Check mobility of tongue (CN XII). Test gag, swallow reflexes (CN IX, X).</p>	<p>Lips, oral mucosa, gingivae pink and without lesions. Teeth all present and clean; dental work present; no obvious caries. Pharynx pink, tonsils +1, palate intact. Tongue smooth, pink, symmetrical with no lesions; taste intact. Full range of motion (ROM). Symmetrical rise of the uvula, positive gag and swallow reflex. Glands nonpalpable.</p>
<p>Palpate parotid and submandibular glands.</p> <p><i>Neck</i> INSPECT/PALPATE/AUSCULTATE Inspect/palpate thyroid gland. Palpate/auscultate carotid pulses. Inspect/measure jugular venous pressure (JVP). Palpate nodes in head, neck. Note deviation of trachea, abnormal masses, pulsations. Note ROM of neck. Test neck muscle strength (CN XI).</p>	<p>Thyroid nonpalpable. Carotid pulses equal, +2. No bruits, thrills, or jugular venous distension (JVD). JVP 2 cm at 45-degree angle. No lymphadenopathy of head, neck. Trachea midline, no abnormal masses or pulsations. Full ROM of neck. +5 muscle strength.</p>

(continued)

Performing a Head-to-Toe Physical Assessment (continued)

AREA/REGION EXAMINED AND TECHNIQUE USED	FINDINGS FOR MRS. MALONE
<p>Posterior Thorax <i>Lungs</i> INSPECT/PALPATE/PERCUSS/AUSCULTATE Inspect and palpate excursion. Palpate fremitus. Percuss the chest and diaphragmatic excursion.</p>	<p>Equal excursion, no lag; equal fremitus. Resonance percussed bilaterally. Posteriorly, diaphragmatic excursion 2 inches (5 cm) separate. Lungs clear, no adventitious sounds.</p>
<p>Auscultate breath sounds. <i>Spine</i> INSPECT/PALPATE/PERCUSS Inspect normal curves of spine (cervical, thoracic, lumbar, sacral). Inspect and test for scoliosis, kyphosis, lordosis. Check ROM of spine. Palpate paravertebral muscles and posterior thorax for tenderness, masses. Fist/blunt percuss costovertebral angle (CVA) tenderness.</p>	<p>Normal curves of spine. No scoliosis, kyphosis, lordosis. Full ROM. Paravertebral muscles nontender. No CVA tenderness.</p>
<p>Anterior Thorax INSPECT/PALPATE Breasts and lymph nodes (axillary and epitrochlear) with patient in various positions. INSPECT/PALPATE/PERCUSS/AUSCULTATE Precordium <i>Lungs</i> Inspect and palpate excursion. Inspect AP:lateral ratio. Palpate fremitus. Percuss lungs. AUSCULTATE Lungs, heart.</p>	<p>Breasts symmetrical, no masses or retraction, no nipple discharge, no lymphadenopathy. PMI 1 cm at fifth intercostal space at midclavicular line. Chest expansion symmetrical, no tenderness, scars, masses, lesions. Anteroposterior transverse diameter 1:2; respirations 16/min unlabored. Equal excursion; equal tactile fremitus. Resonant percussion sound over lung fields. Lungs clear; no adventitious breath sounds. Heart rate 85 BPM and regular, S₁, S₂; no S₃, no S₄, murmurs, gallops, or thrills present.</p>
<p>Upper Extremities INSPECT/PALPATE Palpate brachial, radial, ulnar pulses. Perform Allen's test, if indicated. Perform Tinel or Phalen test for carpal tunnel syndrome, if indicated. Check nails for color, temperature, capillary refill, deformities, clubbing. Inspect hands for joint deformities. Palpate skin temperature and test hand grip. Measure arm lengths and circumferences, note dominant side. Check ROM and strength of upper extremities and hands. Test for pronator drift. Test for coordination with rapid alternating movement (RAM) and finger-thumb opposition.</p>	<p>+2 pulses. Negative Phalen test. Nails pink, brisk capillary refill, no clubbing. No deformities, full ROM. Hands warm, equal +5 hand grip. Patient right-handed, arm lengths and circumferences equal. Full ROM of upper extremities, +5 muscle strength. Negative pronator drift. Positive RAM of hands and finger-thumb opposition.</p>

AREA/REGION EXAMINED AND TECHNIQUE USED	FINDINGS FOR MRS. MALONE
Test for accuracy of movements with point-to-point localization.	Accuracy of movements intact.
Test superficial and deep sensations of arms and hands. Test biceps, triceps brachioradialis DTR.	Superficial and deep sensations intact. +2 DTR of upper extremities.
Abdomen	
<i>INSPECT/AUSCULTATE/PERCUSS/PALPATE</i>	
Inspect for shape, scars, movements, respirations, pulsations, peristalsis, abnormalities. Test for hernias. Auscultate for bowel sounds, vascular sounds. Percuss abdomen and organs. If ascites is suspected, percuss for shifting dullness. Palpate abdomen, liver, kidneys, spleen, aorta.	Abdomen flat, no masses or pulsations, surgical scar on RLQ. Tympany in all four quadrants. Bowel sounds present, no vascular sounds heard. Tympany in all four quadrants. Liver 8 cm at right midclavicular line (RMCL). Abdomen soft, no hepatomegaly, splenomegaly, masses, tenderness. kidneys not palpable, aorta 2 cm.
Test superficial abdominal reflexes.	Negative abdominal reflexes.
Palpate inguinal nodes.	No palpable lymph nodes in inguinal area.
Palpate/auscultate femoral arteries.	No femoral bruits; pulses 2+ and equal.
Lower Extremities	
<i>INSPECT/PALPATE</i>	
Inspect skin color, hair distribution, varicose veins.	Skin warm, hair on both lower legs, no varicose veins.
Perform Trendelenburg test or manual compression test to check venous circulation, if indicated.	
Palpate pedal pulses, temperature.	+2 pedal pulses.
Inspect feet for deformities, toenail condition, lesions.	
Test ROM in lower extremities.	Full ROM in lower extremities, no crepitus.
Measure leg lengths and circumferences.	Leg lengths and circumferences equal.
Perform straight leg test for herniated disc, if indicated.	
Perform patellar tap or bulge sign if fluid is suspected.	
Check for draw sign if torn anterior cruciate ligament suspected.	
Perform Apley's or McMurray's test if torn meniscus suspected.	
Test muscle strength and superficial and deep sensations.	Muscles well developed, +5 muscle strength, superficial and deep sensations intact.
Inspect gait, toe walk, heel walk, deep knee bend.	Gait steady and coordinated, heel-and-toe walk, deep knee bend without difficulty.
Perform Romberg test.	Negative Romberg.
Have patient toe tap and run heel down shin.	Movements coordinated.
Test patellar and Achilles DTR and plantar reflex.	+2 DTR, positive plantar flexion.
	Joints and muscles symmetrical.
Female Genitalia/Rectum	
<i>INSPECT/PALPATE</i>	
Inspect and palpate external and internal genitalia.	External genitalia without lesions or abnormalities.
	Vaginal mucosa pale pink/dry with few rugae.
	Cervix pale pink, no lesions or discharge.
	Nontender, uterus midline, normal size.
	Adnexa without masses or tenderness.
	Perianal area intact without hemorrhoids.
	Rectal wall smooth without masses or tenderness.
Inspect/palpate rectum for masses.	Stool soft, brown, occult blood negative.
Test stool for occult blood.	

Documenting the Physical Assessment Findings

Once you have completed your assessment, document

your findings. You performed the physical assessment by region; now document your findings by system.

Documenting the Physical Assessment Findings

AREA/SYSTEM	FINDINGS
General Health Survey	<ul style="list-style-type: none"> ■ Appears younger than stated age (40 years). ■ Well-developed, well-nourished woman, neatly dressed and well groomed. ■ Oriented to person, place and time; appropriate affect. ■ Speech clear and responds appropriately. ■ No acute distress at this time. ■ Moves all extremities well, gait balanced and coordinated. ■ Height 5'5"; weight 145 lb.
Integumentary	<ul style="list-style-type: none"> ■ Vital signs: Temperature, 98.8°F; pulse, 74; respirations, 18; BP, 128/80. ■ Skin even in color, warm, dry, positive turgor, no suspicious lesions. ■ White scar from appendectomy on RLO. ■ Hair clean, coarse, slightly graying, evenly distributed. ■ Nails pink, brisk capillary refill, no clubbing.
Head/Face/Neck	<ul style="list-style-type: none"> ■ Normocephalic, erect, midline. ■ Scalp mobile, no lesions, tenderness, or masses. ■ Facial features symmetrical. ■ Thyroid not palpable.
Eyes	<ul style="list-style-type: none"> ■ No palpable lymph nodes. ■ <i>Snellen</i>: R: 20/20; L: 20/20; Both: 20/20. ■ Color vision intact. ■ Difficulty noted with near vision. ■ Visual fields normal by confrontation. ■ EOMs intact, no nystagmus. ■ Corneal light reflex symmetrical bilaterally. ■ No drifting. ■ Eyes clear and bright; positive blink reflex; no lid lag, ectropion/entropion, or lesions noted on lids; cornea, iris intact; anterior chamber clear. ■ Sclera white, conjunctivae clear and glossy. ■ Lacrimal glands and ducts nontender. ■ Pupils 3 mm, PERRLA direct and consensual. ■ Positive constriction and convergence. ■ Red reflex present bilaterally, discs flat with sharp margins, vessels present without crossing defects, retina even in color without hemorrhages or exudates, macula even in color.
Ears/Nose/Throat	<ul style="list-style-type: none"> ■ Skin of external ear intact with no masses, lesions, or discharge. ■ Angle of attachment <10 degrees. ■ Tragus, mastoid, helix nontender. ■ Positive whisper test. ■ <i>Weber test</i>: No lateralization. ■ <i>Rinne test</i>: ac > bc bilaterally. ■ External ear canals clear without redness, swelling, lesions, or discharge. ■ Tympanic membranes intact, pearly gray with light reflex and landmarks visible. ■ Frontal and maxillary sinuses nontender. ■ Nares patent; patient recognizes familiar odor. ■ Nasal mucosa pink, septum intact with no deviation.

AREA/SYSTEM	FINDINGS
Mouth	<ul style="list-style-type: none"> ■ Lips, oral mucosa, gingivae pink and without lesions. ■ Teeth all present and clean; dental work present, no obvious caries. ■ Pharynx pink, tonsils +1, palate intact. ■ Symmetrical rise of uvula, positive gag and swallow reflex. ■ Tongue smooth, pink, symmetrical, no lesions, taste intact, full ROM. ■ Glands nonpalpable.
Respiratory	<ul style="list-style-type: none"> ■ Respirations 16/min unlabored, trachea midline. ■ AP < transverse diameter. ■ Chest expansion symmetrical; no tenderness, scars, masses, lesions. ■ Equal excursion, no lag, equal tactile fremitus. ■ Resonant percussion sound over lung fields, diaphragmatic excursion 2 inches (5 cm). ■ Lungs clear; no adventitious breath sounds.
Cardiovascular	<ul style="list-style-type: none"> ■ PMI 1 cm at fifth intercostal space at midclavicular line. ■ Heart rate 85 BPM and regular; S₁, S₂; no S₃, S₄, murmurs, gallops, or thrills present. ■ Pulses +2, no bruits or thrills, no varicose veins. ■ JVP 2 cm at 45-degree angle.
Breasts	<ul style="list-style-type: none"> ■ Symmetrical, no masses, nipple retraction, nipple discharge, lymphadenopathy.
Abdomen	<ul style="list-style-type: none"> ■ Abdomen flat, no masses or pulsations. ■ Surgical scar on right lower abdomen. ■ Bowel sounds present, no vascular sounds heard. ■ Tympany in all four quadrants. ■ Liver 8 cm at right midclavicular line. ■ Abdomen soft, no hepatomegaly, splenomegaly, masses, or tenderness. ■ Negative abdominal reflexes. ■ Kidneys nonpalpable; no palpable lymph nodes in inguinal area. ■ Aorta 2 cm. ■ No CVA tenderness.
Reproductive	<ul style="list-style-type: none"> ■ External genitalia without lesions or abnormalities. ■ Vaginal mucosa pale pink, dry with few rugae. ■ Cervix pale pink, no lesions or discharge. ■ No cervical motion tenderness; uterus midline, normal size. ■ Adnexa without masses or tenderness. ■ Perianal area intact, without hemorrhoids. ■ Rectal wall smooth without masses or tenderness. ■ Stool brown, occult blood negative.
Musculoskeletal	<ul style="list-style-type: none"> ■ Normal curves of spine; no scoliosis, kyphosis, lordosis. ■ Joints and muscles symmetrical, nontender; no deformities. ■ Right side dominant, arm and leg lengths and circumferences equal, equal hand grasp. ■ Muscles well developed, +5 muscle strength.
Neurologic	<ul style="list-style-type: none"> ■ Full ROM in upper and lower extremities, no crepitus. ■ Awake, alert, oriented to time, place, person; cooperative; responds appropriately. ■ CNs I through XII intact. ■ Gait steady and coordinated; no pronator drift; negative Romberg; able to heel-and-toe walk and do deep knee bends without difficulty. ■ Point-to-point localization intact. ■ Superficial and deep sensations intact. ■ +2 DTR, +plantar flexion, negative abdominal reflexes.



Case Study Analysis and Plan

Now that you have completed Mrs. Malone's assessment, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mrs. Malone. Now use this information to identify a list of nursing diagnoses. Some possible ones are provided below; cluster the supporting data.

1. Imbalanced Nutrition: more than body requirements
2. Knowledge deficit related to need for age-appropriate health activities



CRITICAL THINKING ACTIVITY #4

Now that you have identified some nursing diagnoses for Mrs. Malone, select one from the list above and develop a nursing care plan and a brief teaching plan for her.

Collaborative Nursing Diagnoses

Identify any collaborative nursing diagnoses, and be specific as to what the desired outcome of each would be.

Referrals

Mrs. Malone's health history and physical examination findings warrant several referrals to either a physician or a practitioner. Identify what needs to be done at each referral and the reasons behind it.

RESEARCH TELLS US

Osteoporosis is a major health problem for the older population in this country, especially women. It places people at risk for fractures as a result of low bone mass and deterioration of bone tissue. The first manifestation of the disease is usually a fracture, most often in the hip, spine, or wrist. Loss of height over 1 inch and spinal deformity are also indicators of osteoporosis. Fractures cause morbidity in the elder population and have psychological effects such as depression, anxiety, fear, and anger.

Osteoporosis has both modifiable and nonmodifiable risk factors. Modifiable risk factors include a sedentary lifestyle, low calcium intake, low body weight, high alcohol consumption, smoking, and estrogen deficiency. Nonmodifiable risk factors include being a white or Asian female over age 65 and having a slight build, a family history of osteo-

porosis, premature or surgical menopause, and endocrine or chronic disease.

Treatments and techniques for diagnosing osteoporosis have improved in recent years. Techniques include dual x-ray absorptiometry, quantitative computed tomography, and quantitative ultrasound. Treatments include antiresorptive therapies, such as hormone replacement therapy, bisphosphonates, and selective estrogen receptor modulators. Encourage your patients to discuss evaluation and treatments with their healthcare providers.

Prevention continues to focus on health education. Nurses can encourage women to modify risk factors through adequate diet and weight control, adequate calcium intake, exercise, and avoidance of smoking and alcohol.

S U M M A R Y


- A comprehensive health history and a physical examination are important components of a health maintenance program.
- Time intervals for a health assessment vary depending on age and the presence of chronic health problems. Standard recommendations are made by various organizations and the U.S. Preventive Services Task Force (2005).
- The content of periodic health examinations varies with age because of the risks of health problems in different age categories.
- In addition, certain age-specific health screenings, patient counseling, and preventive health activities occur during the health assessment. Much like nursing diagnoses and care plans, these screenings are highly

individualized, based on patient need and presentation in the clinical setting.

- Begin your assessment by obtaining a health history. The health history guides your physical assessment.
- After you have completed the health history, perform the physical examination. For a complete assessment, work head to toe by regions.
- Once you have completed your assessment, document your findings. When possible, organize and document the physical assessment findings by systems.
- Remember to share the information with your patient so that you can develop an effective plan of care.

R E V I E W Q U E S T I O N S

1. You are performing a complete history and physical assessment. Which approach is best for performing a complete assessment?
 - a. By system
 - b. By region
2. Which approach is best when performing a focused assessment?
 - a. By system
 - b. By region
3. Which approach is best for documenting your complete assessment findings?
 - a. By system
 - b. By region




UNIT

4

Adapting Assessment to Special Populations

CHAPTER 23



Assessing the Mother-to-Be

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify normal physiological and psychological changes that occur during pregnancy
- Discuss various cultural influences on motherhood
- Identify pertinent history data included in a maternal history (antepartal and postpartal)
- Obtain a maternal history (antepartal and postpartal)
- Describe additional physical assessment components specific to a maternal assessment
- Differentiate between antepartal and postpartal assessment
- Perform a maternal physical assessment (antepartal and postpartal)
- Document maternal assessment findings
- Identify actual/potential health problems stated as nursing diagnoses with supporting data

Pregnancy is usually an exciting and special time in a woman's life. The duration of human pregnancy is 9 calendar months, 10 lunar months, 40 weeks, or 280 days. The length of pregnancy is divided into three **trimesters**, or 3-month periods. Each trimester is characterized by its own unique and predictable developments for the mother and her baby. To accommodate the changes taking place throughout the gestational period, the pregnant woman's body undergoes changes in size and shape, and all of her organ systems modify their various functions to create an environment that is protective and nurturing for the growing fetus.

Anatomy and Physiology Review

The female body undergoes many physiological and anatomic changes during pregnancy. Several factors are

responsible for the woman's adaptation to pregnancy. Hormonal influences, mechanical pressure arising from growth of the fetus inside the uterus, and the mother's physical adaptation to her changing body all account for the changes that take place during pregnancy. The majority of these changes are brought about by the hormones of pregnancy, primarily estrogen and progesterone. Although the most dramatic changes occur in the reproductive system, every other body system is also affected by pregnancy.

Although pregnancy is a normal event, problems can occur. Therefore, you need an understanding of normal maternal physiology so that you can recognize potential or actual problems that warrant attention. Also, your understanding of the normal physiological and psychological events that take place during pregnancy will assist you in teaching your patient and her family about changes that are normal and expected and how to identify signs and symptoms that should be reported to her healthcare provider.

Normal Changes Associated With Pregnancy

SYSTEM/STRUCTURE/CHANGES

CLINICAL SIGNS AND SYMPTOMS

Integumentary

Skin

- Changes result from hormones and mechanical stretching. Increased skin thickness and hyperpigmentation are caused by increased secretion of melanotropin, an anterior pituitary hormone.
- Increased action of adrenocorticosteroids occurs in 50 to 90 percent of pregnant women, causing cutaneous elastic tissue to become more fragile.
- Increased estrogen results in color and vascular changes.
- Increased sebaceous gland secretions.
- Increased blood supply to skin, increased basal metabolic rate (BMR), and progesterone-induced increased body temperature.

- Chloasma, the "mask of pregnancy," is a brownish hyperpigmentation of the skin over the cheeks, nose, and forehead. It appears in 50 to 70 percent of pregnant women and occurs most often in those with dark complexions, usually after the 16th week. Sunlight enhances the heightened pigmentation, which generally fades after delivery.
- Darkening of the nipples, areolae, axillae, and vulva also occur, and scars and moles may darken.
- The linea alba may become pigmented (linea nigra) and extends from the symphysis pubis to the umbilicus.
- Stretch marks (*striae gravidarum*)—pinkish-red streaks with slight depressions in the skin—may appear over the abdomen, thighs, breasts, and buttocks, fading to silvery or white after pregnancy.
- **Angiomas**, or *vascular spiders*, are tiny, branched, pulsating end-arterioles on the neck, chest, face, and arms. These skin lesions are bluish, do not blanch with pressure, and usually disappear after the baby's birth.
- **Palmar erythema** is characterized by a pinkish-red, diffuse mottling over the palms of the hands.
- The skin develops increased oiliness and acne or takes on a "healthy glow."
- Increased perspiration and feeling "hotter."

SYSTEM/STRUCTURE/CHANGES	CLINICAL SIGNS AND SYMPTOMS
<i>Hair</i>	
<ul style="list-style-type: none"> ■ Increased hair growth during pregnancy as a result of hormonal influences. 	<ul style="list-style-type: none"> ■ Some women may have excessive hair growth in unusual places (hirsutism). Increase in fine body hair growth also occurs but disappears after delivery. Increase in brittle hair growth usually does not. Excessive scalp oiliness or dryness may also occur.
<i>Nails</i>	
<ul style="list-style-type: none"> ■ Changes in nail growth and texture as a result of hormonal influences. 	<ul style="list-style-type: none"> ■ Nails may grow longer, soften, or thin.
HEENT	
<i>Ear</i>	
<ul style="list-style-type: none"> ■ Increased vascularity of upper respiratory tract may cause swelling of the tympanic membrane and eustachian tube. 	<ul style="list-style-type: none"> ■ Decreased hearing, a sense of fullness in the ears, or earaches.
<i>Nose</i>	
<ul style="list-style-type: none"> ■ Estrogen-induced edema and vascular congestion of the nasal mucosa and sinuses. 	<ul style="list-style-type: none"> ■ Nasal stuffiness and epistaxis (nosebleed).
<i>Mouth/Throat</i>	
<ul style="list-style-type: none"> ■ Edema of the larynx. ■ Higher estrogen levels increase vascularity and connective tissue proliferation. 	<ul style="list-style-type: none"> ■ Some women may experience vocal changes. ■ Gum hypertrophy and bleeding of gums while brushing teeth is common. Epulis—raised, red nodules on gums that bleed easily—may develop but generally regress after delivery.
Respiratory	
<ul style="list-style-type: none"> ■ Estrogen promotes relaxation of the ligaments and joints of ribs. 	<ul style="list-style-type: none"> ■ Increase in transverse diameter by 2 cm with a total circumference increase of 6 cm. Increase in the costal angle > 90 degrees. ■ Thoracic breathing as pregnancy progresses. ■ The diaphragm becomes displaced as pregnancy progresses. ■ Slight increase in respiratory rate. ■ 30 to 40 percent increase in tidal volume. ■ Increase in inspiratory capacity. ■ Decrease in expiratory volume. ■ Total lung capacity slightly decreased. ■ Decrease in Pco₂ (27 to 32 mm Hg) leads to increase in pH (more alkaline) and decrease in bicarbonate (18 to 21 mEq/L).
<ul style="list-style-type: none"> ■ Increase in oxygen consumption by 15 to 20 percent. 	
<ul style="list-style-type: none"> ■ Higher levels of progesterone increase sensitivity of respiratory receptors, increasing tidal volume, which results in respiratory alkalosis with compensated mild metabolic acidosis. 	
Cardiovascular	
<ul style="list-style-type: none"> ■ Increase in cardiac output and maternal blood volume by approximately 40 to 50 percent. Because the heart must pump harder, it actually increases in size. The body adapts to increase in blood volume with peripheral dilation to maintain BP. Hormones cause peripheral dilation. 	<ul style="list-style-type: none"> ■ Heart rate may increase by 10 to 15 beats per minute, and systolic murmurs may be heard. Increase in blood volume may cause physiological or “pseudo” anemia because the plasma increase exceeds red blood cell production. Sinus arrhythmias and premature atrial or ventricular contractions may occur. BP is normal in first and third trimesters; systolic and diastolic BP drops 5 to 10 mm Hg during second trimester.
<ul style="list-style-type: none"> ■ Compression of vena cava impairs venous return and results in decreased cardiac output when woman is supine during second half of pregnancy. 	<ul style="list-style-type: none"> ■ Orthostatic hypotension can occur as a result of decreased venous return and decreased cardiac output.

(continued)

Normal Changes Associated With Pregnancy (continued)

SYSTEM/STRUCTURE/CHANGES

CLINICAL SIGNS AND SYMPTOMS



ALERT

Watch for signs of supine hypotension (vena caval syndrome), which may occur when pregnant patient lies in a supine position. Weight of abdominal contents may compress vena cava and aorta, causing a 30 mm Hg drop in BP with reflex tachycardia. Signs and symptoms include pallor, dizziness, faintness, breathlessness, tachycardia, nausea, and clammy skin. Reposition patient on left side until signs and symptoms subside and vital signs return to within normal limits.

- Compression of iliac veins and inferior vena cava increases venous pressure and decreases blood flow to extremities.

Breasts

- Increase in estrogen and progesterone soon after conception causes many changes in mammary glands.
- Increased blood supply to breasts.
- Increased growth of mammary glands.
- Increase in luteal and placental hormones leads to increase in lactiferous ducts and lobule-alveolar tissue.

Gastrointestinal

- Increased levels of hCG and altered carbohydrate metabolism.
- Change in senses of taste and smell.
- Decreased swallowing and increased stimulation of salivary glands by starch ingestion.
- After the 7th month of pregnancy, the upper portion of the stomach may herniate.
- Increased progesterone decreases tone and motility of smooth muscles.
- Increased estrogen leads to decrease in hydrochloric acid.
- Increased progesterone decreases muscle tone and peristalsis.
- Gallbladder becomes increasingly distended because of decreased muscle tone. Emptying time is prolonged and bile thickens.
- Displacement of intestines by uterus.

- Dependent edema, varicose veins in legs and vulva, hemorrhoids.

- Breasts may feel full, with increased sensitivity, tingling, and heaviness. Increased nipple erectility and hypertrophy of Montgomery's tubercles (glands).
- Blood vessels become more visible.
- Increase in breast size.
- Striae gravidarum (stretch marks) on breasts.
- Breasts become softer, looser, and nodular. Colostrum produced by 16th week.

- Morning sickness during first trimester.

- May result in decreased appetite or unusual nonfood cravings (pica).
- Nausea.
- Some women develop **ptyalism**, or excessive salivation.
- Hiatal hernia is more likely to occur in older, obese, or multiparous women.
- Esophageal regurgitation, reverse peristalsis, and delayed stomach emptying result in heartburn (pyrosis).
- Peptic ulcers rarely occur.

- Constipation can be caused by hypoperistalsis, increased water absorption from large intestines, decreased physical activity, displacement of intestines, abdominal distension, and iron supplements.
- Increased risk for gallstones.

- Abdominal changes include round ligament tension, flatulence, distension, cramping, pelvic heaviness, and contractions.

Normal Changes Associated With Pregnancy (continued)	
SYSTEM/STRUCTURE/CHANGES	CLINICAL SIGNS AND SYMPTOMS
<p>Endocrine</p> <p><i>Pituitary and Placental Hormones</i></p> <ul style="list-style-type: none"> ■ Decreased follicle-stimulating hormone. ■ Increased progesterone relaxes smooth muscles, which results in decreased uterine contractions and prevents spontaneous abortion. ■ Increased estrogen. ■ Increased prolactin. ■ Increased oxytocin. ■ Human placental lactogen or hCG acts as a growth hormone and decreases glucose metabolism and increases fatty acids. <p><i>Thyroid</i></p> <ul style="list-style-type: none"> ■ BMR gradually increases throughout pregnancy. ■ Hyperplasia and increased vascularity of thyroid gland. <p><i>Parathyroid</i></p> <ul style="list-style-type: none"> ■ Increased parathyroid hormone peaks between 15 and 35 weeks' gestation to meet increased requirements for calcium and vitamin D for fetal skeletal growth. <p><i>Pancreas</i></p> <ul style="list-style-type: none"> ■ As fetus grows, it requires increasing amounts of glucose. <p><i>Adrenals</i></p> <ul style="list-style-type: none"> ■ Increase in cortisol. <p>Immunologic/Hematologic</p> <ul style="list-style-type: none"> ■ Increased coagulability results from increases in clotting factors VII, VIII, IX, X, and fibrinogen. Fibrinolytic activity is depressed to minimize risk of bleeding. ■ Increase in blood volume 40 to 50 percent > nonpregnant state, about 1500 mL; 1000 mL is plasma, 500 red blood cells. ■ Increase in white blood cells during second and third trimesters. 	<ul style="list-style-type: none"> ■ Amenorrhea. ■ Fat deposits in subcutaneous tissue over abdomen, back, and thighs. ■ Maintains pregnancy. ■ Enlarges uterus, breasts, and genitals; increases vascularity. ■ Causes lactation. ■ Causes uterine contractions at time of delivery and let-down milk reflex. ■ Contributes to breast development. ■ May cause heat intolerance, fatigue, and lassitude. ■ Enlargement of thyroid gland. ■ Slight hyperparathyroidism develops. ■ As mother's glucose stores are depleted, she experiences decreasing blood glucose levels. ■ Increases production of insulin and mother's resistance to insulin. ■ Increased risk for thrombus formation. ■ Hemodilution causes physiological anemia. ■ Increase is seen in granulocytes.

Developmental, Cultural, and Ethnic Variations

Adolescents

Physical assessment of the teenage mother-to-be is essentially the same as for the adult pregnant woman. However, careful attention to the baseline blood pressure (BP) is necessary because teenagers have lower systolic and diastolic pressures than older women. Also, when conducting the physical examination, look for signs of physical and sexual abuse in the young adolescent,

who may be ashamed or afraid to share this important information.



HELPFUL HINT

Remember that adolescent girls are very modest and usually anxious during pelvic examinations. Conducting the examination in a calm, unhurried manner, along with instructions for relaxation techniques, should be helpful.

The diets of adolescents are often lacking in essential nutrients, such as calcium and iron, needed during pregnancy. Additional problems associated with teen pregnancy are related to growth changes occurring in the teenager during the pregnancy. Because the mother and the fetus are both growing, they compete for nutrients. Cephalopelvic disproportion is a common problem for teen pregnancy because the growth of the pelvis lags behind growth in stature. For these reasons, pregnant teens are at increased risk for complications.

Older Women

In general, the risk for maternal or fetal complications increases with the patient's age. Pregnant women over age 35 have an increased risk of developing gestational diabetes, pregnancy-induced hypertension (PIH), gestational bleeding, abruptio placentae, and intrapartum fetal distress. They also have an increased risk for conceiving a child with chromosomal abnormalities. In addition, older women are more likely to have pre-existing medical conditions, such as diabetes and hypertension (HTN). In fact, pre-existing conditions appear to pose a greater risk for

maternal well-being and pregnancy outcome than the mother's age.

People of Different Cultures or Ethnic Groups

When obtaining the biographical information, it is important to establish your patient's ethnicity because certain ethnic populations are more likely to experience race-related genetic disorders. African Americans and Southeast Asians are at risk for sickle cell disease; Jewish women are at risk for Tay-Sachs disease; Mediterranean, Italian, and Greek women are at risk for beta-thalassemia, and Asians are at risk for alpha-thalassemia. Women from these ethnic groups may need to be referred for prenatal screening. Aside from screening for genetic disorders related to specific ethnic groups, assessment of your patient's cultural background may identify pregnancy and childbearing practices specific to a particular cultural/ethnic group. You will need to explore and identify your patient's cultural views and practices on fertility and pregnancy. Specific prescriptive, restrictive, and taboo practices related to pregnancy, birthing, and postpartum care need to be identified and considered when caring for your patient.

Pregnancy and Childbearing Practices of Various Ethnic Groups

CULTURAL GROUP	PRACTICES/BELIEFS
African American	<ul style="list-style-type: none"> ■ Oral contraceptives most popular type of birth control. ■ High rate of teen pregnancy. ■ Primary advisors are grandmother and maternal relatives. ■ Geophagia (eating of earth or clay) common practice. ■ <i>Taboos/beliefs:</i> If food cravings not met, baby is marked. ■ Labor is induced by a bumpy ride, a big meal, castor oil, or sniffing pepper. Photographing pregnant woman causes stillbirth or captures baby's soul. Lifting hands over head causes cord to wrap around baby's neck. Amniotic sac over baby's face denotes that baby has special powers. Child born after set of twins, the seventh child, or a child born with a physical condition has special powers from God.
Amish	<ul style="list-style-type: none"> ■ Children are a gift from God. ■ Average number of children per family is seven. ■ Women have high status because of their role as child producer. ■ Birth control seen as interfering with God's will. ■ Prenatal care by Amish lay-midwives. ■ Fathers expected to be involved and present during delivery and to participate in prenatal classes. ■ No major taboos, but showing emotions is considered inappropriate, so woman labors quietly without expressing discomfort. ■ Childbirth is not viewed as medical condition, so few have health insurance, keeping hospitalizations very short.
Appalachian	<ul style="list-style-type: none"> ■ Family and community are expected to care for mother and baby. ■ Contraceptive practices similar to those of general public, but a common belief is that laxatives facilitate abortion. ■ Believe that healthy living during pregnancy leads to a healthy baby.

(continued)

Pregnancy and Childbearing Practices of Various Ethnic Groups (continued)

CULTURAL GROUP	PRACTICES/BELIEFS
Arab American	<ul style="list-style-type: none"> ■ <i>Taboos/beliefs:</i> Boys are carried high, girls low; photo taking causes stillbirth; reaching over head causes cord strangulation; wearing an opal ring may harm baby; birth marks are caused by eating strawberries or citrus fruit; experiencing a tragedy during pregnancy leads to congenital anomalies. ■ Birthing is considered a natural process to be endured. ■ Family members are expected to care for mother and child during postpartum period. ■ A band may be placed around the baby's abdomen to prevent umbilical hernias and an asafetida bag around the neck to prevent contagious diseases. ■ Believe that God decides family size and God provides. ■ Procreation is the purpose of marriage, resulting in high fertility rates. ■ Irreversible means of contraception and abortions are considered unlawful. Sterility may lead to rejection or divorce. ■ With emphasis on fertility and bearing a son, pregnancy occurs early in marriages. ■ Pregnant women are pampered. Unmet cravings are thought to result in birthmarks. Pregnant women are excused from fasting during sacred days. ■ Men are excluded from labor and delivery. Deliveries are often performed at home by midwives because of limited access to hospitals. Breathing and relaxation techniques are not practiced, and labor pain is openly expressed. ■ Baby's stomach is wrapped after birth to prevent exposure to cold. Male circumcision is a Muslim requirement. ■ Postpartum bathing is avoided for fear of mother being exposed to cold. ■ Breastfeeding is delayed until second or third day to allow mother to rest. ■ Postpartum foods include lentil soup to increase milk production and teas to cleanse the body.
Chinese American	<ul style="list-style-type: none"> ■ To control population, Chinese government has one-child law. Contraception is free; abortion common. ■ Pregnancy is seen as positive experience. Men not very involved in process. Female midwives often preferred because of modesty of Chinese women. ■ <i>Dietary beliefs:</i> Adding meat to diet makes blood "strong" for fetus. Shellfish causes allergies. Iron makes delivery more difficult. Fruits and vegetables should be avoided during postpartum period because they are "cold" foods. Mother drinks rice wine to increase strength and has 5 to 6 meals a day with rice, soups, and 7 to 8 eggs.
Cuban American	<div style="display: flex; align-items: center;">  <div style="text-align: center;"> <p>ALERT</p> <p style="color: red;">Wine may increase bleeding time.</p> </div> </div>
	<ul style="list-style-type: none"> ■ 5- to 7-day hospital stay after delivery, 1 month for recovery. Chinese government allows 6 months' maternity leave with full pay. If mother is breastfeeding, work time is allowed for feedings. ■ Many mothers avoid "cold"—do not go outside or bathe, and wear many layers of clothes even in warm weather. ■ Cuba has lowest birth rate in Latin America because of high labor-force participation for women and high divorce rate. ■ Childbirth is a celebration. ■ <i>Beliefs:</i> Must eat for two during pregnancy (leads to excessive weight gain); eating coffee grounds cures morning sickness; eating fruit ensures that baby will have smooth complexion; wearing necklaces causes umbilical cord to wrap around neck. ■ During postpartum period, family members care for mother for 4 weeks. Mother should avoid ambulation and exposure to cold because this is seen as risk for infection.

CULTURAL GROUP	PRACTICES/BELIEFS
Egyptian American	<ul style="list-style-type: none"> ■ Prefer breastfeeding but wean at 3 months and start solid foods. Believe that a fat child is a healthy child. ■ Must have children to make family complete. Three children ideal number. Woman expected to conceive within first year of marriage, and there is much stress until pregnancy occurs. Inability to conceive is grounds for divorce. Pregnancy brings security and respect. Birthing, especially of a son, has status and power. ■ <i>Beliefs:</i> Curtail activities during pregnancy to prevent miscarriage; eat for two; cravings must be met or baby will be marked with the shape of the craved food. ■ Maternal mother is expectant mother's source of support during labor and delivery. Men excluded from birthing process. In postpartum period, mother should avoid cold, such as bathing. Postpartum period is 40 days, and family tends to mother and baby.
Filipino American	<ul style="list-style-type: none"> ■ Fertility practices influenced by Catholic faith; rhythm is only acceptable method of contraception; abortion is a sin. ■ Child-centered culture. Pregnancy is normal; expectant mothers are pampered. Maternal mother is great source of support and often serves as labor coach. ■ <i>Beliefs:</i> Uses healthcare providers but also a massage therapist. Cravings should be met to avoid harm to baby. Baby takes on characteristics of craved foods, (e.g., dark-colored foods, dark skin). If mother has a sudden scare or stress, it can harm fetus. ■ <i>Postpartum beliefs:</i> Chicken soup stimulates milk production. Showers may cause arthritis, but sponge baths are allowed.
French Canadian	<ul style="list-style-type: none"> ■ Family cares for new mother and baby during postpartum period. ■ Fertility practices influenced by Catholic faith. Abortion considered morally wrong by many. ■ Fear of labor and delivery. Fathers encouraged to be present during delivery. Use of analgesia high. ■ Breastfeeding has regained popularity. ■ Both maternity and paternity leaves are available. ■ <i>Beliefs:</i> Washing floors can trigger labor; so can full moon. Hyperglycemia during pregnancy means that mother is likely to give birth to a boy. Lack of salt during pregnancy means that mother is likely to give birth to a girl.
Greek American	<ul style="list-style-type: none"> ■ Family size is limited because of desire to provide for family and ensure education for children. ■ Greek Orthodox faith condemns birth control and abortion. ■ Infertility causes great stress. ■ Pregnant woman is greatly respected and protected. ■ Birthing usually by midwife or female relatives; fathers remain uninvolved. ■ Mother considered impure and susceptible to illness for 40 days postpartum. Remains at home. Breastfeeding common. ■ <i>Beliefs:</i> Encouraged to eat foods high in iron and protein. Child will be marked if food cravings are unmet. Breastfeeding mothers who shower may cause diarrhea or milk allergy in baby. Silver objects or coins placed in crib bring good luck.
Iranian American	<ul style="list-style-type: none"> ■ Hot and cold influence pregnancy practices. ■ Women are expected to have children early in marriage. ■ Infertility blamed on woman. ■ Breastfeeding used as method of contraception. ■ Bearing a child, especially a boy, is prestigious. ■ <i>Beliefs:</i> Cravings must be met with balance for hot and cold foods. Heavy work causes miscarriages. ■ Much support from female relatives. Father usually not present during delivery. Postpartum period 30 to 40 days.

(continued)

Pregnancy and Childbearing Practices of Various Ethnic Groups (continued)

CULTURAL GROUP	PRACTICES/BELIEFS
Irish American	<ul style="list-style-type: none"> ■ <i>Postpartum beliefs:</i> Baby boys are “hotter” than girls. Baby may be kept at home for first 40 days until strong enough to defend against environmental pathogens. Ritual baby bath given between 10th and 40th day. ■ Fertility practices influenced by Catholicism. Sexual relationships often seen as a duty. Abstinence and rhythm only acceptable methods of birth control, and abortion considered morally wrong. ■ <i>Beliefs:</i> Eating well-balanced diet during pregnancy important; not eating right leads to deformities. Lifting hands above head wraps cord around baby's neck. Experiencing a tragedy during pregnancy results in congenital anomalies. Going to bed with wet hair or wet feet causes illness in pregnant woman.
Jewish American	<ul style="list-style-type: none"> ■ Children are a gift and duty; males are more important. Sterility is a curse. ■ Birth control pill is acceptable form of contraception. ■ Orthodox Jews condone abortion if mother's health is at risk. Reform Jews believe that women have control over their own bodies, therefore it is their decision. ■ Hasidic husband not allowed to touch mother during delivery or view her genitals, therefore may give only verbal support or choose not to be present during delivery. ■ Male circumcision performed on the 8th day by a mohel, a person trained in circumcision, asepsis, and the religious ritual.
Mexican American	<ul style="list-style-type: none"> ■ Fertility practices influenced by Catholicism. Abstinence and rhythm acceptable methods of birth control. Breastfeeding also seen as method of birth control. Abortion morally wrong. ■ Multiple births common; childbearing age between 19 and 24 is seen as best time; may be considered too old if over 24. ■ Large family equates with man's virility. Pregnancy is natural and desirable condition, so many do not seek prenatal care; extended family provides advice and support. ■ <i>Beliefs:</i> Hot food provides fetus with warmth; cold foods should be eaten during postpartum period. Pregnant women sleep on back to prevent harm to baby. ■ Frequent intercourse during pregnancy keeps vaginal canal lubricated and eases delivery. Walking in moonlight and viewing lunar eclipses leads to deformities. ■ Avoid becoming cold. Lifting hands over head causes cord to wrap around baby's neck. Pregnant mother may wear safety pin or metal object to prevent deformities. ■ Father not included in delivery. Mothers are very verbal during labor and do not practice breathing techniques. Mother puts legs together after delivery and wears cotton binder or girdle to prevent air from entering womb. ■ Postpartum period 40 days, warmth preferred, showers avoided. ■ Lactating mothers exposed to pesticides have decreased milk production and increased risk of breast cancer.
Navajo Native American	<ul style="list-style-type: none"> ■ <i>Postpartum beliefs:</i> Cutting baby's nails causes blindness and deafness. ■ Large families are favorable; many do not practice birth control. Having twins is not favorable and traditionally one must die; now, however, one may be placed for adoption. ■ <i>Beliefs:</i> Reluctant to deliver in hospital because hospitals are associated with sickness and death. Birthing necklaces worn during labor to ensure a safe birth. Mother holds onto woven sash when ready to push. Chanting may occur during delivery. Mother does not buy baby clothes before the baby's birth. ■ <i>Postpartum beliefs:</i> Placenta is buried after birth to protect baby from evil spirits. Baby is given special mixtures after birth to rid mouth of mucus.

CULTURAL GROUP	PRACTICES/BELIEFS
Vietnamese American	<ul style="list-style-type: none"> ■ High fertility rate because of long period of childbearing (up to age 44). ■ <i>Taboos/beliefs:</i> Dietary practices influenced by hot/cold, wind/tonic. Wind foods are cold, tonic foods are hot and sweet. Maintaining balance between hot and cold restores bodily balance. In first trimester, woman is considered weak and cold, so she eats hot foods. Second trimester is neutral state. In third trimester, woman is considered hot, so she eats cold foods. Woman remains physically active but avoids heavy lifting and raising hands above head (thought to pull on placenta). Sexual intercourse late in pregnancy thought to cause respiratory distress. Taboo to attend weddings and funerals while pregnant. ■ Early prenatal care is not the norm. Labor usually short, woman may prefer squatting or walking while in labor. ■ <i>Prenatal beliefs:</i> Head of mother and baby considered sacred, so should not be touched. Placenta is buried to protect baby's health. Ritual cleansing of mother without water. Mother avoids showers because of cold influence for a month after delivery. Family helps care for mother and child. Breastfeeding common practice, but colostrum may be discarded. May alternate breast with bottle because of hot/cold influence (hot foods benefit mother's health; cold foods promote healthy breast milk).

Source: Adapted from Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.



CRITICAL THINKING ACTIVITY #1

Consider each of the cultural perspectives just described. How might they influence your approach in planning your patient's care during the antepartal, intrapartal, and postpartal periods?



CRITICAL THINKING ACTIVITY #2

For each of the cultures described above, identify an intervention that incorporates your patient's cultural beliefs into her care.



CRITICAL THINKING ACTIVITY #3

What questions should you ask your patient to identify the cultural impact on her health?

Performing the Assessment

Your patient's prenatal workup includes a health history and physical assessment. A complete health history is essential to providing optimal care for the pregnant woman. If there is no recent complete health history available, you should perform one before proceeding with the specific pregnancy-related questions.

Introducing the Case Study



Mrs. Rosa Ramirez arrives at the Women's Clinic for confirmation of pregnancy. She is a 24-year-old Hispanic American who has missed two menstrual periods. During the interview, Mrs. Ramirez states that she has experienced fatigue, nausea, and occasional vomiting over the past 3 weeks and has noticed an increase in breast size as well as breast tenderness and tingling. She also mentions that she has urinary frequency. Mrs. Ramirez is 65 inches tall and weighs 125 pounds. She states that she usually weighs between 126 and 128 pounds.

After the health history comes the physical examination. Keep the key history findings in mind as you perform it. Taken together, the history and physical examination form a complete picture of your patient's prenatal health.

Health History

This section focuses specifically on the current pregnancy. The first prenatal visit involves collection of baseline information about your patient and her partner and identification of risk factors.

Key points to remember when obtaining a prenatal history:

- Focus on the current pregnancy and the presenting presumptive symptoms. Take a detailed obstetric/gynecologic history.
- Use the past medical history to identify anything that would affect or be affected by pregnancy.
- Pay special attention to the nutritional history.
- Pay special attention to the use of prescribed, over-the-counter (OTC), and illegal drugs; it may have a major impact on the developing fetus.
- Determine the patient's reaction to pregnancy—was it planned?
- Identify major supports—family, spouse, significant other.
- Assess for history or risk of physical abuse.
- After you have completed your questions, ask the patient if she has any problems or concerns that have not been covered, and give her an opportunity to discuss them.

If time is an issue and you are unable to perform a complete health history, perform a focused history on the sensory-neurological system.

Biographical Data

A careful review of the biographical data will be helpful in identifying actual or potential problems. Collecting this data also allows your patient to answer uncomplicated questions comfortably and sets the tone for the remainder of the interview.

First, clarify your patient's name, address, and date of birth. Geographic location may have a bearing on pregnancy outcome because women residing in the southern and western regions of the United States have a

higher incidence of preeclampsia. Women who will be age 35 or older at the time of delivery should be offered genetic counseling and testing. Determine what effect the patient's occupation may have on her pregnancy. Also identify the patient's religious preference and cultural/ethnic group and incorporate them into her care, if appropriate. Biographical data will also be helpful in identifying your patient's supports.

Current Health Status

The current health status includes verifying the patient's pregnancy, performing a symptom assessment, and calculating the estimated date of birth (EDB) or "due date."

Documenting Pregnancy

It is useful to document the patient's pregnancy before proceeding with the initial comprehensive prenatal evaluation. Both urine and serum pregnancy tests are based on levels of human chorionic gonadotropin (hCG), which are secreted into the mother's bloodstream and then excreted into the urine.

Urine pregnancy tests are 95 to 98 percent accurate and are sensitive within 7 days after implantation. The test is inexpensive and widely available without a prescription, so many women test themselves at home. The first voided specimen of the morning is best to use for testing because concentrated urine improves the pregnancy detection rate. Serum pregnancy tests do not indicate pregnancy until levels rise above baseline values—usually around 25 to 30 metric International Units (mIU)/mL. The hCG is detectable in serum as early as 7 to 9 days after ovulation, or just after implantation. During the first

Focused Obstetric History

If your patient's condition or time prohibits a detailed obstetric history, and the current pregnancy has been confirmed, just ask these key questions. They focus on significant past pregnancy events, current pregnancy status, identification of risk factors, and health promotion activities. The patient's response will help to direct your assessment.

- How many times have you been pregnant? What were the outcomes of these pregnancies? Did you experience any problems during these pregnancies?
- When was your LMP? Was this a normal period?
- When is this baby due? How was your due date determined?
- Since you became pregnant this time, have you:
 - Had any x-rays? (If yes, when and why?)
 - Been exposed to any viral illnesses, such as German measles (rubella)? (If yes, when?)
 - Experienced any other childhood illnesses? (If yes, describe.)
 - Had a fever? (If yes, when, and how high was your temperature?)
 - Have you used any medications: prescribed, OTC, home remedies/herbal preparations, vitamin or mineral supplements?
 - Have you had any bleeding or leakage of fluid from your vagina? When?
 - Have you had any prenatal care? If yes, where and when?
 - Do you have any medical conditions (HTN, diabetes, etc.)?
 - How much weight have you gained so far during this pregnancy?

Signs and Symptoms of Pregnancy

SIGNS AND SYMPTOMS

DESCRIPTION/TIME FRAME

Presumptive Signs

- Cessation of menses (amenorrhea).
- Nausea, vomiting.
- Frequent urination.
- Breast tenderness.
- Perception of fetal movement (quickening).
- Skin changes
- Fatigue

Probable Signs

- Abdominal enlargement.
- Piskacek's sign.
- Hegar's sign.
- Goodell's sign.
- Chadwick's sign.

- Braxton Hicks contractions.

- Pregnancy test.
- Ballottement.

Positive Signs

- Fetal heartbeat.

- Visualization of the fetus.

- Uterine lining does not slough off; women may experience spotting during implantation.
- From weeks 2 to 12; usually subsides after 12 weeks.
- Bladder irritability caused by enlarging uterus.
- Starts at 2 to 3 weeks; soreness; tingling.
- Occurs at 16 to 18 weeks; sensation of "fluttering" in abdomen perceived by mother-to-be.
- Increased pigmentation; striae gravidarum.
- Starts at 12 weeks.

- Palpated at 12 weeks.
- Palpated at 4 to 6 weeks; uterus asymmetrical with soft prominence on implantation side.
- 6 weeks; palpable softening of the lower uterine segment.
- Palpated at 8 weeks; softening of the cervix.
- Seen at 6 to 8 weeks; bluish hue on vulva, vagina, cervix from increased venous congestion.
- Painless, irregular, intermittent uterine contractions that typically start after the 4th month and last through remainder of pregnancy.
- Positive 7 to 10 days after conception.
- Occurs at 16 to 18 weeks, passive movement of the unengaged fetus.

- By ultrasound, fetal heart motion is noted by 4 to 8 weeks after conception; by Doppler, auscultated by 10 to 12 weeks; by fetal stethoscope, auscultated by 17 to 19 weeks.

- By ultrasound, at 5 to 6 weeks; by radiograph, by 16 weeks (but rarely used to diagnose pregnancy because of teratogenic effects).

3 to 4 weeks after implantation, the hCG level doubles every 2 days, then peaks at 60 to 70 days.

The diagnosis of pregnancy is based on the following indicators:

- Presumptive signs (experienced by the patient).
- Probable signs (observed by the examiner).
- Positive signs (attributed only to the presence of the fetus).

Symptom Assessment

The presenting symptoms usually relate to the presumptive signs of pregnancy. Your patient may present with multiple symptoms and vague complaints, all related to the pregnancy. Because pregnancy affects every system of the body, the review of systems (ROS) will address every presenting sign and symptom. Remember to perform a symptom analysis (PQRST) for all presenting symptoms.

Calculating the Estimated Date of Birth

Once the pregnancy has been confirmed, the EDB, also termed the *estimated date of confinement* (EDC), is cal-

culated. Establishing the baby's due date involves obtaining accurate information regarding the mother's menstrual history, including the last menstrual period (LMP).

To calculate the EDB, apply **Naegele's rule**: Add 7 days to the first day of the LMP, and then subtract 3 months from that date. Considerations in calculating the EDB include:

- Find out the first day of the LMP. Make sure that the patient is sure of the date because the EDC is based on the LMP. Conception usually occurs around 2 weeks after the LMP in a 28-day cycle.
- Review the patient's menstrual history, including frequency of menses, length of flow, normalcy of the LMP, and contraceptive use.
- Ultrasound studies may also be used to estimate the gestational age.

Past Health History

The purpose of the past health history is to uncover diseases or other risk factors that could affect the woman's

health or the fetus's well-being during pregnancy. Allergies to food, drugs, or environmental factors need to be noted because they can be exacerbated during pregnancy. Ask about exposure to toxins (e.g., radiation or chemicals) in the environment or at work, because this can affect fetal health. Is your patient on any medications? Identify all prescribed and OTC drugs (including alcohol and tobacco) your patient took before and during her pregnancy for their potential effects on the developing fetus.

Also take an obstetric history. Has your patient had previous pregnancies? If so, ask how many and if complications occurred during pregnancy or labor. Also ask about neonatal complications, such as birth defects, jaundice, infection, or death. Be sure to follow up on unclear or vague answers, and remember that sometimes rewording the question may help the patient find a relevant response.

Ask your patient if she has any of the diseases listed in the following paragraphs, which pose a particular risk to the expectant mother and/or fetus.

Diabetes

If your patient has diabetes, ask about the age of onset. If she is insulin dependent, ask what type and amount of insulin she takes. If her diabetes is diet controlled, ask about use of oral hypoglycemics. If she has had other pregnancies, did she have gestational diabetes? Uncontrolled diabetes in pregnancy can cause congenital anomalies, fetal overgrowth (macrosomia), intrauterine fetal death, delayed fetal lung maturation, and neonatal death. Oral hypoglycemics may cause fetal damage and are contraindicated. Women with a history of gestational diabetes are more likely to develop it again with subsequent pregnancies.

Hypertension

If your patient has chronic HTN, ask how she controls it. Does she take antihypertensive medication? If so, explain that these drugs may be contraindicated in pregnancy. HTN may result in decreased placental perfusion and intrauterine fetal growth restriction. PIH may recur.

Cardiac Disease

Patients with mitral valve prolapse (MVP) may need prophylactic antibiotics during labor to prevent streptococcal infections and subsequent bacterial endocarditis and valve disease.

Liver Disease

If your patient has hepatitis B, the infant may require treatment (hepatitis B immunoglobulin and hepatitis vaccine) after birth.

Cancer

Patients with cervical cancer who were treated with cone biopsy (cone-shaped section of the cervix is removed for examination) are at risk for preterm labor.

Infectious Diseases

Rubella, mononucleosis, and other viral infections in the first trimester can cause fetal abnormalities.

Pulmonary Disease

Medications and inhalers used for asthma may be harmful to the fetus or may affect anesthesia used during labor. Inhalants and general anesthesia may be contraindicated for patients with asthma.

Gastrointestinal Disease

Ask about previous abdominal surgery and note the type of scarring; this may influence the type of delivery. Colitis and other bowel problems may be exacerbated in pregnancy.

Other Medical Problems

Varicosities and renal, gallbladder, genitourinary, autoimmune, neurological, and psychiatric conditions may be exacerbated in pregnancy.

Gynecologic Diseases and Sexually Transmitted Diseases

Vaginitis should be identified and treated early to prevent intrauterine complications. Untreated sexually transmitted diseases (STDs), such as genital herpes and gonorrhea, can be transmitted to the fetus during passage through the birth canal. The transmission rate for babies born to human immunodeficiency virus (HIV)-infected mothers is 20 to 35 percent.

Family History

The purpose of the family history is to identify potential physical and emotional complications of pregnancy and familial patterns of health or illness. Ask specific questions to pinpoint inherited diseases. Some questions to ask include:

- “Was anyone in your family diagnosed with heart disease before age 50?” (Cardiovascular disease or heart defects may be inherited.)
- “Does anyone in your family have lung disease, tuberculosis, or asthma?” (Pulmonary disorders may be familial; tuberculosis is contagious.)
- “Do any family members have diabetes?” (Endocrine problems are genetically linked.)
- “Does anyone in your family have cancer?” (There is a genetic component with certain types of cancer.)
- “Is there a history of birth defects, inherited genetic disorders, blood disorders, or mental retardation in your family?” (There is a genetic risk for Down syndrome, spina bifida, brain defects, anencephaly, heart defects, muscular dystrophy, cystic fibrosis, hemophilia, thalassemia, and other disorders.)
- “Did your mother or sisters have complications during pregnancy or labor?” (Daughters and sisters of preeclamptic women have a higher tendency toward preeclampsia.)

Be sure to consider your patient's race/ethnic background when taking a family history. For example, children of African American women are at risk for sickle cell disease. Identification of sickle cell disease will prevent a crisis; testing the mother-to-be is appropriate if status is uncertain.

Review of Systems

Normal changes that occur during pregnancy have an impact on every body system. The ROS will help you to identify normal physiological changes as well as alert you to abnormal findings.

Psychosocial Profile

The psychosocial profile is an important component of the assessment because it lays the groundwork for a trusting nurse-patient relationship. The profile provides an opportunity to explore the patient's reactions to the pregnancy and to identify lifestyle patterns that may pose a threat to her or her baby's well-being.

Start by asking your patient about her health practices and beliefs. Is she proactive (getting regular preventive healthcare) or reactive (seeking healthcare only when ill)? Also inquire about self-care, such as breast self-examinations (BSEs). Determine the patient's acceptance

Review of Systems

AREA/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- How have you been feeling?

- Feelings of fatigue and ambivalence are normal during the first trimester.
- During the second trimester, mothers-to-be are introspective and energetic.
- The third trimester is characterized by restlessness, mood swings, and interest in preparing for the baby.



ALERT

Denial of the pregnancy, withdrawal, depression, or psychosis signal psychological problems that warrant referral.

Body Weight

- What is your normal weight (before pregnancy)?
- Have you lost or gained weight since a year ago? How much?

- Optimal weight gain during pregnancy depends on patient's height and normal weight. Recommended weight gain in pregnancy is:
 - Underweight patient: 28 to 40 lb.
 - Normal weight patient: 25 to 35 lb.
 - Overweight patient: 15 to 25 lb.
 - Twin gestation: 35 to 45 lb.



ALERT

Low pregnancy weight and inadequate weight gain during pregnancy contribute to fetal growth restriction and low birth weight.

Integumentary

- Have you noticed any changes in your skin, hair, or nails?

- Hormonal changes cause hyperpigmentation of skin (chloasma, linea nigra), thin nails, oily hair.

HEENT

Eyes

- Do you have any vision problems?

- Excessive tearing may be associated with allergies; blurred vision or spots before the eyes may indicate preeclampsia.

(continued)

Review of Systems (continued)

AREA/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p><i>Ears</i></p> <ul style="list-style-type: none"> ■ Do you have any hearing problems? 	<ul style="list-style-type: none"> ■ Decreased hearing, earaches, or sense of fullness in ears occurs because tympanic membranes swell as a result of increased vascularity.
<p><i>Nose</i></p> <ul style="list-style-type: none"> ■ Do you have nasal stuffiness? ■ Nosebleeds? 	<ul style="list-style-type: none"> ■ Increased vascularity from increased estrogen causes nasal edema.
<p><i>Neck</i></p> <ul style="list-style-type: none"> ■ Have you noticed any masses in your neck? 	<ul style="list-style-type: none"> ■ Slight thyroid enlargement is normal; marked enlargement may indicate hyperthyroidism.
<p><i>Mouth/Throat</i></p> <ul style="list-style-type: none"> ■ Do you have any trouble with your throat? ■ Since your LMP, have you had a fever or chills without a cold? ■ Do you have a cough that doesn't go away or frequent chest infections? <ul style="list-style-type: none"> ■ Do your gums bleed? When was your last dental exam? <ul style="list-style-type: none"> ■ Do you have increased saliva? 	<ul style="list-style-type: none"> ■ Prolonged nasal congestion with sore throat, fever, and chills may be an upper respiratory infection. ■ Fetal exposure to viral illnesses is associated with fetal growth restriction, developmental delays, hearing impairment, and mental retardation. ■ Persistent cough and frequent chest infections may indicate pneumonia or tuberculosis. ■ Gum hypertrophy is common; bleeding during tooth brushing may be associated with gum disease and warrants further dental evaluation. ■ Ptyalism (excessive saliva) often occurs within 2 to 3 weeks after the first missed period and is not associated with pathology.
<p><i>Respiratory</i></p> <ul style="list-style-type: none"> ■ Do you have shortness of breath? Dyspnea? Other breathing problems? 	<ul style="list-style-type: none"> ■ Thoracic breathing, slight hyperventilation, and shortness of breath occur in late pregnancy. ■ Dyspnea may be associated with respiratory distress; dyspnea with markedly decreased activity tolerance may indicate cardiovascular disease.
<p><i>Cardiovascular</i></p> <ul style="list-style-type: none"> ■ Do you have a history of cardiovascular disease? Palpitations? Dizziness? 	<ul style="list-style-type: none"> ■ Pregnant women with pre-existing cardiovascular disease, such as MVP, can decompensate as a result of increased workload of the heart. Be alert for cardiovascular changes associated with PIH and eclampsia. ■ Supine hypotension can occur from vena caval compression.
<ul style="list-style-type: none"> ■ Do your ankles swell? 	<div data-bbox="699 1451 1310 1661" style="border: 1px solid red; padding: 5px; margin-bottom: 10px;">  <p>ALERT</p> <p>Lying supine compresses vena cava and aorta, decreasing cardiac output. Advise patient to lie on left side to increase renal perfusion and output and reduce edema.</p> </div> <ul style="list-style-type: none"> ■ Dependent edema and varicose veins in legs frequently occur in pregnancy, but may also be associated with PIH and eclampsia.
<p><i>Breasts</i></p> <ul style="list-style-type: none"> ■ Have you noticed pain, lumps, or fluid leaking from your breasts? 	<ul style="list-style-type: none"> ■ Fullness, increased sensitivity, tingling, and heaviness are common early in pregnancy; however, lumps and pain may also indicate breast disease. ■ Colostrum secretion from breasts is normal during pregnancy and varies in color.

AREA/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Gastrointestinal <ul style="list-style-type: none"> ■ Do you have nausea and vomiting that do not go away? ■ Are you more thirsty than usual? 	<ul style="list-style-type: none"> ■ Excessive nausea and vomiting may be associated with hyperemesis gravidarum; sudden, excessive weight gain could indicate a multifetal gestation or fluid retention associated with PIH. ■ Abdominal pain or cramping may be related to round ligament pain or may signal impending miscarriage. ■ Hydration must be maintained because the patient may be at risk for hypovolemia, cholecystitis, or cholelithiasis.
<ul style="list-style-type: none"> ■ Do you ever notice black or bloody stools? ■ Do you have diarrhea or trouble passing stools? 	<ul style="list-style-type: none"> ■ Blood in the stools or a change in bowel habits may indicate constipation or hemorrhoids.
Genitourinary <ul style="list-style-type: none"> ■ Do you ever have burning or pain when you urinate? ■ Do you have to urinate more often than normal? 	<ul style="list-style-type: none"> ■ Urinary urgency and frequency are common during pregnancy and are not cause for concern unless accompanied by pain or burning, which may signal a UTI. ■ During pregnancy, women may have asymptomatic bacteriuria. UTIs must be promptly diagnosed and treated because untreated UTIs predispose patient to complications such as preterm labor, pyelonephritis, and sepsis.
Reproductive <ul style="list-style-type: none"> ■ When was your LMP? ■ Do you have increased vaginal discharge? <ul style="list-style-type: none"> ■ Have you experienced any vaginal bleeding, leakage of fluid, or unusual vaginal discharge? 	<ul style="list-style-type: none"> ■ Needed to determine EDC. ■ Increased white vaginal discharge (leukorrhea) is normal during pregnancy. Discharge accompanied by a foul odor, itching, or burning may indicate infection. ■ Vaginal bleeding, fluid leakage, or vaginal discharge may indicate placenta previa, rupture of membranes, or vaginal infection. Untreated vaginal infections predispose patient to preterm labor or fetal infections.
Musculoskeletal <ul style="list-style-type: none"> ■ Do you have leg cramps? ■ Do you have back pain? 	<ul style="list-style-type: none"> ■ Leg cramps may indicate calcium deficiency. ■ Curvature of the lumbar spine may be accentuated during pregnancy, resulting in backache. ■ Severe back pain may be associated with disc disease.
Neurologic <ul style="list-style-type: none"> ■ Do you have a history of depression, difficulty sleeping, loss of appetite? 	<ul style="list-style-type: none"> ■ Emotional lability can occur during pregnancy; however, these symptoms also may indicate psychological disorders.
	<div style="border: 1px solid red; padding: 5px;"> <p style="text-align: center;"> ALERT</p> <p style="text-align: center;">• Patients with a history of psychological disorders must be continually monitored for signs and symptoms and referred when appropriate.</p> </div>

(continued)

Review of Systems (continued)

AREA/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Have you experienced light-headedness, dizziness, or fainting? ■ Do you have wrist pain, numbness, or tingling? 	<ul style="list-style-type: none"> ■ Fainting may indicate anemia. ■ Wrist pain, especially in the dominant hand, may indicate carpal tunnel syndrome.
<p>Endocrine</p> <ul style="list-style-type: none"> ■ Do you have increased fatigue or heat intolerance? ■ Do you have a history of diabetes or gestational diabetes? 	<ul style="list-style-type: none"> ■ Common symptoms associated with increase in BMR and hormonal changes. ■ Positive history of diabetes calls for close monitoring.
<p>Immunological/Hematological</p> <ul style="list-style-type: none"> ■ Do you have a history of anemia? 	<ul style="list-style-type: none"> ■ Physiological anemia may occur during pregnancy. As a result, pre-existing anemia may worsen.
<ul style="list-style-type: none"> ■ History of thrombophlebitis? 	<ul style="list-style-type: none"> ■ Increase of clotting factors increases risk of thrombus formation.

of the prenatal care plan and understanding of recommended actions, including the need for regular prenatal visits.

Next, ask about your patient's typical day. What constitutes a usual day's activities? How has pregnancy affected her daily lifestyle pattern?

Inquire about nutrition and weight loss or gain. Weight loss often occurs in the first trimester as a result of morning sickness. Failure to gain enough weight in the second and third trimesters may be associated with fetal growth restriction. Rapid, sudden weight gain may indicate PIH.

Ask if the patient takes vitamins and supplements. Supplementation may be necessary to ensure adequate intake and should be started before conception. The Centers for Disease Control and Prevention (CDC) recommends 0.4 mg of folic acid per day for the prevention of neural tube defects; it is best to begin 1 month before conception and continue through the first 3 months of pregnancy. Pregnant women should also receive prenatal vitamins with 30 mg of ferrous iron daily.

Question the patient about the amount and type of exercise she does. Daily exercise is highly recommended as long as it is well tolerated by the patient. Women in good physical condition tend to have easier labors than those who are not physically fit.

Ask about pets. If the patient has a cat, warn her to wear gloves when changing litter boxes because of the risk of toxoplasmosis infection. Suggest that other family members change the cat litter, and encourage the patient to wash her hands well after petting the cat.

Ask about sleep and rest. How much uninterrupted sleep does the patient get every night? Does she take naps? Does she suffer from insomnia? Adequate rest is important in ensuring optimal health for the mother and

fetus. Difficulty sleeping during the third trimester is common and may be treated with massage, tub baths, or warm milk.

Inquire about personal habits. Does the patient currently smoke? Does she take recreational drugs, such as cocaine, speed, or marijuana? Does she drink alcohol? Cigarette smoking during pregnancy is associated with increased perinatal mortality, preterm delivery, premature rupture of membranes, placental abruption, stillbirth, and bleeding during pregnancy. Cocaine use is associated with a higher rate of spontaneous abortions and placental abruption. Alcohol use during pregnancy is associated with low birth weight, stillbirth, and fetal alcohol syndrome, the leading cause of mental retardation. There is no known safety level for alcohol use during pregnancy.

Find out where the patient works and what she does for a living. Does it involve heavy lifting, toxic exposure, or anything else that might harm the fetus? Can she take rest periods? Physiological changes during pregnancy place a tremendous amount of stress on the body, and frequent rest periods are encouraged.

Ask about environmental hazards such as second-hand smoke and paternal exposure to toxins and industrial pollutants. There is an increased incidence of preeclampsia in women residing in southern and western states. Smoke and industrial chemicals are hazardous to the fetus. Paternal exposure to workplace toxins has been associated with an increased risk of spontaneous abortion. Also ask about living conditions. Ascertain whether the patient's home is adequate for the expanding family, and work with her to improve the situation, if needed. Refer to community agencies, if appropriate.

Ask about cultural and religious influences in the patient's life. How do they affect her healthcare practices, especially during pregnancy and delivery?

Ask about the patient's family relationships. Does she have a spouse or partner? Does she have any other children? How old are they? Education and preparation for the baby can offset potential problems with sibling rivalry. Encourage sibling classes. Also ask about support systems outside the immediate family. Does she have relatives or friends to rely on during pregnancy and the postpartum period?

Ask the patient about stress and coping. How has the pregnancy affected her emotional well-being? Many women experience mood swings during pregnancy, but signs of clinical depression must be evaluated and referred.



Case Study Findings

Reviewing Mrs. Ramirez's

Biographical data:

- 24 years old, married, first pregnancy.
- College graduate, full-time social worker.
- Born and reared in the United States, Hispanic descent, Catholic religion.
- Health insurance through own and husband's work plan.
- Referral: Routine prenatal care.
- Source: Self, seems reliable.

Current health status and presenting symptoms:

- Fatigue.
- Nausea and vomiting for 3 weeks.
- Breast enlargement, tingling, and tenderness.
- Urinary frequency.
- Amenorrhea.

Past health history:

- Usual childhood illnesses.
- No surgeries, hospitalizations, or allergies.
- Immunizations current.
- Takes one multivitamin with iron daily; no other medications.
- Physical examinations yearly since age 18; normal Pap smears.
- Used oral contraceptives for 5 years without problems; discontinued them 8 months ago to attempt pregnancy.
- No previous pregnancies.

Family history:

- Grandmother, age 64, takes medication to control high BP.
- All other relatives alive and well; two older sisters married with children and experienced no complications during pregnancies; one younger sister is single.

Review of systems:

- *General Health Survey:* Very tired for past couple of months, needs to take a nap every

day when she gets home from work. Body weight 125 lb, on the low end of her normal reported range of 126 to 128 lb. Has lost 2 lb since becoming pregnant.

- *Integumentary:* Noticed nipples getting darker and line on abdomen. Hair very oily.
- *HEENT:* No changes in head/neck. Has nasal stuffiness but no colds, allergies, or flu. Gums bleed with brushing; last dental exam 6 months ago. No eye changes.
- *Respiratory:* No changes.
- *Cardiovascular:* No history of MVP, palpitations, or edema.
- *Breasts:* Enlargement, tenderness, and tingling sensation.
- *Gastrointestinal:* Nausea and vomiting in the morning, usually subsides by noon; says certain foods "just turn my stomach"; reports constipation.
- *Genitourinary:* Reports frequency of urination, no burning.
- *Reproductive:* LMP 4/10/06.
- *Musculoskeletal:* No changes.
- *Neurological:* Occasional "lightheadedness"; no history of depression; excited at possibility of being pregnant.
- *Endocrine:* No history of gestational diabetes; reports heat intolerance.
- *Immunologic/Hematologic:* No history of anemia.

Psychosocial profile:

- Patient is proactive regarding healthcare. Has yearly physical examination and Pap smear. Performs monthly BSE. Believes people are responsible for maintaining their own health.
- Typical day consists of arising at 6:30 A.M. and eating breakfast with husband. (Lately unable to enjoy usual breakfast of cold cereal and juice because of nausea.) Drives 4 miles to office 5 days a week and works from 8 A.M. to 5 P.M. at desk part of time and interviews patients in a clinic located on the ground floor of her office building for rest of time. Loves her work. Evenings she and husband prepare and eat supper, clean up dishes, take a walk in the neighborhood, then relax by watching television, playing cards, or reading. Patient showers and goes to bed around 10 P.M.
- Usually eats a healthy diet, but lately has been skipping breakfast because of nausea. As day progresses, is gradually able to eat bland food, such as baked chicken, potatoes, and steamed vegetables.
- Walks daily for 45 minutes with husband.

(continued on page 832)



Case Study Findings (continued)

- Has no pets. Recreation includes daily walk with husband, dinner at restaurant or movie once a week, family get-togethers. Hobbies include reading and needlework.
- Sleeps around 8½ hours normally, but lately gets very sleepy at work, especially in mid-afternoon. Combats sleepiness with a walk outside her office and an afternoon snack of fruit and yogurt or crackers. Takes a short nap after work. No problems falling asleep.
- Does not smoke, use drugs, or drink alcoholic beverages.
- No chemical exposure at work.
- Lives in a one-story, two-bedroom single-family home in suburbs; has hot water, heating, and air conditioning; negative for radon gas. Will convert guest room into baby's room. No recent travel outside local community.
- Catholic religion, Hispanic nationality (third generation). No religious or cultural influences that affect healthcare, pregnancy, or delivery.
- Supportive, caring husband (nonsmoker) who is computer analyst for national company; married 5 years; attempted pregnancy for 6 months.
- Mother, mother-in-law, and three sisters live in same community and will assist with infant care. Patient plans to breastfeed and stay at home for 6 weeks after birth and then return to job.
- Copes with stress by walking, talking with husband and family members, doing needlework.



CRITICAL THINKING ACTIVITY #7

Based on the subjective information you have obtained, identify positive lifestyle factors that will enhance Mrs. Ramirez's pregnancy and birthing experience.



Case Study Evaluation

Before performing the physical examination, document what you have learned about Mrs. Ramirez so far.

Physical Assessment

Now that you have completed the subjective data collection, begin to collect objective data by means of the physical examination. Keep the history findings in mind. These findings, along with those from the physical examination, will complete the assessment picture. Then you can analyze the data, formulate nursing diagnoses, and develop a plan of care.

Approach

Begin the physical examination by asking the patient for a clean-catch midstream urine specimen. Then proceed to inspection, palpation, percussion, and finally, auscultation. Remember that normal findings may vary somewhat from one pregnant patient to another; thus, a systematic approach is essential. Explain to your patient that after the examination is complete, she will go to the laboratory for initial prenatal blood tests.



CRITICAL THINKING ACTIVITY #4

From what you have learned about Mrs. Ramirez, calculate her EDB using Naegele's rule.



CRITICAL THINKING ACTIVITY #5

Identify the presumptive signs of pregnancy Mrs. Ramirez has experienced.



CRITICAL THINKING ACTIVITY #6

List strategies that may help combat Mrs. Ramirez's morning sickness.



TOOLBOX

The tools for the obstetric assessment include a stethoscope, light for the pelvic examination, tape measure, fetoscope or fetal Doppler, and equipment for the pelvic exam: speculum, gloves, lubricant, glass slides, potassium hydroxide (KOH), normal saline, and cytology fixative. Also, sharpen your senses because you will use all of them during the obstetric examination.

Performing a General Survey

The first part of the physical assessment focuses on the general survey, including a measurement of height and weight (Fig. 23.1). At the initial visit, it is important to



FIGURE 23.1. Establishing baseline weight.

establish a baseline height, weight, pulse, respirations, and BP; on subsequent prenatal visits, the weight, pulse, respirations, and BP (taken in a sitting position) are also recorded.

Throughout each prenatal examination, observe your patient's behavior. During the first trimester, she is likely to appear tired and even ambivalent about her pregnancy. The second trimester is usually characterized by

excitement, enthusiasm, and energy. The third trimester is marked by periods of restlessness, self-doubts, and concerns about the upcoming birth experience.



HELPFUL HINT

Consider your patient's position and anxiety level when assessing BP. BP is highest when sitting and lowest when supine. So always take BP in the same arm and in the same position.

Performing a Head-to-Toe Physical Assessment

Because physiological changes occurring during pregnancy affect every other body system, physical signs may be seen in any system. You will need to perform a full head-to-toe exam, including a pelvic exam. Pelvic cultures obtained with this exam include a Pap smear and gonorrhea and chlamydial cultures.

For the next component of the assessment, the patient should be comfortably seated on the examining table. She should be wearing a gown and adequately draped to ensure privacy. Explain to her what you will be doing and why and what she may expect to feel just before and during each part of the exam. As you progress through your initial physical assessment, remember that pregnancy affects every other body system and that you will need to be able to differentiate normal changes from abnormal changes. (See *Performing a Head-to-Toe Physical Examination*.)

Prenatal Laboratory Tests

Urine Tests

One of the first laboratory tests to obtain (preferably before beginning the physical exam) is the clean-catch midstream urine specimen, which will be tested for glucose (to assess for diabetes), protein (to assess for PIH), and nitrites and leukocytes (to assess for infection).

Blood Tests

Initial prenatal blood tests include complete blood count, blood type and screen, Rh status, rubella titer, serologic test for syphilis, and hepatitis B surface antigen. Patients of African ancestry are also referred for a sickle cell anemia screen. In addition, patients who are at high risk for infection with HIV should be screened for this disease.

Between 16 and 18 weeks' gestation, a multiple marker or "triple screen" is usually obtained. This

blood test measures the maternal serum level of alpha-fetoprotein (MSAFP), hCG, and unconjugated estriol (uE3). High levels of alpha-fetoprotein are associated with neural tube defects, low triple screen values with Down syndrome, and other chromosomal abnormalities.

MSAFP screening can detect approximately 80 to 85 percent of all open neural tube defects (NTDs) and open abdominal wall defects early in pregnancy. AFP is produced by the fetal liver, gastrointestinal tract, and embryonic yolk sac and readily crosses the placenta and fetal membranes into the mother's bloodstream, resulting in a rise in the maternal serum. The AFP level can be measured by taking a sample of the mother's blood between 15 and 21 weeks' gestation; measurement at 17 weeks is ideal.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

1. Integumentary

SYSTEM/NORMAL VARIATIONS

Skin, Hair, Nails

Inspect

- Linea nigra, striae gravidarum, chloasma, spider nevi, palmar erythema.
- Increased growth, softening, thinning of hair and nails.



Chloasma



Palmar erythema

ABNORMAL FINDINGS/RATIONALE

- *Pale skin:* Anemia.



ALERT

Your patient is anemic if hemoglobin drops below 10 g/dL.

2. HEENT

SYSTEM/NORMAL VARIATIONS

Head and Neck

Palpate

- Palpable smooth, nontender, small cervical chain lymph nodes. Slight thyroid enlargement.

Ear

Inspect

- Tympanic membranes clear, landmarks visible.

Nose

Inspect

- Mucosal swelling and redness, epistaxis (nose-bleeds) common because of increased estrogen.

Mouth, Throat

Inspect

- *Gums:* Gingival hypertrophy and epulis usually regress spontaneously after delivery.
- *Throat:* Pink, no redness or exudates.



Gingival hypertrophy



Epulis

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

2. HEENT (continued)

ABNORMAL FINDINGS/RATIONALE

- *Hard, tender, fixed or prominent cervical nodes:* Cancer.
- *Marked thyroid enlargement:* Hyperthyroidism.
- *Tympanic membranes red and bulging with pus:* Infection.
- *Purulent discharge:* Upper respiratory infection.
- *Bleeding gums:* Gingivitis.
- *Redness:* Exudate present.
- *Enlarged tonsils:* Infection.

3. Respiratory

SYSTEM/NORMAL VARIATIONS

Inspect/Palpate/Percuss/Auscultate

- Increased anteroposterior chest diameter, thoracic breathing, slight hyperventilation, shortness of breath in late pregnancy, lung sounds clear bilaterally.

ABNORMAL FINDINGS/RATIONALE

- *Dyspnea, crackles, rhonchi, wheezes, rubs, absence of breath sounds, unequal breath sounds, respiratory distress:* May indicate pulmonary complications, such as pulmonary edema or acute respiratory distress syndrome.

4. Cardiovascular

SYSTEM/NORMAL VARIATIONS

Heart

Palpate/Auscultate

- Point of maximal impulse (PMI) may be displaced upward and laterally in the latter stages of pregnancy.
- Normal sinus rhythm.
- Soft systolic murmur caused by increased blood volume.

ABNORMAL FINDINGS/RATIONALE

- *Enlarged PMI:* May be associated with HTN.
- *Irregular rhythm:* Cardiac disease.
- *Dyspnea, palpitations, markedly decreased activity tolerance:* Cardiovascular disease.
- *Midsystolic click and late systolic murmur:* MVP.

5. Breasts

SYSTEM/NORMAL VARIATIONS

Inspect/Palpate

- Venous congestion with prominence of Montgomery's tubercles. Increased size and nodularity; increased sensitivity; colostrum secretion in the third trimester.
- Hyperpigmentation of nipples and areolar tissue.

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

5. Breasts (continued)

ABNORMAL FINDINGS/RATIONALE

- Nipple inversion may be problematic for breast-feeding women.
- *Localized redness, pain, and warmth:* Mastitis.
- *Bloody nipple discharge and skin retraction:* Cancer.

6. Gastrointestinal

SYSTEM/NORMAL VARIATIONS

Abdomen

Inspect/Auscultate/Palpate

- Cesarean scars and location; obtain previous pregnancy records to confirm type and location of uterine incision.
- Note linea nigra, striae.
- Abdominal enlargement caused by fetus; in later pregnancy, uterine shape may suggest fetal presentation and position. Palpable uterus at 10 to 12 weeks.
- Fetal movement noticed by mother at 18 to 20 weeks (earlier for multipara).
- Uterine contractions may be present; intensity is described as mild, moderate, or firm to palpation.



HELPFUL HINT

How long are the patient's contractions? Time them from beginning to end of same contraction. How frequent are they? Time them from beginning of one contraction to beginning of next

Measuring fundal height.

Place zero point of tape measure on symphysis pubis and measure to top of fundus.

- Fundal measurement should approximately equal number of weeks gestation; measurements may vary by 2 cm; measurements by different examiners should be approximately the same.




Fundal height measurement



Linea nigra



Striae



Fundal height assessment

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

6. Gastrointestinal (continued)

SYSTEM/NORMAL VARIATIONS

Fetal Heart Tones

FHTs are best auscultated through the back of the fetus.

A fetal Doppler can be used after 10 to 12 weeks' gestation; a fetoscope may be used after 18 weeks' gestation.



Auscultating FHT with fetoscope




Auscultating FHT with Doppler

- *Fetal heart rate (FHR) range:* 120 to 160 beats per minute. In third trimester, FHR accelerates with fetal movement.

Fetal Position

- Use Leopold's maneuvers to palpate the fundus, lateral aspects of the abdomen, and lower pelvic area. Leopold's maneuvers assist in determining: fetal lie (body of fetus in relation to mother's back), presentation (fetal presenting part into the maternal pelvis), size, and position.
- A longitudinal lie (fetal spine axis is parallel to mother's spine axis) is expected. Fetal presentation may be cephalic, breech, or shoulder. Fetal size is estimated by measuring fundal height and by palpation.



FHTs. Intensity varies according to fetal position. RSA, right sacrum anterior; LSA, left sacrum anterior; ROP, right occipitoposterior; LOP, left occipitoposterior; RMA, right mentum anterior; LMA, left mentum anterior; ROA, right occipitoanterior; LOA, left occipitoanterior.

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

6. Gastrointestinal (continued)

SYSTEM/NORMAL VARIATIONS

First Maneuver

Face the patient's head.

Place your hands on the fundal area and palpate.

- You should palpate a soft, irregular mass in the upper quadrant of the mother's abdomen. The soft mass is the fetal buttocks; the round, hard part is the fetal head.



First Leopold's maneuver

Second Maneuver

- Move your hands down to the lateral sides of the mother's abdomen and palpate.
- On one side of the abdomen you will palpate round, irregular nodules—the fists and feet of the fetus. Expect to feel kicking and movement. The other side of the mother's abdomen feels smooth—this is the fetus's back.



Second Leopold's maneuver

Third Maneuver

- Move your hands down to the mother's lower pelvic area and palpate the area just above the symphysis pubis to determine the presenting part of the fetus. Grasp the presenting part with your thumb and third finger.
- If the presenting part is the head, it will be round, firm, and ballotable; if it is the buttocks, it will be soft and irregular.



Third Leopold's maneuver

Fourth Maneuver

- Place your hands on her upper and lower abdomen and try to move your hands toward each other while applying downward pressure.
- If your hands move together easily, the fetal head is not descended into the mother's pelvic inlet. If your hands do not move together and stop because of resistance, the fetal head is engaged into the mother's pelvic inlet.



Fourth Leopold's maneuver



Fetal presentation/position

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

6. Gastrointestinal (continued)

ABNORMAL FINDINGS/RATIONALE

- Note any scars that may indicate previous abdominal surgery and influence type of delivery.
- *Abnormal palpable masses*: Uterine fibroids or hepatosplenomegaly.
- *No fetal movement felt*: Wrong EDC or fetal demise.
- *Regular contractions before 37 completed weeks of gestation*: Preterm labor.
- Measurements greater than 4 cm from the estimated gestational age warrant further evaluation.
- *Greater than expected*: Multiple gestation; polyhydramnios; fetal anomalies; macrosomia.
- *Smaller than expected*: Fetal growth restriction.
- *Diastasis recti abdominis*: Separation of abdominal muscles from pregnancy.
- *Inability to auscultate FHT with a fetal Doppler at 12 weeks*: Retroverted uterus; uncertain dates; fetal demise; false pregnancy.
- *Fetal heart rate decelerations*: Poor placental perfusion.
- *Oblique or transverse lie*: Breech presentation.



HELPFUL HINT

If you hear fetal heart sounds above the umbilicus, it is a breech presentation; below the umbilicus, a vertex presentation.

7. Extremities

SYSTEM/NORMAL VARIATIONS

Inspect/Palpate

- In third trimester, dependent edema is normal; varicose veins may also appear.

ABNORMAL FINDINGS/RATIONALE

- *Calf pain, positive Homans' sign, generalized edema, diminished pedal pulses*: DVT.



Homans' assessment

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

8. Genitourinary

SYSTEM/NORMAL VARIATIONS

Inspect/Palpate

External Genitalia

- Labial and clitoral enlargement; parous relaxation of introitus; scars from episiotomy or perineal lacerations (multiparous women).

Bartholin's and Skene's Glands

- No discomfort or discharge.

Vaginal Orifice

- Small amount of whitish discharge (leukorrhea).

Cervix

- Smooth, pink, or bluish, long, thick, closed; 2.3 to 3 cm long. Softening of lower uterine segment (Hegar's sign) should be present. Bluish color (Chadwick's sign) indicates increased blood flow to pelvic area and is probable sign of pregnancy.



Chadwick's sign



Circular cervical opening: nulliparous



Slitlike cervical opening: multiparous

Uterus

- Uterus is size of orange at 10 weeks; grapefruit at 12 weeks.
- No masses in left and right adnexa; some discomfort caused by stretching of round ligaments.

ABNORMAL FINDINGS/RATIONALE

- *Labial varicosities*: Venous congestion.
- *Discharge and tenderness*: infection.
- *Thick, purulent vaginal discharge*: Gonorrheal infection.
- *Thick, white, cheesy discharge*: Yeast infection.
- *Gray-white discharge, sweet smell, positive clue cells*: Bacterial vaginosis infection.
- *Effaced, opened cervix*: Preterm labor or incompetent cervix if not a term gestation.
- *Uterine size not consistent with dates*: Wrong dates, fibroids, multiple gestation.
- *Palpable masses*: Ectopic pregnancy.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

9. Musculoskeletal

SYSTEM/NORMAL VARIATIONS

Inspect

- Accentuated lumbar curve (lordosis).
- Wider base of support.



Accentuated lumbar curve

10. Neurologic

SYSTEM/NORMAL VARIATIONS

Inspect

- +1 to 2 DTR.
- Negative Phalen's or Tinel's sign.



DTR assessment

ABNORMAL FINDINGS/RATIONALE

- *Hyperreflexia, clonus*: Preeclampsia, eclampsia.
- *Positive Phalen's or Tinel's sign*: Carpal tunnel syndrome.



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Ramirez have concluded. The following are your findings:

- **General Health Survey**
 - AAO × 3, affect pleasant, excited about pregnancy.
 - Neatly dressed and well groomed.
 - No signs of distress.
 - Urine pregnancy test positive today.
- **Vital Signs**
 - BP, 130/80.
 - Pulse, 80 and regular.
 - Respirations, 22/min.
 - Temperature, 98.8 F
 - Weight, 125 lb.
- **Integumentary**
 - Skin pink, warm, dry, good turgor.
 - Mucous membranes pink, moist.
- **HEENT**
 - No palpable cervical chain lymph nodes.
 - Slight thyroid enlargement, no nodularity or tenderness.
- **Respiratory**
 - Lung sounds clear to auscultation bilaterally.
- **Cardiovascular**
 - Normal sinus rhythm.
- **Breasts**
 - Tenderness.
 - Venous congestion.
 - Prominence of Montgomery's tubercles.
 - Darkening of nipples and areolae.
 - No nipple discharge.
- **Gastrointestinal**
 - Linea nigra present.
 - No masses, no enlargement, no fetal heart tones (FHTs) auscultated with Doppler.
- **Musculoskeletal**
 - Extremities normal color
 - No edema, no varicosities, no tenderness, no increased warmth.
 - Deep tendon reflexes (DTRs) 1+.
- **Genitourinary**
 - *External genitalia*: Normal hair distribution, no varicosities, slight labial enlargement.
 - *Bartholin's and Skene's glands*: Nontender, no discharge.
 - *Cervix*: Long (3.0 cm), thick, and closed; Chadwick's sign; slight leukorrhea; positive Hegar's sign.
 - *Uterus*: Approximately orange-sized, no adnexal masses.
 - *Rectal mucosa*: Pink, intact, no hemorrhoids.



Case Study Analysis and Plan

Now that you have completed a thorough assessment of Mrs. Ramirez, document the key history and physical exam findings that will help you formulate your nursing diagnoses.



CRITICAL THINKING ACTIVITY # 8

What are Mrs. Ramirez's strengths? Are there any areas of concern?



CRITICAL THINKING ACTIVITY # 9

As with most pregnant women, Mrs. Ramirez's readiness to learn is high. What information about her pregnancy would be important to share with her at this time?



CRITICAL THINKING ACTIVITY # 10

How often do you expect Mrs. Ramirez to be seen in the prenatal clinic? Which components of the exam will be repeated at each prenatal visit?



CRITICAL THINKING ACTIVITY # 11

What questions will you ask Mrs. Ramirez on each visit?

Nursing Diagnoses

Your next step is to analyze the data from Mrs. Ramirez's history and physical assessment and develop nursing diagnoses. The following are possible nursing diagnoses for this patient. Cluster the supporting data for each diagnosis.

1. Knowledge deficit related to plan for prenatal visits during the remainder of pregnancy
2. Imbalanced Nutrition: less than body requirements, related to nausea
3. Risk for Fluid Volume Imbalance related to vomiting and weight loss associated with early pregnancy
4. Fatigue, related to early pregnancy
5. Health-Seeking Behaviors

Identify any possible additional nursing diagnoses.



CRITICAL THINKING ACTIVITY # 12

Now that you have identified some nursing diagnoses for Mrs. Ramirez, develop a brief teaching plan for her, selecting a diagnosis from the previous section. Include learning outcomes and teaching strategies.



Case Study Follow-Up

Subsequent Prenatal Visits

Mrs. Ramirez is seen at regular intervals throughout her pregnancy. As described previously, her routine prenatal assessments include an interview to determine her general physiological and emotional well-being, to answer questions, and to revise the birth plan, if needed. At each visit, Mrs. Ramirez is reminded of danger signs and symptoms and encouraged to continue her physical activity. Her physical exam consists of measurement of weight, pulse, respirations, BP (sitting), and fundal height; and abdominal inspection and palpation. A clean-catch urine to test for glucose, protein, nitrites, and leukocytes is obtained. A hematocrit is obtained frequently. At 25 weeks, a routine glucose challenge screening for gestational diabetes is conducted and is within normal limits. Fundal height measurements show normal fetal growth and an ultrasound examination at 26 weeks reveals a healthy fetus.

Childbirth

In her 39th week, Mrs. Ramirez begins to experience regular uterine contractions that intensify with activity. Two hours after she arrives at the hospital, her membranes rupture; the following morning, she gives birth to a healthy baby girl, Carmen Lucretia, who weighs

7 pounds, 6 ounces. After the birth, Mrs. Ramirez and Carmen experienced no problems and were discharged from the hospital 2 days later.

The Postpartal Assessment

The postpartal assessment includes a history and physical exam with focus on changes that occur after childbirth. The postpartal assessment immediately after delivery and 6-week follow-up is presented. In addition to reviewing your patient's prenatal records, ask the following questions:

- How many weeks' gestation? Were you full-term?
- How many times have you been pregnant? How many deliveries (children) do you have? How many pregnancies have you had (gravida)? How many pregnancies have you had that reached viability (parity)?
- What type of delivery? Vaginal or C-section?
- What type of anesthesia did you have?
- Do you have episiotomy? Tears?
- When was the last time you went to the bathroom? Last void?
- How do you feel? Pain?
- Are you breast- or bottlefeeding?
- What is your blood type? Rh factor? Predelivery hemoglobin and hematocrit?

(continued on page 845)

Postpartal Physical Assessment

AREA/NORMAL VARIATIONS

Vital Signs

- Temperature may increase to 38°C within first 24 hours owing to dehydration.
- Pulse initially elevated few hours after delivery. Returns to prepregnancy rate within 8 to 10 weeks.
- Respirations return to prepregnancy rate within 6 to 8 weeks.
- *Blood pressure:* Orthostatic hypotension for the first 48 hours owing to splenchnic engorgement; slight transient rise in BP for 4 days after delivery.

HEENT

Neck

- Thyroid may be palpable during pregnancy, returns to prepregnant state after pregnancy.

Respiratory

- *Lungs:* Clear.

Cardiovascular

- PMI returns to normal as cardiac axis is restored.

ABNORMAL VARIATIONS/RATIONALE

- *Elevated temperature after 24 hours:* May indicate puerperal sepsis, mastitis, endometritis, or UTI.
- *Elevated pulse:* May be associated with hypovolemia secondary to hemorrhage.
- *Decreased respiratory rate:* May be due to spinal block, pain medication.
- *Hypoventilation:* May occur with C-sections.
- *Hypotension:* may be associated with shock.
- *HTN:* May be associated with PIH.
- Thyromegaly.
- *Palpable nodules:* Thyroiditis, hypothyroidism.
- *Hypoventilation:* possible with C-section.
- *Tachycardia:* May signal shock.


(continued)

Postpartal Physical Assessment (continued)

AREA/NORMAL VARIATIONS

Abdomen

- Uterus firm and midline.
- 12 hours postpartum 1 cm above or at umbilicus.
- Descends (involutes) 1 to 2 cm (fingerbreadths) every 24 hours. After 9th postpartal day, not palpable.



Fundal heights postpartum

- Afterpains are uterine contractions as involution occurs. Afterpains increase with breast feedings, common in multipara patients.
- *Bowels*: positive bowel sounds, bowel movement within 2 to 3 days

Vaginal Delivery

- Pain with defecation secondary to episiotomy.

C-Section

- Hypoactive bowel sounds; incision clean, dry, and intact.

Genitourinary

- Bladder nondistended.
- Voids within 8 hours postpartum.
- *Perineum*: Episiotomy; hemorrhoids usually regress after birth.
- Sutures intact.
- Perineum tender and edematous.

Lochia

- Rubra (bloody) first few days postpartum. Rubra contains blood, decidual and thromboplastic debris.
- Serous (pink, brown) 3 to 4 days to 10 days postpartum. Serous contains old blood, serum, leukocytes, and tissue debris.
- Alba (yellow, white) 10 days up to 6 weeks' postpartum. Alba contains leukocytes, decidua, epithelial cells, mucus, serum, and bacteria.

Vascular

- *Extremities*: Increase for thromboemboli owing to hypercoagulative state.

ABNORMAL VARIATIONS/RATIONALE



ALERT

Boggy fundus may result in hemorrhage.

- *Displaced uterus*: May be due to distended bladder.
- *Costovertebral angle tenderness*: Kidney infection.
- Subinvolution.

- *Absent bowel sounds*: Sign of paralytic ileus (possibly due to C-section).

- *Inability to void*: Secondary to birth-induced trauma and anesthesia, pain, episiotomy. Overdistended bladder prevents involution, which can lead to hemorrhage.
- *Severe perineal pain*: Hematoma.

- *Increased rubra*: Hemorrhage, cervical or vaginal tears.

- *Positive Homans' sign*: May indicate phlebitis, DVT, varicosities.

AREA/NORMAL VARIATIONS

- Negative Homan's.
 - Varicosities usually regress after delivery.
 - Edema usually decreases after delivery.
- Neurologic**
- Carpal tunnel abates after delivery.
 - Affect depends on outcome of delivery.
 - Emotionally labile, postpartum blues common.
 - Bonding well with baby.
- Musculoskeletal**
- Joint stability returns within 6 to 8 weeks.
 - Increase in shoe size is permanent.

ABNORMAL VARIATIONS/RATIONALE

- **Headaches:** May be caused by PIH, stress, or spinal fluid leaks secondary to spinal anesthesia. Can last 1 to 3 days to weeks depending on cause.
- Postpartum depression or psychosis.



Case Study Follow-Up (continued)

The 6-Week Postpartal Assessment

Mrs. Ramirez, her mother, and Carmen arrive at the Women's Clinic 6 weeks later for the postpartal examination. Mrs. Ramirez is breastfeeding without difficulty, and reports that Carmen is gaining weight appropriately. She has had no problems and is smiling and cooing over the baby. Measurement of her weight reveals a 15-pound loss (her total weight gain during pregnancy was 26 pounds); her pulse, respirations, and BP are all within normal limits. A clean-catch urine specimen is obtained and is negative. Her hemoglobin/hematocrit levels are consistent with her prenatal values. During the interview, she has no complaints, and states that she plans to return to work in 4 more weeks. Her mother and older sister will share child care, and she will continue to breastfeed Carmen. She is still taking prenatal vitamins. She is interested in the diaphragm as a method of contraception. She has not had a menstrual period since her delivery, and reports that her lochia progressed through the normal stages of lochia rubra, lochia serosa, and lochia alba, and now has ceased. She displays no signs of postpartum depression.

Mrs. Ramirez's physical findings include thyroid nonpalpable, breast firm and full, milk expressible,

lungs clear, heart regular rhythm, no murmurs, abdomen soft, nontender, uterus nonpalpable, perineum intact.

Mrs. Ramirez is fitted with a diaphragm and instructed in its use and insertion and removal technique, which she is able to complete without difficulty. Because her prenatal Pap smear was normal, she is given a return appointment for 6 months for a routine follow-up visit.

Risk Factors for Postpartum Depression

- History of mood disorder
- Few or no social supports
- Stress
- Low self-esteem
- Poor marital relationship
- Unplanned pregnancy
- Depressed father
- Bottle feeding
- Mother unemployed or unemployed head of household

Source: Lowdermilk, D.L., Perry, S.E., and Bobak, I.M. (1999). *Maternity Nursing*, ed. 5. St. Louis: Mosby.

RESEARCH TELLS US

Nutritional deficiencies during pregnancy are known to result in poor birth outcomes and pregnancy complications. Use of multivitamins during pregnancy is recommended to improve the nutritional status of the patient and improve pregnancy outcomes. Because the exact timing of a nutrient's effect

on fetal development varies, a delay of using multivitamins until pregnancy is diagnosed may affect the course and outcome of the pregnancy.

The Pregnancy, Infection, and Nutrition Study (n = 2010) specifically compared women who used multivitamins prior to and during pregnancy with those

(continued)

RESEARCH TELLS US (continued)

who did not to determine risk for preterm birth (<37 weeks), and data were analyzed. The women were between 24 and 29 weeks' gestation from four prenatal clinics in North Carolina.

The findings revealed that preconceptual multivitamin use was inversely associated with early (<35 weeks adjusted odds ratio = 0.59, 95 percent confidence interval: 0.12, 2.76) and late (35 to 36 weeks;

adjusted odds ratio = 0.40, 95 percent confidence interval: 0.12, 1.40) preterm births. These findings suggest that women who take multivitamins prior to conception have a decreased risk of preterm birth.

Nurses have a key role in promoting women's health. Although further study is needed, the potential benefits of multivitamin use during reproductive years is apparent.

Source: Vahratian, A., Siega-Riz, A.M., Savitz, D., and Thorp, J. (2004). Multivitamin use and risk of pre-term birth. *American Journal of Epidemiology* 160(9):886-892.

HEALTH CONCERNS

Complications can occur during each trimester of pregnancy. The following is a list of signs and symptoms of complications and their causes.

First Trimester

- **Severe vomiting:** Hyperemesis gravidarum.
- **Chills, fever:** Infection.
- **Burning on urination:** Infection.
- **Abdominal cramping, bloating, vaginal bleeding:** Spontaneous abortion, miscarriage.

Second and Third Trimesters

- **Severe vomiting:** Hyperemesis gravidarum.
- **Leakage of amniotic fluid from vagina before labor begins:** Premature rupture of membranes.
- **Vaginal bleeding, severe abdominal pain:** Miscarriage, placental separation.

- **Chills, fever, diarrhea, burning on urination:** Infection.
- **Change in fetal activity:** Fetal distress, intrauterine fetal demise.
- **Uterine contractions before due date:** Preterm labor.
- **Visual disturbances (blurring, double vision, spots):** Hypertensive disorders (PIH).
- **Swelling of face, fingers, eye orbits, sacral area:** PIH.
- **Severe, frequent, or continuous headaches:** PIH.
- **Muscular irritability or convulsions (seizures):** PIH.
- **Severe stomachache (epigastric pain):** PIH.
- **Glucosuria, positive glucose tolerance test result:** Gestational diabetes mellitus.

SUMMARY

- A thorough prenatal assessment provides invaluable data regarding the patient's overall health status.
- Remember that the initial evaluation includes a thorough review of the patient's health history as well as a careful physical exam. This first visit provides an excellent opportunity to screen for possible complications of pregnancy, as well as to lay the groundwork for a trusting relationship with the pregnant woman.
- Also keep in mind that pregnancy affects every body system. Before you begin the assessment, review normal anatomy and physiology, as well as changes that routinely occur during pregnancy, so that you will be able to recognize abnormal findings.
- Always work in a systematic manner from apex to base and side to side. Establish a baseline, using the patient


as her own control (i.e., as a comparison throughout the pregnancy).

- After completing the assessment, document your findings; the plan for prenatal care should flow from your assessment and your findings with your patient. This provides an excellent opportunity for teaching and is an important beginning of the development of an individualized birth plan that meets the mother's needs and promotes a healthy pregnancy.
- Once the initial prenatal comprehensive assessment has been completed, routine follow-up prenatal visits usually include assessment of maternal weight, pulse, respirations, BP (same arm, with the patient sitting), urinalysis (glucose, protein, nitrites, and leukocytes), edema, fundal height, FHTs, and fetal movement.

R E V I E W Q U E S T I O N S

1. Janet Clark, age 24, has missed two periods and had a positive home pregnancy test. She is complaining of “morning sickness,” breast tenderness, and increased urinary frequency. Her last period was 11/30/06. Calculate her EDB.
 - a. 8/30/07
 - b. 7/23/07
 - c. 7/30/07
 - d. 8/23/07
2. Which sign would be considered a presumptive sign of pregnancy?
 - a. Cessation of menses
 - b. Abdominal enlargement
 - c. Fetal heart beats (FHB)
 - d. Positive pregnancy test
3. Which sign would be considered a probable sign of pregnancy?
 - a. Cessation of menses
 - b. Breast tenderness
 - c. FHB
 - d. Positive pregnancy test
4. Which sign would be considered a positive sign of pregnancy?
 - a. Cessation of menses
 - b. Breast tenderness
 - c. Fetal heart beats (FHB)
 - d. Positive pregnancy test
5. You perform a physical exam on Mrs. Clark. You note a bluish discoloration of the cervix. This sign is referred to as:
 - a. Goodell’s sign.
 - b. Hegar’s sign.
 - c. Chadwick’s sign.
 - d. Piskacek’s sign.
6. Mrs. Clark prepregnant weight was 130 pounds; her height is 5’6”. How much weight she should expect to gain during this pregnancy?
 - a. 28 to 40 lb
 - b. 15 to 25 lb
 - c. 25 to 35 lb
 - d. 35 to 45 lb
7. Your patient, Dawn Stubick, is beginning her third trimester of her pregnancy. As she lies on the examination table, she complains of feeling dizzy. You check her BP and note that she is hypotensive. How should you position your patient?
 - a. Left-side lying
 - b. Right-side lying
 - c. Trendelenburg’s position
 - d. Fowler’s position
8. You are assessing the FHT. What is the normal FHT rate?
 - a. 60 to 100
 - b. 100 to 120
 - c. 120 to 160
 - d. 160 to 200
9. You are listening for FHT with a Doppler. FHT can first be heard with a Doppler at:
 - a. Week 4.
 - b. Week 10.
 - c. Week 18.
 - d. Week 20.
10. Which technique is useful in assessing fetal position and presentation?
 - a. Pelvic exam
 - b. Leopold’s maneuver
 - c. Ballotement
 - d. Abdominal percussion

CHAPTER 24



Assessing the Newborn and Infant

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify normal physiological changes of newborns and infants
- Discuss various cultural influences on children
- Identify developmental tasks for newborn and infant
- Identify health issues and risks for newborn and infant
- Identify pertinent history data included in a newborn and infant history
- Obtain a newborn and infant health history
- Describe additional physical assessment components specific to a newborn and infant assessment
- Perform a newborn and infant assessment
- Document newborn and infant assessment findings
- Identify actual/potential health problems stated as nursing diagnoses with supporting data

Each newborn (birth to 30 days) arrives as a unique person with the energetic desire to grow and learn. For approximately 40 weeks, the fetus has enjoyed a warm, comfortable uterine environment with all needs met. At birth, he or she is totally dependent on the caretaker. Because the newborn is unable to directly communicate his or her needs, the nurse must learn assessment skills to identify abnormal findings and promote a healthy environment.

Infants (1–11 months) have few communication skills. As they grow, they can smile, frown, point, and even say “no,” but it will be several years before they can communicate well enough to provide information during a patient history. Until then, parents, siblings, and extended family are fine sources of information. Healthy families usually raise healthy babies, so take every opportunity to offer education and support during your assessment.

Anatomy and Physiology Review

Before you begin to examine a newborn or infant, you must have an understanding of the basic characteristics of this age group. These findings may vary slightly from child to child and change according to age and development. Use of standard measurement guidelines and physiological assessments will assist you in your task.

Many physiological changes occur at birth. Newborns begin their own oxygenation and circulatory efforts, and they must thermoregulate their bodies, ingest food, process nutrients, and expel wastes. They begin to interact mentally and physically with their surroundings.

Circulatory Adaptation

Major changes occur at birth to transition blood flow from fetal to newborn circulation. Aortic pressure increases and venous pressure decreases. Systemic pressure increases and pulmonary pressure decreases. Closure of the foramen ovale, ductus arteriosus, and ductus venosus occurs. Any variations in these changes may cause hypoxia and cyanosis. Figure 24.1 shows fetal circulation.

Hepatic Adaptation

Newborns have immature livers that cannot break down products of hemolysis easily. Bilirubin, a byproduct of this process, consequently builds up in the blood and cannot be excreted well through urine and feces. This results in jaundice—yellow-tinged skin, sclera, or mucous membranes. Prolonged high levels of bilirubin can cause mental and developmental delays. Physiological jaundice does not become apparent until after the newborn is 24 hours old.

Developmental, Cultural, and Ethnic Variations

As you perform your assessment, keep in mind the rapid developmental changes that occur during the first year of life and note any variations from the norm. Also remember that infants of different races and ethnic groups have different physical characteristics. Last, be aware of cultural variations that may affect parenting.

Developmental Considerations

During the first year of life, the infant experiences rapid growth and developmental changes. Many physical, gross motor, fine motor, primitive reflex, sensory,

Normal Newborn Physiological Changes

AREA/SYSTEM CHANGE	CAUSES/RATIONALE
Integumentary <ul style="list-style-type: none"> ■ <i>Acrocyanosis (peripheral cyanosis)</i>: Normal for 12 hours after birth. ■ <i>Vernix caseosa</i>: White, cheesy substance of sebum and desquamated epithelium, seen in small amounts in skin folds of full-term babies. ■ Desquamation (peeling of skin). ■ <i>Cutis marmorata (mottled skin)</i>: Seen in early infancy. If it persists, it may indicate congenital problems. ■ <i>Harlequin sign</i>: Dependent side red and nondependent side pale. ■ <i>Lanugo</i>: Fine, downy hair on face, shoulders, and back. Head <ul style="list-style-type: none"> ■ Bruises, lacerations, and temporary scalp and skull swelling may occur. 	<ul style="list-style-type: none"> ■ Caused by thermoregulation adjustments. ■ Protects skin in utero. ■ As skin dries after delivery, peeling is normally seen within 2 days. ■ Dilation of superficial vessels, not chilling. ■ Positional color changes seen in low-birth-weight newborns. ■ More common in premature infants. ■ Caused by physical stress of birth process. Increase in swelling may be caused by a hematoma or hydrocephalus.

AREA/SYSTEM CHANGE	CAUSES/RATIONALE
Respiratory <ul style="list-style-type: none"> ■ Rapid, periodic breathing required for extrauterine life. Newborns begin nose breathing immediately at birth. Breathing helps remove birth fluids from respiratory tract. Respiration permits oxygenation. 	<ul style="list-style-type: none"> ■ Respiration initiated by chemical and mechanical events of birth.
Circulatory <ul style="list-style-type: none"> ■ Onset of breathing stimulates transportation of oxygen. Shortly after first cry, heart rate accelerates. 	<ul style="list-style-type: none"> ■ BP is highest immediately after birth.
Hepatic <ul style="list-style-type: none"> ■ At birth, liver takes a major role in iron storage, red cell production, carbohydrate metabolism, bilirubin conjugation, and coagulation. 	<ul style="list-style-type: none"> ■ Immature liver often does not function well, resulting in high levels of bilirubin.
Gastrointestinal <ul style="list-style-type: none"> ■ Newborn has necessary enzyme activity and peristalsis to digest simple carbohydrates, fat, and protein. Stomach capacity is 50 to 60 mL. Stools are passed, often after a feeding. 	<ul style="list-style-type: none"> ■ Feeding promotes peristalsis. Cardiac sphincter of stomach is immature and may be cause of regurgitation after feedings.
Urinary <ul style="list-style-type: none"> ■ Urination often occurs at birth or shortly after. 	<ul style="list-style-type: none"> ■ Newborns are less able than adults to concentrate urine. Fluid intake enhances urination.
Musculoskeletal <ul style="list-style-type: none"> ■ Increased body movements are noted during reactive, awake phases. 	<ul style="list-style-type: none"> ■ Behavioral and sensory abilities trigger movement and exploration.
Thermoregulation <ul style="list-style-type: none"> ■ Newborn strives to maintain body temperature. The large body mass enhances heat loss, which consumes oxygen and may cause respiratory distress. 	<ul style="list-style-type: none"> ■ Body heat loss is caused by: <ul style="list-style-type: none"> ■ <i>Convection</i>: From body to cooler air. ■ <i>Radiation</i>: From body to cooler surfaces not in direct contact. ■ <i>Evaporation</i>: From body moisture to vapor. ■ <i>Conduction</i>: From direct skin to cooler surface.
Neurologic <ul style="list-style-type: none"> ■ During first period of reactivity, newborns are alert and active. ■ A sleep phase occurs after 30 minutes. ■ During second period of reactivity, newborn is awake and alert. 	<ul style="list-style-type: none"> ■ This phase lasts 30 minutes and is the natural inclination to search for a feeding. ■ This restorative phase lasts 2 to 4 hours. ■ Readiness for feeding is observed.


communication, and socialization changes are seen. During the first year of life, psychosocial development focuses on establishing trust. Trust is established when the infant senses that her or his needs are being met.

Cognitive development during the first year of life is also called the **sensorimotor phase** of development. The three major tasks during this phase are separation, realizing that self is separate from other objects; object permanence, realizing that objects are permanent even when not in sight; and mental representation, recognizing symbols of objects without actually experiencing the object. The infant also is developing a body image by exploring and playing with different parts of his or her body.

Social development at this age includes developing attachment to parents or caregivers and experiencing separation anxiety and fear of strangers.

Communication and language development also occurs during the first year. The Denver Developmental Screening Test-II (DDST-II) and growth charts are frequently used to assess and track the growth and development of the infant during well-baby checkups.

As you assess the infant, be sure to note if her or his physical development is appropriate for her or his age and whether she or he is performing appropriate developmental tasks for that age. Because growth and development are so rapid during the first year of life, even the



FETAL CIRCULATION

FIGURE 24.1. Fetal circulation.

slightest developmental delay may signal an underlying problem and warrant further investigation.

Key physical changes include:

- Birth weight doubles by 6 months, triples by 12 months.
- Height increases by 1 inch per month for first 6 months.
- Fontanels are closing.
- Lumbar curve develops with a lordosis once the infant begins to walk.
- Drooling and teething occur.
- Primitive reflexes disappear as the neurological system matures.

Gross motor changes include:

- Rolls, crawls.
- Pulls self up to sit.
- Begins to walk.
- Achieves head control.

Fine motor changes include:

- Grasps objects.
- Puts objects in mouth.
- Holds bottle.
- Plays with toes.
- Develops pincer grasp.

Sensory changes include:

- Develops better vision.

- Follows objects with eyes.
- Responds to sounds.

Communication changes include:

- Initially cries to convey needs.
- Babbles.
- Laughs.
- Says three to five words by 12 months.
- Begins to comprehend simple directions.
- Imitates sounds.

Socialization changes include:

- Identifies parents.
- Develops social smile.
- Is aware of strange situations.
- Has increasing difficulty separating from parents.
- Becomes more fearful of strangers.
- Begins to develop memory.
- Shows emotions.

Cultural and Ethnic Considerations

Cultural or ethnic influences may affect your assessment findings. You need to be aware of these normal variations so that you do not mistake them for abnormal findings. Cultural or ethnic influences may also affect the relationship between child and parent and define the roles of both parent and child within the family.

Cultural and Ethnic Variations in Infants

CULTURAL GROUP	INFLUENCE
African American	<ul style="list-style-type: none"> ■ Mongolian spots and other birthmarks more prevalent than in other ethnic groups.
Amish	<ul style="list-style-type: none"> ■ Babies seen as gifts from God. Have high birth rates, large families.
Appalachian	<ul style="list-style-type: none"> ■ Newborns wear bands around abdomen to prevent umbilical hernias and asafetida bags around neck to prevent contagious diseases.
Arab American	<ul style="list-style-type: none"> ■ Children “dearly loved.”
Chinese American	<ul style="list-style-type: none"> ■ Male circumcision is a religious requirement. ■ Children highly valued because of one-child rule in China. ■ Mongolian spots occur in about 80 percent of infants. ■ Bilirubin levels higher in Chinese newborns than others, with highest levels seen on day 5 or 6.
Cuban American	<ul style="list-style-type: none"> ■ Childbirth is a celebration. Family takes care of both mother and infant for the first 4 weeks. ■ Tend to bottle-feed rather than breastfeed. If breastfeeding, weaning is early, around 3 months. If bottle feeding, weaning is late, around 4 years.
Egyptian American	<ul style="list-style-type: none"> ■ Children very important.
Filipino American	<ul style="list-style-type: none"> ■ Mother and infant cared for by family for the first 50 days. ■ Eyes almond shaped, low to flat nose bridge with mildly flared nostrils. Mongolian spots common.
French Canadian	<ul style="list-style-type: none"> ■ Five mutations account for 90 percent of phenylketonuria (PKU) in French Canadians. High incidence of cystic fibrosis and muscular dystrophy.
Greek American	<ul style="list-style-type: none"> ■ High incidence of two genetic conditions: thalassemia and glucose-6-phosphate dehydrogenase (G6PD).
Iranian American	<ul style="list-style-type: none"> ■ Believe in hot/cold influences, with baby boys “hotter” than baby girls. Infant may be confined to home for first 40 days. Ritual bath between 10th and 40th day.
Jewish American	<ul style="list-style-type: none"> ■ Children seen as valued treasure. ■ High incidence of Tay-Sachs disease. ■ Male circumcision is a religious ritual.
Mexican American	<ul style="list-style-type: none"> ■ Wears stomach belt (<i>ombliguero</i>) to prevent umbilicus from popping out when infant cries. ■ Believe that cutting nails in the first 3 months after birth causes blindness and deafness.
Navajo Native American	<ul style="list-style-type: none"> ■ Infants kept in cradle boards until they can walk.
Vietnamese American	<ul style="list-style-type: none"> ■ Mongolian spots common.

Source: Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.

Introducing the Case Study

As you recall from the previous chapter, Mrs. Ramirez gave birth to a baby girl, Carmen. Carmen is now 2 hours old. Her mother had a normal vaginal delivery, and you are admitting them both to the postpartum unit. You assessed Mrs. Ramirez and noted normal vital signs, firm uterus, intact perineum, and moderate lochia rubra. She breastfed Carmen in the delivery room. You also obtained Apgar scores for Carmen of 8 at 1 minute and 10 at 5 minutes. Now you will assess Carmen more thoroughly. When you change her diaper, you note that she has voided and passed a meconium stool.



Performing the Initial Newborn Assessment

The initial newborn assessment consists of obtaining an Apgar score immediately after birth, taking a health history, and then performing a physical assessment that includes a general survey, head-to-toe assessment, and tests for reflexes. Depending on the newborn's condition, certain diagnostic and screening tests may also be performed.

Apgar Score

Heart Rate	0 = absent	1 \leq 100 BPM	2 \geq 100 BPM
Respirations	0 = absent	1 \leq 30, irregular	2 = strong cry, regular
Muscle Tone	0 = flaccid	1 = some flexion in arms and legs	2 = full flexion, active movement
Reflex Irritability	0 = no response	1 = grimace, weak cry	2 = vigorous cry
Color	0 = pale, blue	1 = body pink, extremities blue	2 = totally pink

Apgar Score

Assess the newborn's physiological and cardiopulmonary status immediately after birth. The Apgar score assesses five vital areas: heart rate, respiration, color, muscle tone, and reflex irritability. A score from 0 to 2 is assigned to each category, for a score range of 0 to 10. This score is obtained at 1 and 5 minutes after birth. A score of 8 to 10 is excellent, 4 to 7 is guarded, and 0 to 3 is critical.

Newborn Health History

The newborn's health history addresses two main areas: prenatal health and labor and delivery. The focus of prenatal health is on the mother's prenatal care, obstetric history; use of medications; and use of tobacco, alcohol, and drugs. Labor and delivery address gestational week at time of delivery, duration of labor, type of delivery, and type of anesthesia.



Case Study Findings

Newborn Carmen's health history reveals:

- 39 weeks' gestation, gravida 1/para 1.
- Positive prenatal care.

- Mother generally healthy, no history of medical problems.
- No prenatal diseases.
- No history of smoking, drug or alcohol abuse in mother.
- Prenatal vitamins once a day.
- 16-hour labor, uncomplicated vaginal delivery.
- Epidural anesthesia.
- First-degree laceration and midline episiotomy.
- Parents Cuban American, bilingual.

Newborn Physical Assessment

Now, proceed to the newborn physical assessment. Keep your history findings in mind and combine them with the physical examination findings to form a total picture of the newborn.

Approach

Use your senses of sight, smell, hearing, and touch along with all four techniques of physical assessment. Remember that your equipment and techniques will have to be adjusted for the size of your patient. In

Newborn Health History

HEALTH HISTORY

- General health, prenatal diseases or conditions, prenatal care, number of pregnancies.
- Use of prescribed or over-the-counter medications, tobacco, alcohol, illegal drugs.
- Duration of pregnancy and labor, type of anesthesia, type of delivery, complications.

SIGNIFICANCE

- Maternal health problems (e.g., gestational diabetes, cardiac or kidney disease) may cause potential risk factors in newborn.
- Medications and other agents may affect physiological systems (e.g., smoking during pregnancy related to low birth weights, alcohol use related to fetal alcohol syndrome [FAS]).
- Details of labor and delivery alert nurse to observe for potential newborn problems.

addition, instead of proceeding from head to toe, adapt your assessment sequence to fit the newborn's needs. For example, move techniques that may evoke crying—such as the otoscopic examination—to the end of the assessment. Also be sure to keep the room and the baby warm.



TOOLBOX

Gather the following equipment: Tape measure, stethoscope, thermometer, blood pressure cuff, penlight, otoscope, and ophthalmoscope. You will also need a baby scale.

Performing a General Survey

Before you perform your head-to-toe assessment of the newborn, ask yourself the following questions and record your observations as baseline data.

- **Integumentary:** Are there any abrasions, lacerations, or birthmarks? If so, describe them, and monitor for infection, bleeding, and trauma.
- **Head and Neck:** Are there masses on the head or neck? These may indicate cephalohematoma or a fractured clavicle.
- **Eyes, Ears, Nose, Mouth, and Throat:** Is there exudate in the eyes? If so, monitor for infection. Is the baby blinking? Do his or her eyes follow an object within 8 inches? Is there discharge from the ears, nose, and throat or nose congestion? Fluid may be from delivery or could indicate infection. Does the newborn respond to sound? A reaction to sound should occur.



ALERT

Babies are obligatory nose breathers, so clear secretions from nose to ease breathing.

- **Respiratory Characteristics, Lungs, and Breathing:** Is the newborn congested or gasping for breath? Fetal fluid that remains in the lungs can cause airway obstruction.
- **Cardiovascular Characteristics:** Is there cyanosis? It may denote poor vascular perfusion. Is the newborn alert? A newborn in difficulty will demonstrate irritability or be unarousable.
- **Temperature Regulation:** Is body temperature maintained? Newborns are poor regulators of body temperature and are dependent on their environment for warmth. Cold stress can cause respiratory distress.
- **Hepatic Regulation:** Does the newborn have jaundice? Increasing jaundice denotes increased blood bilirubin, which can cause mental deterioration if untreated.



ALERT

Jaundice within 24 hours of birth is pathological jaundice; jaundice after 24 hours is probably physiological jaundice.

- **Gastrointestinal Adaptation:** Has the newborn passed stool? This proves that an anus is present. Has she or he vomited? Persistent vomiting suggests intestinal obstruction.
- **Genitourinary Adaptation:** Has the baby voided? Voiding denotes kidney function.
- **Neurological Characteristics:** How are extremities moving? Muscle tone should be symmetrical and not flaccid, extremities should be partially flexed; hand-mouth behavior should be evident; and when awake, newborn should demonstrate random, purposeless, bilateral movements. How does cry sound? It should be loud, not high pitched or weak.
- **Endocrine Characteristics:** Is there evidence of fetal or maternal endocrine disease? Is the newborn jittery? Maternal diabetes can cause severe hypoglycemia in the newborn. Low glucose levels often manifest as jittery extremities.
- **Immunologic Adaptation:** Are defenses maintained? Because the newborn's immune response is not well established, protect portals of entry (umbilical stump and breaks in skin) from infection.
- **Sleep and Rest Patterns:** What is the sleep pattern? Newborns sleep up to 18 hours a day. The long sleep period promotes growth and development. They should awaken for feedings every 3 to 4 hours.
- **Relationships, Psychosocial Profile, and Cultural/Ethnic Variations:** How are family members relating to the newborn? The newborn is dependent on the family for its well-being. Parents should demonstrate touching and holding of infant. Assist in the bonding process. Responses may vary depending on cultural/ethnic variations.

Performing a Head-to-Toe Physical Assessment

Next, perform a head-to-toe physical assessment of the newborn.

Performing Barlow-Ortolani Maneuver

Place the infant supine on a flat surface. Flex his or her knees, holding your thumbs in the inner thighs and your fingers on the outside of the greater trochanters of the hips. Move the legs inward until your thumbs touch. Gently but firmly rotate the hips outward so that the knees touch the flat surface. There should be no clicking or crepitus noted during this procedure. If there is, the joint is unstable.

(Text continued on page 867)

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

1. Posture

AREA/SYSTEM/NORMAL FINDINGS

- Inspect posture.
- *Head and extremities flexed.*

ABNORMAL FINDINGS/RATIONALE

- *Limp posture with extension of extremities:* Associated with birth injuries, anesthesia, acidosis, hypoglycemia, hypothermia, or congenital problems.

2. Head Circumference

AREA/SYSTEM/NORMAL FINDINGS

- Measure head circumference.



Measuring head circumference

- *33 to 35 cm.*



HELPFUL HINT

Measure head circumference from occiput to forehead.

ABNORMAL FINDINGS/RATIONALE

- *Head circumference < 10% of normal:* Microcephaly related to congenital malformation or infection.
- *Head circumference > 90% of normal:* Macrocephaly related to hydrocephalus.

3. Chest Circumference

AREA/SYSTEM/NORMAL FINDINGS

- Measure chest circumference.



Measuring chest circumference

- *30.5 to 33 cm (2 to 3 cm less than head).*
- *Breast engorgement can affect measurement.*



HELPFUL HINT

Measure chest at nipple line.

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

4. Abdominal Circumference

AREA/SYSTEM/NORMAL FINDINGS

- Measure abdominal circumference.



Measuring abdominal circumference



HELPFUL HINT

Measure abdomen above the umbilicus.

- *Similar to chest measurement. Should not be distended.*

5. Length

AREA/SYSTEM/NORMAL FINDINGS

- Measure length.



Measuring length

- **Crown to rump:** 31 to 35 cm (about equal to head circumference).
- **Head to heel:** 45 to 55 cm (18 to 22 inches) at birth.
- *Molding can affect measurement.*

6. Weight

AREA/SYSTEM/NORMAL FINDINGS

- Weigh newborn.



Weighing the infant

- *Newborn weight is usually between 2500 and 4000 g (5 lb, 8 oz, and 8 lb, 13 oz).*

(continued)

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

6. Weight (continued)

ABNORMAL FINDINGS/RATIONALE

- Birth weights < 10 or > 90 percent are abnormal.
- *Low birth weight (small for gestational age)*: Associated with prematurity.
- *Macrosomic infant (large for gestational age)*: Associated with gestational diabetes in mother.

7. Temperature

AREA/SYSTEM/NORMAL FINDINGS

- Take newborn's temperature.
- **Axillary**: 36.5 to 37.2°C

ABNORMAL FINDINGS/RATIONALE

- Hypothermia leads to cold stress. Sepsis, environmental extremes, and neurological problems can cause hypothermia or hyperthermia.

8. Pulse

AREA/SYSTEM/NORMAL FINDINGS

- Auscultate heart rate.
- *Apical rate 120 to 160 BPM.*
- *Rate increases with crying and decreases with sleep.*

ABNORMAL FINDINGS/RATIONALE

- Irregular rhythms such as bradycardia (<100 BPM) and tachycardia (>160 BPM).
- Most murmurs are not pathological and disappear by age 6 months.

9. Respirations

AREA/SYSTEM/NORMAL FINDINGS

- Take newborn's respirations.
- *30 to 60 breaths a minute; irregular.*



HELPFUL HINT

Anesthesia during labor and delivery can affect respirations.

ABNORMAL FINDINGS/RATIONALE

- Respirations < 30 or > 60 breaths a minute.
- Periods of apnea > 15 seconds.

10. Blood Pressure

AREA/SYSTEM/NORMAL FINDINGS

- Take newborn's BP.
- **Systolic**: 50 to 75 mm Hg.
- **Diastolic**: 30 to 45 mm Hg.



HELPFUL HINT

Crying and moving increase systolic pressure.

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

10. Blood Pressure (continued)

ABNORMAL FINDINGS/RATIONALE

- *Low BP:* May be caused by hypovolemia.
- Late clamping of umbilical cord can increase BP because of expanded blood volume from the “placental transfusion.”

11. Integumentary

AREA/SYSTEM/NORMAL FINDINGS

Skin

- Inspect skin, note, color, lesions.
- *Skin may be red, smooth, edematous, mottled (cutis marmorata).*
- *Hands and feet may be cyanotic (acrocyanosis).*
- *Physiological jaundice occurs after 24 hours.*
- *Color may change with position (harlequin sign).*
- *Cheesy substance (vernix caseosa) decreases as baby’s gestational age increases to term.*
- *Desquamation (peeling), ecchymosis, and petechiae may occur from trauma during delivery.*
- *Milia (white papules) may occur on face.*

- *Miliaria or audamina (papules or vesicles on face) are caused by blocked sweat ducts.*
- *Mongolian spots (bluish discoloration in sacral area) are commonly seen in African, Asian, Latin, and Native American babies.*
- *Telangiectatic nevi.*
- *Flat hemangiomas (“stork bites”) may be present at nape of neck.*

Hair

- Inspect hair, and note distribution.
- *Some lanugo is normal.*

Nails

- Inspect neonate’s nails.



Milia



Stork bite

(continued)

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

11. Integumentary (continued)

ABNORMAL FINDINGS/RATIONALE

- *Persistent acrocyanosis*: May indicate thermoregulation problem or hypoglycemia.
- *Extensive desquamation*: Seen in post-term baby
- Pathological jaundice occurs within first 24 hours.
- *Plethora*: May indicate polycythemia.
- *Pallor*: May indicate anemia, hypothermia, shock, or sepsis.
- *Persistent ecchymosis or petechiae*: May be caused by thrombocytopenia, sepsis, or congenital infection.
- *Poor turgor*: Intrauterine growth retardation or hypoglycemia.
- *Café-au-lait spots (light brown spots)*: If more than six or larger than 4 x 6 cm, may indicate neurofibromatosis and can become precancerous with age.
- *Nevus flammeus (port-wine stain)*: Disfigures face and may be associated with cerebral vascular malformation.
- Giant hemangiomas and nevus vasculosus ("strawberry marks") tend to trap platelets and lower circulating platelet counts. They usually disappear by age 5.
- Reddish-blue round mass of blood vessels (cavernous hemangioma) must be monitored, and if size increases, surgery may be necessary.
- Erythema toxicum, a common newborn rash of red macules and papules, usually disappears in 1 week.
- *Bullae or pustules*: May indicate infections such as syphilis or staphylococcus.
- Thin, translucent skin and vernix caseosa are signs of prematurity.
- Genetic disorders may cause extra skin folds.
- Abundant lanugo is a sign of prematurity.
- Genetic disorders may cause abnormal hair distribution unrelated to gestational age.
- Long nails are seen in post-term babies.

12. HEENT

AREA/SYSTEM/NORMAL FINDINGS

Head/Face

- Inspect head and fontanels.
- Gently palpate fontanels.
- Inspect facial features and movements.
- *Molding in birth canal may cause asymmetry of face and skull and should resolve within 1 week.*
- **Anterior fontanel:** *Diamond shaped, 2.5 to 4 cm.*
- **Posterior fontanel:** *Triangle shaped, 0.5 to 1 cm.*
- *Soft and flat.*
- *Symmetrical facial movements.*

Neck

- Inspect and palpate the neck.
- Test tonic neck reflex.
- *Short neck.*
- *Positive tonic reflex.*
- *Able to hold head up with "pull-to-sit" test.*



Palpating fontanel




Transilluminating fontanel

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

12. HEENT (continued)

Eyes

- Inspect eyes, position, edema, exudates, color of sclera, parallel alignment, pupil size, and equality.
- Test corneal/blink reflex.
- Test red light reflex.
- *Eyes may be edematous after vaginal delivery.*
- *Eyes equal and symmetrical.*
- *Blue-gray or brown iris; white or bluish-white sclera.*
- *Antimongolian slant; Mongolian slant seen in Asian infants.*
- *Positive red light reflex.*
- *Positive blink reflex.*
- *Positive corneal reflex.*
- *No tears (tear production begins by 2 months).*
- *Positive fixation on close objects.*
- *Positive pupillary reaction to light.*
- *Strabismus and searching nystagmus caused by immature muscular control.*



Normal eye line



HELPFUL HINT

Avoid bright light because it will cause the newborn to avoid opening her or his eyes and make assessment difficult.

Ears

- Inspect shape, position, and drainage.
- Test hearing.
- *Pinna flexible, without deformity, aligns with external canthus of eyes.*
- *Positive startle reflex.*



HELPFUL HINT

American Academy of Pediatrics recommends hearing screening by auditory brainstem response or evoked otoacoustic emissions on newborns before discharge.

Nose

- Inspect nares.
- Nares patent.
- *Small amount of thin white mucus.*
- *Nose may be flattened and bruised from birth.*



ALERT

Natal teeth must be removed by a specialist because they usually fall out and can cause choking.

Mouth/Throat

- Inspect mucous membranes, lips, tongue, and palate.
- Test sucking, rooting, gag, extrusion and swallowing reflexes.
- *Mucous membranes pink and moist.*
- *Frenulum of tongue and lip intact.*
- *Palate intact, uvula midline.*
- *Strong sucking reflex, positive rooting, gag, extrusion, and swallowing reflexes.*
- *Minimal saliva.*
- *Strong cry.*
- *Natal teeth may be benign or associated with congenital defects.*



Natal teeth

(continued)

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

12. HEENT (continued)

- *Small white, pearl-like epithelial cysts on the palate (Epstein's pearls), disappear within a few weeks.*



Epstein's pearls

ABNORMAL FINDINGS/RATIONALE

- Fused sutures.
- *Large fontanels:* Associated with hydrocephaly, osteogenesis imperfecta, congenital hypothyroidism.
- *Small fontanels:* Associated with microcephaly.
- *Bulging fontanels:* May indicate increased intracranial pressure.
- *Depressed fontanels:* Associated with dehydration.
- Craniosynostosis (premature closure of the sutures).
- Cephalohematoma (hematoma between periosteum and skull with unilateral swelling).
- Most uncomplicated cephalohematomas totally resolve within 2 weeks to 3 months.
- Caput succedaneum (edema of soft scalp tissue from birth trauma) decreases gradually in several days.
- *Asymmetrical facial movements:* May result from damage to facial nerve during forceps delivery.
- *Absent tonic reflex:* Erb's palsy if unilateral or dislocation of cervical spine or fractured clavicle.
- *Head lag with "pull-to-sit" test:* Muscle weakness.
- Torticollis (wry neck).
- *Subconjunctival hemorrhage:* Trauma during delivery.
- *Brushfield spots (speckling of iris), epicanthal fold, and Mongolian slant:* Down syndrome.
- *Absent red light reflex:* May indicate congenital cataract.
- *Ptosis:* Neuromuscular weakness.
- *Sun-setting (crescent of sclera over iris caused by retraction of upper lid):* Hydrocephalus.
- *Yellow sclera:* Jaundice.
- *Blue sclera:* Osteogenesis imperfecta.
- *Persistent nystagmus, absent blink reflex, inability to follow objects:* May indicate vision problem, such as blindness.
- *Dilated or fixed pupil:* May indicate anoxia or neurological damage.
- Chemical conjunctivitis from eye prophylaxis may occur during first 24 hours.
- *Low-set ears:* Down syndrome.
- *Absent startle reflex:* Possible hearing problem.
- Because infants are obligatory nose breathers, large amounts of mucus drainage may obstruct nostrils and cause respiratory difficulty.
- *Nasal flaring:* Sign of distress.
- *Cyanotic mucous membranes:* Hypoxia.
- *Candida albicans (thrush):* Contracted during vaginal delivery.
- *Weak sucking, swallowing reflex:* May be caused by maternal anesthesia or perinatal asphyxia.
- *Opening in palate or lips:* Cleft palate or lip. Any opening is abnormal. A series of surgical interventions will be necessary.
- *Weak cry:* May indicate neuromuscular problem, hypotonia, and prematurity.



ALERT

The ears and kidneys develop at the same time in utero, so malformed ears may be accompanied by renal problems.



ALERT

Cleft lip or palate will cause newborn to have difficulty with feeding



Thrush

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

13. Chest

AREA/SYSTEM/NORMAL FINDINGS

- Inspect shape, symmetry, and chest excursion.
- Inspect breast size, drainage.
- *Anteroposterior:lateral (1:1).*
- *Equal chest excursion.*
- *Breast engorgement.*
- *Clear or milky liquid from nipples (“witch’s milk”) develops from maternal hormones in utero.*
- *Supernumerary nipples are a benign finding.*

ABNORMAL FINDINGS/RATIONALE

- *Funnel chest (pectus excavatum):* Congenital anomaly.
- *Pigeon chest (pectus carinatum):* Obstructed respiration in infancy.
- *Asymmetrical excursion, retraction:* Respiratory distress.
- Red, firm nipples.

14. Respiratory

AREA/SYSTEM/NORMAL FINDINGS

- Auscultate breath sounds.
- *Lungs clear, bronchial to bronchovesicular breath sounds audible.*
- *Cough reflex absent at birth, but present 1 to 2 days later.*
- *Scattered crackles a few hours after birth.*



A. Auscultating lungs posteriorly.



B. Auscultating lungs anteriorly.

ABNORMAL FINDINGS/RATIONALE

- Persistent crackles, wheezes, stridor, grunting, paradoxical breathing, decreased breath sounds, prolonged periods of apnea (>15 to 20 sec) are signs of respiratory problems.

(continued)

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

15. Cardiac

AREA/SYSTEM/NORMAL FINDINGS

- Auscultate heart sounds.
- S_1 , S_2 normal rhythm with respiratory variations.
- PMI fourth left intercostal space midcostal line.



HELPFUL HINT

Quiet but clearly audible murmurs occur in 30 percent of newborns but should disappear in 2 days.



Auscultating the heart

ABNORMAL FINDINGS/RATIONALE

- Dextrocardia (heart on right side).
- *Cardiomegaly*: Displaced point of maximal impulse (PMI).
- Murmurs are often heard at base or along left sternal border, and are usually benign, but need to be evaluated to rule out cardiac disorder.
- Thrills.

16. Abdomen

AREA/SYSTEM/NORMAL FINDINGS

- Inspect abdomen and umbilical cord.
- Note hernias or diastasis recti.
- Gently palpate abdomen and femoral pulse.
- Auscultate bowel sounds.
- *Abdomen round.*
- *Positive bowel sounds.*
- *Liver edge palpable 2 to -3 cm.*
- *Tip of spleen, kidneys palpable.*
- *Cord bluish white with two arteries and one vein.*
- *Positive femoral pulses.*
- *Umbilical hernias and diastasis recti (separation of rectus muscles) more common in African American infants and often resolve within a year.*



Inspecting the cord



Palpating femoral pulse



Auscultating bowel sounds

ABNORMAL FINDINGS/RATIONALE

- *Abdominal distension, ascites, distended veins*: May indicate portal hypertension.
- *Green umbilical cord*: May indicate infection.
- *Absence of umbilical vessels*: Associated with heart and kidney malformations.

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

17. Rectum

AREA/SYSTEM/NORMAL FINDINGS

- Inspect anus.
- *Anus patent.*
- *Passage of meconium stool within 48 hours.*
- *Positive anal reflex (“anal wink”).*

ABNORMAL FINDINGS/RATIONALE

- Anal fissures or fistulas.
- *No stools:* May indicate malformation in gastrointestinal (GI) tract.



ALERT

Imperforate anus (absent anus) requires immediate surgical repair.

18. Female Genitalia

AREA/SYSTEM/NORMAL FINDINGS

- Inspect genitalia. Place thumbs on either side of labia and gently separate tissues to visualize perineum.
- Note presence of clitoris, vagina, and hymen.
- *Urination within 24 hours.*
- *Urinary meatus is midline and an uninterrupted stream is noted on voiding.*
- *Labia majora and minora may be edematous.*
- *Blood-tinged vaginal fluid may be noted (pseudomenstruation).*

ABNORMAL FINDINGS/RATIONALE

- Fused labia or absent vaginal opening.
- Ambiguous genitalia.
- Meconium from vaginal opening.
- Inability to urinate within 24 hours.



ALERT

A newborn clitoris larger than 0.5 cm is abnormal.

19. Male Genitalia

AREA/SYSTEM/NORMAL FINDINGS

- Inspect penis, and note position of urinary meatus.
- Retract foreskin.
- Palpate scrotum, and note undescended testicle if present.
- *Urination within 24 hours.*
- *Foreskin retracts.*
- *Urethral opening at tip of penis.*
- *Scrotum edematous.*
- *Smegma.*
- *Palpable testes.*



Palpating the scrotum

ABNORMAL FINDINGS/RATIONALE

- Hypospadias (urethral opening on ventral surface of penis).
- Epispadias (urethral opening on dorsal side of penis).

(continued)

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

19. Male Genitalia (continued)

ABNORMAL FINDINGS/RATIONALE

- Chordee (ventrally curved penis).
- Hydrocele (fluid in scrotum).
- Inability to urinate within 24 hours.
- Inability to retract foreskin.
- Undescended testicles.
- Inguinal hernia.
- Ambiguous genitalia.
- Meconium from scrotum.

20. Musculoskeletal

AREA/SYSTEM/NORMAL FINDINGS

- Inspect extremities, and note number of digits and deformities.
- Inspect spine, and note shape.
- Inspect gluteal folds.
- Perform Barlow-Ortolani maneuvers.
- *10 fingers and 10 toes.*
- *Full ROM.*
- *No clicks in joints.*
- *Equal gluteal folds.*
- *C curve of spine, no dimpling.*
- *When arms and legs are extended:*
 - *Muscles symmetrical and with equal muscle tone.*
 - *Arms and legs symmetrical in size and movement.*
- *Hands held as fists until after 1 month, when grasp becomes strong and equal.*
- *Position in utero may affect appearance.*



Checking gluteal folds



Checking for C curve



A. Barlow-Ortolani maneuver #1



B. Barlow-Ortolani maneuver #2



C. Barlow-Ortolani maneuver #3

PERFORMING A NEWBORN HEAD-TO-TOE PHYSICAL ASSESSMENT

20. Musculoskeletal (continued)

ABNORMAL FINDINGS/RATIONALE

- *Polydactyly*: Extra digits.
- *Syndactyly*: Webbed digits.
- *Phocomelia*: Hands and feet attached close to chest.
- *Hemimelia*: Absence of distal part of extremity.
- *Talipes (clubfoot)*: Foot permanently twisted out of shape.
- Severe bowing of legs is abnormal.
- *Unequal gluteal folds and positive Barlow-Ortolani maneuver*: Associated with congenital hip dislocation. Requires immediate referral.
- Decreased ROM and muscle tone.
- *Swelling, crepitus, neck tenderness*: Possible broken clavicle.
- *Simian (transverse palmar) creases*: Down syndrome.

21. Neurological

AREA/SYSTEM/NORMAL FINDINGS

- Test the newborn's reflexes.
- *Positive newborn reflexes*.
- *Positive knee reflex*.



Checking knee reflex

ABNORMAL FINDINGS/RATIONALE

- *Hypotonia*: Floppy, limp extremities.
- Paralysis.
- Marked head lag.
- Tremors.
- Asymmetrical posture.
- *Hypertonia*: Tightly flexed arms and stiffly extended legs with quivering.
- *Opisthotonic posture*: Arched back.
- *Dimpling of spine, tuft of hair*: May indicate spina bifida or pilonidal cyst.

Testing Reflexes

Infant reflexes are often present at birth and occur because the neurological system is immature. Many of these reflexes disappear as the neurological system develops. Some reflexes, such as rooting, sucking, and swallowing, are critical for survival. Other reflexes are protective reflexes and last throughout life. These include the blink or corneal, papillary reaction, sneezing, gag, cough, and yawn reflexes. (See *Newborn/Infant Reflexes*.)



HELPFUL HINT

In premature babies, newborn reflexes may persist longer.

Performing Diagnostic/Screening Tests

Depending on the newborn's condition, various diagnostic and screening tests are performed shortly after birth. Mandatory testing varies from state to state. (See *Diagnostic/Screening Tests for Newborns*.)

(Text continued on page 874)

NEWBORN/INFANT REFLEXES

1. Moro

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth and lasts 1 to 4 months
- *Technique:* Startle infant by suddenly jarring bassinet, or with infant in semisitting position, let head drop back slightly.
- *Quickly abducts and extends arms and legs symmetrically.*
- *Makes "C" with index finger and thumb. Legs flex up against trunk.*



Moro reflex

ABNORMAL RESPONSE

- Premature or ill infants may have sluggish response.
- Positive response beyond 6 months indicates neurological problem.
- Asymmetrical response may be caused by injury to clavicle, humerus, or brachial plexus during delivery.

2. Startle

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth and lasts 4 months
- *Technique:* Startle infant by making loud noise.
- *Hands clenched, arms abducted, flexion at elbow.*



Startle reflex

ABNORMAL RESPONSE

- Same as Moro.

3. Tonic Neck

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present between birth and 6 weeks; disappears at 4 to 6 months.
- *Technique:* With infant supine, rotate head to one side so that chin is over shoulder.
- *Infant assumes "fencing position," with arm and leg extended in direction to which head was turned.*



Tonic neck reflex

NEWBORN/INFANT REFLEXES

3. Tonic Neck (continued)

ABNORMAL RESPONSE

- Response after 6 months may indicate cerebral palsy.

4. Palmar Grasp

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 3 to 4 months.
- *Technique:* Place object or finger in palm of infant's hand.
- *Infant grasps object tightly. If he or she grasps your fingers with both hands, infant can be pulled to a sitting position.*



Palmar grasp reflex

ABNORMAL RESPONSE

- Negative grasp seen with hypotonia or prenatal asphyxia.

5. Plantar Grasp

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 3 to 4 months.
- *Technique:* Place thumb firmly against ball of infant's foot.
- *Toes flex tightly downward in a grasping motion.*



Plantar grasp reflex

ABNORMAL RESPONSE

- Negative grasp seen with hypotonia or spinal cord injury.

6. Babinski

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 1 year.
- *Technique:* Stroke lateral surface of sole of infant's foot.
- *Toes should fan.*



Babinski reflex

(continued)

NEWBORN/INFANT REFLEXES

6. Babinski (continued)

ABNORMAL RESPONSE

- Diminished response associated with neurological problem.

7. Stepping or Dancing

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 3 to 4 weeks.
- *Technique:* Hold infant upright with her or his feet touching a flat surface.
- *Infant steps up and down in place.*



Stepping or dancing reflex

ABNORMAL RESPONSE

- Poor response caused by hypotonia.

8. Rooting

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 3 to 6 months.
- *Technique:* Brush cheek near corner of mouth.
- *Infant turns head in direction of stimulus and opens mouth.*



Rooting reflex

ABNORMAL RESPONSE

- Prematurity or neurological problem may cause weak or absent response.

NEWBORN/INFANT REFLEXES

9. Sucking

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 10 to 12 months.
- *Technique:* Touch lips.



HELPFUL HINT

Don't check for rooting or sucking responses immediately after a feeding—they will be difficult to elicit.

- *Sucking motion occurs.*



Sucking reflex

ABNORMAL RESPONSE

- Weak or absent response associated with prematurity or neurological defect.

10. Swallowing

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth and lasts throughout life.
- *Technique:* Automatically follows sucking response during feeding.
- *Sucking and swallowing should occur without coughing, gagging, or vomiting.*

ABNORMAL RESPONSE

- Weak or absent response associated with prematurity or neurological problem.

11. Extrusion

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth and lasts 3 to 4 months.
- *Technique:* Touch tip of tongue.
- *Tongue protrudes outward.*



Extrusion reflex

ABNORMAL RESPONSE

- Absence may indicate neurological problem.
- Continued extrusion of large tongue associated with Down syndrome.

(continued)

NEWBORN/INFANT REFLEXES

12. Glabellar

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth.
- *Technique:* Tap on forehead.
- *Newborn blinks for first few taps.*



Glabellar reflex

ABNORMAL RESPONSE

- Persistent blinking with repeated taps indicates extrapyramidal problem.

13. Crawling

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 6 weeks.
- *Technique:* Place infant on abdomen.
- *Newborn attempts to crawl.*



Crawling reflex

14. Crossed Extension

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears at 2 months.
- *Technique:* Infant supine with leg extended. Stimulate foot.



A. Crossed extension



B. Stimulate the foot

- *Flexion, adduction, then extension of opposite leg.*

ABNORMAL RESPONSE

- Peripheral nerve damage causes weak response.
- Spinal cord lesion causes absent response.

NEWBORN/INFANT REFLEXES

15. Pull-to-Sit

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth.
- *Technique:* Pull infant to sitting position.
- *Head lags as infant is pulled to sitting position, but then infant is able to hold up head temporarily.*



Pull-to-sit reflex

ABNORMAL RESPONSE

- Inability to hold up head suggests prematurity or hypotonia.

16. Trunk Incurvation

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth; disappears in a few days to 4 weeks.
- *Technique:* With infant prone, run finger down either side of spine.
- *Flexion of trunk with hip moving toward stimulated side.*



Trunk incurvation reflex

ABNORMAL FINDINGS/RATIONALE

- Absent response indicates neurological or spinal cord problem.

17. Magnet

REFLEX/TECHNIQUE/NORMAL RESPONSE

- Present at birth.
- *Technique:* With infant supine, flex leg and apply pressure to soles of feet.
- *Extends legs against pressure.*



A. Flex the legs



B. Apply pressure to the soles of the feet

ABNORMAL FINDINGS/RATIONALE

- Breech birth may diminish reflex.
- Absent response caused by spinal cord problem.

Diagnostic/Screening Tests for Newborns

TEST	SIGNIFICANCE
<p><i>Phenylketonuria</i>: Measures amount of phenylalanine amino acids in blood.</p> <p><i>Galactosemia</i>: Transferase deficiency.</p>	<ul style="list-style-type: none"> ■ Test is collected after several days of feedings. ■ High phenylalanine levels can cause brain damage. ■ Elevated galactose and low fluorescence may result in mental retardation, blindness, or death from dehydration and sepsis. ■ Elevated leucine can result in acidosis, seizures, mental retardation, and death. ■ Elevated methionine can lead to mental retardation, seizures, and behavioral disorders. ■ Elevated 17-hydroxyprogesterone can lead to hyponatremia, hypokalemia, hypoglycemia, and ambiguous female genitalia. ■ Decreased activity of biotinidase on colorimetric assay causes mental retardation, skin changes, hearing and vision problems, even death. ■ Thyroxine is a thyroid hormone necessary for growth, development, and metabolism. ■ Low values are associated with brain defects. ■ Rh and ABO incompatibility can cause hemolysis and subsequent jaundice. ■ If mother is Rh– and newborn is Rh+, mother must be treated promptly to avoid sensitization to future Rh+ fetuses. ■ Detects presence of hemoglobin S. ■ Electrophoresis identifies whether infant has the trait or disease. ■ Identifies hearing disorders early. ■ Early virus detection can initiate early treatment that may enhance survival. ■ Accurate estimate of gestational age influences care management and special monitoring needs. ■ Full-term infant should have creases over entire sole of foot, full breast areolas, firm ear cartilage. ■ Males should have pendulous testes and deep scrotal rugae. ■ In females, the labia majora should cover the clitoris and labia minora.
Maple syrup urine disease	
Homocystinuria	
Congenital adrenal hyperplasia	
Biotinidase deficiency	
Thyroxine	
Blood type	
Sickle cell anemia	
<p>Audiometric testing</p> <p><i>Human immunodeficiency virus (HIV)</i>: Newborns of mothers with HIV are tested for virus.</p> <p><i>Gestational age</i>: Normal gestational age is 38 to 42 weeks. A newborn less than 38 weeks is premature. A screening test, such as the Ballard test, estimates fetal gestation.</p>	



Case Study Findings

Newborn Carmen's physical assessment findings include:

- **General Health Survey/Anthropometric Measurements**
 - Flexed head and extremities.
 - Head, 33 cm; chest, 31 cm; abdomen, 32 cm.
 - Length head to heel, 50 cm.
 - Weight, 7 lb, 6 oz.

■ Vital Signs

- Axillary temperature, 98°F
- Pulse, 144 beats per minute (BPM).
- Respirations, 44 and irregular.
- Blood pressure (BP), 64/38.

■ Apgars

- 8 and 10.

■ Integumentary

- Skin pink and warm with no jaundice.
- Lanugo on face and shoulders.
- Vernix caseosa in groin skin folds.
- One 4-cm Mongolian spot on each buttock.

■ Head/Face

- Positive molding, but skull appears symmetrical.
- Fontanels soft and flat.

■ Head, Eyes, Ears, Nose, and Throat (HEENT)

- Intact, positive red light reflex both eyes (OU).
- React to noise.
- Ears align with external canthus of eyes.
- Small amount of clear mucus in nose and mouth.
- Palate intact.
- Strong sucking reflex.

■ Respiratory

- Irregular, unlabored breathing.
- Occasional periods of apnea lasting no more than 5 seconds.
- Lungs clear.

■ Cardiovascular

- No cyanosis.
- Apical pulse, 144 regular.
- No murmurs.
- Positive femoral pulses.

■ Gastrointestinal

- Positive bowel sounds.
- Anus patent.
- Meconium stool.
- Positive anal reflex.
- Umbilical cord white with two arteries and one vein, intact with no drainage.

■ Genitourinary

- Voided.

■ Genitalia

- Pink and edematous.

■ Musculoskeletal

- 10 fingers and 10 toes.
- C curve of spine.
- No dimpling.
- Equal gluteal folds.
- Full range of motion (ROM) of extremities.
- No fractures or dislocations.

■ Neurological

- Good muscle tone.
- Strong cry.
- Positive newborn reflexes.



CRITICAL THINKING ACTIVITY #1

How would you adjust your assessment of Carmen to include the educational needs of her parents? What resources on the hospital unit would you use?



CRITICAL THINKING ACTIVITY #2

Mrs. Ramirez brings 3-day-old Carmen to the pediatric clinic. She states that Carmen nurses for 30 to 40 minutes every 3 to 4 hours but sleeps after each feeding. She is concerned that the baby sleeps 15 to 16 hours a day. Mrs. Ramirez states that she had an uncomplicated pregnancy and delivery and that she feels so good; she cannot understand why Carmen sleeps so much. What information about sleeping and feeding patterns would you offer Mrs. Ramirez?



Case Study Follow-Up

Carmen is now 6 months old. Mrs. Ramirez brings her to the pediatric clinic for her routine checkup. She states that last month Carmen was diagnosed with otitis media and was on an antibiotic for 1 week. Mrs. Ramirez is concerned that Carmen does not appear to be well fed and wonders if formula would help her gain weight. She states that she continues to enjoy her job as a social worker and appreciates her mother's and sister's help with child care while she works.

COMMON HEALTH PROBLEMS

As you gather information, consider some common health problems that affect newborns and infants.

Newborns

- Heart murmurs
- Respiratory distress syndrome
- Mother-to-child disease transmission (e.g., sexually transmitted diseases [STDs] such as HIV and syphilis)

Infants

- Otitis media
- Upper respiratory infections
- Diaper rash
- Allergies
- Colic

Performing the Infant Assessment

The initial assessment is performed immediately after birth, while the newborn is still in the hospital. It serves as a baseline. The subsequent history and physical examination are performed when the infant visits the clinic or pediatrician's office for well-baby checkups.

Infant Health History

A complete health history can be obtained from the newborn's chart and from a parent or guardian. Review the chart, paying special attention to prenatal care, gestational or family health problems, and supports. Then, ask the parent or guardian about developmental milestones, and discuss safety issues and immunizations. The parent or guardian is critical in obtaining a full health history. Encourage discussion of concerns and questions. The interview is also an ideal teaching opportunity.

Developmental Milestones

Ask the parent or guardian about milestones, such as when the infant first rolled over. This information will identify delays that must be addressed.

Home Safety

Ask about safety in the home, including fall prevention, fire and burn prevention, gun and poison control, and car seat restraints. Take every opportunity to educate parents on safety issues.



ALERT

Congenital anomalies and prematurity with low birth rate are the top two leading causes of death in infants. Unintentional injuries are the leading cause of death in children over age 1 year, with automobile accidents being the number-one cause.

Immunizations

Immunizations are critical for primary prevention of childhood diseases. Discuss the immunization schedule with the parent or guardian. The following immunizations are recommended:

- *Hepatitis B*: First dose within 2 months of age, second dose 1 month after first dose, and third dose 6 months after second dose.
- *Diphtheria, pertussis, tetanus toxoid (DPT)*: Doses at 2 months, 4 months, and between 6 and 18 months.
- *Haemophilus Influenzae type B (Hib)*: Doses at 2 months, 4 months, and 6 months (check with manufacturer if third dose at 6 months is required).
- *Inactivated polio vaccine (IPV)*: Doses at 2 months, 4 months, and between 6 and 18 months.
- *Pneumococcal conjugate vaccine (PCV)*: Doses at 2 months, 4 months, and between 6 and 18 months.

For the most recent information about vaccines, visit the National Immunization Program Home Page at <http://www.cdc.gov/nip/>, or call the National Immunization Hotline at 800-232-4636 (English and Spanish).

Review of Systems

The last part of your history is a review of systems (ROS). This review will help you identify normal physiological changes as well as alert you to abnormal findings.

Review of Systems

AREA/QUESTIONS TO ASK

General Health Survey

- How is your infant doing now?

Body Weight

- Has he or she been gaining weight?

Integumentary

- Does the infant have good skin turgor?
- Does the skin appear healthy?

- Do hair and nails appear healthy or brittle?

RATIONALE/SIGNIFICANCE

- Determining the parents' view of their infant's general well-being provides a baseline for you to begin your assessment.

- Weight gain within normal parameters is an indicator of nutritional status.

- Skin tenting denotes dehydration.

- Scaly skin may be caused by an underlying medical condition or by dryness from soaps or lotions.

- Brittle hair and nails may be caused by poor protein intake.

AREA/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p><i>Head and Neck</i></p> <ul style="list-style-type: none"> ■ Is head shape symmetrical? ■ Can infant hold head upright while sitting and move head from side to side? 	<ul style="list-style-type: none"> ■ Symmetry is normal. ■ Head control is a necessary motor development milestone.
<p><i>Eyes, Ears, Nose, and Throat</i></p> <ul style="list-style-type: none"> ■ Can infant recognize toys you give her or him? ■ Do eyes focus symmetrically? ■ Is there eye exudate? <ul style="list-style-type: none"> ■ Does infant turn head when his or her name is called? ■ Can she or he vocalize? <ul style="list-style-type: none"> ■ Is there exudate from the nose? 	<ul style="list-style-type: none"> ■ Refer infants with signs of visual difficulties to specialist. ■ Infants are highly susceptible to eye infections and may require medications. ■ Ability to hear and speak is critical for development. If you notice signs of hearing/speaking problems, refer infant to specialist. ■ Nose exudate may signal infection or allergies or may be normal mucus production.
<p><i>Respiratory</i></p> <ul style="list-style-type: none"> ■ Is breathing labored? <p><i>Cardiovascular</i></p> <ul style="list-style-type: none"> ■ Is infant active? ■ Is skin ever blue tinged? 	<ul style="list-style-type: none"> ■ Monitor abnormal breathing patterns carefully. ■ Infants should be active and lively when awake.
<p><i>Gastrointestinal</i></p> <ul style="list-style-type: none"> ■ How is the infant's appetite? ■ What does he or she usually eat and drink each day? <ul style="list-style-type: none"> ■ How many bowel movements a day does she or he have? 	<ul style="list-style-type: none"> ■ Lethargy may be related to cardiac difficulties, of which cyanosis is a warning sign. ■ Good appetite denotes good health.
<p><i>Genitourinary</i></p> <ul style="list-style-type: none"> ■ Is urination pattern consistent with fluid intake? ■ How many diaper changes a day? 	<ul style="list-style-type: none"> ■ Adequate daily nutrient intake is necessary for growth. ■ Regular bowel movements suggest normal GI activity. ■ Adequate urination pattern denotes normal kidney function. ■ Usually about eight diapers a day is normal, depending on fluid intake.
<p><i>Musculoskeletal</i></p> <ul style="list-style-type: none"> ■ Does infant roll over, sit unsupported, transfer toy from one hand to the other? ■ Can she or he creep forward or backward on tummy? 	<ul style="list-style-type: none"> ■ Body movements should be within developmental norms.
<p><i>Neurologic</i></p> <ul style="list-style-type: none"> ■ Have you noticed any unusual movements such as tremors? ■ Have you noticed any problems with sucking, swallowing? 	<ul style="list-style-type: none"> ■ Tremors or other unusual movements may indicate seizures. ■ Diminished sucking or swallowing reflexes may indicate an underlying neurological problem.
<p><i>Infections</i></p> <ul style="list-style-type: none"> ■ Has infant had any colds, fevers, infections? 	<ul style="list-style-type: none"> ■ Usual infant infections may not be preventable. However, teach parents infection control methods.
<p><i>Development</i></p> <ul style="list-style-type: none"> ■ Does infant prefer to play with people rather than toys? ■ Does he or she search for an object that is out of sight? ■ Is she or he afraid of strangers? 	<ul style="list-style-type: none"> ■ Deviations from these developmental milestones may impair social activities.
<p><i>Relationships</i></p> <ul style="list-style-type: none"> ■ How do you feel about the new baby and about being a mother or father? 	<ul style="list-style-type: none"> ■ Mother may cry, feel irritable, and have loss of appetite and sleeplessness for first 10 days or so after delivery. If these problems persist for more than 2 weeks, they may signal postpartum depression and warrant referral.



Case Study Findings

Your health history of infant Carmen has concluded. Her developmental, safety, and immunization findings include:

■ Developmental

- Sits.
- Bears weight on hands, stands with assistance.
- Rolls over.
- Controls head.
- Grasps objects.
- Is teething.
- Imitates sounds.
- Is attached to parents.
- Has separation anxiety.

■ Safety

- Smoke and carbon monoxide detectors.
- Safety gates.
- Cabinet locks.
- Electrical outlet caps.
- Car seats.
- Medications and cleaning solutions out of reach.

■ Immunizations

- Hepatitis B 1 and 2.
- DPT 1 and 2.
- Polio 1 and 2.
- HiB 1 and 2.
- PCV 1 and 2.

Infant Carmen's ROS reveals:

■ General Health Survey

- Very robust and active infant.
- Healthy appearance.

■ Body Weight

- 16 lb, gained 2 lb since last month's visit for otitis media.

■ Integumentary

- No scaly skin.

■ Head and Neck

- Easily holds head upright and in alignment.

■ Eyes

- Normal response to visual stimuli, no drainage.

■ Ears

- Normal sound recognition and making sounds.
- No apparent abnormalities.
- Otoscope examination indicated because of recent otitis media.

■ Respiratory

- No breathing problems.

■ Cardiovascular

- No apparent distress.
- Active when awake.

■ Gastrointestinal

- Exclusively breastfed and nurses every 3 to 4 hours during the day, with daily bowel movements, yellow in color.

■ Genitourinary

- Normal urination pattern noted.
- Eight diapers a day.

■ Musculoskeletal

- Attempts to crawl (normal finding).
- Rolled over and sat up on her own (normal finding).

■ Neurological

- No abnormal movements.
- Nurses well.
- No problems.

■ Immunologic

- No infection aside from otitis media.

■ Developmental

- Laughs easily with mother but is fearful of examiner.
- Normal age-related behavior observed.

■ Relationships

- Warm, caring responses noted toward baby from parents.
- Both parents say they enjoy parenthood very much.



Case Study Evaluation

Before you proceed to the physical assessment, record what you have learned about infant Carmen's health history so far.

Infant Physical Assessment

After the history, perform the physical examination on the infant, considering the important role that the parent or guardian plays in this process. Physical examinations should be conducted in a warm environment. The alert, well-fed newborn is usually a most cooperative patient. Involve the family whenever possible because this is a fine opportunity for education and support. Infants after the age of 6 months will most likely fear the practitioner. Therefore, conduct as much of the examination as possible with the child in the mother's lap.

Performing a Head-to-Toe Physical Assessment

Next, perform a head-to-toe physical assessment of the infant.

(Text continued on page 883)

Performing an Infant Head-to-Toe Physical Assessment

AREA/SYSTEM/NORMAL VARIATIONS ABNORMAL FINDINGS/RATIONALE

Anthropometric Measurements

Head and Chest

- Apply tape measure around widest part of head, just above eyebrows.
- Measure chest at nipple line.

Height and Weight

- Measure length from head to heels.
- Weigh without clothing or diaper.
- Increases by 1.5 cm each month for first 6 months and by 0.5 cm per month until age 12 months. Chest circumference equals head circumference by 12 months. Height should increase 50 percent by age 12 months. Birth weight should double by 6 months and triple by 12 months.

Integumentary

Skin

- Inspect skin appearance and skin folds.
- Test skin turgor.
- Take axillary temperature.
- Inspect skin appearance throughout your physical exam.
- Skin pigmentation varies. Newborns of dark-skinned parents may appear light-skinned at birth. True skin color develops by 3 months of age.
- Good skin turgor in newborns and infants is a sign of adequate hydration.

- Axillary temperature range for newborns and infants is 35.9°C to 36.7°C (96.6°F to 98.0°F).



ALERT

Never use a mercury thermometer on a newborn or infant. Mercury is highly toxic.

- Greater-than-normal head circumference may indicate hydrocephalus. Smaller-than-normal head circumference is microcephaly.

- Anthropometric measurements deviating from normal may be caused by underlying disease or inadequate eating or nutritional pattern.

- Persistent cyanosis in a warm infant is never normal and requires immediate referral. Other deviations include:

- *Diaper rash (diaper dermatitis)*: Red, excoriated skin. Secondary yeast infection (*Candida*) causes bright, round, scaling patches.

- *Eczema (atopic dermatitis)*: Dry, itchy patches on face and skin folds in infants with allergies.

- *“Cradle cap” (seborrheic dermatitis)*: Flat, greasy scales on scalp that may be caused by infrequent shampooing. Some cultures fear touching fontanel area.

- *Depressed fontanels and skin tenting*: Signs of severe dehydration.

- Low temperature suggests hypothermia.

- High temperatures can cause seizures in newborns and infants.

HEENT

Head/Face/Neck

- Inspect face, fontanels.
- Gently palpate fontanels.
- *Face and skull symmetrical. Fontanels flat and soft. At birth, anterior fontanel is 2.5 to 4 cm across; it closes between 12 and 18 months of age. Posterior fontanel is initially 2 cm and closes by 3 months.*
- Neck is short and should rotate left to right.

- Sunken fontanel associated with dehydration. Bulging fontanel associated with intracranial pressure.

(continued)

Performing an Infant Head-to-Toe Physical Assessment (continued)

AREA/SYSTEM/NORMAL VARIATIONS	ABNORMAL FINDINGS/RATIONALE
<p>Eyes</p> <ul style="list-style-type: none"> ■ Inspect external eye, eyelids, lacrimal ducts, conjunctiva, sclera, pupils. ■ Test corneal/blink reflex. ■ Test visual acuity using the DDST II. ■ <i>Bilateral blinking.</i> ■ <i>Corneal reflex positive.</i> ■ <i>Transient strabismus common.</i> ■ <i>No tearing in first month.</i> <ul style="list-style-type: none"> ■ <i>Sclera may be blue tinged.</i> ■ <i>Pupils constrict with light and are round and equal in size.</i> ■ <i>Infants have full binocular vision.</i> <p>Ears</p> <ul style="list-style-type: none"> ■ Inspect external and internal ear. ■ Test hearing. ■ <i>Infant should respond to noise.</i> ■ <i>Pinna flexible.</i> 	<ul style="list-style-type: none"> ■ Continued strabismus after 6 months is abnormal. ■ Lack of tears after 2 months may be caused by clogged lacrimal ducts and requires medical attention. ■ Fixed or dilated pupils indicate neurological problem. <ul style="list-style-type: none"> ■ Lack of response to noise may indicate hearing problem.
<div style="border: 1px solid black; padding: 5px;">  <p>HELPFUL HINT</p> <p>To perform otoscopic exam, place infant flat and restrain arms above head. Pull auricle down and back to observe tympanic membrane.</p> </div>	
<p>Nose</p> <ul style="list-style-type: none"> ■ Infants are nose breathers. Check patency by closing one nostril, then the other. ■ <i>Thin white mucus discharge and sneezing are normal.</i> 	<ul style="list-style-type: none"> ■ Flaring of nares is sign of respiratory distress. ■ Bloody discharge or large amount of nasal secretions may obstruct nares.
<p>Mouth/Throat</p> <ul style="list-style-type: none"> ■ Inspect oral cavity: palate, tongue, gums, tonsils (use light and tongue blade). ■ <i>White nodules (Epstein's pearls) may be found on hard palate and usually disappear by 3 months.</i> ■ <i>Tongue should not protrude from mouth.</i> ■ <i>No coating in mouth.</i> ■ <i>Suck should be strong.</i> ■ <i>First primary teeth usually appear at 6 to 8 months.</i> 	<ul style="list-style-type: none"> ■ Protruding tongue associated with congenital disorders, such as Down syndrome or hypothyroidism.
<p>Chest/Back/Shoulders</p> <ul style="list-style-type: none"> ■ Inspect back and spine. ■ <i>Convex spinal curvature (C curve).</i> ■ <i>No blemishes or skin openings.</i> ■ <i>By 12 to 18 months, lumbar curve develops and lordosis is present as walking begins.</i> 	<ul style="list-style-type: none"> ■ Limited ROM may result from injury. ■ Dimpling in spine may be associated with neural tube defects.
<p>Respiratory</p> <p>Chest</p> <ul style="list-style-type: none"> ■ Percuss chest lightly. ■ Palpate back with two fingers while baby is crying. ■ Auscultate lungs. 	

AREA/SYSTEM/NORMAL VARIATIONS

ABNORMAL FINDINGS/RATIONALE

**HELPFUL HINT**

Use direct percussion.

- Normal respiratory rate 25 to 50 breaths per minute. Apnea is less common in infants than in newborns. Normal breath sounds are more bronchial to bronchovesicular in infants. Infants are abdominal breathers. Anteroposterior:lateral diameter is equal.

Cardiovascular*Heart/Peripheral Pulses*

- Auscultate heart sounds, heart rate.
- Test capillary refill.
- Palpate peripheral pulses.

**HELPFUL HINT**

Use a stethoscope with a small diaphragm and perform exam when infant is quiet.

- Apical pulse felt in fourth or fifth intercostal space just medial to midclavicular line.
- Heart rate range 80 to 160 BPM.
- Capillary refill < 1 second.
- Peripheral pulses present.

Gastrointestinal**Abdomen**

- Inspect abdomen, note hernias.
- Auscultate bowel sounds.
- Gently palpate abdomen.
- Gently percuss (direct) abdomen.
- Bowel sounds present.
- Bowel soft.
- Tympany may be heard because of air swallowing.
- Umbilicus flat.
- Liver edge 1 to 2 cm below right rib cage (costal margin).

**ALERT**

Apnea > 15 seconds accompanied by decreased heart rate is abnormal.

- Stressful breathing with flaring nares and sighing with each breath are signs of respiratory distress and require immediate attention.
- Inspiratory stridor, expiratory grunts, retractions, paradoxical (seesaw) breathing, asymmetrical or decreased breath sounds, wheezing, and crackles are abnormal.
- Depressed sternum may affect normal respiration.

- Abnormal heart rate range requires attention.
- Murmurs accompanied by cyanosis may indicate congenital heart defects.
- Capillary refill times longer than 2 seconds may indicate dehydration or hypovolemic shock.
- Evaluate newly discovered murmurs.
- Infant eating poorly may have cardiovascular problem.

- Umbilical hernias > 2 cm wide may require further evaluation.
- Abdominal pain may indicate childhood illnesses.
- Enlarged liver or palpable spleen may indicate disease.

(continued)

Performing an Infant Head-to-Toe Physical Assessment (continued)

AREA/SYSTEM/NORMAL VARIATIONS	ABNORMAL FINDINGS/RATIONALE
Extremities	
<ul style="list-style-type: none"> ■ Inspect (hands, arms, feet, legs, hips)/test ROM. 	<ul style="list-style-type: none"> ■ Inadequate ROM may indicate congenital malformation or birth injury or may result from pulling or lifting infant.
<ul style="list-style-type: none"> ■ <i>Feet flat.</i> 	
<ul style="list-style-type: none"> ■ <i>Legs equal in length.</i> 	
<ul style="list-style-type: none"> ■ <i>Transfers objects from one hand to other by 7 months.</i> 	
<ul style="list-style-type: none"> ■ <i>Crawls and sits unsupported by 7 months.</i> 	<ul style="list-style-type: none"> ■ Inability to meet developmental milestones may indicate neurological or environmental deficits.
<ul style="list-style-type: none"> ■ <i>Pulls to standing position and holds on to furniture by 11 months.</i> 	
Genitourinary	
Male	
<ul style="list-style-type: none"> ■ Note infant's sex. Inspect penis, urethra 	<ul style="list-style-type: none"> ■ Ambiguous genitalia abnormal.
<ul style="list-style-type: none"> ■ <i>In uncircumcised male infant, foreskin may not be fully retractable until 1 year.</i> 	<ul style="list-style-type: none"> ■ Phimosis (tight foreskin) can constrict penis. Instruct parents on gentle retraction of foreskin to prevent phimosis.
<ul style="list-style-type: none"> ■ <i>Urethra midline.</i> 	
<ul style="list-style-type: none"> ■ <i>Urine flow straight and strong.</i> 	<ul style="list-style-type: none"> ■ Weak urine stream or dribbling suggests stricture at urinary meatus.
<ul style="list-style-type: none"> ■ Palpate testes. 	<ul style="list-style-type: none"> ■ Solid scrotal masses are abnormal. Hernias present as scrotal masses.
<ul style="list-style-type: none"> ■ <i>No palpable masses in testes.</i> 	<ul style="list-style-type: none"> ■ Testes not palpable indicates undescended testicles.
<ul style="list-style-type: none"> ■ <i>Two testes should be palpated in scrotum or brought down from inguinal ring.</i> 	<ul style="list-style-type: none"> ■ If scrotum is swollen, transilluminate to determine if fluid (hydrocele) present.
Female	
<ul style="list-style-type: none"> ■ Inspect external genitalia. 	<ul style="list-style-type: none"> ■ Ambiguous genitalia abnormal.
<ul style="list-style-type: none"> ■ <i>Note infant's sex.</i> 	<ul style="list-style-type: none"> ■ Blood-tinged fluid from vagina abnormal after 1 week.
<ul style="list-style-type: none"> ■ <i>External genitalia of female infant should be pink and moist.</i> 	<ul style="list-style-type: none"> ■ Vaginal discharge or labial redness or itching may be caused by diaper or soap irritation or sexual abuse.
Anus and Rectum	
<ul style="list-style-type: none"> ■ Inspect anus. 	
<ul style="list-style-type: none"> ■ <i>In infants, bowel movements may occur with each feeding. By 1 year, they occur once or twice a day.</i> 	<ul style="list-style-type: none"> ■ Watery stools and explosive diarrhea indicate infection. Constipation or hard stools indicate inadequate hydration or nutrition.
<ul style="list-style-type: none"> ■ <i>Breastfed babies have stools that are mustard colored and soft.</i> 	
<ul style="list-style-type: none"> ■ <i>Formula-fed babies have stools that are yellowish-green and more formed.</i> 	
Neurological	
<ul style="list-style-type: none"> ■ Inspect and test muscle strength, function, sensory function, reflex movements. 	<ul style="list-style-type: none"> ■ Delays in motor or sensory activity may indicate brain damage, mental retardation, illness, malnutrition, or neglect.
<ul style="list-style-type: none"> ■ <i>Infant's body stays erect when you support him or her with both hands under axilla. Motor control develops from head to toe. Infant opens eyes to noise and responds to touch.</i> 	<ul style="list-style-type: none"> ■ Asymmetrical posture or spastic movements need further evaluation.
<ul style="list-style-type: none"> ■ <i>Presence of infantile reflexes denotes healthy neurological system.</i> 	<ul style="list-style-type: none"> ■ Maintenance of infant reflexes past usual age is abnormal.

Performing Diagnostic/Screening Tests

Depending on the infant's condition, certain diagnostic and screening tests are performed. Mandatory screening varies from state to state.

- **Developmental screening:** Should occur regularly throughout childhood. DDST-2 assesses skills in areas of personal/social, fine motor, language, and gross motor function. If delays are noted, refer the child to a specialist for further testing. Medications, malnutrition, neurological, and emotional conditions affect test results.
- **Urine screening:** Dipstick screening test is usually sufficient. Any abnormalities require a full urinalysis.
- **Hemoglobin or hematocrit screening:** A hemoglobin level below 10 g/dL is low and the infant is considered anemic. A hematocrit level below 29 percent is low.
- **Lead screening:** Perform only if environmental exposure is suspected. Elevated blood lead levels can cause neurological damage.



Case Study Findings

Infant Carmen's physical assessment findings include:

- **General Health Survey**
 - Well-developed, robust, alert 6-month-old girl sitting upright in mother's lap.
- **Anthropometric Measurements**
 - Length, 66 cm (26 inches).
 - Weight, 7.25 kg (16 lb).
 - Head circumference, 40.5 cm (16 inches).
- **Vital Signs**
 - Temperature, 36.7°C (98.0°F).
 - BP, 113.
 - Respirations, 30, lungs clear.
- **Integumentary**
 - Skin pink and warm to touch.
 - Hair shiny and clean.
- **Head and Face**
 - Head and skull symmetrical.
 - Posterior fontanel closed.
 - Anterior fontanel flat and soft.
- **HEENT**
 - Eyes clear, follows objects.
 - External ears intact, piercing for earrings noted.
 - Internal ear intact; no redness or exudates.
 - Responds to noise and when name is called.
 - Mouth and palate intact, two teeth noted.
- **Chest, Neck, Shoulders, and Back**
 - Intact.
- **Cardiovascular**
 - Positive peripheral pulses.

Gastrointestinal

- Abdomen soft.
- Positive bowel sounds (BS+).
- Several soft, yellow stools daily.

Extremities

- Legs, feet, arms, and hands intact with symmetrical movement.

Genitourinary

- Voids with every feeding.
- Labia and buttocks excoriated (diaper rash).

Musculoskeletal

- Barlow-Ortolani maneuver negative.
- Equal gluteal skin folds.
- Sits unassisted.
- Attempting to crawl.
- Transfers object from one hand to other.

Reflexes

- Tonic neck present.
- Plantar grasp present.
- Babinski present.
- Suck present.

Relationships

- Smiling baby, comfortable with mother.

Additional Data

- Immunization and assessment visits on schedule.
- Nutrition status is good and within normal for age.
- Mother is breastfeeding but wonders if Carmen's weight is normal.



CRITICAL THINKING ACTIVITY #3

Unfortunately, Mrs. Ramirez's mother and sister are moving to another state. They have both helped with child care and are a major source of support. List community resources that would be of assistance to the family.



Case Study Analysis and Plan

Now that you have completed a thorough assessment of infant Carmen, document the key history and physical examination findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Analyze the data on infant Carmen and develop nursing diagnoses. Then cluster the supporting data for each diagnosis. Because the caregiver affects the infant, be sure to include the family in the plan.

1. Impaired Tissue Integrity
2. Risk for Infection
3. Risk for Imbalanced Nutrition: more than body requirements
4. Potential for Social Isolation
5. Health-Seeking Behaviors



CRITICAL THINKING ACTIVITY #4

Now that you have identified some nursing diagnoses for infant Carmen, select one and develop a nursing care plan and a brief teaching plan for her mother, including learning outcomes and teaching strategies.

RESEARCH TELLS US

Kernicterus, a form of icterus neonatorum, may occur the 2nd to 8th day after birth. It results from the breakdown of hemoglobin with an increase in bilirubin and can lead to permanent brain damage. In the past, kernicterus and bilirubin encephalopathy were used interchangeably, but they are now distinguished as acute, reversible, bilirubin encephalopathy, and chronic, irreversible, kernicterus.

In 2001, the Centers for Disease Control and Prevention (CDC) called for rigorous detection and management of hyperbilirubinemia (jaundice) in newborns. Breastfed newborns less than 38 weeks' gestation are at higher risk for hyperbilirubinemia, and visual determination of jaundice is unreliable. The American Academy of Pediatrics has developed guidelines designed to identify and manage hyperbilirubinemia in newborns (≥ 35 weeks' gestation).

Four key recommendations are aimed at reducing the risk of severe hyperbilirubinemia, while at the same time reducing maternal anxiety, maintaining breastfeeding, and reducing unnecessary cost and treatments. Major points of the recommendations include:

1. *Promote and support breastfeeding.* Because inadequate calories or fluid from inadequate breastfeeding may contribute to hyperbilirubinemia, breastfeeding is encouraged 8 to 12 times a day for the first few days after birth.

2. *Perform a thorough newborn assessment prior to discharge to identify potential cases of severe hyperbilirubinemia.* Assess for jaundice every 8 to 12 hours; if present, test TcB (transcutaneous bilirubin) or TSB (total serum bilirubin). Develop nursery protocol to identify hyperbilirubinemia. Identify risk for severe hyperbilirubinemia.
3. *Perform a systemic assessment of all infants prior to discharge.* Give written and verbal instructions to parents on how to monitor and identify jaundice. Schedule follow-up after discharge within 3 to 5 days after birth (bilirubin likely to be at highest level).
4. *Assess and determine risk for severe hyperbilirubinemia on newborns prior to discharge.* Follow-up is dependent on identified risk for hyperbilirubinemia. During the follow-up visit, assess the infant's weight, percent change in weight from birth, adequacy of intake, output (wet diapers and stools), and presence or absence of jaundice. If hyperbilirubinemia is present, phototherapy or exchange transfusions may be used to prevent severe hyperbilirubinemia and kernicterus.

Nurses have a key role not only in assessing hyperbilirubinemia but also in educating parents in the assessment and treatment of jaundice.

Source: Holcomb, S. (2005). Managing jaundice in full-term infants. *The Nurse Practitioner The American Journal of Primary Health Care* 30(1):6-12.

HEALTH CONCERNS

Health concerns for infants focus on health promotion, such as maintenance of regular well-baby assessments and immunization schedules, supporting and

educating parents in their critical role as infant caregivers, and using community resources to enhance the family unit.


S U M M A R Y

- A thorough newborn and infant assessment is a fundamental component in health promotion and disease prevention for the entire family.
- The baby clinic visit may be the only access point to the healthcare system for the family.
- Because infant health depends on family health, incorporating the total family is critical to your approach to health assessment.
- Encourage parents to keep all scheduled child assessment visits, and take every opportunity to make the necessary referrals for family members.

R E V I E W Q U E S T I O N S

1. Jane Johnson, age 30, delivered an 8-pound healthy baby boy. You note on the assessment that the baby has acrocyanosis. Where would you expect to see acrocyanosis?
 - a. Periphery
 - b. Oral mucosa
 - c. Conjunctiva
 - d. Tongue
2. You check baby Johnson's Apgar scores. Which of the following ranges reflects a healthy state?
 - a. 0–3
 - b. 4–7
 - c. 8–10
3. As you perform the initial physical assessment of baby Johnson, which finding would be cause for concern and warrant medical follow-up?
 - a. Vernix caseosa
 - b. Lanugo hair
 - c. Desquamation of skin
 - d. Jaundice
4. As you auscultate baby Johnson's heart, which heart rate would be cause for concern and warrant medical follow-up?
 - a. 100 BPM
 - b. 120 BPM
 - c. 140 BPM
 - d. 160 BPM
5. You note color changes on Baby Johnson associated with position changes. These color changes are referred to as:
 - a. Harlequin sign.
 - b. Cutis marmorata.
 - c. Acrocyanosis.
 - d. Plethora.
6. You assess the fontanel. At what age would you expect the anterior fontanel to close?
 - a. At birth
 - b. 1 month
 - c. 6–12 months
 - d. 12–18 months
7. You are testing the newborn's reflexes. How would you test the glabellar reflex?
 - a. Stroke lateral aspect of the foot
 - b. Tap on forehead
 - c. Touch the tip of the tongue
 - d. Stroke the newborn's cheek
8. You are testing the newborn's reflexes. What would be the normal response for the magnet reflex?
 - a. Eye blinking
 - b. Toe flexion
 - c. Legs extend
 - d. Tongue protrudes
9. Baby Johnson has a positive Ortolani maneuver. What does this finding suggest?
 - a. Congenital hip dislocation
 - b. Spina bifida
 - c. Scoliosis
 - d. Talipes
10. You auscultate baby Johnson's lungs. Which type of breath sound is normally heard in newborns?
 - a. Tracheal
 - b. Bronchial
 - c. Vesicular

CHAPTER 25



Assessing the Toddler and Preschooler

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify normal physiological changes of toddler and preschooler
- Discuss various cultural influences on children
- Identify developmental tasks for toddler and preschooler
- Identify health issues and risks for toddler and preschooler
- Identify pertinent history data included in a toddler and preschooler history
- Obtain a toddler and preschooler health history
- Describe additional physical assessment components specific to a toddler and preschooler assessment
- Perform a toddler and preschooler assessment
- Document toddler and preschooler assessment findings
- Identify actual/potential health problems stated as nursing diagnoses with supporting data

The terms **toddler** and **preschooler** refer to children between the ages of 1 and 5. This is a time of challenge, discovery, and learning control. Specifically, the toddler period extends from 12 to 36 months, and the preschool period from 3 to 5 years. The toddler years are often referred to as the *terrible 2s* as the child struggles to gain control. During the preschool years, the child matures cognitively and emotionally in preparation for entering school. This chapter will help you effectively assess this age group by familiarizing you with the physical, psychosocial, and cognitive developmental changes that occur.

Before you begin the health assessment, ask yourself:

- How does assessment of the toddler and preschooler differ from assessment of other patients?
- What approach is best for assessing the toddler and preschooler?
- What health concerns should I keep in mind as I proceed with the assessment?

Developmental Changes

Physical Changes

During the toddler and preschool years, physical growth continues at a steadier pace than in infancy. Body systems also develop more mature functioning. The specific physical changes that occur are presented during the physical examination section of this chapter.

Psychosocial and Emotional Changes

During the toddler years, the child struggles to develop a sense of autonomy. He or she works to gain control over bodily functions, tolerate separation from parents, differentiate self from others, develop socially acceptable behavior and verbal communication skills, become less egocentric, and tolerate delay in gratifying needs. Unsuccessful attempts at developing autonomy result in doubt and shame. The negativism associated with this period explains the term the *terrible 2s*, when “no” seems to be the only word in the child’s vocabulary. But even though the child is trying to achieve control, he or she also looks for consistency within the environment. This gives the child a sense of safety and comfort. During the toddler and preschool years, the child also recognizes sexual differences and roles. Sex role imitation, such as “dressing up,” is common during these years. Once the toddler has developed autonomy, he or she moves to the next stage, which Eric Erikson calls *initiative versus guilt*. This is a time of playing and learning. If the child goes beyond set limits, guilt results. The superego, or con-

science, is in its early developmental stages, and the ability to tolerate a delay in satisfying needs reflects ego development. During the preschool years, the superego continues to develop. Behavior is seen as good or bad based on the outcome—reward or punishment.

Rapid cognitive development also occurs during this time. The child develops new skills, constantly building on what she or he has learned. The toddler begins to see causal and spatial relationships, develops object permanence, and imitates household activities and sex-role behavior. Preoperational thought processes develop. During the preschool years, language skills continue to develop, and play becomes an important means of expression and learning.

As the child approaches the preschool years, he or she begins to have some concept of a Supreme Being, having been introduced to the idea at an early age through bedtime or mealtime prayers. Preschoolers’ perceptions of the Supreme Being are concrete and based on parental influence.

Health Issues and Risk Factors

As a professional nurse, you are in an ideal position to screen for risk factors and teach parents health prevention. As you proceed with the assessment, stay alert for the following problems that are common in toddlers and preschoolers.

Toilet Training

The child needs to be physiologically and psychologically ready for toilet training, and this usually occurs between 18 and 24 months. Usually, the child gains bowel control before bladder control, and full bladder control may not be achieved until age 5.

Sibling Rivalry

Jealousy is a normal reaction in a toddler or preschooler, who may have trouble understanding and accepting a new baby. Talking to children and letting them help prepare for the new baby helps minimize conflict. Because the toddler or preschooler may act out, close supervision is needed once the baby comes home.

Temper Tantrums and Negativism

As the child tries to assert her or his independence, she or he may act out by having temper tantrums and holding breath. Although ignoring this behavior is usually the best approach, emphasize to parents that the child needs to be protected from injury. Negativism (saying “no”) can be minimized by avoiding yes-or-no choices.

Fears, Stress, and Regression

A child's fears may be real or imagined. Helping the child deal with fears may help resolve them. Usually, imaginary fears disappear by the time the child reaches school age. Even young children can experience stress. A variety of situations, such as a new baby, being hospitalized, or fighting among family members, can cause stress. But because young children have not yet developed effective skills to deal with stress, they may regress (return to an earlier reaction) when confronted with stressful situations. Regression is often seen when the child is hospitalized.

Communicable Diseases

Many communicable diseases can be prevented by proper immunizations. Provide parents with written information on immunization schedules. The hand-mouth activity common with toddlers increases their risk for various intestinal parasitic diseases and stomatitis. Emphasize the importance of handwashing.

Injuries and Poisoning

Accidents are the leading cause of deaths for all children over the age of 1. Motor vehicle injuries are the leading cause. Car seats or restraints should be used by all children. Car restraints need to be appropriate for the age and size of the child, and children under the age of 12 should never be in the front seat.

The inquisitiveness of toddlers and preschoolers increases the risk for accidents and ingestion of toxic substances. Safety is of the utmost importance. Make sure that parents have the number for the Poison Control Center. Also advise them to have syrup of ipecac in their medicine cabinets in case Poison Control advises them to induce vomiting. Discuss how to childproof the home to decrease risks. Also discuss the use of bicycle helmets. Emphasize that supervision is the best prevention.

Child Abuse and Neglect

Abuse or neglect can occur at any age, but younger children are dependent on parents for all their needs, both physical and psychological, and are therefore more vulnerable. See Guidelines for Assessing Abuse in Appendix C.



ALERT

A sudden change in a child's behavior may reflect an underlying problem that warrants further investigation.

Recognizing Abuse

Abuse can occur at any age and comes in many different forms. It can take the form of neglect, physical injury, sexual abuse, or psychological abuse. Be alert for the following signs and symptoms of abuse or neglect in children:

- Obvious physical signs of abuse or neglect.
- Repeated emergency room visits.
- Siblings blamed for injury.
- Inconsistent account of how injury occurred.
- Report of abuse by child.
- Inappropriate response by child or parent to injury.
- Inconsistency between physical findings and cause of injury.
- Inconsistency between injury and child's developmental level.
- Previous history of abuse.

Introducing the Case Study



Travis Harrison, 3 years old, is scheduled for his annual checkup. He sits quietly on his mother's lap but looks wide eyed and anxious as you approach to bring him into the exam room

Performing the Toddler and Preschooler Assessment

At this age, the assessment is performed with the parent or guardian present. Even so, you need to explain what you are doing and why. Because the toddler thinks in a very concrete way, keep your explanations very simple. Toys, such as hand puppets, or games can be helpful when examining the toddler or preschooler. If some procedures might cause discomfort—such as the otoscopic examination if the child has an earache—leave them until the end.

As you perform the exam, keep in mind the normal developmental changes affecting this age group. As with all assessments, the process involves subjective and objective data collection, nursing diagnoses, planning, interventions, and evaluation.

Health History

Although the parent will provide most of the history, you may want the child's response to a few questions. For instance, if there is an acute problem, ask the child to

describe it in his or her own words. As you perform the assessment, also look for changes in the child's behavior that might reflect his or her health status.

Biographical Data

Begin with the biographical data. Take note of where and with whom the child resides. Also note who is listed as contact person—it may not be the parent. Do not assume anything.

Current Health Status

Ask about the health status of the child. Has she or he been healthy, or is there an acute problem? Be sure to ask about all current prescription and over-the-counter (OTC) medications the child may be taking.

Past Health History

Besides asking specific health history questions, ask the mother about her pregnancy and the delivery of the child. Problems at this time may affect the child's growth and development. Also, be sure to ask whether the child's immunizations are current. (See Health Concerns, p. 906, for required immunizations.)

Family History

Ask about heart disease, stroke, cancer, and other serious health conditions within the family.

Review of Systems

Go through every system, asking the parent about changes or problems specific to the toddler or preschool child.

Review of Systems

SYSTEM/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- How has the child been feeling?
- Has his or her behavior changed? If so, how?
- How much does the child weigh?

- Screens for any obvious problems.
- Good indicator of how child is feeling.
- Monitor weight closely to evaluate normal growth and development.

Integumentary

- Does the child have any skin problems, such as rashes or excessive itching?

- Tinea capitis, ringworm, and contagious disorders often occur in toddlers and preschoolers, especially if the child is in day care or preschool.
- Atopic dermatitis (eczema) is often associated with allergies and has a familial tendency.
- Diaper dermatitis (diaper rash) occurs from prolonged exposure to an irritant, such as urine or feces.
- Pediculosis capitis (head lice) often occurs in preschoolers in day care or preschool. Playing dress-up or sharing hats increases transmission.
- Excessive sun exposure (ultraviolet rays) without sunblock increases risk for skin cancer later in life.

- Is the child exposed to the sun?

- Does she or he wear sunblock?

HEENT

Eyes

- Does the child have vision problems?
- Does he or she have crossed eyes?
- When was the child's last eye exam?

- Normal visual acuity is 20/40 for toddlers. Depth perception is still developing.
- Test vision routinely during physical exams.

Ears

- Does the child have hearing problems?
- Does she or he respond to sounds?
- Is he or she talking or making sounds?
- When was the child's last hearing exam?
- Does the child have frequent ear infections?
- Is the child irritable?
- Does she or he have a poor appetite or sleep poorly?

- Test hearing routinely during physical exam.

- Toddlers and preschoolers have a high incidence of otitis media, with the highest incidence between 6 months and 2 years.

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p><i>Teeth</i></p> <ul style="list-style-type: none"> ■ How many teeth does the child have? ■ Has the child had his or her first dental exam? ■ Does the child brush her or his own teeth? How often? ■ Is your water fluorinated? ■ Does the child go to bed with a bottle? 	<ul style="list-style-type: none"> ■ All 20 primary teeth should be in by age 2½. ■ The first dental exam should occur shortly after primary teeth start erupting. At least one dental exam per year recommended. ■ Review toothbrushing technique with child and parent. ■ Fluoride decreases tooth decay. ■ Bedtime bottles, nocturnal breast feeding, or coating pacifiers with a sweet substance can cause baby bottle caries, which occurs most frequently between 18 months and 3 years. <div style="border: 1px solid red; padding: 5px; margin-top: 10px;">  <p>ALERT</p> <p>Fluoride is toxic if ingested. Advise parents to supervise toothbrushing and not allow the child to eat or swallow toothpaste.</p> </div>
<p><i>Respiratory</i></p> <ul style="list-style-type: none"> ■ Does the child have a cough, sneezing, runny nose, or fever? ■ Does he or she wheeze or have trouble breathing? ■ Is the child exposed to air pollutants, such as smoke or second-hand smoke? 	<ul style="list-style-type: none"> ■ Upper respiratory infections (URIs) are common in toddlers and preschoolers. ■ Respiratory syncytial virus (RSV) causes one-half of all bronchiolitis and usually occurs in children under age 2. ■ Asthma is the most common disease among children. Onset is usually between ages 3 and 8. ■ Exposure to second-hand smoke increases risk for respiratory and ear infections and asthma attacks.
<p><i>Cardiovascular</i></p> <ul style="list-style-type: none"> ■ Were you ever told that the child had a heart murmur? ■ Has the child ever turned blue? 	<ul style="list-style-type: none"> ■ Although cardiovascular disease is uncommon in children, murmurs are a common finding. Most are innocent, functional murmurs, but follow-up is indicated to rule out pathology. ■ Change in skin color may indicate cardiopulmonary problem.
<p><i>Gastrointestinal</i></p> <ul style="list-style-type: none"> ■ Is the child potty trained? ■ What is her or his bowel pattern? 	<ul style="list-style-type: none"> ■ Bowel control is usually achieved before age 3.
<p><i>Genitourinary</i></p> <ul style="list-style-type: none"> ■ Does the child have bladder control? 	<ul style="list-style-type: none"> ■ Full bladder control may not occur until age 5.
<p><i>Musculoskeletal</i></p> <ul style="list-style-type: none"> ■ Is the child walking? 	<ul style="list-style-type: none"> ■ The child should be able to walk without difficulty by 12 or 13 months.
	<div style="border: 1px solid red; padding: 5px;">  <p>ALERT</p> <p>Even though the child can walk, he or she is still at risk for falls because he or she does not have refined coordination or full depth perception.</p> </div>

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Neurologic</p> <ul style="list-style-type: none"> ■ Is the child speaking? ■ Does she or he use complete sentences? ■ Does he or she have any fears? <p>■ Does the child have temper tantrums or aggressive behavior?</p> <p>Lymphatic/Hematologic</p> <ul style="list-style-type: none"> ■ Does child have any lumps in the neck, groin, or underarms? 	<ul style="list-style-type: none"> ■ Cognitive skills continue to develop and speech normally develops between ages 2 and 4. ■ Fear of real and imaginary things is common (e.g., fear of dogs or the dark). ■ May need to educate parents about effective ways to deal with child's behavior. <ul style="list-style-type: none"> ■ May indicate infection or malignancy.

Psychosocial Profile

The psychosocial profile should focus on the child's health practices and behaviors that affect health and well-being.

Health Practices

When was the child's last checkup? (Checkups are recommended every year.) Keep in mind that the child may have no control over frequency of health screenings. The parent or guardian usually assumes this responsibility, and health insurance or lack of it may be an influencing factor. You may need to make appropriate referrals or provide information on available health resources.

Typical Day

What is the child's typical day like? Does he or she have a babysitter or go to day care or preschool? If so, does he or she enjoy it? What kind of day-care situation is it (e.g., day-care center, child care in private home, babysitter in your own home)?

Nutritional Patterns

If necessary, provide nutritional education to children and families.

Ask: Is child still nursing or bottle feeding? If yes, how often? Because breast milk contains immunoglobulin A (IgA), and breastfeeding minimizes reflux of milk into the eustachian tube, breastfeeding decreases the risk of otitis media and respiratory viruses and allergies. Ask for a 24-hour diet recall, inquiring about how often child eats breakfast and what she or he usually eats at each meal. Ask the child to name her or his favorite foods and snacks. Also ask about junk food and fast food, which amount to empty calories and contribute to the high incidence of obesity among children.

Toddlers between 12 and 18 months still have high protein and calorie requirements for growth and high

calcium, iron, and phosphorous requirements for bone growth. After 18 months, the toddler's appetite decreases (**physiological anorexia**) in response to decreased nutritional needs. Preschoolers need an average of 1800 calories a day, as well as calcium and minerals for bone growth. They should eat limited fats. During the toddler and preschool years, the child develops taste preferences and may become a picky eater.

Activity and Exercise Patterns

Is the child very active? What does he or she enjoy doing? Is he or she supervised during activities? Toddlers and preschoolers are normally very active and need close supervision.

Recreation and Hobbies

Safety is a major concern for the toddler and preschooler. If recreational activities or hobbies increase the risk for injury, ask if protective measures, such as bike helmets, are used. Also ask about the parents' hobbies. For example, are there guns in the home? Play for toddlers is parallel (playing next to each other but not together); play for preschoolers is associative, group play, with more social interaction. In both age groups, safety is paramount, supervision is needed, and toys must be appropriate for the child's age.

Sleep and Rest Patterns

Ask about sleep patterns and naps, bedtime rituals, and sleep problems (nightmares, night terrors, somnambulism, enuresis). How many hours does the child sleep? Twelve hours a day is normal. Toddlers usually need a nap, but preschoolers do not.

Personal Habits

Ask parents about their and other caregiver's personal habits. Do they smoke, drink alcohol, or use drugs? Exposure to these substances places the child at risk for

health problems. If drug and alcohol abuse is an issue, the caregiver's ability and the home environment need to be assessed to ensure the child's safety. Referrals may be warranted.

Roles and Relationships

If there are siblings, ask the parent about the relationship. Sibling rivalry is not uncommon.

Religion and Spirituality

The toddler and preschooler's perception of a Supreme Being is concrete and based on the beliefs and practices of parents and caregivers. Prayers are often part of bedtime rituals and can be a source of comfort for a child.

Sexuality

Toddlers and preschoolers are curious by nature and exhibit sex-role imitation. They are also curious about their own bodies, and it is not uncommon for them to masturbate. How parents address this behavior is important and can have lasting effects.

Stress and Coping

Does the child behave aggressively or have temper tantrums? If the child hurts herself or himself or others, she or he needs to learn effective coping strategies, and a referral may be necessary.

Physical Assessment

After you complete the health history, perform the physical examination. You will use all four assessment techniques (for more information, see Chapter 3, Approach to the Physical Assessment). When assessing the toddler and preschooler, keep in mind the specific variations that are unique to this developmental period.



FIGURE 25.1. Taking temperature.

Begin with a general survey, obtaining height, weight, and vital signs such as temperature and blood pressure (BP) (Figs. 25.1 and 25.2). Keep track of the child's growth and development by plotting height and weight on growth charts (see Appendix B). The toddler usually gains 4 to 6 pounds and grows 3 inches a year. Head and chest circumferences are usually equal by age 2. The preschooler gains 5 pounds and grows 2½ to 3 inches a year. Changes in vital signs include a slight, gradual increase in BP and a slight decrease in temperature, pulse, and respirations.

Once you have completed your assessment, document your findings, identify any nursing diagnoses, and formulate a plan of care.

(Text continued on page 897)



FIGURE 25.2. Taking BP.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

1. General Health Survey

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Inspect overall appearance, noting appropriate growth and development for child's age.
- *Toddler has potbelly and wide base of support. Preschooler loses potbelly and becomes taller and leaner.*
- Note delays or premature maturation.

ABNORMAL FINDINGS/RATIONALE

- Note obvious weight problems.
- Inspect skin for lesions.

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

2. Integumentary

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Inspect hair and scalp for lice.
- *Skin intact, no lesions or bruising.*
- *Head free of pediculosis.*

ABNORMAL FINDINGS/RATIONALE

- Lesions, such as tinea capitis or ringworm, need treatment.
- Pediculosis is common among preschoolers.



ALERT

Suspect abuse if you find unexplained bruising or injury (see Recognizing Abuse, page 889).

3. HEENT

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

Head and Face

- Inspect head and face.
- Measure head circumference.
- Palpate anterior fontanel.
- *Head circumference appropriate for age of child (plot on graph).*
- *Anterior fontanel closes by 18 months.*

Eyes

- Test visual acuity.
- Test for “lazy eye” (strabismus) with corneal light reflex or cover-uncover test.
- *Visual acuity is normally 20/40 during the toddler years.*
- *Begin vision screening between ages 3 and 4.*

Ears

- Test hearing by age 3 or 4.
- Test hearing with pure tone audiometer.
- Inspect external ear canal and tympanic membrane.

- *Hearing intact.*
- *No drainage or foreign objects, tympanic membrane intact.*

Nose

- Inspect nasal septum and mucosa.
- *Nasal mucosa pink, no drainage or foreign objects.*



ALERT

When inspecting nares or external ear canal, be alert for foreign objects.

Mouth

- Inspect oral mucosa and pharynx.
- Inspect number and condition of teeth.
- *Generally, tonsils are large.*
- *Eruption of primary teeth is usually complete by age 2½ years.*
- Review dental hygiene with parent and child.

Neck

- Palpate for lymph nodes.
- *Lymph nodes not palpable.*



Inspecting ear with otoscope

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

3. HEENT (continued)

ABNORMAL FINDINGS/RATIONALE

- Head size growth slows to 1 inch a year by end of age 2; then ½ inch a year until age 5.
- Premature or delay in closure of fontanel.
- Visual deficits warrant follow-up.
- Referral needed for strabismus to prevent **amblyopia**.
- Hearing deficits warrant follow-up.
- Toddlers and preschoolers have a high incidence of otitis media.
- Toddlers and preschoolers have a tendency to put foreign objects into ears.
- *Boggy, bluish-purple, or gray turbinates:* Chronic rhinorrhea, which can result from allergic rhinitis.
- Toddlers and preschoolers have a tendency to put foreign objects into nose.
- Enlarged, red tonsils with exudates may indicate an infection.



Baby bottle caries

- Note baby bottle caries.
- *Enlarged lymph nodes:* Infection or lymphoma.

4. Respiratory

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Inspect and measure size and shape of chest.
- Auscultate lungs.
- *Chest circumference appropriate for age of child.*
- *Anteroposterior to lateral diameter should be 1:2 by end of 2nd year.*
- *Lungs clear.*



Auscultating the lungs

ABNORMAL FINDINGS/RATIONALE

- Abnormal chest shape or circumference warrants referral.
- Toddlers and preschoolers have a high incidence of URIs.

5. Cardiovascular

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Auscultate heart, noting rate and rhythm.
- *Children often have a sinus arrhythmia and a split second heart sound that both change with respiration. This is a normal variation.*
- *Systolic innocent murmurs and venous hum are common findings.*



ALERT

If you detect a murmur, refer the patient for follow-up to rule out pathology.



Auscultating the heart

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

5. Cardiovascular (continued)

ABNORMAL FINDINGS/RATIONALE

- Murmurs should be evaluated.

6. Gastrointestinal

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Inspect, palpate, and auscultate the abdomen.
- *Potbelly normal for toddler; disappears as abdominal muscles strengthen.*



Auscultating the abdomen

ABNORMAL FINDINGS/RATIONALE

- Umbilical hernias.

7. Genitourinary

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Inspect external genitalia.
- *Skin and genitalia intact.*



HELPFUL HINT

If child is still in diapers, inspect for diaper dermatitis.

ABNORMAL FINDINGS/RATIONALE

- Diaper dermatitis.

8. Musculoskeletal

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Inspect gait.
- Test muscle strength.
- *Toddlers usually can walk alone by 12 to 13 months. Balance is unsteady with wide base of support. Valgus (knock knees) or varus (bowlegs) may be present. The preschooler's gait is more balanced, smaller base of support; child walks, jumps, climbs by age 3.*
- *Strength increases during preschool years.*

ABNORMAL FINDINGS/RATIONALE

- Persistent gait and balance problems beyond age 3 need referral.
- Muscle weakness.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

9. Neurologic

NORMAL VARIATIONS/SYSTEM/ASSESSMENT

- Test balance, coordination, and accuracy of movements.
 - *Balance and coordination improve with refinement of fine motor skills.*

ABNORMAL FINDINGS/RATIONALE

- Balance and coordination problems may indicate developmental delays.



Back to the Case Study

Now that you understand the physical and psychosocial issues surrounding toddlers and preschoolers, apply your knowledge to our case study subject, 3-year-old Travis Harrison, by answering the following questions.



CRITICAL THINKING ACTIVITY #1

How does assessment of the toddler and preschooler differ from assessment of other patients?



CRITICAL THINKING ACTIVITY #2

What approach is best for assessing the toddler and preschooler?



CRITICAL THINKING ACTIVITY #3

What health concerns should you keep in mind as you proceed with the assessment?



CRITICAL THINKING ACTIVITY #4

List some nursing diagnoses that are appropriate for the toddler and preschooler.

RESEARCH TELLS US

Obesity is a growing national health problem that is now affecting children, and TV viewing is a contributing factor. While interventions to reduce TV viewing time have been developed for older children, they have yet to be developed for younger ones. To combat this, a recent study was performed to determine the effects of a program to reduce viewing time for younger children.

The study focused on randomized control of 16 preschools and day-care facilities, with children ranging in age from 2.6 to 5.5 years. The control group had safety prevention programs, whereas the intervention group had seven sessions designed to reduce TV viewing. Over the course of the school year, changes in TV viewing and growth measurements were used to determine the outcome.

The findings revealed a statistically significant difference in viewing time between the groups. Prior to the intervention, TV viewing time for both groups ranged from 11.9 to 14 hr/wk. After the intervention, the TV viewing time decreased in the intervention group by 3.1 hr/wk, whereas the control group increased their

view time by 1.6 hr/wk, for an adjusted difference between groups of -4.7 hr/wk (95 percent confidence interval, -8.4 to -1.0 hr/wk; $p = .02$). And, children who watched TV more than 2 hr/wk decreased significantly in the intervention group from 33 to 18 percent, whereas the control group increased TV view time from 41 to 47 percent ($p = .046$). No statistically significant changes were seen in growth measurements.

Although no significant changes were seen in growth measurements, the reduction in TV viewing time was significant and warrants further study. Because TV viewing habits develop at an early age, an intervention geared at reducing viewing time seems valuable. The American Academy of Pediatrics recommended that children limit TV viewing to a maximum of 2 hr/day, with no TV viewing for children under the age of 2. Less TV viewing time allows more time for children to engage in physical activity.

Nurses working in primary care and schools have the opportunity to promote such behaviors, which may in turn reverse the trend of childhood obesity.

HEALTH CONCERNS

Immunizations needed between ages 1 and 5 include:

- *Polio*: Dose three between 12 and 18 months; dose four between ages 4 and 6.
- *Diphtheria, tetanus, pertussis (DTP)*: Dose four between 12 and 18 months; dose five between ages 4 and 6.
- *Measles, mumps, rubella (MMR)*: Dose one between 12 and 15 months; dose two between ages 2 and 6.

- *Varicella zoster (chickenpox) (VZV)*: Between 12 and 18 months or at any age after that if the child has never had chickenpox.
- *Hepatitis B (HepB)*: Dose three between 6 and 18 months.
- *Haemophilus influenzae (Hib)*: Dose four between 12 and 15 months.

COMMON PROBLEMS

Problems occurring frequently in toddlers and preschoolers include toilet training issues, sleep problems, temper tantrums, sibling rivalry, fears,

stress, regression, child abuse and neglect, poisoning, injuries, otitis media, baby bottle caries, nutritional problems, and asthma.

S U M M A R Y

- Toddlers and preschoolers have their own unique developmental and psychosocial issues.
- Understanding these issues and knowing how to communicate honestly and effectively with children and


parents will help you conduct a thorough assessment and develop an effective plan of care.

R E V I E W Q U E S T I O N S

1. You are performing a vision assessment on Robert Daniels, age 4. Which far-vision finding would be considered normal for a 4-year-old child?
 - a. 20/20
 - b. 20/40
 - c. 20/100
 - d. 20/200
2. Which musculoskeletal finding is often seen in toddlers as they are learning to walk?
 - a. Wide base of support
 - b. Wide stride length
 - c. Toeing in
 - d. Toeing out
3. Toddlers often have potbellies. Which spinal change would be considered normal in toddlers?
 - a. Scoliosis
 - b. Kyphosis
 - c. Lordosis
4. At what age does the child usually gain bowel control?
 - a. 2 years
 - b. 3 years
 - c. 4 years
 - d. 5 years
5. To what age might it take for a child to gain full bladder control?
 - a. 2 years
 - b. 3 years
 - c. 4 years
 - d. 5 years
6. Mark Benner, age 5, is hospitalized for appendicitis. Mrs. Benner is concerned that Mark wet the hospital bed and states he doesn't wet the bed at home. How would you respond to Mrs. Benner?
 - a. "It is not uncommon for children to regress when hospitalized."
 - b. "He will need to be retrained."
 - c. "There is an underlying problem that will need further evaluation."
 - d. "This is his way of showing that he is angry for being in the hospital."

7. According to Erikson, what is the developmental task for the toddler, ages 1 to 3 years?
- Trust vs. mistrust
 - Autonomy vs. shame
 - Initiative vs. guilt
 - Identity vs. role diffusion
8. According to Erikson, what is the developmental task for the preschooler, ages 3 to 6 years?
- Trust vs. mistrust
 - Autonomy vs. shame
 - Initiative vs. guilt
 - Identity vs. role diffusion

CHAPTER 26



Assessing the School-Age Child and Adolescent

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify normal physiological changes of the school-age child and adolescent
- Discuss various cultural influences on children
- Identify developmental tasks for the school-age child and adolescent
- Identify health issues and risks for the school-age child and adolescent
- Identify pertinent history data included in a school-age child and adolescent history
- Obtain a school-age child and adolescent health history
- Describe additional physical assessment components specific to a school-age child and adolescent assessment
- Perform a school-age child and adolescent assessment
- Document school-age child and adolescent assessment findings
- Identify actual/potential health problems stated as nursing diagnoses with supporting data

During the period from school age through adolescence, the patient grows and matures from a child to a young adult. Dealing with this age group is a challenging, yet potentially very rewarding aspect of nursing care. To effectively assess and intervene, you must be familiar with the significant physical, psychosocial, and cognitive developmental changes that occur before, during, and after puberty. Although each stage has its own set of milestones, the assessment techniques you use are essentially the same. This chapter covers the complete health history and physical assessment for the school-age patient first, and then moves on to the adolescent.

Assessing the School-Age Child

The school-age child is between the ages of 6 and 12 years. This age span (called the *latency period* by Freud) is like the calm before the storm and both follows and precedes a time of great growth and development. Although this is a slower, more stable period, some major changes do occur. For example, when the child begins school, a whole new world opens up to him or her, and the child begins to develop relationships other than those within the family.

Before beginning your health assessment, ask yourself:

- How does assessment of the school-age child differ from assessment of other patients?
- What approach is best for the school-age child?
- What health concerns should I keep in mind as I proceed?

As you proceed with the assessment, stay alert for the following health problems and risk factors common in school-age children:

- **Injuries and accidents:** These are the most frequent cause of morbidity and mortality in children. The school-age child is very active and involved in many activities. Participation in sports especially increases the risk for injury. Injury can result from overuse or from engaging in activities not yet suitable for the child's age. Common problems include tendinitis, osteochondritis, Osgood-Schlatter disease (osteochondritis of the epiphysis of the tibial tuberosity), and stress fractures.
- **Drug, alcohol, and nicotine use:** Although the use of drugs, alcohol, and nicotine occurs more frequently among adolescents, experimentation often begins during the school-age years and may continue into adolescence. Remember, peer pressure exerts great influence on a child's behavior. Prevention works best, so education needs to begin at an early age.

- **Eating disorders:** Obesity is an increasing problem among American children. Eating habits are established at an early age, and psychosocial and cultural factors can influence them. Eating may be associated with nurturing. The effects of obesity are not only physical; they also can affect the child's self-image. If obesity continues through adolescence into adulthood, the effects can be long lasting, increasing the person's risk for cardiovascular disease, hypertension (HTN), hyperlipidemia, diabetes, degenerative joint disease, sleep apnea, and colon cancer.
- **Learning problems:** During the school-age years, learning problems become apparent. Attention deficit hyperactivity disorder (ADHD) and learning disability (LD) are common. ADHD is characterized by developmentally inappropriate inattention, impulsiveness, and hyperactivity; LD is difficulty with reading, writing, listening, reasoning, speaking, mathematical skills, and social skills. Care must be taken not to prematurely label a child as having ADHD or LD because many children exhibit one or more of these behaviors. The symptoms of ADHD begin before age 7, and children exhibit six or more symptoms for at least 6 months. As a nurse, you will need to make appropriate referrals, as indicated, and help with the education and therapeutic management of children with ADHD and LD.
- **Lying, cheating, and stealing:** Children may lie to avoid punishment or cheat to win a game. These common behaviors usually occur in this age group and diminish as the child becomes older and more responsible for her or his behavior.
- **Emotional problems:** Children commonly experience stress. The source can be the home, school, or community. Children also experience various types of emotional problems, such as school phobia, recurrent abdominal pain, conversion reactions, depression, and even schizophrenia. Often, the emotional problem is triggered by a traumatic event, such as an illness, the loss of a parent, or a major family crisis such as divorce. Be alert for signs of stress that warrant further assessment such as headaches; stomachaches; sleep problems; enuresis; changes in eating habits; changes in behavior, such as aggressiveness; poor performance in school; isolation or withdrawal from participation; and regression. The key is to identify the problem and respond promptly to avoid further ill effects.

Developmental Changes

The changes that take place during the school-age years are subtle. Growth and development are both physical and psychosocial. Before you can detect any abnormalities, you need to understand the normal changes that occur in this age period.

Physical Changes

As the school-age child grows, his or her measurements become more proportional as baby fat disappears. The neuromuscular system continues to develop, and the child becomes more agile and coordinated. For specific physical changes that occur, see the Physical Assessment section.

Psychosocial and Emotional Changes

Psychosocial changes also occur at an even pace. Erikson describes this stage as industry versus inferiority. The child becomes more independent, developing skills and competencies. Social skills also develop as the child becomes more involved in school and community activities. With each success at school, at home, or in the community, confidence and independence grow. The child learns to cooperate, work, and compete with others. It is important that both adults and peers recognize the child's accomplishments.

Peer acceptance becomes especially important. The child needs a sense of belonging and of "fitting in." Peers often have a greater influence on a child than her or his parents do. Sex roles are established through peer relationships, and same-sex friendships develop. School provides the child with the opportunity to see different perspectives and to argue, negotiate, resolve conflicts, work together, and develop friendships. Belonging to clubs, teams, and peer groups is important.

Even though peer influence seems greater than parental influence, parents still play a major role in the child's development. Children test parents, so it is impor-

tant that parents set limits to give the child a sense of security.

Team play becomes important, either on sport teams or in quiet games. Both involve rules and rituals that children strongly adhere to. Watching sports, creating artwork, reading, collecting things, cooking, sewing, and similar activities are also important activities that often carry through adolescence into adulthood. As the child progresses through this stage, her or his self-concept further develops. For a positive self-concept to develop, the child needs to feel special.

The child also moves from preconceptual thinking to conceptual thinking. Piaget calls this stage of cognitive development *concrete operations*, when the child begins using thought processes to perceive the world. The child masters the tasks of conservation and classifications. Reading is the most important skill that develops during this stage.

During the school-age years, the child becomes less egocentric and begins to develop a conscience. Initially, his or her views were black and white, right or wrong based on the parents' moral values. But as the child grows and matures, he or she incorporates different points of views before making judgments. The Supreme Being is seen in concrete terms. Illness or injury may be seen as a punishment.

Cultural and Ethnic Variations

Various cultures have different perspectives on children and their role in the family. You need to understand any cultural influences that might have an impact on your patient's plan of care.

Cultural and Ethnic Variations in School-Age Children

CULTURAL GROUP INFLUENCE

Amish	<ul style="list-style-type: none"> ■ Children are seen as a gift from God. ■ Parents are very directive of children throughout school.
Appalachian	<ul style="list-style-type: none"> ■ Parents believe in physical punishment. ■ Having children is associated with sense of importance.
Arab American	<ul style="list-style-type: none"> ■ Child's character and success depend on parental influence. ■ Children are taught conformity and cooperation. ■ A "good" child is obedient; behavior that would bring dishonor is avoided.
Chinese American	<ul style="list-style-type: none"> ■ Parents believe in physical punishment. ■ Boys and girls play together when young but are separated when older. ■ Children are pressured to succeed. ■ Males are more valued than females.
Cuban American	<ul style="list-style-type: none"> ■ Children are expected to help out at home.
Egyptian American	<ul style="list-style-type: none"> ■ Children are expected to study and respect their parents.
Filipino American	<ul style="list-style-type: none"> ■ Children are expected to be studious and goal oriented as well as respectful and loyal to family.
	<ul style="list-style-type: none"> ■ Children are expected to honor and respect their parents.

(continued)

Cultural and Ethnic Variations in School-Age Children (continued)

CULTURAL GROUP INFLUENCE

French American	<ul style="list-style-type: none"> Children are well educated and a source of pride for the family.
Greek American	<ul style="list-style-type: none"> Children are the center of the family. Parents use teasing as a form of discipline to toughen children and make them aware of public opinion.
Iranian American	<ul style="list-style-type: none"> This is a child-oriented society. Children are expected to be respectful.
Irish American	<ul style="list-style-type: none"> Boys are expected to be more aggressive than girls. Children are expected to show self-restraint, discipline, respect, and obedience.
Jewish American	<ul style="list-style-type: none"> Children are seen as a valued treasure. Children are expected to respect and honor parents.
Mexican American	<ul style="list-style-type: none"> Children are highly valued but must respect parents.
Navajo Native American	<ul style="list-style-type: none"> Children are allowed to make own decisions, such as taking medicine, which might seem irresponsible on part of parents.
Vietnamese American	<ul style="list-style-type: none"> Children are prized and valued. Children are expected to be obedient and devoted.

Introducing the Case Study



Joshua Goldfarb, age 6, is starting first grade this fall. He is in the pediatrician's office for his required physical exam. Joshua is sitting close to his mother and seems a little anxious. He asks you, "Do I have to get any shots?"

if there is an acute problem, you should ask the child to describe it in her or his own words. Because most of the health history is identical to an adult's, the sections that follow contain only information that is especially important to ask the school-age child or that is specific to this age group.

Biographical Data

Begin with the biographical data, noting where and with whom your patient resides. Also note who is listed as contact person—it may not be a parent. Do not assume anything.

Current Health Status

Ask the parent or guardian about all prescription and over-the-counter (OTC) medications.

Past Health History

Ask the parent or guardian specific past health history questions. An immunization history is especially important (see Health Concerns, p. 912, for required immunizations).

Family History

Ask the parent or guardian about heart disease, strokes, cancer, and other serious health conditions within the family. Be sure to ask if anyone died at an early age (under 35) from a sudden heart attack. This may be a risk factor for sports participation and may require additional investigation.

Review of Systems

Go through every system, asking about changes or problems specific to the school-age child. Although the questions that follow are addressed to the child, you will need to question the parent or guardian in the case of younger children.

Performing the School-Age Child Assessment

Now that you understand the major issues affecting this age group, you can begin the actual assessment. The process is similar to that for an adult. To communicate with this age group, be honest, open, and nonjudgmental. School-age children often need an adult to look up to, and you are in an ideal position to serve as a role model. You are also in an ideal position to screen for risk factors and teach healthy lifestyle behaviors.

As with all assessments, the process involves data collection (both subjective and objective), diagnosis, planning, interventions, and evaluation.

Health History

Follow the history format used for an adult, but keep in mind the normal developmental changes and the issues confronting the school-age child. Although the parent or guardian usually provides most of the history, you may want the child's response to a few questions. For example,

Review of Systems

SYSTEM/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- How have you been feeling?
- Have you lost or gained weight recently?

- Screens for obvious problems.
- Changes in weight, especially gains, may indicate poor eating habits.

Integumentary

- Do you have any skin rashes or lesions?
- Do you have excessive itching?

- Tinea capitis; ringworm; verruca (warts); contact dermatitis; and poison ivy, oak, and sumac frequently seen in school-age children.
- Pediculosis capitis (head lice) common in school-age children.
- Excessive sun exposure (ultraviolet rays) without sunblock increases risk for skin cancer later in life.

- Are you exposed to the sun?
- Do you use sunblock?

HEENT

Eyes

- Do you have any visual problems?
- Do you wear glasses? If yes, do you wear them consistently, especially while at school?
- Do you wear contact lenses?

- Visual acuity reaches 20/20 by school age.
- Children may not consistently wear glasses for fear of being teased by other children.
- May identify teaching needs regarding care of contact lenses.
- Eyes should be tested routinely during physical exams.

- When was your last eye exam?

Ears

- Do you have any hearing problems?
- Do you listen to loud music?

- Cochlear damage can result from excessive exposure to loud noise, such as CD players and rock concerts.
- May identify health teaching need regarding noise pollution's effect on hearing.
- Hearing should be tested routinely during physical exams.

- When was your last hearing exam?

Teeth

- How many second teeth do you have?

- Primary teeth are lost during school-age years and replaced by secondary teeth.
- These years are called the "age of loose tooth" or "ugly duckling stage," because secondary teeth are initially too big for the face.
- Should have at least an annual dental exam.
- Common dental problems include dental caries, periodontal disease, malocclusion, and dental injury.

- When was your last visit to the dentist?
- Do you see an orthodontist?



ALERT

In secondary tooth evulsion, the tooth should be replanted as soon as possible. If it is replanted within 30 minutes, replantation is 70 percent successful.

Respiratory

- Do you have asthma?
- Do you ever have trouble breathing or do you wheeze when exercising or running?
- Are you exposed to air pollutants, smoke, or second-hand smoke?

- Asthma is the most common respiratory disease in children.
- Allergies and asthma often become apparent during school-age years.
- Exercise-induced bronchospasm: Acute, reversible, self-terminating airway obstruction that occurs after vigorous exercise, peaks 5 to 10 minutes once activity stops, and then ceases within 30 minutes.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Cardiovascular</p> <ul style="list-style-type: none"> ■ Do you have any chest pain? ■ Does your heart ever skip a beat? 	<ul style="list-style-type: none"> ■ Although cardiovascular disease is not common in children and adolescents, it can occur. Prolonged QT syndrome can cause sudden cardiac death in a seemingly healthy child.
<p>Gastrointestinal</p> <ul style="list-style-type: none"> ■ Do you have any stomach problems? ■ How often do you have a bowel movement? 	<ul style="list-style-type: none"> ■ Encopresis (fecal incontinence and voluntary or involuntary loss of bowel control) is more common in boys, usually secondary to constipation. ■ Recurrent abdominal pain is common during childhood and is often psychosomatic. Child usually has poor self-image. School can precipitate attacks.
<div style="border: 1px solid red; padding: 5px; background-color: #f0f0f0;">  <p>ALERT</p> <p>If child presents with recurrent abdominal pain, rule out physiological causes first.</p> </div>	
<p>Genitourinary</p> <ul style="list-style-type: none"> ■ Do you have bladder control? ■ Do you wet the bed? 	<ul style="list-style-type: none"> ■ Nocturnal bedwetting is more common in boys and often ends by ages 6 to 8.
<p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Do you have any back problems? ■ Have you ever been told you had a spinal problem? 	<ul style="list-style-type: none"> ■ Heavy backpacks are associated with low back problems in children. ■ Scoliosis (lateral curvature of spine) occurs more often in girls than in boys. Screening usually occurs in middle school and high school. If present, determine whether scoliosis is structural or postural (see Chapter 20, Assessing the Motor-Musculoskeletal System). ■ Physical maturity varies and does not always correlate with emotional or social maturity.
<p>Neurologic</p> <ul style="list-style-type: none"> ■ How would you describe your mood? 	<ul style="list-style-type: none"> ■ Stress, anxiety, and fear are seen in children. ■ Depression, conversion reactions, and even schizophrenia can occur during childhood and can be difficult to detect.
<p>Lymphatic/Hematological</p> <ul style="list-style-type: none"> ■ Have you been tired? ■ Do you have any lumps in your neck, underarms, or groin? 	<ul style="list-style-type: none"> ■ Non-Hodgkin's lymphomas occur most frequently in children before age 15

Psychosocial Profile

The psychosocial profile focuses on the child's health practices and on behaviors that affect health and well-being. Although the questions below are addressed to the child, you will need to question the parent/guardian with younger patients.

Health Practices

When did the child have his or her last checkup? Checkups are recommended every 2 years, more fre-

quently if child has an existing health problem or is involved in sports. Keep in mind that the child may have no control over frequency of health screenings. The parent or guardian usually assumes this responsibility, and health insurance or lack of it may be an influencing factor. You may need to make appropriate referrals or provide information on available health resources.

Typical Day

What is a typical day like? How is the child doing in school? This allows you to see the patient through her

or his own eyes and possibly identify stresses confronting her or him in everyday life. A change in academic performance or a sudden disinterest in school may reflect a more serious problem. What subjects does the child like and dislike? If she or he is having problems with a particular subject, has she or he sought help? How many days of school does she or he usually miss every term? ADHD and LD may become more apparent once the child starts school, and you should make appropriate referrals. School phobias are common at age 10, with somatic complaints such as nausea, headache, or abdominal pain. These complaints quickly subside once the child is assured that she or he doesn't have to go to school. If the patient is a "latchkey child," ask what she or he does with the time until the parent or guardian returns home. Children who are left home alone are at a higher risk for injury or delinquent behavior.

Nutritional Patterns

Ask for a 24-hour diet recall. Inquire about eating patterns, asking, "How often do you eat breakfast?" "What do you usually have for lunch?" "What are your favorite snacks?" Junk food and fast food amount to empty calories and contribute to the high incidence of obesity among children. Take this opportunity to provide nutritional education to children and families.

Activity and Exercise Patterns

Ask what the child enjoys doing outside the home. Does he or she play any sports? If so, what? Contact sports increase the risk for injury. Does he or she wear protective equipment? What does the child like to do with his or her friends? How often does he or she watch TV during the week? How much time does he or she spend on the computer? You may identify a need for health education about the importance of routine exercise.



ALERT

If a child is active in sports, she or he should engage in those that are appropriate for her or his age.

Recreation/Hobbies

What does the child do for fun? What hobbies does he or she have? If recreational activities or hobbies increase the risk for injury, ask if he or she uses protective measures such as a bike helmet. Ask whether there are weapons in the home. If there are guns, ask if child has taken gun or hunting safety courses.

Sleep/Rest Patterns

Ask when the child usually goes to bed and when she or he awakens. Children from ages 8 to 11 may resist going

to bed. Does the child have a bedtime routine? Does she or he wet the bed? Enuresis can continue into adolescence.

Personal Habits

Substance abuse is occurring at younger and younger ages; so many school-age children have already tried smoking or alcohol. Experimentation is common in this age group, and peer pressure can be great. So inquire about drug, alcohol, and nicotine use. Ask: "If you smoke cigarettes, how many do you smoke in an average day?" "If you use alcohol, how much do you usually drink during the week?" "If you use drugs, what type and what method?" (Methods include smoking, snorting, huffing, ingesting, and injecting.) Asking questions in this way is less threatening to the child; so he or she will be more likely to give an honest response.

The choking game is a dangerous game for school-age children and adolescents to get a "high," but the result can often be deadly. Remember, prevention is best. Educating parents and children as to the effects of such behaviors is important. (See Chapter 5, Approach to the Mental Health Assessment.)

Roles/Relationships/Self-Concept

Ask the child, "Who lives at home with you?" If she or he has siblings, ask about their relationship. Also ask, "Is there an adult in your life whom you feel comfortable talking with?" Teachers can have a major influence on a child's development, either positively or negatively. Ask about peer relationships. Does the child have a special friend, and what do they enjoy doing together? During the school-age years, a child develops close same-sex friendships.

Ask the child to describe herself or himself by saying, "Tell me about yourself." Self-image and body image are often influenced by a child's perception of how she or he is seen by others. Anything out of the norm may become an easy target for teasing, and such teasing can have lasting effects.

Religious and Spiritual Influences

The school-age child's values and beliefs are often guided by those of the parents. The Supreme Being is thought of in concrete terms. Illness may be perceived as a punishment. Prayer can be a source of comfort for a child; for example, prayers are often part of bedtime rituals.

Sexuality Patterns

The school-age child is curious by nature, and it is not unusual for school-age children to incorporate some form of sexuality into play. This is the best time to begin sex education, and you are in an ideal position to educate both children and parents.

Stress and Coping

Ask questions such as: "What makes you angry?" "What do you do when you get angry?" "What do you do to have

fun or relax?” This identifies if the child has healthy coping strategies to deal with life’s stresses. Violence is ever present in our society, at home, at school, and within the community. If the child uses violence as a coping mechanism, you need to help him or her to develop more effective coping strategies or make appropriate referrals.

Physical Assessment

After you complete the history, perform the physical exam. Physical assessment of the school-age child does not differ greatly from that of the adult—you use all four assessment techniques and the same tools (for more information, see Chapter 3, Approach to the Physical Assessment). However, there are specific variations that are unique to this developmental period.

Begin with a general survey, obtaining height and weight. Plot the patient’s height and weight on growth

charts to note growth and development (see Appendix B for growth charts). School-age children grow about 5 cm a year and gain about 4½ to 6½ lb a year. Boys and girls are relatively equal in size until they reach preadolescence; then girls usually surpass boys in both height and weight. Until the boys catch up during adolescence, this growth discrepancy can be distressing for both boys and girls.

Also take vital signs (Figs. 26.1 and 26.2). Be aware that as children age, their pulse and respiration rates decrease and their blood pressure (BP) increases compared with what they were during infancy.

Once you have completed your assessment, document your findings, identify any nursing diagnoses, and formulate a plan of care. If you identify problem areas, actively involve both the child and her or his parents in the plan of care.



FIGURE 26.1. Taking temperature.



FIGURE 26.2. Taking BP.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

1. General Health Survey

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect overall appearance, noting appropriate growth and development for child’s age.
- *General appearance becomes more slender; baby fat disappears.*

ABNORMAL FINDINGS/RATIONALE

- Note delays or premature maturation.
- Note obvious weight problems.

2. Integumentary

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect skin for lesions.
- Inspect hair and scalp for lice.
- *Skin intact, no lesions.*
- *No pediculosis.*



Inspecting hair and scalp

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

2. Integumentary (continued)

ABNORMAL FINDINGS/RATIONALE

- Lesions, such as tinea capitis or ringworm, need treatment.
- Pediculosis is common among school-age children.

3. HEENT


SYSTEM/ASSESSMENT/NORMAL VARIATIONS

Head and Face

- Inspect head and face.
- *Head size becomes more proportionate to body.*

Eyes

- Test visual acuity.
- *Visual acuity should be 20/20 by age 6.*



Testing visual acuity

Ears

- Test hearing with pure tone audiometer.
- Examine ear with otoscope.
- *Hearing intact.*
- *Tympanic membrane intact, no drainage or abnormal findings.*



Examining ear with otoscope

Nose

- Inspect nasal septum and mucosa.
- *Nasal mucosa pink and intact.*



Testing hearing

Mouth and Throat

- Inspect oral mucosa and pharynx.
- Inspect occlusion, number of teeth.
- Also assess for orthodontic devices and brushing technique around them.
- *Oral mucosa pink, no lesions.*
- *Tonsils are usually large.*
- *Child loses first teeth during this time.*
- *Secondary teeth begin to erupt around age 6 (6-year molars), and most secondary teeth are in by the time 12-year molars erupt.*



Examining mouth and throat

Neck

- Palpate lymph nodes.
- *Lymph nodes not palpable.*



Palpating thyroid nodes

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

3. HEENT (continued)

ABNORMAL FINDINGS/RATIONALE

- Abnormal head sizes warrant referral.
- Visual deficits warrant follow-up.
- Hearing deficits warrant follow-up.
- Take this opportunity to teach hazards of listening to loud music.
- *Discharge, red tympanic membrane*: Otitis media (inflammation of middle ear). Common in children.
- Foreign bodies in ear are common.
- *Boggy, bluish-purple, or gray turbinates*: Allergic rhinitis. Can result in chronic rhinorrhea.
- *Enlarged, red tonsils with exudates*: Associated with infection.
- If malocclusion present, may need to make referral to orthodontist.
- *Enlarged lymph nodes*: Infection or lymphoma.

4. Respiratory

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Auscultate lungs.
- *Lungs clear.*

ABNORMAL FINDINGS/RATIONALE

- Children with EIA require bronchodilators before activity. Usually, albuterol via a metered-dose inhaler is given 20 to 30 minutes before exercise.

5. Cardiovascular

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Auscultate heart, noting rate and rhythm.
- Auscultate for venous hum and murmurs.
- *Heart regular rhythm, no extra sounds or physiological S₃ and innocent murmur may be normal.*
- *Children often have sinus arrhythmia and split S₂, which change with respiration. This is a normal variation.*
- *Jugular venous hum is a normal, common finding in children.*



Auscultating venous hum



HELPFUL HINT

Be careful not to confuse a venous hum with a murmur. Remember: Jugular venous pulsations are easily obliterated.

ABNORMAL FINDINGS/RATIONALE

- Murmurs are also common in children. They are often innocent and functional, but should be followed up to rule out pathology.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

6. Gastrointestinal

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect, auscultate, and palpate the abdomen.
- *Abdomen soft, nontender, no organomegaly.*

ABNORMAL FINDINGS/RATIONALE

- Appendicitis is the most common illness during childhood that requires surgery.

7. Genitourinary/Reproductive

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect external genitalia.
- *External genitalia intact, appropriate development for age of child.*

ABNORMAL FINDINGS/RATIONALE

- *Precocious puberty:* Sexual development before age 8 in girls and age 9 in boys warrants follow-up evaluation.

8. Musculoskeletal

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect and palpate spinal curves.
- Test for spinal deformities (see Chapter 20, Assessing the Motor-Musculoskeletal System).
- Test muscle strength.
- *Normal curves noted, no abnormalities.*
- *No weakness, = muscle strength.*
- *Strength doubles during school-age years.*

ABNORMAL FINDINGS/RATIONALE

- Scoliosis is a major variation within musculoskeletal system.
- Screening should be done in the preadolescent period, generally when the child is in the fifth to sixth grade. Significant curvature should be referred for evaluation and follow-up.
- Muscle weakness warrants referral.

9. Neurological

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Test balance, coordination, and accuracy of movements.
- *Balance and coordination greatly improve, with refinement of fine motor skills.*

ABNORMAL FINDINGS/RATIONALE

- Balance and coordination delays need follow-up



Back to the Case Study

Now that you understand the physical and psychosocial issues surrounding school-age children, apply your knowledge to our case study subject, 6-year-old Joshua Goldfarb, by answering the following questions.



CRITICAL THINKING ACTIVITY #1

How does assessment of the school-age child differ from assessment of other patients?



CRITICAL THINKING ACTIVITY #2

What approach is best for assessing the school-age child?



CRITICAL THINKING ACTIVITY #4

List some nursing diagnoses that are appropriate for the school-age child.



CRITICAL THINKING ACTIVITY #3

What health concerns should you keep in mind as you proceed with the assessment?

RESEARCH TELLS US

More and more families in the United States are becoming homeless. An estimated 40 percent of children live below the poverty level and account for 25 percent of all homeless people. One recent study compared the health status of children using a school-based health center (SBHC) for comprehensive care. Of the 308 children seen at the SBHC, 76 lived in shelters and 232 lived in homes. The 308 charts were reviewed to identify health problems and medical coverage. The findings revealed

that homeless children had significantly more health problems than children who lived at home. The authors concluded that homeless children were at higher risk for health problems and lack of medical coverage and that SBHC is an effective means for providing health services to an increasingly underserved population. These findings are not surprising and have great implications for nursing. Support of SBHC would allow nurses to provide health care to this group.

Source: Berti, L., Zylbert, S., and Rolnitzky, L. (2001). Comparison of health status of children using a school-based health center for comprehensive care. *Journal of Pediatric Health Care* 15(5):244–250.

HEALTH CONCERNS

Immunizations needed between ages 6 and 12 include:

- *Polio*: Dose four by age 6.
- *Diphtheria, tetanus, pertussis (DTP)*: Between ages 4 and 6.
- *Tetanus-diphtheria “booster” (Td)*: After age 7 for children who have not gotten at least three doses of any tetanus and diphtheria vaccine; whenever child suffers a wound, if booster has not been received in last 5 years.
- *Measles, mumps, rubella (MMR)*: Between ages 11 and 12 if second MMR shot was not received between ages 4 and 6.
- *Varicella zoster (chickenpox) (VZV)*: Any time between ages 6 and 12 if child has never had chickenpox.
- *Hepatitis B (HepB)*: Any time if child was not immunized during infancy or school-age years. Children between ages 11 and 15 may need only two doses, 4 to 6 months apart.
- *Hemophilus influenzae (Hib)*: Not necessary unless child has special health concerns, such as sickle cell disease, human immunodeficiency virus/acquired immunodeficiency syndrome (HIV/AIDS), removal of spleen, bone marrow transplant, or cancer treatment with drugs.

COMMON PROBLEMS

Problems occurring frequently in school-age children include behavioral problems (e.g., school phobias, learning problems, and lying, cheating, or stealing); injuries and accidents; skin problems, such

as ringworm, pediculosis, and contact dermatitis; asthma; nocturnal bedwetting (mainly boys); malocclusion; and eating disorders (including poor nutrition).

S U M M A R Y

- School-age children have their own unique developmental and psychosocial issues.
- Understanding these issues and knowing how to communicate honestly and effectively with children and parents will help you conduct a thorough assessment and develop an effective plan of care.

Assessing the Adolescent

The term **adolescence** literally means “growing to maturity.” In the past, the terms *adolescent* and *teenager* were considered synonymous. Today, however, the span of adolescence is considered not so much a specific range of years (13 to 19), but rather a time period marked by specific developmental tasks, both physical and psychosocial. With this definition, adolescence can extend from age 8 to age 22. Before assessing an adolescent, ask yourself these questions:

- How does assessment of the adolescent differ from assessment of other patients?
- What approach is best for assessing the adolescent?
- What health concerns should I keep in mind as I proceed with the assessment?

As you proceed with the assessment, stay alert for the following health problems and risk factors common in adolescents:

- **Injuries and accidents:** Accidents, especially motor vehicle accidents, are the leading cause of death among adolescents. Alcohol is often a contributing factor. Sports injuries are also common.
- **Eating disorders:** Body-image issues contribute to eating disorders. Anorexia and bulimia are more common in girls, but they are also seen in boys. Anorexia usually involves almost totally abstaining from food, resulting in extreme weight loss, electrolyte imbalance, and loss of lean muscle tissue (including organ tissue such as heart muscle). It can also include obsessive exercise and is the patient’s way of taking control over her or his life. Bulimia involves bingeing on large quantities of food and then purging, through either vomiting or laxative use. This also results in electrolyte imbalance as well as dental erosion. Boys with eating disorders are usually involved in sports that strictly limit weight categories, such as wrestling or boxing. Girls are more affected by the media, in which the emaciated model is the desired female ideal. Family or self-pressure to be successful (e.g., to be a straight-A student) can also be a factor. Early identification and referral is critical to prevent serious sequelae.
- **Obesity:** Now seen in over 30 percent of U.S. adolescents, obesity is largely caused by decreased physical activity, increased TV and computer time, and too many high-fat foods.
- **Pregnancy:** The United States has the highest rate of adolescent pregnancy of any developed country in the world, although rates have decreased slightly. One cause includes the “it can’t happen to me” mind-set common in this age group. Pregnancy may also be intentional or an outcome of sexual abuse. All states provide confidential referral of adolescents to family planning centers for counseling and education on abstinence and contraception. Services are generally free and completely confidential, regardless of the patient’s age.
- **Sexually transmitted diseases (STDs):** Thousands of sexually active adolescents have STDs, which may cause long-term negative effects. Chlamydia, the most common and least recognized STD, is the major cause of pelvic inflammatory disease (PID) and can result in fertility problems. Human papillomavirus (HPV) is the most common cause of abnormal Pap tests in adolescent females and is a major cause of cervical cancer. Confidential services are available nationwide for the diagnosis and treatment of STDs and can be located through county or city health departments. Generally free of charge, these services also provide counseling related to safe sex and supplies such as condoms.
- **Emotional disorders:** Depression, bipolar disorders, conduct disorders, and obsessive-compulsive disorders can have serious outcomes, including suicide, if undiagnosed and untreated. Early detection and prompt referral for treatment is crucial. In many states, adolescents can “self-refer” for diagnosis and treatment of mental illness.
- **Substance abuse:** Alcohol is the most commonly reported substance of abuse, but marijuana, cocaine, methamphetamine, and some “designer drugs” such as “ecstasy” are also widely used. More than 30 percent of teens also report using cigarettes or smokeless tobacco. Education is a valuable tool, but to be effective, information must be presented factually and scare tactics avoided. In most states, adolescents can self-refer for treatment.
- **Violence:** Violent acts involving adolescents continue to increase. School shootings and random gang violence give some young people a fatalistic view. They think, “I’m going to die anyway, so what’s the big deal?” Those without a strong sense of self-worth and purpose are most likely to be involved in violence, either as recipients or as perpetrators. Although recent studies fail to reveal any one identifiable personality trait of violent adolescents, those who are withdrawn, noncommunicative, angry,

and “loners” may be at greater risk. Talking to the patient about behaviors of this type may open the door for therapeutic interventions.

Developmental Changes

Many physical changes take place during adolescence. Height, weight, and muscle mass increase rapidly. Sexual maturation also occurs. You need to understand the normal changes that occur in order to detect abnormalities.

Stages of Adolescence

Many authorities divide adolescence into early, middle, and late (based on specific developmental tasks), or prepubescence, puberty, and postpubescence (based on physical changes). Early adolescence begins at puberty (as early as age 8 but more typically age 11 to 14 for girls and age 12 to 16 for boys). This period of rapid physical growth corresponds with the onset of menstruation in girls and sperm production in boys. Middle adolescence typically ranges from age 14 to 16 in girls and age 16 to 18 in boys. Females have generally achieved their adult height by this stage, but males may continue their linear growth. Late adolescence typically starts around age 17 and can continue into the early 20s. This rapid period of both physical growth and psychosocial development is what makes the adolescent period so complex and challenging for healthcare providers.

Prepubescence is defined as the 2-year period before puberty in which preliminary physical changes occur. Puberty is the point when sexual maturity occurs, reproductive organs begin to function, and secondary sexual characteristics develop. Postpubescence occurs 1 to 2 years after puberty, when skeletal growth is complete and reproductive function is regular.

Physical Changes

For girls, the first sign of sexual maturation is breast development. Menstruation generally begins within 2 years after the onset of breast development. For boys, sperm production (spermatogenesis) corresponds with increased testicular size and penile enlargement. Nocturnal emissions (“wet dreams”) typically start about 1 year after the penis begins to enlarge in size. Sexual maturity can be tracked using the Tanner scale. The achievement of Tanner stage 5 for both girls and boys corresponds with adult sexual maturity. (See Chapter 16, Assessing the Breasts, and Chapter 19, Assessing the Male Genitourinary System, for more on Tanner scales.)

Rapid physical growth is responsible for much of the clumsiness and awkwardness associated with this age period. Adolescents are faced with adjusting to rapidly changing physical bodies that may not always have the agility and grace associated with younger years. They often feel uncoordinated and ill at ease with their rapidly

changing physiques. A number of other physical changes occur that will be addressed later in this chapter.

Psychosocial and Emotional Changes

From a psychosocial perspective, Eric Erikson describes this period as one of “identity versus identity diffusion.” Adolescents become preoccupied with others’ opinions of them in an attempt to be accepted. The focus of influence shifts from family to peer group, and the adolescent begins to forge a sense of identity with the peer group. This is a normal developmental progression but one with potential conflicts, depending on the chosen peer group. If the peer group supports a sense of self-identity and self-worth, the outcome tends to be a positive one. But if the peer group discourages such internalization of values (such as with gangs), the adolescent does not progress in the direction of independent thought and decision making. Ideally, the adolescent moves through this stage developing a set of personal values and a sense of self-competency while beginning to plan for a future career.

Jean Piaget describes cognitive development during adolescence as the “formal operational stage.” This means that the adolescent begins a process of more logical thinking. He or she begins to hypothesize, wondering “what if?” and starts to reflect on outcomes to specific hypotheses. Previously, thought was based on very concrete concepts—what could be readily visualized or experienced. Concepts such as right versus wrong and good versus bad now become more abstract and situational. Other people’s concepts and views become more important, which only adds to the decision-making dilemma. This is a time when conflicts with parents begin to occur. Traditional family values may be challenged by the exposure to new ideas and values from peers.

During this time, the adolescent also struggles to develop her or his own set of moral principles by questioning established moral codes. Peers often have more influence than parents, but often preestablished values and morals persist. This is also a time for questioning spiritual beliefs, and many adolescents turn away from formal religion, eventually resolving the questions and identifying their sense of spirituality.

Adolescence is also a time of increased risk taking. This is a result of several factors: The desire to separate from parental influence, peer pressure and the need to belong, and a feeling of invulnerability or “it can’t happen to me.” Unprotected sexual activity, substance use, and unsafe driving are examples of risky behavior. Understanding this phenomenon is essential to effective counseling and education in the area of risk reduction.

Cultural and Ethnic Variations

Cultural influences may affect adolescent behaviors by establishing expectations.

Cultural and Ethnic Variations in Adolescents

CULTURAL GROUP INFLUENCE

Amish	<ul style="list-style-type: none"> Adolescent may want to break away from cultural norms (e.g., dress), but this is considered experimental. Teen expected to eventually assume adult role and adhere to prescribed norms.
Appalachian	<ul style="list-style-type: none"> Formal education not stressed, but teen expected to work and support family. Marriage at ages as young as 13 to 15 common. Having children associated with sense of importance. Underage alcohol abuse common.
Arab American	<ul style="list-style-type: none"> Chastity and decency expected.
Chinese American	<ul style="list-style-type: none"> Expected to score well in national tests by age 18. Must make career choice during adolescent years. Teenage pregnancy not common. Teens expected to respect elders.
Cuban American	<ul style="list-style-type: none"> Boys expected to learn a trade; girls expected to keep themselves honorable and prepare for marriage. “Rite of passage” (ready for courting) celebration at age 15 for girls. Unmarried couples may have chaperones.
Egyptian American	<ul style="list-style-type: none"> Loss of virginity affects marriageability, so parents are very strict.
Filipino American	<ul style="list-style-type: none"> Short courtships.
Greek American	<ul style="list-style-type: none"> Girls have less dating freedom than boys. Dating often prohibited until late adolescence.
Iranian American	<ul style="list-style-type: none"> Girls expected to maintain virginity.
Irish American	<ul style="list-style-type: none"> Expected to remain loyal to family.
Jewish American	<ul style="list-style-type: none"> Rite of passage at age 13 for boys, age 12 for girls.
Navajo Native American	<ul style="list-style-type: none"> Menarche for teenage girl seen as passage to adulthood. Older children taught to be stoic and not complain.
Vietnamese American	<ul style="list-style-type: none"> Teens expected to respect elders.

Introducing the Case Study



Margaret Chung, a pretty 14-year-old, has been in your school health office four times in the last 2 weeks, each time with a different vague complaint. So far, you have been unable to assess any specific malady. Today, when you ask what brings her there, she says that she just wants to talk. Several other students are waiting, so you escort Margaret to one of the private examination rooms and explain that you will return shortly, after seeing the students who were there first. After attending to them, you place an “Exam In Progress—Do Not Disturb” sign on the door, and then you begin your assessment.

Communicating during the assessment can be a challenge, but many adolescents are looking for an adult who will listen and not preach or talk down. For tips on talking to adolescents, see *Communicating with Adolescents*, page 924.

As with all assessments, the process involves subjective and objective data collection, diagnosis, planning, interventions, and evaluation.

Health History

Interview the adolescent alone, at least for some part of the examination. If a parent is present, explain that it is appropriate for young people to begin having independent relationships with their healthcare providers in preparation for managing their healthcare as adults. Because the history is essentially the same as for an adult, in this section differences will be presented as they relate specifically to the adolescent.

Biographical Data

Begin with standard biographical data, but also take note where the patient lives and with whom. Note who is listed the contact person—it may not be the parent. Do not assume anything.

Performing the Adolescent Assessment

Assessing an adolescent is similar to assessing an adult. As you begin the process, keep in mind the normal developmental changes and issues affecting adolescents.

Communicating with Adolescents

As a professional nurse, you may be the only receptive adult in an adolescent's life. Many young people feel that they have no adult to communicate with. Or they may be separating themselves from parental authority and moving toward self-autonomy, but still want and need adult guidance. The guidelines below will help you communicate with adolescents:

1. *Listen actively*: This may be the most effective strategy of all. Adolescents understand that adults will not always agree with them, but they desperately want a solid "sounding board." Listening can help you understand the issues at hand, even if you do not condone them. Here are some tips for active listening:
 - Start with less sensitive subjects and work your way up to more sensitive ones.
 - Make sure you and your patient understand each other, clarify as needed, and never assume.
 - Use gender-neutral terms and open-ended questions, and restate if necessary.
 - Show true concern and respect for the adolescent's perspective and opinion.
 - Be nonjudgmental.
2. *Help with decision making*: Listening actively puts you in a position to summarize the issues or questions and then help the adolescent verbalize the pros and cons of each side. This way, she or he can more objectively deal with the issue at hand and is more likely to make a logical, informed decision.
3. *Avoid scare tactics*: Adults often use these in an attempt to control behavior. Most young people know that one puff on a cigarette will not addict them for life, but adults often make statements like this. Such comments can destroy your credibility and make the patient less likely to follow your advice on other health issues.
4. *Maintain confidentiality*: This is a major issue. Make sure the female patient knows she is guaranteed confidentiality in pregnancy testing, contraceptive services, and diagnosis and treatment of STDs. Then keep your promises. If the patient has expressed the desire to harm himself or herself or others, be sure that he or she knows this is one problem you *cannot* keep confidential.

Current Health Status

Be sure to ask about all prescription and OTC medications. Many adolescents (especially those involved in athletics) take vitamin or protein supplements, so assessing for these is important. Ask girls if they are on birth control pills (oral contraceptives), take monthly "shots" (Depoprovera), or have a contraceptive implant (Norplant). Also ask girls if they take the acne medication Acutane. If they do, tell them that taking Acutane while pregnant can cause serious birth defects, and counsel them about pregnancy prevention.

Past Health History

Because the adolescent may not be aware of all his or her past health history, you may need to check with a parent or guardian for specifics such as immunizations (see Health Concerns, p. 924, for required immunizations).

Family History

If the adolescent is being interviewed alone, she or he may have limited knowledge in this area. Many, however, will be aware of heart disease, stroke, cancer, and other serious health conditions within the family. It is also important to ask if anyone in the family died before age

35 from a sudden heart attack. This may be a risk factor for sports participation and may require additional investigation.

Review of Systems

Review every system, keeping in mind changes or problems specific to the adolescent.

Psychosocial Profile

The psychosocial profile focuses on how the adolescent's health practices and behaviors affect his or her health and well-being.

Health Practices and Beliefs and Self-Care Activities

When did the patient have her or his last health examination (every 2 years is recommended). More frequent examinations may be needed if she or he has an existing health problem or is involved in sports. Keep in mind that adolescents may have no control over the frequency of their health screenings. The parent or guardian often assumes this responsibility, and health insurance or the lack of it may be an influencing factor. You may need to make appropriate referrals or provide information on available health resources.

Review of Systems

SYSTEM/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

General Health Survey

- How have you been feeling?
- Have you had any weight changes?

- Screens for any obvious problems.
- *Weight loss or gain*: May indicate eating disorder.

Integumentary

- Do you have any skin problems, such as acne?
- Have you had changes in body hair growth?
- Do you have piercings or tattoos?
- Do you use sunscreen?
- Do you lie in the sun or go to tanning salons?

- Adolescents are very self-conscious about appearance, and acne may affect self-image.
- Secondary sexual changes include increase in body hair—pubic hair, axilla and leg hair, and facial hair for boys.
- *Body piercing and tattooing*: Increased risk for HIV or hepB.
- *Excessive exposure to sun (ultraviolet rays) without sunscreen*: Increased risk for skin cancer later in life.

HEENT

Eyes

- Do you have any vision problems?
- Do you wear glasses or contact lenses?
- When was your last eye exam?

- Visual refractive problems peak during adolescence.
- Identifies teaching need regarding care of contact lenses.
- Test visual acuity routinely during physical exams.

Ears

- Do you have any hearing problems?
- Do you listen to loud music?
- When was your last hearing exam?

- Identifies teaching need regarding noise pollution's affect on hearing.
- Cochlear damage can result from excessive exposure to loud noise such as CD players and rock concerts.
- Test hearing routinely during physical exams.

Mouth

- When was your last visit to the dentist?
- Do you go to an orthodontist?

- Should have at least annual dental exam.
- Need for orthodontia usually becomes apparent during adolescence and can make patient even more self-conscious about appearance.

Respiratory

- Do you have asthma?
- Do you ever have trouble breathing or wheeze when you exercise or run?
- Are you exposed to air pollutants, such as smoke and second-hand smoke?

- Asthma is the most common disease among children.
- Exercise-induced bronchospasm is an acute, reversible, self-terminating airway obstruction that occurs after vigorous exercise. It peaks 5 to 10 minutes after activity stops, and then stops within 30 minutes.

Cardiovascular

- Do you have chest pain?
- Does your heart ever skip a beat?

- Although cardiovascular disease is not common in adolescents, it can occur.
- Prolonged QT syndrome can cause sudden cardiac death in a seemingly healthy adolescent.

Breasts

Girls

- Are your breasts tender?
- Have they increased in size?
- Are they growing equally?

- Determines whether normal breast development is occurring. May identify areas for health teaching. Initially, breasts may develop asymmetrically, but they should even out eventually.

Boys

- Have your breasts enlarged?

- During early puberty, boys may experience temporary gynecomastia (breast enlargement), which usually disappears within 2 years. Because their appearance is of great concern, gynecomastia can be very alarming for boys and make them very self-conscious.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
Gastrointestinal <ul style="list-style-type: none"> ■ Do you have any stomach problems? ■ How often do you have a bowel movement? ■ Do you use laxatives? 	<ul style="list-style-type: none"> ■ Identifies underlying eating disorder. ■ Excessive use of laxatives associated with eating disorders.
Genitourinary/Reproductive <i>Girls</i> <ul style="list-style-type: none"> ■ Did your periods start? When? How often do you have them? How heavy is the flow? ■ Do you use pads or tampons? ■ Do you have burning when you go to the bathroom? ■ Have you had urinary tract infections (UTIs)? 	<ul style="list-style-type: none"> ■ Identifies start of menarche and determines if cycle is normal. ■ Tampon use has been associated with risk for toxic shock syndrome (TSS). ■ May identify health teaching needs. ■ Frequent UTIs in girls can be caused by sexual activity and require follow-up.
<i>Girls and Boys</i> <ul style="list-style-type: none"> ■ Are you sexually active? ■ Have you ever had sex? ■ Are you practicing safe sex? 	<ul style="list-style-type: none"> ■ Identifies health teaching need.
Musculoskeletal <ul style="list-style-type: none"> ■ Do you have any back problems? ■ Have you ever been told you had a spinal problem? 	<ul style="list-style-type: none"> ■ Heavy backpacks are associated with low back problems in children and adolescents. ■ Scoliosis (lateral curvature of spine) occurs more frequently in girls than in boys. Screening usually occurs in middle school and high school. If it is present, determine if problem is structural or postural (see Chapter 20, Assessing the Motor-Musculoskeletal System).
Neurologic <ul style="list-style-type: none"> ■ How would you describe your usual mood? ■ Do you feel sad a lot? 	<ul style="list-style-type: none"> ■ Adolescence is a time of change, and emotional response to these changes varies. ■ Depression or suicidal ideations must be identified and treated.
Endocrine <ul style="list-style-type: none"> ■ Do you have any swelling in your neck? Difficulty swallowing? Hoarseness? 	<ul style="list-style-type: none"> ■ The thyroid is more active during puberty. ■ <i>Enlarged thyroid:</i> Hypo- or hyperthyroid disease. Lymphocytic thyroiditis (Hashimoto's disease or juvenile autoimmune thyroiditis) is most common in children after age 6 and reaches peak occurrence in young people during adolescence. ■ Graves' disease occurs in young people between ages 12 and 14. ■ If sexual maturation is not progressing, consider endocrine problems.
Lymphatic/Hematologic <ul style="list-style-type: none"> ■ Have you been tired? ■ Do you have any lumps in your neck, underarms, or groin? 	<ul style="list-style-type: none"> ■ Rapid growth and poor dietary habits increase risk of iron deficiency anemia. ■ Hodgkin's lymphoma is prevalent between ages 15 and 19. ■ Non-Hodgkin's lymphomas occur more frequently before age 15.

Typical Day

What constitutes a usual day's activities? How is the adolescent doing in school? What subjects does he or she like and dislike? Has the adolescent sought help for subjects that he or she is having problems with? This allows you to identify stresses confronting the patient in everyday life. A change in academic performance or a sudden disinterest in school may reflect a more serious problem. How many days of school has he or she missed this term? High absenteeism (except for serious illness) correlates with other risky behaviors.

Nutritional Patterns, Weight Loss, and Weight Gain

Ask for a 24-hour diet recall. Then ask questions about body image, such as "How much do you think you should weigh?" Further explore this issue if the response is unrealistic. Ask about eating patterns, such as "How often do you eat breakfast?" "What is a usual lunch for you?" and "What are your favorite snacks?" These open-ended questions will provide a much clearer picture of the adolescent's nutritional status and potential risks.

Vitamin and Mineral Supplements

At this age, dietary needs for calcium, iron, and zinc are greater. Calcium is needed for bone growth and bone mass development during adolescence; it also influences the development of osteoporosis later in life. Assess both overeating and undereating problems. Girls are usually more concerned about losing weight, whereas boys may want to "bulk-up" and improve strength.

Activity and Exercise Patterns

Ask what the patient enjoys doing outside of the home. What sports does she or he play? What does she or he like to do with her or his friends? How much TV does she or he watch? How much time does she or he spend on the computer? You may identify a need for health education about the importance of routine exercise. If the teen does participate in sports, ask if she or he uses protective equipment and if it is a contact sport. Contact sports may increase the risk for injury.

Recreation and Hobbies

What does the patient do for fun? If activities increase the risk for injury, ask if protective measures, such as bike helmets, are used. Also ask about weapons in the home and hunting safety courses, if appropriate.

Sleep and Rest Patterns

What time does the patient usually go to bed and wake up? Does he or she feel refreshed in the morning or still tired? How does this pattern change on weekends? Even though sleep needs increase during growth spurts, teens often get less-than-ideal amounts of sleep during the week and sleep late on weekends. This can disrupt

healthy sleep patterns. Also ask about bedwetting, which is not uncommon at this age, especially in boys. Phrase your question in a nonthreatening way to decrease embarrassment. For example, say: "Many young people find they sometimes wet themselves during their sleep. Does this ever happen to you?"

Personal Habits

Ask if the patient uses drugs, alcohol, or nicotine, and phrase your questions in a nonthreatening way to increase the likelihood of an honest response. For example, say: "If you smoke cigarettes, how many do you smoke in an average day?" "If you use alcohol, how much do you usually drink during the week?" "If you use drugs, what type and what method do you use to take them?" Methods for taking drugs include smoking, snorting, huffing, ingesting, and injecting.

Risky Behavior

Accidents are the leading cause of death among teenagers, especially motor vehicle accidents. If the patient drives, ask if she or he attended a driver's education program. Does she or he wear a seat belt? Is she or he aware of the hazards of driving under the influence of drugs or alcohol? Take this opportunity to reinforce the importance of not driving under the influence or riding with someone under the influence. If drug or alcohol use is identified, make appropriate referrals.

Roles, Relationships, and Self-Concept

Ask the young person to tell you what he or she likes best about himself or herself and what he or she would like to change. Remember that adolescence is a time of great change. These questions will help you identify how satisfied the patient is with the changes that have occurred. Also ask about family relationships. For example, ask: "Who lives at home with you?" "How do you get along with your family?" "Is there an adult in your life whom you feel comfortable talking with?" Ask about peer relationships. Does the patient have a special friend, and what things does he or she enjoy doing with that friend? Does he or she date?

Religious and Spiritual Influences

Ask the adolescent what is important to her or him and whether she or he believes in God or a higher power. Established spiritual beliefs may be a source of support and influence behavior.

Sexuality Patterns

Ask girls if they have any questions or concerns about menstrual issues. Ask boys if they have any questions about physical changes. Inquire about sexual activity. You might ask: "If you are sexually active, how do you protect yourself against STDs and unwanted pregnancy?" Keep in

mind that sexual preferences become apparent during adolescence, and gay or bisexual teens may be confronted with homophobic attitudes.

Stress and Coping

Ask: “What do you do when you get angry?” “What makes you angry?” “What do you do to relax?” Identify whether the adolescent has healthy coping strategies to deal with life’s stressors. Violence is on the rise among teenagers. If your patient uses violence as a coping mechanism, you may need to make referrals to help him or her develop more effective coping strategies.

Physical Assessment

After you complete the history, perform the physical examination. Physical assessment of the adolescent does not differ greatly from that of the adult—you use all four

assessment techniques and the same tools (for more information, see Chapter 3, Approach to the Physical Assessment).

Begin with a general survey, obtaining height and weight (Figs. 26.3 and 26.4). Also take vital signs, which are essentially the same as for adults. Then perform a head-to-toe physical examination.



HELPFUL HINT

Always weigh adolescents in private. When weighing is done in front of other people—for example, in school settings—it is a source of anxiety for teens, especially those who are overweight or think they are.

(Text continued on page 924)



FIGURE 26.3. Weighing the patient.



FIGURE 26.4. Measuring the patient.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

1. General Health Survey

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect overall appearance, noting appropriate growth and development for age of patient.
- *Growth and development appropriate.*

ABNORMAL FINDINGS/RATIONALE

- Detects delays or premature sexual maturation.
- Detects obvious weight problems.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

2. Integumentary

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect skin for lesions.
- Note piercings and tattoos.
- Note body hair distribution.
- *Skin intact, no lesions.*
- *Hair distribution appropriate for developmental age of child.*

ABNORMAL FINDINGS/RATIONALE

- Acne is a common problem and a major concern. This condition should be treated and not dismissed as insignificant. Severe acne can result in permanent scarring. Many effective are now available.
- Piercings and tattoos increase risk for HIV and hepB.
- Absent or excessive body hair may relate sexual maturation problem.

3. HEENT

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

Eyes

- Test visual acuity.
- *Vision 20/20.*

Ears

- Test hearing with pure tone audiometer.
- *Hearing intact.*

Nose

- Inspect nasal septum and mucosa.
- *Nasal mucosa pink, intact.*

Mouth

- Inspect oral mucosa and pharynx.
- Inspect occlusion.

- Inspect number of teeth and check for wisdom teeth.
- Assess for orthodontic devices and brushing technique around them.
- *Hypertrophied tonsils of early childhood have usually shrunk to more normal proportions.*
- *No malocclusion.*
- *Should have 28 permanent teeth. Third molars (“wisdom teeth”) may erupt during this period.*

Neck

- Palpate thyroid.
- *Thyroid nonpalpable.*

ABNORMAL FINDINGS/RATIONALE

- Many teenagers have some degree of myopia (near-sightedness) but are reluctant to wear prescribed eyeglasses.
- Listening to loud music, especially with earphones, can cause early hearing loss. Use opportunity to teach hazards of listening to loud music.
- *Boggy, bluish-purple turbinates:* Chronic rhinorrhea caused by allergic rhinitis.
- *Mucosal damage:* Possible cocaine use, which also causes rhinorrhea.
- Enlarged, red tonsils with exudates may indicate an infection, such as strep throat or mononucleosis.
- If malocclusion is present, refer to orthodontist.
- May need referral if wisdom teeth are impacted.
- *Improper brushing:* Increased risk for gum problems and cavities.
- *Tender, enlarged thyroid gland:* Acute thyroiditis.

4. Respiratory

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Auscultate lungs.
- *Lungs clear.*

ABNORMAL FINDINGS/RATIONALE

- Adolescents with EIA require bronchodilators before activity (usually albuterol via a metered-dose inhaler 20 to 30 minutes before exercise).

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

5. Cardiovascular

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Auscultate heart and note rate and rhythm.
- *Heart regular rhythm, no extra sounds.*
- *Physiological S_3 , sinus arrhythmia and split S_2 that change with respiration are normal variations.*



Auscultating the heart

ABNORMAL FINDINGS/RATIONALE

- Murmurs warrant evaluation.

6. Breasts

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect and palpate breasts.
- *Breast development appropriate for age and gender of child.*
- *Breast examinations should begin during adolescence, along with instruction in breast self-examination.*

ABNORMAL FINDINGS/RATIONALE

- Abnormal breast development should be followed-up.
- Gynecomastia occurs in about one-third of boys.

7. Gastrointestinal

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect, auscultate, and palpate the abdomen.
- *Abdomen soft, nontender, no organomegaly.*

ABNORMAL FINDINGS/RATIONALE

- Except for pregnancy, no specific variations occur with the abdomen. It is not unusual to detect pregnancy during a routine exam in an adolescent who is denying it. By 20 weeks' gestation, the fundus is at level of the umbilicus.

8. Genitourinary/Reproductive

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect external genitalia.
- Genitalia intact, no lesions or discharge.
- *Boys:* Perform testicular exam.
- *Girls:* Perform pelvic exam at age 21 or 3 years after onset of sexual activity.
- Note Tanner stage for boys and girls.
- Instruct boys in testicular self-exam.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

8. Genitourinary/Reproductive (continued)

ABNORMAL FINDINGS/RATIONALE

- Examine sexually active adolescents for STDs.
- Testicular mass.
- *Pubertal delay for boys:* No changes in testes or scrotum by age 14 or incomplete genital growth within 4 years of initial change.
- *Pubertal delay for girls:* No breast development by age 13 or no menarche within 4 years of initial breast changes.

9. Musculoskeletal

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect and palpate spinal curves.
- Test for spinal deformities.
- If patient is active in sports, test knee strength and stability.
- Test muscle strength grip.
- *No spinal deformities.*
- *Knee joints stable.*
- *+ 5 muscle strength, no weakness.*
- *For more information on testing for scoliosis and muscle strength, see Chapter 20, Assessing the Motor-Musculoskeletal System.*



Inspecting for scoliosis



Testing muscle strength



Testing knee strength and stability

ABNORMAL FINDINGS/RATIONALE

- Scoliosis is a major variation within the musculoskeletal system. Screening should be done in the preadolescent period, generally during fifth and sixth grades. Refer children with significant curvature for evaluation and follow-up.
- Muscle weakness, joint instability warrant follow-up.

10. Neurological

SYSTEM/ASSESSMENT/NORMAL VARIATIONS

- Inspect the patient, focusing on affect and cognitive functioning.
- *Affect appropriate, doing well in school, cognitive functioning intact.*

ABNORMAL FINDINGS/RATIONALE

- Depression and suicidal ideations warrant immediate intervention.

Now that your assessment is over, analyze your data and formulate nursing diagnoses. If you identify problem areas, actively involve both the adolescent and her or his parents in the plan of care.



Back to the Case Study

Now that you understand the physical and psychosocial issues surrounding adolescents, apply your knowledge to our case study subject, 14-year-old Margaret Chung, by answering the following questions:



CRITICAL THINKING ACTIVITY #5

How does assessment of the adolescent differ from assessment of other patients?



CRITICAL THINKING ACTIVITY #6

What approach is best for assessing the adolescent?



CRITICAL THINKING ACTIVITY #7

What health concerns should you keep in mind as you proceed with the assessment?



CRITICAL THINKING ACTIVITY #8

Some nursing diagnoses are appropriate for all adolescents. What are they?

RESEARCH TELLS US

Childhood obesity poses a major health problem in the United States, with 10 to 15 percent of children being affected. *Overweight* is the preferred term to be used with children rather than obesity. A child with a body mass index (BMI) between the 85th and the 95th percentiles places him or her at risk for being overweight. A child with a BMI greater than the 95th percentile is considered overweight.

Two critical growth and development periods have been identified for overweight development. The first occurs between ages 4 to 6, when the child's BMI rises owing to growth spurts. The second occurs during adolescence owing to the increase in body fat secondary to puberty.

Researchers have also linked cultural and ethnic backgrounds and socioeconomic status to overweight risk. American Indian, Hispanic, and African American children have the highest overweight rates. A genetic link associated with neuropeptide Y has been identified with overweight Mexican American children. Socioeconomic status is inversely proportional to weight; the lower socioeconomic

status the greater the risk for higher being overweight.

Additional contributing factors have also been identified:

- Sedentary lifestyle, including increased TV viewing.
- Food preferences, availability, and family eating patterns, including an increase in consumption of fat foods and decreased consumption of fruits and vegetables.

The effects of childhood overweight are immediate and long-lasting. Immediate effects of childhood overweight include the development of metabolic syndrome, type 2 diabetes, orthopedic problems, and psychosocial disorders. Half of overweight adolescents will become obese as an adult. The long-term effects include the development of cerebrovascular disease, HTN, and hyperlipidemia.

Nurses are well positioned in the community to screen children for overweight problems, develop programs, and educate children and their families in preventing and treating this major health problem.

Source: Hardy, L., Harrell, J., and Bell, R. (2004). Overweight children: Definitions, measurements, confounding factors and health consequences. *Journal of Pediatric Nursing* 19(6):376-384.

HEALTH CONCERNS

Immunizations recommended during adolescence include:

- *Tetanus-diphtheria "booster" (Td)*: Between ages 11 and 12 and every 10 years thereafter; whenever person suffers a wound, if booster not received

within past 5 years.

- *MMR*: Between ages 11 and 12 if second MMR not given between ages 4 and 6.
- *VZV*: At any age if person has never had chickenpox; if vaccine not given until age 13 or older, should get two doses, 4 to 8 weeks apart.

HEALTH CONCERNS (continued)

- *HepB*: Any time between ages 6 and 12, if not immunized during infancy. Children between ages 11 and 15 may need only two doses, 4 to 6 months apart.
- *HiB*: Not recommended for healthy adolescents. Children over age 5 usually do not need this vaccine unless they have special health concerns, such as sickle cell disease, HIV/AIDS, removal of spleen, bone marrow transplant, or cancer treatment with drugs.
- *Meningitis (Menomune)*: Before starting college.
- *Gardasil vaccine*: for four types of HPV (which cause 70% cervical cancer and 90% genital warts) between ages 9 and 26. Series of three shots over six month period.

COMMON PROBLEMS

Common problems in this age group include acne, eating disorders, scoliosis, pregnancy, STDs, injuries

and accidents, substance abuse, malocclusion, exercise-induced asthma (EIA), and pubertal delay.


S U M M A R Y

- Managing adolescent healthcare presents challenging issues for the nurse.
- The key to success lies in effective communication and understanding the developmental and psychosocial issues related to adolescence.
- Having a foundational understanding of these concepts will help you develop an effective plan of care.

R E V I E W Q U E S T I O N S

1. You are performing a vision assessment on Regina Davis, age 14. Which vision change is frequently seen in adolescence?
 - a. Hyperopia
 - b. Myopia
 - c. Strabismus
 - d. Amblyopia
2. As you proceed with Regina's physical exam and examine the head, face, and neck, what structure may be palpable owing to hormonal changes associated with puberty?
 - a. Submaxillary glands
 - b. Parotid glands
 - c. Thyroid gland
 - d. Lymph nodes
3. According to Erikson, what is the developmental task for the school-age child, ages 6 to 12 years?
 - a. Industry vs. inferiority
 - b. Intimacy vs. isolation
 - c. Initiative vs. guilt
 - d. Identity vs. role diffusion
4. According to Erikson, what is the developmental task for the adolescent, ages 12 to 18 years?
 - a. Industry vs. inferiority
 - b. Intimacy vs. isolation
 - c. Initiative vs. guilt
 - d. Identity vs. role diffusion
5. What common spinal deformity is seen in school-age children and adolescents?
 - a. Kyphosis
 - b. Lordosis
 - c. Scoliosis
6. Jack Warner, age 18, will be starting college in the fall. Which immunization should be given to adolescents prior to college?
 - a. HepB
 - b. Tetanus
 - c. Chickenpox
 - d. Meningitis
7. What common breast problem is seen in adolescent boys as they proceed through puberty?
 - a. Adenomas
 - b. Gynecomastia
 - c. Mastitis
 - d. Galactorrhea

CHAPTER 27



Assessing the Older Adult

Learning Outcomes

After you have successfully completed this chapter, you should be able to:

- Identify normal physiological changes of the older adult
- Discuss various cultural influences on the older adult
- Identify developmental tasks for the older adult
- Identify health issues and risks for the older adult
- Identify pertinent history data included in an older adult history
- Obtain an older adult health history
- Describe additional physical assessment components specific to an older adult assessment
- Perform an older adult assessment
- Document older adult assessment findings
- Identify actual/potential health problems stated as nursing diagnoses with supporting data

Obtaining an accurate and useful history and physical assessment is a vital skill in working with all adults, but it is particularly challenging in working with older adults. This challenge is a result of the normal changes that occur with aging, the tendency of older people to present with disease in an atypical fashion, their long medical histories, and common communication problems caused of hearing loss, aphasia, or cognitive impairment. The actual assessment process does not differ from that used with any other adult. You just need to focus your attention on the changes that occur with aging. Understanding the normal changes that occur with age and how they interrelate with disease and its presentation will improve your assessment skills and facilitate the diag-

nosis and management of common problems in older patients.

Anatomy and Physiology Review

In evaluating the older adult, it is often difficult to differentiate between normal changes and those caused by environmental factors, lifestyle, or disease. For example, skin changes such as wrinkling, which were once believed to be related exclusively to aging, are now believed to be influenced by sun exposure. Understanding the normal changes associated with aging in each body system and being familiar with them are essential to assessing older adults.

(Text continued on page 932)

Normal Changes Associated With Aging

SYSTEM OR STRUCTURE/CAUSES	CHANGES
Integumentary <i>Skin</i> <ul style="list-style-type: none"> ■ Decreased collagen and subcutaneous fat ■ Atrophy of sweat glands and decreased function ■ Decline in fibroblast proliferation, cell production, and epidermal turnover ■ Capillary fragility and decreased vascularity ■ Decreased sensory receptors and increased thresholds 	<ul style="list-style-type: none"> ■ Increased wrinkling ■ Decreased elasticity ■ Increased dryness ■ Pruritus ■ Thinning ■ Increased healing time ■ Bruising ■ Decreased sensory perception ■ Decreased vitamin D production ■ Increased skin lesions
<i>Hair</i> <ul style="list-style-type: none"> ■ Decreased melanocytes ■ Decreased hair follicle density 	<ul style="list-style-type: none"> ■ Graying of body hair ■ Uneven skin color ■ Loss and thinning of hair
<i>Nails</i> <ul style="list-style-type: none"> ■ Hypo/hyperplasia of nail matrix ■ Decreased blood supply to nails 	<ul style="list-style-type: none"> ■ Increased longitudinal ridges ■ Nails thick and brittle ■ Growth slow
HEENT <i>Eyes</i> <ul style="list-style-type: none"> ■ Decreased orbital fat ■ Decreased elasticity of lids ■ Decreased tears ■ Decreased corneal sensitivity ■ Increased lipid deposits around cornea ■ Decreased aqueous humor ■ Atrophy of ciliary muscles ■ Decreased elasticity of lens ■ Increased density of lens ■ Decreased color of iris ■ Decreased pupil size ■ Increased vitreous debris 	<ul style="list-style-type: none"> ■ Sunken eyes ■ Ectropion or entropion ■ Dry eyes ■ Decreased corneal reflex ■ Arcus senilis ■ Decreased lens accommodation ■ Decreased peripheral vision ■ Decreased ability to adapt to light and dark ■ Glare intolerance ■ Impaired night vision ■ Decreased visual acuity ■ Floaters

SYSTEM OR STRUCTURE/CAUSES	CHANGES
<i>Ears</i>	
<ul style="list-style-type: none"> ■ Increased external canal hair in men ■ Decreased cerumen ■ Degeneration of middle ear bones ■ Thickened tympanic membrane ■ Decreased hair in inner ear ■ Atrophy of cochlea and organ of Corti 	<ul style="list-style-type: none"> ■ Conductive hearing loss ■ Decreased speech discrimination ■ Difficulty hearing higher-frequency sound
<i>Nose</i>	
<ul style="list-style-type: none"> ■ Atrophic changes 	<ul style="list-style-type: none"> ■ Vasomotor rhinitis ■ Decrease in sense of smell and ability to distinguish odors
Respiratory	
<ul style="list-style-type: none"> ■ Rigid ribs and thoracic wall ■ Decreased muscle strength ■ Atrophy of cilia ■ Decreased elastic recoil ■ Decreased pulmonary bed ■ Thickening and decrease in number of alveoli ■ Decreased response to hypoxia/hypercarbia 	<ul style="list-style-type: none"> ■ Increased anterior-posterior diameter ■ Senile kyphosis ■ Decreased vital capacity ■ Increased residual lung capacity ■ Reduced cough and clearing ■ Decreased lung compliance ■ Decreased ventilation and perfusion ■ Decreased PaO₂ and O₂ saturation ■ More difficulty in maintaining acid-base balance
Cardiovascular	
<i>Heart</i>	
<ul style="list-style-type: none"> ■ Decreased cardiac output and cardiac index ■ Decreased response to beta-adrenergic stimulation ■ Decreased heart muscle with increase in fat and collagen ■ Thickening of ventricular walls ■ Decreased compliance ■ Increased dependence on atrial contraction ■ Calcification of valves ■ Decreased sinoatrial node pacer cells and bundle of His fibers 	<ul style="list-style-type: none"> ■ Decreased stroke volume and output ■ Increased myocardial oxygen demands ■ Ventricular hypertrophy ■ S₄ (see Chapter 14, Assessing the Cardiovascular System) ■ Murmurs ■ Arrhythmias ■ Slower rates in response to stress
<i>Arteries</i>	
<ul style="list-style-type: none"> ■ Decreased elastin and smooth muscle ■ Decreased compliance and stiffness of vessels ■ Increased peripheral vascular resistance ■ Aortic dilatation ■ Decreased baroreceptor response ■ Rigidity of arteries leading to decreased peripheral circulation 	<ul style="list-style-type: none"> ■ Increased BP ■ Orthostatic hypotension ■ Decreased pulses ■ Cool temperature
<i>Veins</i>	
<ul style="list-style-type: none"> ■ Increased tortuosity 	<ul style="list-style-type: none"> ■ Varicosities
Gastrointestinal	
<i>Mouth and Teeth</i>	
<ul style="list-style-type: none"> ■ Decreased dentine ■ Gingival recession ■ Decreased papillae on tongue ■ Increased threshold for tasting salt and sugar ■ Decreased saliva 	<ul style="list-style-type: none"> ■ Potential loss of teeth ■ Decreased sense of taste ■ Dry oral mucous membranes

(continued)

Normal Changes Associated With Aging <i>(continued)</i>	
SYSTEM OR STRUCTURE/CAUSES	CHANGES
Esophagus <ul style="list-style-type: none"> ■ Decreased sphincter pressure ■ Decreased motility 	<ul style="list-style-type: none"> ■ Heartburn ■ Dysphagia ■ Increased risk for hiatal hernia, gastroesophageal reflux disease (GERD), and aspiration
Stomach <ul style="list-style-type: none"> ■ Decreased gastric acid and hydrochloric acid ■ Atrophy of mucosa ■ Decreased blood flow ■ Delayed emptying 	<ul style="list-style-type: none"> ■ Decreased absorption of iron, B12, and calcium ■ Food intolerance ■ Decreased hunger ■ Weight changes
Small Intestine <ul style="list-style-type: none"> ■ Decreased villae, enzymes, and motility 	<ul style="list-style-type: none"> ■ Decreased absorption of nutrients and fat-soluble vitamins
Large Intestine <ul style="list-style-type: none"> ■ Decreased blood flow and motility ■ Decreased sensation of need to defecate 	<ul style="list-style-type: none"> ■ Constipation ■ Increased risk for diverticular disease
Liver <ul style="list-style-type: none"> ■ Decrease in number and size of cells ■ Decreased protein synthesis ■ Decreased regeneration 	<ul style="list-style-type: none"> ■ Decreased drug metabolism and ability to detoxify
Pancreas <ul style="list-style-type: none"> ■ Decreased lipase and reserve 	<ul style="list-style-type: none"> ■ Impaired fat absorption ■ Possible glucose intolerance
Kidneys <ul style="list-style-type: none"> ■ Decreased renal mass, nephrons, glomerular filtration rate, blood flow 	<ul style="list-style-type: none"> ■ Decreased ability to concentrate urine, resulting in loss of free water and increased sensitivity to salt ■ Decreased creatinine clearance ■ Increased blood urea nitrogen ■ Decreased toxins and drug clearance
Bladder <ul style="list-style-type: none"> ■ Decreased smooth muscle and elastic tissue ■ Decreased sphincter control 	<ul style="list-style-type: none"> ■ Decreased control and possible incontinence ■ Decreased capacity ■ Increased frequency, urgency, and nocturia
Female Reproductive <ul style="list-style-type: none"> ■ Decreased hormones ■ Decreased size of ovaries and uterus ■ Decreased pelvic elasticity ■ Atrophy and fibrosis of cervical and uterine walls ■ Decreased elasticity of vagina ■ Vaginal secretions pH alkaline ■ Involution of mammary gland tissue ■ Decreased elasticity and subcutaneous tissue ■ Increased adipose tissue 	<ul style="list-style-type: none"> ■ Thin, pale vaginal mucosa ■ Decreased vaginal secretions ■ Decreased intensity of sexual response ■ Potential for prolapses and infections ■ Sagging of breasts ■ Possible stringy feeling of mammary ducts
Male Reproductive <ul style="list-style-type: none"> ■ Enlarged prostate ■ Decreased sperm count and seminal fluid volume ■ Seminal vesicles atrophy ■ Increased estrogen levels ■ Decreased testosterone ■ Reduced elevation and decreased size of testes 	<ul style="list-style-type: none"> ■ Prostatic hypertrophy ■ Decreased intensity of sexual response ■ Increased time to achieve erection ■ Decreased force of ejaculation ■ Tendency of testes to hang lower ■ Gynecomastia

SYSTEM OR STRUCTURE/CAUSES	CHANGES
Musculoskeletal	
<i>Bones</i>	
<ul style="list-style-type: none"> ■ Narrow intervertebral discs ■ Increased cartilage in nose and ears ■ Decreased bone mass, bone growth, and osteoblastic activity 	<ul style="list-style-type: none"> ■ Loss of height (1–4 inches) ■ Kyphosis ■ Wider pelvis ■ Increased length of nose and ears ■ Increased risk for osteoporotic fractures
<i>Muscles</i>	
<ul style="list-style-type: none"> ■ Decreased number of muscle fibers ■ Muscle atrophy ■ Increased fat in muscles ■ Slow muscle regeneration ■ Stiffening of ligaments and tendons ■ Increased contraction and latency time 	<ul style="list-style-type: none"> ■ Decreased strength ■ Decreased agility
<i>Joints</i>	
<ul style="list-style-type: none"> ■ Decreased cartilage ■ Increased erosion and calcium deposits 	<ul style="list-style-type: none"> ■ Decreased ROM and mobility ■ Osteoarthritis
Neurological	
<i>Brain</i>	
<ul style="list-style-type: none"> ■ Decreased brain size, weight, and volume ■ Decreased neurons, glial cells, and conduction of nerve fibers ■ Neurofibrillary tangles ■ Hypoperfusion ■ Atrophy ■ Decreased neurotransmitters, dopamine, norepinephrine, serotonin, and acetylcholine ■ Elevated cortisol, sodium, and monoamine oxidase levels ■ Decreased deep sleep and rapid eye movement (REM) sleep 	<ul style="list-style-type: none"> ■ Decreased processing and reflexes ■ Delayed reaction time ■ Decreased psychomotor performance ■ Depression ■ Altered pain response ■ Decreased proprioception ■ Increased balance problems ■ Decreased sensory input ■ Increased periods of being awake and difficulty falling asleep ■ Decreased dreaming
Endocrine	
<ul style="list-style-type: none"> ■ Decreased BMR ■ Decreased sensitivity to hormones ■ Decreased febrile response ■ Decrease in hormones (e. g., growth, thyroid) 	<ul style="list-style-type: none"> ■ Increased weight ■ Decreased insulin response, glucose response, glucose tolerance and sensitivity of the renal tubules to antidiuretic hormone (ADH) ■ Decreased shivering and sweating ■ Effects of hormonal change
Immunologic/Hematologic	
<ul style="list-style-type: none"> ■ Decreased immunoglobulin IgA ■ Involved thymus ■ Decreased thymopoietin, lymphoid, antibodies, T lymphocytes ■ Increased autoantibodies ■ Decreased memory of previous antigenic stimuli ■ Decreased responsiveness to immunizations ■ Increased energy 	<ul style="list-style-type: none"> ■ Decreased ability to reject foreign substances ■ Increased autoimmune disorders ■ Delayed hypersensitivity reactions ■ Decreased response to acute infection ■ Increased incidence of malignancy ■ Recurrent latent herpes zoster or tuberculosis

Developmental, Cultural, and Ethnic Variations

Assessing your patient's developmental status is part of the assessment process. All humans continue to grow and develop throughout their lifetime. Erikson's psychosocial development theory identifies the final stage of development as ego integrity versus despair. Ideally, adults over age 65 will be able to do a life review with a sense of satisfaction and accomplishment and be able to accept their own mortality. This stage of life correlates well with Maslow's Hierarchy of Human Needs' self-actualization need, in which self-fulfillment of one's potential is actualized. Many factors over a lifetime influence whether the older adult will attain self-actualization, a sense of accomplishment and satisfaction or despair. You need to be sensitive to your patient's

response to aging and assist him or her in achieving successful resolution of this final stage of development.

The definition and value of aging are dependent on culture. Aging can be defined in years, functional abilities, or social mores. Understanding this cultural influence may be helpful in planning care of the older patient.



CRITICAL THINKING ACTIVITY # 1

Consider the different cultural perspectives that follow. What impact do you think each of these perspectives has on the way families provide health-care for their elderly members?

Cultural Perspectives on Aging

CULTURAL GROUP	VIEW OF AGING
African American	<ul style="list-style-type: none"> ■ Elderly people are valued and treated with respect. ■ The grandmother has a central role and often offers economic support and child care. ■ Grandchildren are often raised by the grandmother.
Appalachian	<ul style="list-style-type: none"> ■ Elderly people are respected and honored, live close to their children, and participate in care of grandchildren.
Brazilian American	<ul style="list-style-type: none"> ■ Elderly parents live with children and are included in all activities.
Chinese American	<ul style="list-style-type: none"> ■ Elderly people are highly valued and respected and are considered very wise. ■ Children are expected to care for parents.
Cuban American	<ul style="list-style-type: none"> ■ Elderly parents live with children. Multigenerational households are common.
Egyptian American	<ul style="list-style-type: none"> ■ Elderly people are respected and thought to become wiser with age. ■ Family is expected to take care of elders. ■ Women gain status with age and childbearing; however, elderly women are expected to care for elderly men.
European American	<ul style="list-style-type: none"> ■ Value is on youth and beauty; elderly people are seen as less important than young people, and little attention is paid to their problems.
Filipino American	<ul style="list-style-type: none"> ■ Multigenerational households are common. ■ Grandparents act as surrogate parents while parents work.
Greek American	<ul style="list-style-type: none"> ■ Elders are well respected. ■ Children are expected to care for parents, who actively participate in family activities.
Iranian American	<ul style="list-style-type: none"> ■ Believe that with age comes experience, worldliness, and knowledge; therefore, elderly persons are seen with respect. ■ Caring for elderly parents is children's obligation.
Irish American	<ul style="list-style-type: none"> ■ Elderly people are well respected and opinions are valued. ■ Elderly parents are cared for in home.
Jewish American	<ul style="list-style-type: none"> ■ Elderly people are respected and seen as having wisdom. ■ Honoring and caring for parents is important. ■ Old age is a state of mind.
Korean American	<ul style="list-style-type: none"> ■ 60 is considered old and persons over that age are expected to retire.
Mexican American	<ul style="list-style-type: none"> ■ Elderly parents live with children. Large extended family.
Navajo Native American	<ul style="list-style-type: none"> ■ Elderly people with many children are seen with respect. ■ Elderly people have important role in teaching and keeping rituals to children and grandchildren.
Vietnamese American	<ul style="list-style-type: none"> ■ Elders are honored, have key role in family activities. ■ Elders are consulted on major family decisions.

Introducing the Case Study



Mrs. Nurick, an 80-year-old widow, comes into the health office of the continuing care retirement community where she has lived for the past 8 years. She complains of vague fatigue, not being able to participate in her walking group over the past few weeks, decreased appetite, and general malaise. She says that she just doesn't feel like herself. She has been forcing herself to eat small amounts, take care of her cat, and do other necessary things. She still goes down to the dining room for dinner, and most weekends one of her four daughters takes her to mass and to lunch at a restaurant or dinner at her home. She has no specific complaints of bowel or bladder changes or problems, cough or shortness of breath, indigestion, or pain.

and then simply wait and watch what happens, accept the changes they are experiencing, deny any danger associated with the symptoms, and delay seeking help. Many older adults worry that if they report symptoms to their healthcare provider, they may end up in a nursing home or in a more restrictive level of care. The cost of healthcare is also a major concern for many older adults. Also, when older adults do complain of a particular problem, they are more likely to identify a specific disease than are younger patients.

Multiple Pathologies

Another major challenge in doing a history and physical exam in an older adult is the likelihood of multiple pathologies. Active medical problems frequently interact with one another and can confuse the clinical picture. For example, uncontrolled diabetes in an older woman can result in a urinary tract infection, which causes subsequent urinary incontinence. The urinary incontinence may then result in the patient slipping on the wet bathroom floor and fracturing her hip. When moved to an acute-care setting, this same patient may develop atrial fibrillation.

Alternately, older adults may have unidentified multiple pathologies that interact with the current treatment of a newly diagnosed problem. These are referred to as disease-treatment interactions. An older man who presents with insomnia and is given an antihistamine to help him sleep at night may return to the office with acute urinary retention because of an underlying enlargement of his prostate that was previously undetected.

Performing the Assessment

A complete history and physical exam are essential to providing comprehensive, holistic care for the older adult. If the patient has a long and complicated medical history, you may need to do the history and physical exam on separate visits and schedule an hour for each visit. You may also need to allow more time to help your patient into the exam room and with dressing and undressing. When obtaining a history, ask one question at a time and allow enough time for your patient to respond. You may also need to repeat questions and confirm answers.

Remember that older adults may not present in the same way as younger people when they are ill. When performing your assessment, keep these three factors in mind:

- Older adults may minimize or ignore symptoms.
- They often have several concurrent medical problems.
- They often present with atypical signs and symptoms of disease.

Symptom Reporting

Although some older adults are “health pessimists” who exaggerate their symptoms, the opposite is more often true. Many older adults are “health optimists” who downplay symptoms and give a more positive evaluation of their overall health status in the face of disease and disability. Unfortunately, failing to report symptoms or explaining them away can lead to late recognition of serious medical problems.

Older adults are also the least likely of all patients to take action in response to symptoms of serious illness. Instead, they may attribute symptoms to normal aging

Disease Presentation

You should also recognize when doing a history and physical on an older adult that these patients do not always present with the typical signs and symptoms of disease found in younger patients. Illness in older persons is not likely to be signaled by a single specific symptom or sign that points to the organ with the pathology. Older persons are much more likely to present with nonspecific problems such as a change in cognition or functional performance, a fall, decreased appetite and intake, new-onset incontinence, dizziness, or vague weakness and lethargy.

Vulnerable systems, such as the brain, are likely to decompensate as a result of systemic impact of disease anywhere in the body. Older patients often present with blunted or atypical signs and symptoms of specific disease states. For example, an older patient may have a myocardial infarction without pain or pneumonia without cough, shortness of breath, or fever. She or he may complain only of “not feeling well.”

Health History

The format for history taking with older patients is similar to that used with younger patients. However, the

history should also include a functional assessment and investigation into advanced directives. Even though the formats are similar, some liberalization of the traditional interpretation of these components of the history is necessary to be relevant to the needs of the older adult.

Traditionally, the patient is the primary source of information for the history. However, many older adults are unable to provide comprehensive medical information because of cognitive changes or a lack of interest or knowledge about their medical problems. When working with these patients, it may be necessary to get information from medical records (from an acute-care hospital stay, an outpatient or a nursing home chart) or from family, friends, and hospital or nursing home staff. The choice to have a family member present during the history and exam should be based on the patient's wishes. If a family member is present, it should be made clear that the patient, not the relative, is to answer the questions when possible. The relative can clarify points and augment findings at the end of the history.

Even patients with known cognitive impairment or known to be unreliable in terms of their ability to provide information should be given an opportunity to answer questions and describe their symptoms. It is, however, essential to evaluate the patient's cognitive status to determine his or her ability to provide reliable health information. Even in severe dementia, the patient's response to questions regarding current symptoms may yield useful information, and it is important to establish a caring relationship by listening to what the patient has to say.

To quickly establish an overall sense of the patient's cognitive status, begin the history taking with questions that focus on orientation and past information. For example, ask questions that test time and place orientation and ask about the reasons for the visit, previous healthcare contacts, biographical data, and medication use.

Key points to remember when obtaining a history:

- Realize that age differences between you and your patient may influence her or his response.
- Be aware of your own views and values associated with aging.
- Explain what you are doing and why.
- Allow more time than you do with younger patients.
- Realize that you may need to obtain the history over several visits.
- Ask the patient if he or she can hear you clearly.
- Sit in front of and at the same level as the patient without invading her or his personal space.
- Maintain eye contact.
- Speak slowly and clearly; the lower the pitch the better.
- Set time limits for your interview.
- Redirect the interview as needed, but respect the patient's need to reminisce.
- Allow the patient to respond to each question before asking another.

- Listen to what your patient is saying and be alert for any signs of fatigue or discomfort.
- Use lay terminology that is culturally relevant to elicit more comprehensive information (e.g., "sugar" rather than "glucose" in diabetes).

Biographical Data

Begin by scanning the biographical data. The patient's age and gender will identify any risk factors. Her or his address, contacts, cultural or religious factors, and financial status may signal an underlying problem, explanation of a problem, or need for referral.

Current Health Status

Begin by determining the reason for seeking treatment. Remember that the older patient often has more than one presenting symptom and is more likely to have multiple medical problems with associated multiple symptoms and complaints. The complaints may also be overlapping and nonspecific. It is useful to develop a list of chief complaints rather than focusing on one complaint. Help your patient prioritize the problems according to which ones are the most distressing.

Symptom Assessment

As with chief complaints, there may be a history relevant to numerous lesser complaints, and each of these needs to be explored. In exploring them, it may be possible to find some connections among complaints and help identify the problem and relieve symptoms. For example, the patient may describe a sudden onset of right knee pain and also report increased urinary urgency. On further questioning, you may learn that the knee pain resulted in decreased functional ability, causing the patient to limit fluid intake so that he or she would not have to go to the bathroom very often. Limiting fluids resulted in strong, concentrated urine that caused the sensation of urgency in this patient.

The chronology of events is particularly challenging when assessing older adults who have some cognitive impairment or are retired and less focused on the days, dates, and timing of events. It is sometimes necessary to use a calendar to review the sequence of events or to relate the onset of symptoms to timing of the patient's activities. For example, you might ask if the pain started before the patient went to church this week or after her friend's birthday party.

When expanding on the present illnesses of older adults, it is important to use open-ended questions. Older patients look to the healthcare provider to have the answers and will probably follow your lead. Rather than asking if the pain was throbbing or stabbing, ask the patient to describe the pain. If the patient is unable to describe the pain in her or his own words, provide a choice of adjectives to describe it rather than asking the patient to respond to yes or no questions. It is also

important to repeat questions and confirm findings frequently.

Medications

Older adults often take many prescribed and over-the-counter (OTC) medications, so polypharmacy may account for many of your patient's symptoms. Determine your patient's understanding of the medications and whether additional teaching may be needed. Also determine if the medications are being taken as prescribed, or if use of (OTC) medications is affecting the desired action of prescription medications. Visual impairment may also cause mistakes in medication use. Ensuring safe administration of medication is a primary concern.

Past Health History

It is not uncommon for older adults to have an extensive medical history. Taking the time to obtain a comprehensive medical history is essential to prevent recurrence of past problems. Begin by asking your patient about known medical problems, providing prompts such as, "Do you remember being told you had a problem with your heart?" Next, ask about specific diseases, asking questions such as, "Have you ever been told you have diabetes?" Another way to prompt your patient's memory is to ask about past hospitalizations and the reasons for them. Do not focus on childhood illnesses, with the exception of asking about any childhood illness that kept the patient in bed or out of school for an extended period of time. This question will help identify, for example, a history of rheumatic fever, which may cause rheumatic heart disease later in life.

Surgery

With regard to surgical history, it may also be necessary to prompt older patients' memory with questions about common surgeries that occur with age, such as cataract

surgery, joint replacements, or removal of skin lesions. It is also necessary to ask specifically about the removal of organs such as the gallbladder, appendix, uterus, or prostate. During the physical examination, confirm any obvious scars with the surgical history given and question the patient about any unexplained scars you identify.

Medications

Reviewing medications is a good way to stimulate the older adult to recall previous diagnoses and medical problems. For example, if a patient is on amiodarone (Cordarone) but does not report any cardiac history, it is helpful to ask specifically if the patient knows why he or she is taking that medication and if he or she ever had an irregular heart rate.

Family History

Asking older adults about causes of mortality among parents and other relatives is usually irrelevant. Aside from a known history of Alzheimer's disease or nonspecified dementia, most other familial diseases are present before the person reaches old age. Therefore, don't spend a long time reviewing family health histories.

Review of Systems

In addition to the usual review done with all adult patients, the review of systems (ROS) with older adults should focus on problems that are particularly prevalent in this population. Specifically, these problems include a change in cognition, urinary or bowel incontinence (see *Types of Incontinence* on page 941), falls (see *Evaluating the Risk Factors for Falls* on page 941), immobility, insomnia, dysphagia, and sensory changes. Positive responses to questions about any of these problems can be a useful starting point for a more comprehensive evaluation of the problem and the identification of more serious underlying problems.

(Text continued on page 942)

Review of Systems

SYSTEM/QUESTIONS TO ASK

General Health Survey

- How would you describe your usual state of health? Are you able to do what you usually do?
- Have you noticed any changes in your height or weight?
- Do you notice any difference in how your clothes fit?

RATIONALE/SIGNIFICANCE

- Helps to identify current health status and activity tolerance.
- Loss of height (normally 6–10 cm) occurs with aging.
- Changes in weight distribution occur with age; a decrease of subcutaneous fat is seen on the face and extremities, with an increase on the abdomen and hips.
- Differences in the way clothes fit may signal subtle changes in weight that warrant further investigation.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Integumentary</p> <p><i>Skin</i></p> <ul style="list-style-type: none"> ■ Have you noticed any changes in your skin (e.g., dryness, itching, rashes, blisters)? ■ How often do you bathe? What type of soap do you use? Do you use lotions, sunscreens? ■ Have you been in the sun a great deal? ■ What type of work did you do? ■ Have you noticed any skin changes such as changes in size of growths, or open, sore, cracked, itchy, bleeding areas that won't heal? ■ Have you noticed any new rough areas of your skin that do not seem to go away? <p><i>Hair</i></p> <ul style="list-style-type: none"> ■ Any hair loss, increased hair growth, graying, dry scalp, or other hair changes? ■ What are your usual patterns of hair care? ■ Have you undergone hair replacement treatments? <p><i>Nails</i></p> <ul style="list-style-type: none"> ■ How do you care for your nails? Do you cut them yourself? Do you see a podiatrist? 	<ul style="list-style-type: none"> ■ There are many age-related changes in the skin, some of which are also caused by the environment. They result in uncomfortable symptoms such as pruritus and xerosis (dry skin), which occurs frequently in elderly people during the winter. ■ Skin care practices, sun exposure, and occupational history may influence current skin condition. ■ The incidence of skin cancer increases with age. ■ Remember that visual changes that occur with aging may make early recognition of skin changes more difficult. ■ Changes in hair growth and distribution are commonly associated with aging. ■ Dry scalp is a normal and common complaint. Hair loss may be distressing for both men and women. ■ Women may also be concerned about increased facial hair. ■ You need to differentiate normal changes from possible disease. ■ Nails thicken with age and may be more difficult to trim. Fungal nail infections are common. A podiatry consult may be indicated, especially if the patient has a history of diabetes or vascular disease.
<p>HEENT</p> <ul style="list-style-type: none"> ■ Do you have facial pain? 	<ul style="list-style-type: none"> ■ Increased incidence of temporal arteritis in elderly people may exhibit pain over temporal artery.
<p><i>Head and Neck</i></p> <ul style="list-style-type: none"> ■ Can you move your head easily? 	<ul style="list-style-type: none"> ■ ROM in the head and neck may be limited by musculoskeletal changes or osteoarthritis.
<p><i>Eyes</i></p> <ul style="list-style-type: none"> ■ Have you noticed any changes in your vision? ■ Can you read normal-sized print or large-print materials? ■ Do you have problems going from light to dark areas? ■ Do you drive at night? <ul style="list-style-type: none"> ■ Do your eyes feel dry? 	<ul style="list-style-type: none"> ■ Common problems (e.g., cataracts, macular degeneration, and glaucoma) can affect vision. Normal age changes include increased sensitivity to glare and decreases in visual acuity, lens elasticity, peripheral vision, color intensity (specifically blue, green, and purple), night vision, accommodation to changes in lighting, tear production and viscosity, depth perception, and near vision (presbyopia). ■ Dry eyes are very common. They may be caused by decreased tear production or blocked tear ducts, so further investigation is warranted.

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Do you have floaters? ■ Do you see flashes of light? ■ Does it look like a shade is being pulled over your eye? 	<ul style="list-style-type: none"> ■ There are several eye complaints that require immediate attention, such as a sudden onset of floaters or flashes of light seen peripherally, with decreased visual acuity. The patient may describe this as being like a curtain coming down over the field of vision (amaurosis fugax). These symptoms indicate retinal detachment, transient ischemic attack (TIA), or stroke. A complaint of a sudden onset of painless unilateral loss of vision also requires immediate attention because this may be a result of retinal vein occlusion.
<p><i>Nose</i></p> <ul style="list-style-type: none"> ■ Have you noticed any changes in your sense of smell? ■ Do you have a runny nose (rhinorrhea)? ■ Do you sneeze frequently? ■ Do you have postnasal drip? 	<ul style="list-style-type: none"> ■ Anosomia—a decreased ability to identify and discriminate odors—occurs with aging. ■ Atrophic changes associated with aging may cause vasomotor rhinitis.
<p><i>Mouth and Throat</i></p> <ul style="list-style-type: none"> ■ Do you have difficulty chewing, swallowing, tasting, smelling, or enjoying food? 	<ul style="list-style-type: none"> ■ Age-related changes on the surface of the tongue and atrophy of taste buds affect the ability to eat and enjoy food. In particular, there is a decrease in the ability to taste sweets and salt. Gums become thinner and recede, resulting in loose teeth and exposure of the roots.
<ul style="list-style-type: none"> ■ Do you wear dentures? Do they fit properly? ■ Do you have any sores in your mouth? ■ When was the last time you saw your dentist? ■ When you are eating, do you ever cough or choke? Does this occur with liquids and/or solids? Do you have problems swallowing? ■ Does your mouth always feel dry with a bad taste? ■ Do you have difficulty swallowing dry food or difficulty speaking for long periods? 	<ul style="list-style-type: none"> ■ Poorly fitted dentures may lead to poor nutrition and mouth sores. May identify need for referral. ■ Dysphagia, defined as the mechanical interruption of eating, may be related to a variety of underlying problems and is not a normal part of aging. It warrants further investigation. ■ Dry mouth (xerostomia) is a very common problem in older adults and can be caused by medication, decreased production of saliva, inadequate fluid intake, atrophy of the oral mucosa, vitamin deficiencies, poor nutrition, and poor oral hygiene. ■ Burning mouth syndrome is also common in the older adult. It causes dry mouth, altered taste, thirst, difficulty swallowing, swelling in the face and cheeks, and altered sense of smell.
<p><i>Respiratory</i></p> <ul style="list-style-type: none"> ■ Do you have any difficulty breathing? If yes, when does it occur and with how much exertion? Has it affected your ability to perform usual activities? ■ Do you have a cough? 	<ul style="list-style-type: none"> ■ Respiratory disorders are common in older adults. Patients may describe themselves as having breathing difficulties, trouble getting a deep breath or sufficient air. With chronic obstructive pulmonary disease (COPD), dyspnea is usually insidious in onset and progressive. ■ Lung cancer most frequently occurs between ages 55 and 74. ■ There is a recent increase in tuberculosis among elderly people. ■ If there is a new-onset cough, determine if the patient has been exposed to a change in environment or has taken any new medications, such as angiotensin-converting enzyme (ACE) inhibitors.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<p>Cardiovascular</p> <ul style="list-style-type: none"> ■ Do you get short of breath when walking or making the bed? ■ Do you have swelling in the feet, hands, face, or abdomen; weight gain; or shoes or clothing that no longer fit? ■ Does the swelling get worse as the day goes on and disappear in the morning? ■ Do you get dizzy? When? Does the dizziness get worse when going from lying to standing, or is it worse with exertion? ■ Are there any changes in your energy level? ■ Do you have a cough? Is it worse at night? ■ Do you get more short of breath when lying flat? ■ Do you need to sleep with several pillows? ■ Do you have headaches? ■ Do you have leg pain when walking? ■ Skin changes, swelling, ulcers, or varicose veins in your legs? <p>Gastrointestinal</p> <ul style="list-style-type: none"> ■ What is your typical diet for a day? Any change? Any food intolerance? ■ What are your usual bowel patterns? ■ What is your diet and fluid intake? ■ What medications, prescribed and OTC, do you take? 	<ul style="list-style-type: none"> ■ Cardiovascular disease is not a normal change associated with aging, but it is the most common problem. Unfortunately, it often does not exhibit the typical signs and symptoms found in younger people. For example, an older adult may present with atrial fibrillation but have no symptoms at all or complain vaguely of just not feeling right. If the patient reports swelling, explore if this is related to her or his activity level or a change in medication or diet. ■ Dizziness is a very common complaint and may be described as lightheadedness or a woozy feeling. ■ Vague symptoms of dizziness may indicate postural (orthostatic) hypotension or a more serious underlying medical problems such as atrial fibrillation or congestive heart failure. ■ Cough may be related to congestive heart failure or a cardiac medication. ■ Vascular disease increases with age. ■ Headaches may be associated with HTN, stroke, or temporal artery disease. ■ Intermittent claudication (pain in legs when walking) and skin changes (thin, shiny and hairless) suggest arterial vascular disease. There is a high incidence of venous disease in elderly people. ■ Food intolerance is a common complaint and may be associated with hiatal hernia and esophageal reflux. ■ Constipation is one of the most common digestive complaints in older adults, and accounts for 2.5 million physician visits annually. The prevalence of constipation increases with age, is more common in women than in men, in nonwhites than in whites, and in people with lower family income and less education. Constipation may have dangerous complications in older adults, including acute changes in cognition, urinary retention, urinary incontinence, and fecal impaction. Fecal impaction can result in intestinal obstruction, ulceration, and urinary problems. Chronic straining to defecate can have adverse effects on cerebral, coronary, and peripheral vascular circulation. ■ Constipation can be categorized as functional (slow transit of stool) or rectosigmoid outlet delay (anorectal dysfunction, 10 min or more needed to defecate). Bowel changes associated with bleeding and weight loss suggest a malignancy.
<p>Genitourinary</p> <p><i>Urinary</i></p> <ul style="list-style-type: none"> ■ Do you have trouble getting to the bathroom on time? ■ Do you need to wear a pad? If so, how many times a day do you have to change it? 	<ul style="list-style-type: none"> ■ Urinary incontinence is not a normal aspect of aging, but it is a common problem. Approximately 10 million Americans suffer from urinary incontinence. Report of

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Does the incontinence occur with coughing or sneezing, on the way to the bathroom, or at night? Does it interfere with your ability (or desire) to do daily activities or engage in social activities? ■ Ask men about frequent urination, hesitancy, weak or intermittent stream, a sensation of incomplete emptying of the bladder, dribbling after voiding, and nocturia. 	<p>new-onset incontinence, loss of appetite, vomiting, falls, nocturia, difficulty urinating, or behavioral and cognitive changes should alert you to a possible urinary tract infection.</p> <ul style="list-style-type: none"> ■ Prostate enlargement can cause incontinence. (See <i>Types of Incontinence</i>.)
<p>Female Reproductive</p> <ul style="list-style-type: none"> ■ Do you have any vaginal discharge? If so, what is the type, color, odor, and consistency of the discharge? ■ Do you have vaginal pressure, or an uncomfortable, bearing-down sensation, in addition to symptoms of urinary incontinence? ■ Do you have any vaginal bleeding? ■ Are you taking hormone replacement therapy? ■ When did you go through menopause? ■ When was your last Pap test? ■ Do you do breast self-examinations? ■ When was your last mammogram? Do you get them yearly? <ul style="list-style-type: none"> ■ Are you satisfied with your sexual activity? ■ Do you have pain during intercourse (dyspareunia) or vaginal dryness? <ul style="list-style-type: none"> ■ Have you ever had a sexually transmitted disease? ■ Do you practice safe sex? 	<ul style="list-style-type: none"> ■ Changes in vaginal secretions, amount, and pH increase the risk for vaginal infections. ■ Symptoms may be related to the presence of a uterine prolapse, cystocele, or rectocele. ■ Bleeding may be related to hormone replacement therapy. ■ If patient is not on hormones, bleeding that occurs after 1 yr postmenopause is abnormal and needs follow-up. ■ Incidence of breast cancer increases with age. Yearly mammograms are recommended for women over age 40. May identify teaching needs. ■ Older adults continue to be sexually active, unless they no longer have a sexual partner, have a disease, or are exposed to a treatment that decreases libido or makes intercourse uncomfortable. None of the age-related changes in either men or women preclude the continuation of a satisfying sex life. ■ Decreased vaginal secretions may result in dyspareunia. If this is present, an appropriate plan of care should be developed. ■ Because older women do not fear pregnancy, they are less likely to ask their partners to use condoms as a form of protection.
<p>Male Reproductive</p> <ul style="list-style-type: none"> ■ When was the last time you had your prostate checked? <ul style="list-style-type: none"> ■ Have you ever had a PSA test? ■ Do you have any urinary changes or problems? ■ Are you satisfied with your sexual activity? ■ Have you been feeling more tired than usual? 	<ul style="list-style-type: none"> ■ Increased incidence of prostate cancer in older men. Yearly prostate exams should begin at age 50. May identify need for health teaching. ■ Sexuality does not normally decrease with age. However, the physical act and response may require more time and be less intense. ■ Be alert for vague complaints. Give men the opportunity to talk about impotence and associated feelings so that individualized plans can be developed.
<p>Musculoskeletal</p> <ul style="list-style-type: none"> ■ Do you have pain, stiffness, joint enlargement, decreased ROM, and functional changes? 	<ul style="list-style-type: none"> ■ Osteoarthritis is the most common joint disease in the older adult and affects over 21 million adults. ■ Rheumatoid arthritis also increases with age. ■ Gout is also common, exacerbated by use of diuretics or alcohol.

(continued)

Review of Systems (continued)

SYSTEM/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Do you have pain, stiffness, and decreased ROM in the neck, shoulders, or hips persisting for at least 1 mo? ■ Do you have severe headache, visual loss, scalp tenderness, and mouth pain? ■ Have you had any fractures, bone pain, and loss of height? ■ Do you take calcium and vitamin D? ■ Do you exercise? If so, what type of activity do you do? ■ Are you taking any additional bone building medication? ■ Do you have balance problems or a history of falls? ■ Do you have any foot problems? ■ How do your shoes fit? ■ Do you see a podiatrist? ■ How do you care for your feet? 	<ul style="list-style-type: none"> ■ Common problems in older adults include polymyalgia rheumatica, a syndrome that involves the musculoskeletal system; giant cell arteritis, a vasculitic disorder of the cranial arteries associated with polymyalgia rheumatica; and osteoporosis. ■ Identifies participation in measures to prevent further bone loss and possible teaching needs. ■ Refer to the checklist <i>Evaluating the Risk Factors for Falls</i>. This can help determine whether the fall is caused by gait or balance disorder or another underlying problem for which the patient needs to be evaluated and treated. ■ Foot problems are common in elderly people and may result from poorly fitted shoes or poor foot care. Referrals may be warranted.
<p>Neurologic</p> <ul style="list-style-type: none"> ■ Do you have problems with balance, mobility, coordination, sensory interpretations, level of consciousness, intellectual performance, personality, communication, comprehension, emotional responses, and thoughts? ■ Do you have dizziness? Do you feel that the room is spinning or you're spinning? ■ Do you have a known history of seizures? ■ Do you have repetitive shaking or muscle contractions, brief lapses of consciousness, or any abnormal sensations? 	<ul style="list-style-type: none"> ■ Changes in the neurological system may be normal or caused by disease. ■ Stroke is the most common neurological problem in elderly people. ■ Parkinson's disease is the most common extrapyramidal problem. ■ Vertigo is a common problem, and diagnosis is based mainly on clinical symptoms. ■ Incidence of seizures significantly increases in people over age 65 because of an increase in strokes, tumors, subdural hematomas, metabolic disorders, dementia, and medications.
<p>Endocrine</p> <ul style="list-style-type: none"> ■ Have you been feeling depressed? ■ Experiencing weight loss or gain? ■ Does your heart ever flutter, race or skip beats? ■ Do you have increased thirst, urination, and appetite? 	<ul style="list-style-type: none"> ■ Thyroid disease presents differently in older adults than in younger adults. Signs of hyperthyroidism include apathy, depression, and emaciation rather than hyperactivity. Atrial fibrillation also often occurs. ■ Hypothyroidism is the most frequent thyroid disorder in elderly persons, but it is easy to miss because signs (dry skin and hair, hypotension, slow pulse, sluggishness, depressed muscular activity, goiter, weight gain) are often attributed to aging.
<p>Immunologic/Hematologic</p> <ul style="list-style-type: none"> ■ Have you been feeling more tired than usual? 	<ul style="list-style-type: none"> ■ The incidence of diabetes mellitus increases with age. ■ When an elderly person becomes sick, symptoms may be vague because of changes in the immune system. ■ Fatigue may be associated with anemia, a common problem in elderly people, usually caused by iron deficiency.

Types of Incontinence

TYPE OF INCONTINENCE	DEFINITION	PATHOPHYSIOLOGY	SIGNS AND SYMPTOMS
Stress	<ul style="list-style-type: none"> ■ Involuntary loss of urine caused by urethral sphincter failure with increases in intra-abdominal pressure. 	<ul style="list-style-type: none"> ■ Usually caused by weakness and laxity of pelvic floor musculature or bladder outlet weakness. Also may be caused by urethral hypermobility. 	<ul style="list-style-type: none"> ■ Urine is lost during coughing, sneezing, laughing.
Urge	<ul style="list-style-type: none"> ■ Leakage of urine because of inability to delay voiding after sensation of bladder fullness is perceived. 	<ul style="list-style-type: none"> ■ Associated with detrusor hyperactivity, central nervous system disorders, or local genitourinary conditions. 	<ul style="list-style-type: none"> ■ Urine is lost on the way to the bathroom or as soon as the urge to void is felt.
Overflow	<ul style="list-style-type: none"> ■ Leakage of urine resulting from mechanical forces on an overdistended bladder. 	<ul style="list-style-type: none"> ■ Results from mechanical obstruction or an acontractile bladder. 	<ul style="list-style-type: none"> ■ Variety of symptoms, including frequent or constant dribbling, increased incontinence at night, frequency, and urgency.
Functional	<ul style="list-style-type: none"> ■ Urine leakage caused by inability to get to toilet because of cognitive or physical impairment. 	<ul style="list-style-type: none"> ■ Cognitive and physical functional impairment. 	<ul style="list-style-type: none"> ■ Patient is aware of the need to void, but urine is lost on the way to the bathroom.

Evaluating the Risk Factors for Falls

Patient's name _____

Age _____ Gender _____ Date _____

- History of previous falls:
 - Yes
 - No
- Medications:
 - Four or more prescriptions
 - New prescription in the last 2 weeks
 - Use of any of the following medications: Tranquilizers, sleeping pills, antidepressants, cardiac medications, antidiabetic agents
- Known gait problem or muscular weakness:
 - Yes
 - No
- Dizziness, vertigo, or loss of consciousness at time of fall:
 - Yes
 - No
- Visual changes:
 - Yes
 - No
- Environmental problems:
 - Clutter
 - Dim lighting
 - Uneven flooring
 - Inappropriate footwear or lack of footwear
 - Inappropriate assistive device
- Major illnesses:
 - Neurological: Parkinson's disease, stroke, dementia
 - Musculoskeletal: Arthritis, contracture, fracture
 - Cardiac: Hypotension, arrhythmia, acute infarct
 - New acute illness: Infection
 - Other

Psychosocial Profile

A detailed psychosocial profile is critical when assessing the older adult because it can yield important data about the patient's health habits and health status. Your history should explore several areas, which are described as follows.

Typical Day

For most patients, a description of a typical day is helpful in assessing their ability to perform activities of daily living (ADLs). For elderly patients, this assessment is critical. A thorough functional assessment is indicated so that you and the patient can develop a plan together to meet any deficits that may be identified. (For more information, see the Functional Assessment section.)

Nutritional History

A number of variables can compromise nutritional status in the older adult: oral or gastrointestinal disorders, drug effects, systemic illness, financial problems, and psychiatric disease. A nutritional history is best obtained by asking the patient about her or his typical 24-hour diet, pattern of weight during recent years, and food shopping and preparation habits and concerns. Identify the social context of food and eating for your patient. Does she or he eat alone? Also identify changes in eating patterns. Have they resulted from loss of spouse or financial constraints? Are there physical limitations that affect the person's ability to obtain or eat nutritious foods?

Certain types of foods may affect existing health problems. Ask if the patient eats canned and processed foods more often than fresh foods. These foods tend to have high sodium concentrations and do not provide the same fiber content as fresh fruits and vegetables. The high sodium content may be contraindicated for patients with cardiovascular disease. Because decreases in sense of taste and smell occur with aging, older adults often overuse spices, salt, and sugar, which may be contraindicated for patients with cardiovascular disease or diabetes. In particular, older adults should be asked about involuntary weight gain or loss. Obesity is a common problem in this age group because the basal metabolic rate (BMR) lowers with age, caloric requirements decrease, and physical activity may decrease. The daily caloric requirements are 1280 to 1900 calories for older women and 1530 to 2300 calories for older men. Weight loss can also be a problem. Loss of 5 lb in 6 months or less is indicative of protein-energy malnutrition. Unexplained weight loss in an older person warrants further investigation because it may be a sign of a serious underlying disease, such as a malignancy.

You also need to assess your patient's fluid intake. Older adults often limit fluid intake as a result of decreased thirst and incontinence problems. The daily fluid intake should be at least 1300 mL unless contraindicated for existing cardiovascular disease (congestive heart failure). It is also important to determine the patient's calcium intake because the risk of osteoporosis increases with age, especially in women.

A thorough drug history is essential when evaluating your patient's nutritional status. There are many prescription and OTC medications that can affect your patient's fluid and electrolyte and nutritional status. Ask the patient about *every* medication he or she is currently taking, both prescribed and OTC, as well as those in his or her possession. Determine the dose of the drug if possible, or at least the scheduling (how many times a day). Ask specifically about vitamins, minerals, herbs, laxatives, sleeping pills, and cold preparations because older adults may not consider these medications. See Drug-Nutrient Interactions on page 951 for more information.

Physical Activity and Exercise

Exercise and other forms of physical activity have been associated with both physical and psychosocial benefits to older adults. Physical activity is body movement created by skeletal muscles that expends energy and produces health benefits. Exercise is physical activity that is planned, structured, and repetitive and that enhances or preserves physical fitness. Both physical activity and exercise have been associated with improved functional performance in older adults. Studies show that physical activity and exercise can prevent osteoporosis, coronary artery disease, and non-insulin-dependent diabetes mellitus; decrease the risk of falling; improve sleep; enhance mood and general well-being; improve blood pressure (BP); and decrease relative abdominal fat.

Despite the known benefits of exercise and physical activity, only 30 percent of older adults report regular exercise. Moreover, 50 percent of older adults who are not physically active have no intention of starting a regular exercise program to increase their activity. It is therefore essential to ask about exercise behavior in older patients and to encourage them to exercise regularly. The American Heart Association (2005) recommends that all adults engage in at least 30 minutes of moderate-intensity physical activity on most, or preferably all, days of the week.

Sleep

Sleep changes and disorders are a common complaint of older adults and should be included in the history. Some of these changes may be a result of the normal process of aging or psychosocial issues that affect sleep. Explore

Drug-Nutrient Interactions

MEDICATION	POTENTIAL EFFECT ON NUTRIENTS
Mineral oil	Potential Effect on Nutrients
Anticonvulsants	Decreased absorption of fat-soluble vitamins A, D, E, and K
Antacids with aluminum	Reduced storage of vitamin K and absorption of calcium and vitamin D
Antacids with aluminum or magnesium	Reduced absorption of phosphorous and fluoride and increased calcium excretion
Antacids with sodium bicarbonate	Reduces absorption and increases intestinal elimination of phosphate
Gentamicin	Sodium overload and water retention
Penicillin	Increased potassium and magnesium excretion
Tetracycline	Increased potassium excretion
Aspirin	Decreased absorption of iron, calcium, zinc, and magnesium
Corticosteroids	Decreased folate and ascorbic acid, causing iron deficiency as a result of blood loss
Potassium supplements	Decreased calcium and phosphorous absorption
Laxatives	Increased need for pyridoxine, folate, ascorbic acid, and vitamin D
Cholestyramine	Decreased vitamin B ₁₂ absorption as a result of decreased acidity in the ileum
Cimetidine	Hypokalemia, hypoalbuminemia, decreased calcium absorption, malabsorption with steatorrhea
Neomycin	Decreased absorption of fat-soluble vitamins and calcium
Isoniazid	Decreased absorption of vitamin B ₁₂ as a result of hypochlorhydria
Hydralazine	Decreased absorption of fat, lactose, nitrogen, calcium, iron, potassium, and vitamin B ₁₂
	B ₆ and niacin deficiency
	B ₆ deficiency

Source: Miller, C. (1990). *Geriatric Nursing* 11(6):301–302 with permission.

with your patient when the sleep disorder began, its duration and severity, and how the change in sleep impacts her or his life. Ask about difficulty falling and staying asleep, feeling rested in the morning, snoring, and movement, pain or jerking in the leg that either prevents the patient from falling asleep or wakes her or him during the night.

Also ask if the patient is having any associated symptoms, such as chest pain, indigestion, back pain, or urinary frequency, that are influencing sleep. Review bedtime routines, caffeine use, and exercise/activity as they relate to sleep behavior. Be sure to ask the patient if he or she naps during the day, the approximate length of the

naps, and the amount of time he or she spends in bed not sleeping. Roommates, caregivers, and spouses or significant others should also be asked about the patient's snoring or changes in breathing patterns, which might provide a clue to an underlying problem such as sleep apnea, and excessive leg movement or jerking during the night, which might interrupt normal restful sleep patterns.

Self-Care Responsibilities

The health history should also include questions about health promotion and prevention practices. There is evidence to suggest that older people benefit just as much from primary and secondary health promotion activities as middle-aged people do. Exercise and reducing cholesterol levels improve overall health status and physical fitness, including aerobic power, strength, balance, and flexibility and help prevent acute medical problems such as fractures, myocardial infarctions, and strokes. Mammograms, Pap tests, digital examinations for monitoring prostate size, and yearly evaluations of stool specimens for occult blood can help reduce mortality and morbidity among older adults.

Unfortunately, with age, there is a general decline in primary and secondary preventive behaviors, particularly in regard to mammography, Pap tests, having stools checked for occult blood, and monitoring cholesterol.

How Age Affects Sleep

- Longer time to fall asleep
- Increased time in stages 1 and 2 sleep
- Decreased time in deeper stages of sleep (stages 3 and 4)
- Decreased REM sleep
- Increased and shorter repetition of sleep cycle
- Increased nighttime awakenings
- Altered circadian rhythm with a need to fall asleep earlier and awaken earlier

Reasons include advanced age, unwillingness to undergo treatment if given a positive finding, and not being referred for testing by a primary healthcare provider. During history taking, healthcare providers should take an individualized approach, discussing the pros and cons of primary and secondary prevention activities and helping older adults make decisions about participation in these activities. Once patients have been educated, it is important that healthcare providers listen to the patients' wishes and respect their decisions. For example, if an older woman says that she will not undergo surgery for a breast mass, it may not be appropriate to send her for mammography.

Personal Habits

Ask about the use of nicotine, alcohol, and street drugs. Rather than asking the older adult if she or he drinks alcohol, ask how much alcohol she or he drinks. Assuming that an older adult drinks alcohol regularly reduces the feeling that judgment is being passed. It is also helpful to use screening instruments to explore alcohol use in the older patient. The CAGE is a short four-item questionnaire that helps identify people at risk for alcohol abuse. (See Chapter 4, Approach to the Mental Health Assessment.)

Support Systems

Information on support systems is a relevant part of the patient history because it is necessary to understand what personal and community support the older patient has to help with ADLs. Determining if the patient lives alone or with others and what type of setting he or she lives in is important, as is information about the patient's community at large. Explore all potential social supports with the patient and determine what type of interaction is available (phone calls or visits).

Environmental Structure and Hazards

Ask the patient to describe her or his home, the presence and condition of any stairs, the accessibility of the bathroom and kitchen, and household activities with which she or he currently needs assistance (e.g., grocery shopping, cooking, cleaning, laundry). Assessment of your patient's home environment may identify needs for assistive devices or referrals.

Functional Assessment

A very important component of the history in older adults is exploring functional performance. Functional evaluations should include ADLs such as eating, bathing, dressing, transferring, toileting, and ambulation. Instrumental activities of daily living (IADLs) include activities such as taking medications, paying bills, using the telephone, and using public transportation. Use of a functional status instrument will help you obtain baseline information that you can follow from year to year on your patient. Many such instruments are available, and the decision to use one over another may depend on the patient. For example, the Katz Index is a basic measure that describes the patient as dependent, semidependent, or independent in ADLs but does not identify small changes over time. Conversely, the Barthel Index gives weighted scores for each ADL and allows the nurse to recognize small changes over time. A score of less than 60 on this index indicates that the patient needs assistance with ADLs. Duke University provides an example of a common measure used to review IADLs. Reviewing these items with the patient allows you to determine his or her level of functioning and make appropriate referrals for care.

The Katz Index of ADL

Abbreviations: I = independent; A = assistance; D = dependent

1. Bathing (sponge, shower, or tub)
 - I: Receives no assistance (gets in and out of the tub)
 - A: Receives assistance in bathing only one part of the body
 - D: Receives assistance in bathing more than one part of the body
2. Dressing
 - I: Gets clothes and gets completely dressed without assistance
 - A: Gets clothes and gets dressed without assistance except in tying shoes
 - D: Receives assistance in getting clothes or in getting dressed or stays partly or completely undressed
3. Toileting
 - I: Goes to bathroom, cleans self, and manages clothes without assistance (may use an assistive device)
 - A: Receives assistance in going to bathroom or in cleaning self, managing clothes, or emptying a bedpan
 - D: Doesn't go to bathroom for elimination
4. Transfer
 - I: Moves in and out of bed or chair without assistance (may use assistive device)
 - A: Moves in and out of bed or chair with assistance
 - D: Doesn't get out of bed

5. Continence
 I: Controls urination and bowel movements independently
 A: Has occasional accidents
 D: Urine or bowel control maintained with supervision; patient is incontinent or has catheter
6. Feeding
 I: Feeds self without assistance
 A: Feeds self except for cutting meat or buttering bread
 D: Receives assistance in feeding or is fed partly or completely by tubes or intravenous fluids

Source: Adapted from Katz, S., Ford, A., and Moskowitz, R. (1963). Studies of illness in the aged: The index of ADL. *Journal of the American Medical Association* 185:914–919.

The Barthel Index

LEVEL OF CARE	INTACT	LIMITED	HELPER	NULL
Self-Care				
1. Feed	10	5	3	3
2. Dress (upper extremities)	5	5	3	0
3. Dress (lower extremities)	5	5	2	0
4. Don brace	0	0	−2	0
5. Grooming	4	4	0	0
Cleanse Perineum	4	4	2	0
Sphincters	Completely voluntary	Urgency/appliance	Some help needed	Frequent accidents
6. Bladder	10	10	5	0
7. Bowel	10	10	5	0
Mobility/Transfer	Easy/no device	With difficulty or uses device	Some help needed	Dependent
1. Chair	15	15	7	0
2. Toilet	6	5	3	0
3. Tub	1	1	0	0
4. Walk 50 yd	15	15	10	0
5. Stairs	10	10	5	0
6. Wheelchair 50 yd	15	5	0	0

Source: Adapted from Barthel DW et al., (1965) Functional Evaluation: The Barthel Index. *Maryland State Medical Journal* 14:61–65.

Instrumental Activities of Daily Living (IADL)

Abbreviations: I = independent; A = assistance; and D = dependent.

1. Telephone:
 I: Able to look up numbers, dial, receive, and make calls without help
 A: Able to answer phone or dial operator in an emergency, but needs special phone or help in getting number or dialing
 D: Unable to use the telephone
2. Traveling:
 I: Able to drive own car or travel alone on bus or taxi
 A: Able to travel but not alone
 D: Unable to travel

(continued)

Instrumental Activities of Daily Living (IADL) (continued)

3. Shopping:

I: Able to take care of all shopping with transportation provided

A: Able to shop but not alone

D: Unable to shop

4. Preparing meals:

I: Able to plan and cook full meals

A: Able to prepare light foods, but unable to cook full meals alone

D: Unable to prepare any meals

5. Housework:

I: Able to do heavy housework (e.g., scrubbing floors)

A: Able to do light housework, but needs help with heavy tasks

D: Unable to do any housework

6. Medication:

I: Able to take medications in the right dose at the right time

A: Able to take medications, but needs reminding or someone to prepare it

D: Unable to take medications

7. Money:

I: Able to manage buying needs, write checks, pay bills

A: Able to manage daily buying needs, but needs help managing checkbook and paying bills

D: Unable to manage money

Source: Duke University Center for the Study of Aging and Human Development, pp. 169–170, with permission.

Advance Directives and Preferences for Care

Directives for acute and end-of-life care are not traditionally part of the patient history. But, in the current health-care arena, it is essential to address these issues with older adults. The 1990 Patient Self-Determination Act (PSDA), which requires that patients be asked about the existence of advance directives at the time of enrollment or admission to a healthcare facility, increased awareness of the right to determine acute and end-of-life care options. Unfortunately, discussions and decisions about these care options with older adults have increased only moderately since the PSDA. A large number of older adults still do not have advance directives. Moreover, many written advance directives have care options that are poorly delineated, so the proxy is not aware of the patient's specific wishes.

With regard to directives for care, old-old people (90 years of age and older) usually express a decreased interest in receiving aggressive medical interventions for acute and end-of-life care. It is not uncommon when working with old-old patients to have them report a feeling of satisfaction with a long life lived, yet a desire for this life to end quietly. Certainly these wishes need to be acknowledged. Older adults are also more likely to want to be hospitalized for treatment of an acute illness, but are less likely to want treatment for a potentially life-threatening injury. They do not want life prolonged if their function and quality of life will be impaired.

Unfortunately, 6 to 20 percent of older adults defer decisions regarding acute and end-of-life care to their primary healthcare providers or the people who have power of attorney. It has been shown repeatedly that proxy decisions are different from the patient's wishes. Older adults should be encouraged to exercise their moral and ethical right to determine their care. Healthcare providers have a responsibility to educate them about these rights, to provide accurate, up-to-date information, and to correct misinformation about acute and end-of-life care options. Discussions about care decisions, executing a living will, and designating a proxy decision maker and durable power of attorney for healthcare will help older adults make sure that their wishes are carried out if they can no longer dictate their own care.



Case Study Findings

Before performing the physical examination, document what you've learned about Mrs. Nurick so far. Reviewing Mrs. Nurick's

Biographical data:

- 80-year-old widow with four daughters.
- Widowed since 1988.
- Resident of continuing care retirement community for 8 years.
- Catholic religion, Polish ancestry.

- Not referred by doctor—came to outpatient office of her own accord.
- Source: Self, seems reliable.

Current health status:

- Complains of vague fatigue all day, decreased appetite, and inability to engage in usual activities such as walking group. Fatigue has been present for several weeks and has grown worse in the past week even though the patient has been going to bed around 9 P.M. and getting up at 8 A.M. Is forcing herself to participate in social activities such as going to the dining room with friends for dinner. However, has no desire to eat.
- Is taking Lasix 20 mg PO qd for hypertension (HTN) and Tamoxifen 20 bid to prevent recurrence of breast cancer.

Past health history:

- **Medical History**
 - Breast cancer 20 years ago.
 - HTN controlled by diuretics.
 - Degenerative joint disease (DJD), mostly in hips.
- **Surgical History**
 - Cataract surgery, 1990 (right eye) and 1991 (left eye).
 - Lumpectomy, right breast, 1979.
 - Elective left total hip replacement, 1985.

Family history:

- No known history of cognitive problems. She believes that both of her parents died of heart disease in their 70s and 80s, and does not know of any family history of cancer.

Review of systems:

- **General Health Survey:** Complains of fatigue and of “not feeling like myself.” No changes in height or weight reported.
- **Integumentary:** Fair skin with no abnormal lesions. Complains of dry skin in winter.
- **Head, Eyes, Ears, Nose, and Throat (HEENT):** No vision changes, dizziness, change in smell or taste, or difficulty swallowing. Wears corrective lenses; has yearly eye examination. Reports being satisfied with quality of hearing.
- **Respiratory:** No cough or shortness of breath.
- **Cardiovascular:** No chest pain or shortness of breath during the day or at night. No shortness of breath while walking to the dining room. No leg cramping or pain either at rest or walking. Some swelling in ankles

on arising, but usually disappears in a few hours.

- **Gastrointestinal:** No indigestion or burning. Has a regular bowel movement daily and takes milk of magnesia rarely, only if she has not had a bowel movement in several days. Has not felt much like eating, but has not had stomach pain, nausea, or vomiting.
- **Genitourinary/Reproductive:** Loses urine when she coughs, sneezes, or laughs and therefore wears a sanitary pad, which she changes two or three times a day. This problem has not gotten any worse recently. No vaginal discharge or bleeding. Menopause at age 51; was never on hormone replacement therapy. Last Pap test around 1988; last mammogram in 1999.
- **Musculoskeletal:** Mild morning stiffness usually gone by the time she has breakfast. Takes Tylenol when pain is bothersome. Had two falls since moving into the facility but no falls over past 6 months. No increase in joint pain or warmth or swelling of joints.
- **Neurologic:** No dizziness, change in cognition, problems with balance, tremor, seizure, or difficulty with concentration. Is very happy at continuing care retirement community and feels strongly that this was the best move for her because she could no longer manage her home independently. Enjoys living near friends, participating in activities, and having visits with her daughters. Is glad they don't have to take care of her or worry about her being home alone. No previous history of depression.
- **Endocrine:** No weight changes, no increased thirst or urination. Heart does not flutter, race, or skip beats.
- **Immunologic/Hematologic:** Reports feeling more tired than usual.

Psychosocial history:

- Typical day starts around 8 A.M. She fixes herself breakfast in her efficiency apartment, feeds her cat, then has coffee and watches TV. Does laundry, light housecleaning, and “putters around” apartment. At midmorning usually goes to the activity room to socialize or play cards with other residents. Fixes own lunch. Until recently, walked 20 minutes three times a week with walking group in afternoon. Goes on planned outings with other residents. Sees daughters every weekend for lunch or dinner, shopping, and mass on Sunday. Previously

(continued on page 948)



Case Study Findings (continued)

went to bed at 11 P.M. but has been retiring at 9 P.M. for past few weeks because of fatigue.

- Doesn't smoke and rarely has a glass of wine socially.

Functional assessment:

- Performs all of her personal care independently. Does her own laundry, shopping, finances, and cleaning aside from the vacuuming, which is done by the facility every 2 weeks.
- Walks independently with a rolling walker. Has been participating in a walking group and until recently was walking for 20 minutes three times per week.



CRITICAL THINKING ACTIVITY #2

How might Mrs. Nurick's reactions to medications be affected by the changes associated with aging?



CRITICAL THINKING ACTIVITY #3

What normal changes associated with aging might account for Mrs. Nurick's loss of appetite?



CRITICAL THINKING ACTIVITY #4

What are Mrs. Nurick's strengths? Are there any areas of concern?

Physical Assessment

Once you have completed the health history, collect objective data through your physical exam.

Approach

When examining the older patient, make sure the environment is as safe and appropriate as possible. To ensure a senior-friendly environment, do the following:

- Keep examination rooms warm (between 70°F and 80°F).
- Use bright but nonglaring lights.
- Keep background noise to a minimum.
- Provide higher than standard seating (because patients might have trouble getting up, and if they have DJD or have had joint replacement, they shouldn't flex the joint more than 90 degrees) with arm rests on all chairs.
- Use exam tables that mechanically elevate the patient from lying to sitting and vice versa and a broad-based step stool to help patients get onto the table.

- Use a private exam room, if possible, or at least pull the privacy curtain if there is a roommate.
- Minimize position changes to keep the patient from getting tired.
- Uncover only the area being assessed, making sure that patient is warm and covered with blankets or drapes.
- Provide reading materials with large print.
- Allow more time than usual for the exam. The complete exam may need to be scheduled over several meetings.
- Make safety a priority. If your patient can't tolerate or perform what is expected for the exam, then adapt the exam to meet her or his needs.
- Take the time to explain everything you are doing to your patient.



TOOLBOX

To perform the physical exam, you will need all the tools of assessment.

Performing a General Survey

The physical exam of the older patient begins as soon as you meet. During the patient history, you observed the functional mobility of your patient, his or her overall appearance (particularly hygiene), emotional status, body habitus, and so on. As you begin the physical exam, note the patient's ability to transfer from a chair to the exam table and to dress and undress. Look at your patient's posture. A common problem is senile kyphosis (dowager's hump or widow's hump), an accentuated thoracic curve in which the head tilts back and the neck shortens, giving a rounded "E" profile. A wide base of support may signal balance problems. Note if there are any other obvious problems or signs of distress, such as suspicious skin lesions or shortness of breath. Also obtain vital signs and height and weight in a standardized fashion. If the scale requires a step up, be sure to help the patient. In light of the increased risk and incidence of hypothermia in older adults, it is important to have a reliable low-reading thermometer. To check for orthostatic hypotension, take BPs in the supine position after at least 10 minutes of rest, then immediately upon standing, and then 3 minutes after standing. Evaluating the heart rate response to postural changes at the same time provides important information about the cause of orthostatic hypotension. A rise of less than 10 beats per minute (BPM) with a drop in BP suggests baroreceptor reflex impairment.

Performing a Head-to-Toe Physical Assessment

You will use all four techniques of physical assessment and all of your senses when you perform the exam. As you examine your patient, keep in mind the changes that normally occur with aging so that you can differentiate them from abnormal changes.

(Text continued on page 957)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

1. Integumentary

SYSTEM/NORMAL VARIATIONS

Skin

- Inspect skin.



Inspecting the patient's skin



Seborrheic keratosis



Acrochordon



Senile purpura

**HELPFUL HINT**

An easy way to measure skin lesions is to apply a piece of Scotch tape over them and trace the outline of the lesion onto the tape. Place the tape in the patient's chart and reevaluate over time.

- Palpate temperature, turgor, texture, and hydration.
- *Skin color uneven in areas; increased creases, wrinkle lines, and skin lesions. Some common lesions include seborrheic keratosis, lentigines (liver spots), and acrochordons. Senile purpura, commonly found on hands and forearms, is caused by frail capillaries.*
- **Temperature:** Warm, but hands and feet may be cool.
- **Turgor:** Normally decreased. Not a marker of hydration status.
- **Hydration/texture:** Dry, flaky, and thin.

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

1. Integumentary (continued)

Hair and Scalp

- Inspect hair and scalp.
- Palpate hair and scalp.
- **Hair color/distribution:** *Graying in both sexes; thinning/balding especially in men.*
- *Increased facial hair (hirsutism) in women.*
- *Coarse, dry hair and dry flaky scalp (senile xerosis).*

Nails

- Inspect nails.
- Palpate texture.
- *Yellow, dry, brittle nails with longitudinal ridges.*

ABNORMAL FINDINGS/RATIONALE

- Areas of pressure or pressure sores from immobility or splints/appliances.
- Elderly patients are more apt to have body folds and develop intertrigo (inflammatory, moist erythema and scaling lesions) or fungal infections. See *Common Skin Lesions Associated with Aging*, pages 956 and 957 for more information.
- Cool extremities may signal vascular disease. Unilateral cool temperature may indicate an occlusion and warrants medical attention.
- Decreased turgor increases risk for injury and skin breakdown.
- Excessive dryness may indicate dehydration.
- Changes in hair may also relate to endocrine problems or may occur as a side effect from medications.
- Yellow, thick nails may also be caused by disease or a fungal infection.



ALERT

If you detect excessive bruising, consider the possibility of elder abuse.

2. HEENT

SYSTEM/NORMAL VARIATIONS

Head and Neck

- Inspect ROM of head and neck.
- Palpate neck.
- *May have decreased ROM caused by musculoskeletal changes.*

Lymph Nodes

- Inspect and palpate lymph nodes.
- *Lymph tissue decreases in size with aging and should be nonpalpable.*

Thyroid

- Inspect and palpate thyroid.
- *Nonpalpable.*

Temporal Artery

- Inspect and palpate temporal artery.
- *Palpable and nontender.*

Eyes

- Inspect external eye structures.
- Test visual acuity and extraocular membrane (EOM).
- Perform funduscopic exam.
- *Decreased near vision (presbyopia) may cause difficulty reading.*
- *Far vision may be intact.*
- *Decreased peripheral vision.*
- *Dry eyes.*

- *Enophthalmus (recession of eyeball into orbit).*
- **Arcus senilis:** *White to yellow deposit at outer edge of the cornea, along with xanthelasma (yellowish, raised plaque on upper or lower lids).*



Arcus senilis

- *Senile entropion (inversion of the lashes).*
- *Senile ectropion (eversion of the lashes).*
- *Xanthelasma (lipid deposits) on lids.*
- *Pale or yellow-tinted conjunctiva; pale iris; pingueculae (clear to yellow fleshy lesion on conjunctiva).*
- **Pterygium:** *Similar to pingueculae, but extends over cornea.*
- *Increased stimulation needed to elicit corneal reflex.*
- *Small pupils; reaction equal but may be less brisk.*
- **Fundoscopic—Retina and optic disc become paler with age.**

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

2. HEENT (continued)

Ears

- Inspect external and internal ear structures.
- Test hearing acuity.
- Perform otoscopic exam.

Internal

- *Gross hearing intact but diminished with decreased pitch discrimination.*
- *Potential conductive hearing loss (presbycusis).*
- *Increased difficulty hearing high-pitched sounds, especially s, t, f, and g.*
- *Difficulty understanding speech.*
- *Equilibrium-balance problems.*

External

- *Lobes elongate.*
- *Increased external ear canal hair in men.*

Otoscopic

- *Dry cerumen.*
- *Diminished cone of light.*

Nose

- Inspect the patient's nose and nares.
- *Elongated nose.*
- *Increased nasal hair.*
- *Decreased sense of smell (CN I).*

Mouth and Throat

- Inspect structures of mouth and throat.
- *“Purse-string” appearance of mouth.*
- *Teeth may show staining, chipping, or erosion. Some teeth may be loose or missing.*
- *Buccal mucosa and gums thin and pale.*
- *Oral mucosa dry, causing halitosis.*
- *Decreased sense of taste (CNs VII and IX).*
- *Decreased gag reflex (CNs IX and X).*
- *Decreased papillae on tongue.*
- *Varicose veins under tongue (caviar spots).*



HELPFUL HINT

When assessing the mouth, be sure to remove dentures and partial plates and evaluate for bacterial plaque, food debris, and pressure areas.

ABNORMAL FINDINGS/RATIONALE

- Decreased ROM of the neck may also be associated with DJD.
- Palpable nodes warrant referral.
- Palpable thyroid may signal thyroid disease (common in elderly people) and warrants referral.
- Pain, nodularity, and presence of a pulse may indicate temporal arteritis.
- Loss of peripheral vision, halos, and eye pain may indicate glaucoma and warrant referral. Visual changes may also occur with cerebrovascular disease.
- Basal cell carcinoma frequently found on inner third of lower lid.
- Yellow or opaque lens associated with increased incidence of cataracts.
- Entropion increases the risk of corneal abrasions.
- Ectropion increases risk of dryness and conjunctivitis.
- Macular degeneration.
- Check for diabetic and hypertensive retinal changes because the incidence of these diseases increases with age.
- Balance problems and tinnitus may indicate a neurological problem.
- Balance problems increase risk for falls and injury.
- Dry ears with scratch marks related to senile pruritus.
- Cerumen impaction can decrease hearing acuity by 40 to 45 dB; removal of cerumen corrects the impairment.
- Hearing aids can cause contact dermatitis.
- Vasomotor rhinitis causes pale nasal mucosa and boggy turbinates.
- Gum recession and bleeding (atrophic gingivitis).
- Chewing and swallowing difficulty may be caused by poor dentition or poorly fitted dentures, or more serious problem such as stroke.
- Leukoplakia (white precancerous lesion) in mouth, particularly under tongue.

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

3. Respiratory

SYSTEM/NORMAL VARIATIONS

- Inspect and palpate chest.
- Auscultate breath sounds.
- *Senile kyphosis (increased anterior to posterior diameter) caused by musculoskeletal changes, barrel chest appearance.*
- *Decreased respiratory excursion.*
- *Cheyne-Stokes breathing may occur during sleep.*
- *Breath sounds may be decreased with few crackles at bases.*

ABNORMAL FINDINGS/RATIONALE

- Respiratory changes increase risk for pulmonary problems, such as pneumonia.

4. Cardiovascular

SYSTEM/NORMAL VARIATIONS

- Inspect and palpate pulsations.
- Auscultate heart and arteries.
- *Increase in premature beats and irregular pulse.*
- *Decreased pedal pulses.*
- *Stiffer arteries.*
- *Slight increases in BP and wider pulse pressure.*
- *Orthostatic drops in BP more common in elderly people.*



Checking the pulse

ABNORMAL FINDINGS/RATIONALE

- Increased incidence of vascular disease. If present, carotid bruit present and carotid thrills may be palpable.
- Increased varicosities.
- *Abnormal heart sounds:* Rate irregular with ectopic beats, S_4 common, systolic murmurs associated with aortic stenosis.
- Differentiate arterial insufficiency from venous insufficiency.



HELPFUL HINT

Here's how to differentiate a radiating murmur from a carotid bruit: A murmur is usually heard bilaterally and diminishes as you move away from the heart; a carotid bruit is unilateral and increases in intensity as you move away from the heart.

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

5. Gastrointestinal

SYSTEM/NORMAL VARIATIONS

- Inspect abdomen.
- Auscultate bowel sounds and vascular sounds.
- Palpate and percuss abdomen.
- Perform a rectal examination.
- *Abdomen very soft because of decreased musculature, so organs may be easier to palpate.*
- *Bowel sounds may be slightly decreased.*



HELPFUL HINT

Check for old scars—they may provide information about surgeries the patient forgot to mention in the history.

- *Stool negative for occult blood; no fecal impaction; prostate soft and smooth and not enlarged.*

ABNORMAL FINDINGS/RATIONALE

- Palpable bladder with retention.
- Palpable intestines if filled with stool.
- A pulsatile mass in the abdomen may be an aneurysm. Incidence of abdominal aortic aneurysms increases with age.



HELPFUL HINT

An aneurysm may have lateral as well as antero-posterior pulsation. Aneurysms are usually wider than 3 cm and often have an associated bruit. Surgical evaluation may be appropriate, particularly if aneurysm is greater than 5 cm.

- Dullness over bladder may signal urinary retention; dullness over bowel may signal stool retention. Do a follow-up physical exam to determine that dullness is not caused by a tumor.
- Bruits may be heard over stenotic arteries or aneurysms.
- Incidence of colorectal peaks between ages 85 and 92 and accounts for 20 percent of all cancers found in persons age 90 and above.
- Risk of benign prostatic hypertrophy (BPH) and prostatic cancer increases with age. If prostate feels abnormal, refer patient for urological evaluation.

6. Female Reproductive

SYSTEM/NORMAL VARIATIONS

- Inspect and palpate external genitalia.
- Perform pelvic exam.
- *Decreased elasticity, breast sag, cordlike feel to breasts.*
- *Decrease and graying of pubic hair.*
- *External genitalia decrease in size and skin becomes thin, inelastic, and shiny.*
- *Pelvic exam reveals pale vaginal walls and narrow, thick, glistening cervix as a result of decreased estrogen.*
- *Uterus and ovaries decrease in size. Ovaries should not be palpable.*



HELPFUL HINT

Elderly women may not be able to assume the full lithotomy position because of arthritic or cardiovascular changes. Adjust the position to ensure comfort, and use a smaller speculum for the pelvic exam.

ABNORMAL FINDINGS/RATIONALE

- Palpable ovaries, masses, rectocele, cystocele, or prolapsed uterus require referral.

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

7. Male Reproductive

SYSTEM/NORMAL VARIATIONS

- Inspect and palpate male genitalia.
- Perform rectal exam.
- *Gynecomastia may be seen.*
- *Decrease and graying of pubic hair.*
- *Scrotum and penis decrease in size.*
- *Testes hang lower and have fewer rugae.*
- *Prostate enlargement (BPH).*

ABNORMAL FINDINGS/RATIONALE

- Any mass requires referral.
- Enlarged prostate follow-up to rule out malignancy.

8. Musculoskeletal

SYSTEM/NORMAL VARIATIONS

- Inspect gait.
- Inspect and palpate muscles and joints.
- Test muscle strength.
- *About half of people over 65 have decreased arm swing during gait; a wider base of support; a decline in step length, stride length, and ankle ROM; decreased vertical and increased horizontal head excursions; decrease in spinal rotation and arm swing; increased length of double support phase of walking, and a reduction in propulsive force generalized at the push-off phase. Decrease in sensory input, slowing of motor responses, and musculoskeletal limitations result in an increase in unsteadiness or postural sway under both static and dynamic conditions.*
- *Thoracic curvature (senile kyphosis).*
- *ROM decreased.*
- *Crepitation, stiffness with ROM.*
- *Decreased muscle strength and tone. Muscle strength depends on muscle mass. Strong equal hand grip usually remains intact.*



HELPFUL HINT

An easy way to test upper extremity function is having the patient touch the back of his or her head with the hands, pick up a penny, and shake hands.

- *Shoes fit well, wear evenly. No lesions, calluses or deviations noted. Toenails well groomed.*



Inspecting the gait



Testing strength

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

8. Musculoskeletal (continued)

ABNORMAL FINDINGS/RATIONALE

- Musculoskeletal changes associated with aging increase the risk for falls and injury.
- Heberden's nodes, involving the distal interphalangeal joints, are commonly seen with osteoarthritis or DJD but are rarely inflamed.



Heberden's nodes



DJD (osteoarthritis)

- Asymmetrical decrease in strength and tone may be associated with TIA or stroke.
- Diabetes-related ulcerations, fungal infections of the feet or toenails, calluses, bunions, hallus valgus (a lateral deviation of the large toe), and other deformities are very common and can affect function.



Hallus valgus and bunions

9. Neurological

SYSTEM/NORMAL VARIATIONS

Mental Status

- Test mental status.
- Use the MMSE and Clock Scoring Test to screen for dementia. (See *Clock Scoring*, page 957)



HELPFUL HINT

To assess depression quickly, use the single-item Yale Depression Screening Tool, which simply asks, "Do you often feel sad or depressed?" This tool has been shown to correlate very closely with the 15-item GDS.



ALERT

Know the symptoms of depression. They can be easily misread as normal changes associated with aging.

- *Cognitive ability intact; benign forgetfulness; short-term memory and long-term memory for new information may decrease with age.*

Cranial Nerves

- Inspect and palpate the cranial nerves.
- Slower response time bilaterally.
- *CN I olfactory—decreased.*
- *CN II optic—decreased visual acuity.*
- *Presbyopia.*
- *Fundoscopy changes.*
- *CN III oculomotor—Pupils smaller and reaction to light not as brisk.*
- *CN III oculomotor.*
- *CN IV trochlear—Extraocular movements intact. [CNs III, IV, and VI]*
- *CN VI abducens*
- *CN V trigeminal—Increased stimulation needed to elicit corneal reflex.*
- *CN VII facial—Decreased taste.*
- *CN VIII acoustic—Presbycusis, increased loss of hearing of high-pitched sounds progresses to loss of all frequencies.*
- *CN IX glossopharyngeal—Decreased taste.*
- *CN X vagus—Decreased gag reflex.*
- *CN XI spinoaccessory—No change, but ROM and strength depend on musculoskeletal changes.*
- *CN XII hypoglossal—No change.*

(continued)

PERFORMING A HEAD-TO-TOE PHYSICAL ASSESSMENT

9. Neurological (continued)

Muscle Function Movements

- Test pronator drift and abnormal movements.
- *Positive drifting with minimal weakness.*
- *No abnormal movements.*

Cerebellar Function

- Test reflexes and cerebellar function.



HELPFUL HINTS

- Vibratory testing is more reliable on the malleolus (protuberance on both sides of the ankle joint) than on the great toe.
- Aging may cause decreased vibratory sense in the lower extremities and fingers. To test functioning of the posterior column, check position sense rather than vibratory sense. Keep in mind that position sense of the great toe may also be decreased.

Reflexes

- *Romberg sign—Slight swaying normal.*
- *Loss of balance (positive Romberg sign) may be associated with Parkinson's disease.*
- *Deep tendon reflex (DTR) may be increased or decreased (+1 or +3).*
- *Achilles reflex more difficult to elicit. May need to use reinforcement techniques to elicit response.*
- *Superficial abdominal reflexes disappear with age.*
- *Older adults may demonstrate the release of some primitive reflexes including the snout, glabellar, and palmomental.*

ABNORMAL FINDINGS/RATIONALE

- Confusion may be caused by delirium, an underlying dementia, or depression. These conditions may be difficult to differentiate and may occur independently or together.
- Patients with dementia work hard to answer questions and confabulate answers. Delirium causes difficulty concentrating on the questions and attending to the task. Patients with depression are often unwilling to try to complete the task or answer the question. Detection of dementia, delirium, or depression warrants referral. Several bedside screening tools are available to differentiate between dementia, delirium, and depression and to evaluate cognitive function. See Chapter 5, Approach to the Mental Health Assessment, for Characteristics of Dementia, Delirium, and Depression on page 117, and GDS on page 117 for further information.
- Diminished response time on only one side warrants further investigation.
- Tremors are common in older patients. Types include:
 - Postural or physiological—Benign fine tremors.
 - Intention, essential, familial, and senile—Visible tremor associated with intentional movements, diminishes with rest.
 - Rest—Visible tremor at rest but absent or diminished with movement; “pill-rolling;” associated with Parkinson's disease.
 - Action—Large, irregular tremors of limbs, associated with cerebellar dysfunction (e.g., in multiple sclerosis).
- Asymmetrical reflexes may indicate an underlying problem.
- Positive primitive reflexes may also indicate a severe neurological assault.

Common Skin Lesions Associated With Aging

SKIN LESION

DESCRIPTION

Lentigines	■ Hyperpigmented macular lesions (liver spots)
Ichthyosis	■ Dry, scaly, fishlike skin
Acrochordons	■ Small, benign polyp-growths (skin tags)
Actinic keratosis	■ Rough precancerous skin macule or papule from sun exposure
Seborrheic keratosis	■ Benign pigmented lesions with a waxy surface on face and trunk
Senile purpura	■ Vascular lesion of ecchymoses and petechiae on arms and legs caused by the frail nature of capillaries and decreased collagen support
Venous lakes	■ Bluish-black papular vascular lesions

SKIN LESION

DESCRIPTION

Senile ectasias	■ Red-purple macule or papule lesions (senile or cherry angiomas)
Basal cell carcinoma	■ Pearly, papular, or plaquelike cancerous lesions that may be ulcerated in the center; associated with sun exposure
Squamous cell carcinoma	■ Erythematous, indurated areas that may be scaly or hyperkeratotic, associated with sun exposure, and tend to grow more rapidly than basal cell carcinomas

Clock Scoring

Give the patient a piece of paper and ask her or him to draw a circle. Instruct the patient to draw the face of a clock inside the circle, putting the numbers in the correct positions. Then ask her or him to draw the hands to indicate 10 minutes after 11 or 20 minutes after 8.

Scoring

Assign 1 point for each of the following:

- Draws closed circle
- Places numbers in correct positions
- Includes all 12 numbers in correct positions
- Places clock hands in correct positions.

There are no specific cutoff scores. If performance on clock drawing is impaired, consider a complete diagnostic evaluation for dementia.

normal-sized print. Able to hear normal tone of voice but unable to repeat three whispered words. Ear canals bilaterally without evidence of impacted cerumen and tympanic membrane normal. Neck has full range of motion (ROM) without evidence of dizziness during range.

- **Respiratory:** Respirations, 16 and unlabored.
- **Cardiovascular:** Rate, 72 and regular without S₃, S₄, or murmur. Carotid upstroke normal. No bruits bilaterally.

	Right	Left
PULSES	2+	2+
RADIAL	2+	2+
BRACHIAL	2+	2+
DORSALIS PEDIS	2+	2+
POSTERIOR TIBIALIS	2+	2+
POPLITEAL	2+	2+
FEMORAL	2+	2+

- **Gastrointestinal:** Normal bowel sounds in all four quadrants. Abdomen soft with slight tenderness in right upper quadrant. Liver edge palpable and tender. No bruits noted.
- **Genitourinary/Reproductive:** Vaginal mucosa dry and pale but skin intact. No evidence of discharge. Speculum exam/Pap test refused because of age, and she would not do anything even if the test were abnormal. No evidence of prolapse. Has urinary leakage with bearing down when lying and also when standing and coughing or bearing down. Incisional scar on left breast. Breasts symmetrical and no masses or tenderness noted. No nipple retraction or discharge.
- **Musculoskeletal:** Slight enlargement of distal interphalanges (Heberden's nodules), but no evidence of warmth or tenderness. Full ROM in upper extremities with strength 5/5 in upper proximal and distal extremities and 4/5 in lower proximal and distal extremities. Increased curvature of thoracic spine. Gait steady with walker with 6-inch base of

(continued on page 958)



Case Study Findings

Your general health survey and head-to-toe assessment of Mrs. Nurick have concluded. The following are your findings:

- **General Health Survey:** 80-year-old woman who is well groomed and dressed, pleasant, and appropriate in behavior during the interview.
- **Vital Signs:** Temperature, 97°F; BP, 130/70; height, 5'2"; weight, 140 lb. Body mass index of 22.
- **Integumentary:** Skin intact throughout without evidence of growths or lesions. Skin is dry, especially on extremities. Toenails are thick and yellow on all nails bilaterally.
- **HEENT:** Hair is gray and thin. Pupils equal and react to light and accommodation. Conjunctiva white and no drainage or tearing noted. Fundoscopic exam revealed intact retina with no signs of hemorrhage. Vision 20/30 in the right eye and 20/20 in the left eye with corrective lenses. Able to read



Case Study Findings (continued)

support. Positive accentuated thoracic curve (kyphosis).

- **Neurologic:** Sensation intact to light touch, localization, and pin prick. Position sense intact. Cerebellar intact finger to nose, heel to shin, and rapid hand movement. Romberg negative. Reflexes symmetrical and 2+ throughout with Babinski downgoing. Scored a 29/30 on the Mini-Mental Status Exam (MMSE). Scored a 1/15 on the Geriatric Depression Scale (GDS).



Case Study Analysis and Plan

Now that your examination of Mrs. Nurick is complete, document the key history and physical exam findings that will help you formulate your nursing diagnoses.

Nursing Diagnoses

Consider all of the data you have collected during your assessment of Mrs. Nurick, and then use this information to identify a list of nursing diagnoses. Some possible ones are provided below; cluster the supporting data.

1. Imbalanced Nutrition: less than body requirements, related to loss of appetite
2. Risk for Injury related to altered mobility and fatigue
3. Risk for Activity intolerance related to fatigue
4. Fatigue

Identify any additional nursing diagnoses.



CRITICAL THINKING ACTIVITY #5

Now that you have identified some nursing diagnoses for Mrs. Nurick, select a nursing diagnosis from above and develop a nursing care plan and a brief teaching plan for her, including learning outcomes and teaching strategies.

RESEARCH TELLS US

Cognitive impairment in older adults adversely affects outcomes and may result in an increased hospital stay, increased complications and readmission, poor rehabilitation, increased cost, and increased mortality rates. Cognitive function may not always be assessed in the acute care setting.

A recent exploratory study was designed to determine the extent of cognitive impairment in hospitalized older adults. The study also identified the needs of cognitively impaired older adults and their caregivers during and after the acute illness. Cognitive function was assessed using the MMSE, Brief Dementia Severity Rating Scale, and Confusion Assessment Method. Interviews of both patient and caregivers were used to identify needs.

The sample consisted of older adults hospitalized for common medical/surgical problems at three hospitals over 4 months. A total of 145 completed the screening; 65 percent (n = 94) had no evidence of cognitive

impairment; 35 percent (n = 51) had cognitive impairment, of which 65 percent (n = 33) were identified as cognitively impaired only through screening.

The patients and caregivers identified a need for increase responsiveness. The first and second weeks following discharge were identified as vulnerable periods for the patient and caregiver. This time is the point when the highest risk for relapse to likely occur. The study also identified that the need for support still existed at 6 weeks postdischarge.

The implications for nursing are many. The findings suggest that cognitive impairment is often missed. Nurses should routinely assess all older adults for cognitive impairment. Discharge planning should include the patient and the caregiver. Educate patients and caregivers of the risk and behaviors of cognitive impairment. Supply support or referrals postdischarge, keeping in mind the vulnerable periods following discharge.

Source: Naylor, M., Stephens, C., Bowles, K., and Bixby, B. (2005). Cognitively impaired older adults: From hospital to home: An exploratory study of these patients and their caregivers. *American Journal of Nursing* 105(2):52-61.

HEALTH CONCERNS

The screening tests recommended for adults are especially important for older adults because of their

increased risk of disease. The following are the pros and cons of several of these tests.

Pros and Cons of Health Screening Tests and Preventive Treatments


ACTIVITY SUGGESTED	FREQUENCY	PROS	CONS
Immunization		<ul style="list-style-type: none"> ■ Disease prevention, particularly to prevent an epidemic. 	<ul style="list-style-type: none"> ■ Side effects such as local irritation or flulike symptoms.
Pneumonia Flu Tetanus	Every 6 yr Every yr Every 10 yr	<ul style="list-style-type: none"> ■ Decreased morbidity and mortality associated with pneumonia or flu. 	
Pap test	Every year for high-risk women; otherwise, every 3 yr	<ul style="list-style-type: none"> ■ Risk for cervical cancer increases with age. ■ Older women may not have had the opportunity to get regular Pap tests and may want this early screening. ■ Pursue only if woman is willing to undergo treatment if disease is identified. ■ Noninvasive treatment may improve quality of life by detecting and managing problems such as atrophic and dry vaginal mucosa. 	<ul style="list-style-type: none"> ■ Cervical cancer develops slowly and is unlikely to be cause of death in women over age 90. Less risk of cervical cancer if patient is not sexually active. ■ Testing can be difficult and uncomfortable in older women, particularly those no longer (or never) sexually active.
Mammogram	Every yr	<ul style="list-style-type: none"> ■ Breast cancer risk increases with age. These tumors are generally estrogen receptor positive and treatable. ■ Pursue only if woman is willing to undergo treatment if disease is identified. 	<ul style="list-style-type: none"> ■ New-onset breast cancer is not likely to be cause of death in women age 90 and older. Tumors in older women tend to grow slowly. ■ Discomfort associated with mammogram.
Prostate exam	Digital rectal exam and prostate-specific antigen (PSA) every yr	<ul style="list-style-type: none"> ■ Risk for prostate cancer increases with age. ■ Pursue only if man is willing to undergo treatment if disease is identified. Survival rates increase dramatically with early detection. 	<ul style="list-style-type: none"> ■ Controversy persists with regard to effectiveness of treatment options and usefulness of treatment, especially for men with > 10 years' life expectancy.
Fecal Occult Blood Testing (FOBT)	Every yr	<ul style="list-style-type: none"> ■ Early detection of a cancer that could cause discomfort and affect quality of life if left untreated. ■ Easily performed with no discomfort to patient. ■ FOBT has better predictive value in older adults than in young adults. 	<ul style="list-style-type: none"> ■ False positives, which may result in additional testing and anxiety for patient.
Cholesterol Monitoring/ Interventions	Initial and then as appropriate	<ul style="list-style-type: none"> ■ Although not tested in patients age 90 and above, decreasing cholesterol reduces mortality and cardiac events in patients over 65. 	<ul style="list-style-type: none"> ■ Diet restrictions may affect quality of life. May be no advantage to restricting dietary fat in those age 90 and over. Focus should be to eat a healthy diet low in fat and high in fruits, vegetables, and grains. ■ Lipid-lowering medications may have unpleasant side effects and/or interfere with other medications.

S U M M A R Y

- The history and physical exam of the older adult should take into consideration the normal changes associated with aging and focus strongly on function.
- Every system clearly plays a role in the older adult's ability to perform ADLs. As you evaluate the systems, your goal is to identify problems that can be treated to improve function.
- Pay special attention to gait disorders, incontinence, sensory problems, cognitive status, and mood changes because these problems often reflect the presentation of a much more serious condition.

R E V I E W Q U E S T I O N S

1. You are performing a complete assessment on Theresa Wagner, age 75. Considering Mrs. Wagner's age, what nail change would you expect to find?
 - a. Increase in transverse ridges
 - b. Increased thinning of nails
 - c. Thick and brittle nails
 - d. Increase growth of nails
2. When examining Mrs. Wagner's eyes, which finding would be considered a normal finding associated with aging?
 - a. Enlarged pupils
 - b. Arcus senilis
 - c. Macular degeneration
 - d. Cataracts
3. When assessing Mrs. Wagner's skin, which finding would be considered a normal finding associated with aging?
 - a. Increased temperature
 - b. Decreased turgor
 - c. Increased thickness
 - d. Increased moisture
4. When assessing Mrs. Wagner's pupils, what finding would be considered a normal finding associated with aging?
 - a. Increase in pupillary size
 - b. Irregular reaction to light
 - c. Decrease in pupillary size
 - d. Sluggish reaction to light
5. When assessing Mrs. Wagner's spine, what finding would be considered a normal finding associated with aging?
 - a. Kyphosis
 - b. Scoliosis
 - c. Lordosis
6. When assessing Mrs. Wagner's hearing, what finding would be considered a normal finding associated with aging?
 - a. Difficulty hearing high-pitched sounds
 - b. Difficulty hearing low-pitched sounds
7. You are performing a pelvic exam on Mrs. Wagner. What finding would be considered a normal finding associated with aging?
 - a. Pale vaginal walls
 - b. Red vaginal walls
 - c. Pink vaginal walls
8. You are performing a pelvic exam on Mrs. Wagner. What finding would warrant medical follow-up?
 - a. Pale, pink vaginal walls
 - b. Palpable ovary
 - c. Decrease in pubic hair
 - d. Decrease vaginal secretions
9. Mrs. Wagner states that she has a bladder problem and is incontinent every time she coughs. What type of incontinence does Mrs. Wagner have?
 - a. Stress
 - b. Urge
 - c. Overflow
 - d. Functional
10. When assessing Mrs. Wagner's musculoskeletal system, what finding would be considered a normal finding associated with aging?
 - a. Decreased base of support
 - b. Increased stride length
 - c. Decreased height
 - d. Accentuated lumbar curve



APPENDIX

A

Chapter Feedback

CHAPTER 1

PAGE 17: CRITICAL THINKING ACTIVITY #1

- **Headache:** Subjective
- **BP 170/110:** Objective
- **Nausea:** Subjective
- **Diaphoresis:** Objective
- **Equal pupillary reaction:** Objective
- **Tingling sensation:** Subjective
- **Dizziness:** Subjective
- **Decreased muscle strength:** Objective
- **Slurred speech:** Objective
- **Numbness of left arm:** Subjective

PAGE 17: CRITICAL THINKING ACTIVITY #2

- **Primary data source:** Patient, Eileen Ploransky.
- **Secondary data source:** Husband, prenatal records.
- **Subjective data:** Contractions like “severe cramps.” Husband says, “I think her water broke on the way to the hospital.”
- **Objective data:** 29-year-old woman, contractions 5 minutes apart, 6 cm dilated, fetal heart rate (FHR) 120 BPM, VS 140/80, 90, 22, H & H 12.0, 45, blood type AB+.

PAGE 22: CRITICAL THINKING ACTIVITY #3

- **Physiological:** Tissue Perfusion, ineffective; Nutrition: imbalanced; Elimination, alterations in
- **Safety:** Mobility, Impaired Physical; Injury, Risk for
- **Love and belonging:** Social Interaction, impaired; Role Performance, Ineffective
- **Self-esteem:** Self-esteem, Risk for Situational Low; Body Image, Disturbed
- **Self-actualization:** Spiritual Well-Being, Readiness for Enhanced; Spiritual Distress; Health-Seeking Behaviors

PAGE 22: CRITICAL THINKING ACTIVITY #4

- **Bleeding:** Primary
- **Fatigue, pallor, hemoglobin 7.0:** Secondary
- **Nausea and vomiting:** Secondary
- **Grieving:** Tertiary
- **Sleep disturbance, fever, night sweats:** Secondary
- **Knowledge, deficient, on self-administration of insulin:** Tertiary
- **Impaired mobility, fractured hip:** Secondary
- **Irregular heart rate > 140 BPM:** Primary
- **BP 70/40:** Primary
- **New parent role:** Tertiary

PAGE 25: REVIEW QUESTIONS

1. a. Vital signs, height and weight measurements are assessment data. Assessment is the data collection phase of the nursing process.

2. d. Assessment is the data collection phase of the nursing process.
3. a. Preventive healthcare at the primary level focuses on health promotion and prevention. Teaching a drug awareness program provides education about the health risks associated with drug use.
4. c. Tertiary preventive care focuses on extended, rehabilitative care. Hospice nursing is an example of care at the tertiary level of preventive healthcare.
5. b. A focused assessment is problem oriented and may be an initial assessment or an ongoing assessment. A focused assessment is warranted when the patient’s condition is acute or unstable.
6. c. When obtaining a health history, the personal space should be about 3 to 4 feet.
7. b. Open-ended questions are questions that elicit the patient’s perception of the problem or situation. Closed-ended questions are questions that elicit a yes/no response.
8. a. Redirecting is a communication technique that allows you to refocus the conversation to the topic at hand. This technique keeps the communication goal directed.
9. b. The primary purpose of the working phase of the interview is data collection.
10. c. As a beginning practitioner, having a colleague or physician double-check your finding is an ideal means to validate your data.

CHAPTER 2

PAGE 29: CRITICAL THINKING ACTIVITY #1

- Where is the chest pain? Point to where it hurts.
- What does it feel like?
- Did you ever have this before?
- What were you doing before the pain started?
- Does it go anywhere else?
- Does anything make it better?
- When did it start? How long did it last?

PAGE 29: CRITICAL THINKING ACTIVITY #2

- **Past Health History:** Did you ever have this before? Do you have a history of heart disease or high BP?
- **Family History:** Do you have a family history of heart disease, heart attack, or high BP?
- **Review of Systems:** Do you have any breathing problems? Nausea or other GI complaints?
- **Psychosocial Profile:** How has this problem affected your activities of daily living, ability to work, or ability to have fun?

PAGE 36: CRITICAL THINKING ACTIVITY #3

- Cardiovascular disease
- HTN

- MVP
- Breast cancer

PAGE 51: REVIEW QUESTIONS

1. a. The quality of a symptom is best assessed by asking the patient "What does it feel like?"
2. b. The severity of a symptom is very subjective and is best assessed by asking the patient to scale the symptom on a 0 to 10 scale.
3. d. Having the patient localize the symptom is the best way to assess location of a symptom.
4. a. The source and reliability of the history is found in the biographical data.
5. a and c. The chief complaint is a direct quote from the patient as to what the problem is and how long it has occurred.
6. c. The purpose of the family history is to identify familial or genetically linked health problems. In-laws need not be identified since there is no genetic link.
7. c. The purpose of the review of systems is to identify the current and past health of each system.
8. b. One of the purposes of the psychosocial history is to identify environmental health risk factors.
9. c. The genogram is a means of documenting the family history to identify familial or genetically linked health problems.
10. a. Once the chief complaint is identified, a symptom analysis is performed to provide a thorough description of the symptom.

CHAPTER 3

PAGE 74: CRITICAL THINKING ACTIVITY #1

With a blind patient, explaining exactly what you plan to do is even more important. Letting the patient feel the equipment before you use it may also help put him or her at ease.

PAGE 74: CRITICAL THINKING ACTIVITY #2

- Color changes such as cyanosis or pallor
- Diaphoresis
- Grimacing
- Guarding because of pain

PAGE 85: REVIEW QUESTIONS

1. d. A focused physical assessment is indicated when the patient's condition is acute or time does not permit a complete physical assessment. Since the patient has an acute problem, chest pain, a focused physical assessment is indicated.
2. a. The bell portion of the stethoscope is best for detecting low-pitched sound. When using the bell, it should be held lightly on the chest wall.

3. c. The dorsal aspect of the hand is best for assessing temperature changes.
4. a. The balls (bony portion) of your hand are best for detecting vibratory sensations.
5. b. The finger pads are the most discriminatory able to detect fine sensations.
6. a. Tenderness is elicited upon palpation; pain is present whether you are touching the patient or not.
7. d. The normal heart rate for a healthy infant ranges between 120 to 160 BPM.
8. a. The difference between the systolic and the diastolic pressure is referred to as the *pulse pressure*.
9. b. The best technique for taking a BP is to inflate the cuff 30 mm Hg above the point at which the palpable pulse disappears. This prevents falling into the auscultatory gap and obtaining an inaccurate reading.
10. a. Apnea is a period of absent respirations lasting for 15 seconds.

CHAPTER 4

PAGE 102: CRITICAL THINKING ACTIVITY #1

- Questions to ask Mrs. Kane prior to surgery:
 - What do you do to manage your pain?
 - Do you use alternative means of pain control such as meditation?

PAGE 102: CRITICAL THINKING ACTIVITY #2

- Yes, Mrs. Kane could experience acute and chronic pain.
 - She may experience chronic pain associated with DJD.
 - She'll experience acute pain related to surgery

PAGE 102: CRITICAL THINKING ACTIVITY #3

- 0 to 10 pain scale would be appropriate.

PAGE 107: CRITICAL THINKING ACTIVITY #4

- Patient verbalizes cause of pain.
- Guarding surgical site.
- Limited movement.
- Increase in vital signs.

PAGE 107: CRITICAL THINKING ACTIVITY #5

- Pain assessment should be assessed every hour.

PAGE 108: REVIEW QUESTIONS

1. a. Pain is a subjective symptom; therefore, asking the patient what she or he is feeling is the best way to assess pain.
2. b. The best way to assess the quality of pain is to ask the patient what does it feel like.

3. c. The best way to assess location of pain is to have the patient point to where it hurts.
4. d. The best way to assess the severity of pain is to have the patient scale the pain.
5. a. The best question to assess timing of the pain is to ask when the pain began. The other questions addressed different characteristics of pain.
6. d. The best question that identifies precipitating factors of pain is to ask the patient what he or she was doing before the pain started. The other questions identify different characteristics of pain.
7. c. Chronic pain persists for 3 months or longer.
8. a. Caustic chemical burns usually result in a burning sensation.
9. c. Musculoskeletal pain such as joint pain associated with DJD results in aching sensation.
10. a. Acute appendicitis usually results in a sharp pain sensation.

CHAPTER 5

PAGE 126: CRITICAL THINKING ACTIVITY #1

- Significant data:
 - Hypothyroidism and breast cancer
 - Side effects of medications
 - Depression
 - Insomnia
- Additional questions:
 - Questions to determine if she is suicidal
 - Questions to determine if she is taking her medicine

PAGE 126: CRITICAL THINKING ACTIVITY #2

- Priority concern is risk for suicide

PAGE 126: CRITICAL THINKING ACTIVITY #3

- Fatigue
- Inability to sleep
- “No energy. . .”
- Feeling “overwhelmed”
- Depressed
- Recently divorced

PAGE 137: CRITICAL THINKING ACTIVITY #4

- Weight gain
- Poor eye contact
- Slow to respond
- Disheveled appearance
- Slouched position
- Crying
- Difficulty staying focused
- Constipation

PAGE 137: NURSING DIAGNOSES

- **Risk for suicide**
 - Patient states “maybe things would be better if I could disappear.”
 - History of depression
- **Ineffective Individual Coping**
 - Inability to sleep
 - Feeling overwhelmed
- **Fatigue**
 - Inability to sleep
 - No energy
 - Fatigue
 - Side effects of medication

PAGE 142: REVIEW QUESTIONS

1. a. Lithium may cause hypothyroidism. Thyroid function should be assessed periodically.
2. c. Gamma-aminobutyric acid (GABA) is a neurotransmitter associated with anxiety.
3. a. A sudden change in mental status with confusion is often associated with delirium. Dementia, psychosis, and personality disorders that change are not as acute.
4. d. Depression is the most common mental health problem seen in older adults.
5. a. Patients with schizophrenia often experience auditory hallucinations.
6. a. Alcoholic patients who have memory loss often use confabulation to “fill in the blanks” or gaps of memory loss.
7. b. Depression is a side effect of beta-blocker drugs.
8. b. Because tricyclic medications could increase intraocular pressure, use would be contraindicated in patients with glaucoma.
9. c. A patient with bipolar disorder in a manic state often has difficulty staying focused and may exhibit flight of ideas.
10. d. Detaching oneself from a situation is referred to as *depersonalization*.

CHAPTER 6

PAGE 151: CRITICAL THINKING ACTIVITY

At the end of the lesson, Lucas will be able to put on and remove a condom according to the effective condom use checklist.

PAGE 151: REVIEW QUESTIONS

1. a. A patient with hyperthyroidism may have difficulty with temperature. You may need to adjust the room temperature to meet the patient’s needs.
2. b. Since the patient will be self-administering insulin, assessment of vision is important to ensure accurate dosage.

3. a. Older adults often have memory loss (immediate and recent). Assessing memory would be important to ensure retention. Providing written instructions may be helpful for the patient who has memory deficits.
4. a. Patients with CHF often experience fatigue related to decreased cardiac output. Shortening class time may be an effective way of preventing fatigue in this patient.
5. c. Since patients with immunosuppression are at increased risk for infection, one-on-one instruction may minimize the risk.

CHAPTER 7

PAGE 162: CASE STUDY ANALYSIS AND PLAN

Jan (age 16) is a junior in high school. She had screenings for vision, dental, hearing, height, weight, hemoglobin and hematocrit, vital signs, and stress. She was assessed for current vaccinations and received educational materials on exercise, nutrition, weight management, safe sex, accident prevention (such as driving and seat belt use), and drug and alcohol abuse.

Mrs. Chen had her height, weight, body mass index (BMI), vision, cholesterol, pulse, respiration, and BP checked. She was also assessed for history of immunizations such as tetanus and hepatitis B. She received educational materials on menopause, nutrition, prevention of osteoporosis, self-assessment of risk for cancer, and heart disease.

Mrs. Wong had her height, weight, BMI, vision, and vital signs checked. In addition, she had a screening bone mineral density (BMD) test of her heel to detect signs of osteoporosis, which would indicate whether more extensive testing (e.g., hip and spine) was needed. Her immunization history (e.g., tetanus, hepatitis B, influenza, and pneumonia) was also taken. She received educational materials on nutrition, especially the need for calcium and vitamin D, osteoporosis, use and abuse of medications and alternative therapies, safe exercise, fall prevention, and screening for heart disease, cancer, and depression.

PAGE 162: REVIEW QUESTIONS

1. a. Obesity and snoring are risk factors for sleep apnea.
2. b. A side effect of beta-blockers is nightmares, which may affect a person's quality of sleep.
3. c. Older adults often have difficulty falling asleep. Total sleep time is usually the same as that of a younger adult, but often older people have a fragmented sleep pattern.
4. b. To calculate target heart rate while exercising, subtract patient's age from 220, then multiply by 60 to 80 percent.

5. c. Developmentally, adolescents are known risk takers. Accidents are the leading cause in children and young adults.

CHAPTER 8

PAGE 167: CRITICAL THINKING ACTIVITY #1

- **Illnesses that interfere with food intake:** These include dental disease, anorexia nervosa, and diseases or treatments that cause anorexia.
- **Lack of suitable foods:** Grocery stores in urban areas may not stock perishable items such as fresh fruits and vegetables.
- **Intake habits that favor deficiencies:** Busy schedules have engendered a preference for fast foods that are quick and tasty but high in calories, salt, and fat and lacking in nutrients.
- **Inability to afford necessary food for an adequate diet:** This is a principal reason for primary malnutrition among poor and older adults.
- **Knowledge deficiency:** Lack of knowledge about good nutrition.

PAGE 167: CRITICAL THINKING ACTIVITY #2

Chronic and hereditary conditions may impair digestion and absorption of nutrients from the GI tract or cause competition for nutrients at the cellular level. Acute medical conditions (e.g., GI infection or inflammatory bowel disorders), multiple trauma, serious burns, and major abdominal surgery are causes of secondary malnutrition.

PAGE 177: CASE STUDY EVALUATION

Biographical Data

- 78-year-old African American male.
- Widowed, lives alone in urban area.
- Retired postal worker.
- Baptist religion.
- Medicare/Medicaid insurance.
- Daughter is primary contact.

Current Health Status

- Vague complaints of not feeling well, being very tired, and having no energy. Weight loss of 10 lb since wife's death.
- Current medications are Lasix 40 mg od and Vasotec 5 mg bid. No OTC/OTC medications except occasional milk of magnesia (MOM) for constipation. No illegal drug use.

Past Health History

- Positive history of HTN for 10 years.
- Hospitalized last year for uncontrolled HTN.

- Appendectomy at age 12.
- No allergies to food, drugs, or environmental factors.
- Medications listed previously.

Family History

- Daughter, age 50, alive and well.
- Wife deceased age 74, breast cancer.
- Mother deceased age 75, HTN, stroke.
- Father deceased age 80, HTN, MI.
- Sister, age 70, HTN.
- Sister age 73, alive and well.
- Brother, age 75, DM, HTN, CAD.
- Maternal aunt deceased age 80, CAD, MI.
- Maternal uncle deceased age 78, stroke.
- Paternal aunt deceased age 82, colon cancer.
- Paternal uncle deceased age 81, stroke.

Review of Systems

- **General Health Survey:** 10-lb weight loss, fatigue.
- **Integumentary:** Dry, flaky skin.
- **HEENT:** Wears dentures.
- **Respiratory:** No changes.
- **Cardiovascular:** No chest pain; positive history of HTN.
- **Gastrointestinal:** Bowel movement every 2 to 3 days, brown and formed. Takes MOM as needed.
- **Genitourinary/Reproductive:** On Lasix; no problems with urination; voids about eight times a day. Not sexually active since wife died.
- **Musculoskeletal:** Weakness and occasional cramping in legs.
- **Neurological:** No sensory changes. Very lonely since wife's death.
- **Lymphatic:** No allergies or recent infections.

Psychosocial Profile

- Wife used to prepare meals, finds it hard to cook for one person.
- Says dentures are a little loose since he lost weight.
- Visits doctor every 6 months for BP check.
- Typical day: Awakens at 7 A.M., has breakfast, walks to corner for newspaper, takes midmorning nap, usually skips lunch, watches TV, has dinner around 5 P.M., watches TV after dinner, usually falls asleep while watching TV, goes to bed after the 10 P.M. news.
- Has three 8 oz. glasses of water a day, not trying to lose weight. Eats out about once a week, likes fried foods. Not much appetite since wife died. Usually eats alone.
- Daily walk to corner for paper, no other scheduled exercise.
- Sleeps 7 to 8 hours a night. Awakens at least once to go to bathroom. Takes two 20- to 30-minute naps during day.

- Never smoked or used drugs. Occasional alcoholic drink about twice a month.
- Retired postal worker on fixed income with government pension; able to meet financial needs.
- Most close friends have passed away.
- No cultural or religious influence on dietary habits.
- Says life is "pretty routine without much stress."
- Daughter is main source of support; lives 1 hour away, visits once a week, takes patient shopping.

24-Hour Recall

- Breakfast: 8 oz. black coffee, 8 oz. water, bowl of Cheerios with whole milk
- No lunch.
- Afternoon snack: Four pretzels.
- Dinner: Frozen dinner of Salisbury steak, potatoes, and corn, piece of apple pie, 8 oz. water, 8 oz. black coffee.

PAGE 177: CRITICAL THINKING ACTIVITY #3

- Recent loss of wife.
- Fact that he lives alone.
- Weight loss.
- Reports that it is hard to prepare meals for himself and that he skips.
- Diuretic use and urination patterns.
- Fatigue.
- Dry, flaky skin.

PAGE 182: KEY HISTORY FINDINGS

- Wife died 6 months ago.
- Lives and eats alone.
- 10-lb weight loss in past 6 months.
- Says it is hard to cook for one person; skips lunch, eats frozen dinners; not much appetite.
- Dry, flaky skin.
- Wears dentures.

PAGE 182: KEY PHYSICAL EXAMINATION FINDINGS

- 10-lb weight loss since last examination.
- Increased heart rate.
- Sad affect, poor eye contact.
- Dry, flaky skin.
- Dry mucous membranes.
- Dentures poorly fitted.
- Decreased muscle strength.

PAGE 182: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- 24-hour recall, weight loss, dry mucous membranes, patient's admission that he does not eat regularly and that it is hard to cook for one.

- Limited intake, urination patterns, dry mucous membranes, increased heart rate.
- 24-hour recall, weight loss, skipped meals, fluid intake, diuretic use.

PAGE 185: REVIEW QUESTIONS

1. a. Primary malnutrition resulting from inadequate intake of nutrients. Anorexia nervosa is characterized with a preoccupation with being thin and dieting, leading to excessive weight loss.
2. c. There are many different means to perform a nutritional assessment. A 24-hour recall is ideal for screening for nutritional deficits.
3. b. A BMI is calculated using the person's height and weight.
4. a. Normally, difference between mid-arm circumferences should not exceed 1 cm.
5. a. The BMI is a valuable piece of assessment data that reflects body fat.
6. d. Measurements of triceps skinfolds are useful in estimating body fat.

CHAPTER 9

PAGE 190: CRITICAL THINKING ACTIVITY #1

Recognizing the range of different traditions, the best approach would be to ask the patient and/or family how you might best assist in providing comfort to the patient. For example, you may discover that the patient wishes to wash before praying or that she or he would like to have religious symbols pinned to her or his hospital gown or visible in the room.

PAGE 190: CRITICAL THINKING ACTIVITY #2

- Who are your support people?
- What provides you with strength and hope?
- What gives your life meaning and purpose?
- How is your faith important to you?

The answers to these questions will demonstrate the importance of spiritual issues in your patient's care. For example, the patient may mention a clergyman or fellow church (temple, mosque) member as important in his or her spiritual life. If the patient mentions only family members, then connectedness to others rather than organized religion is more important (but still spiritual in nature). The presence of religious symbols, such as the Bible, Qur'an (Koran), or prayer books; wearing of jewelry with religious symbols; presence of other objects such as crystals; or listening to inspirational or religious music will also signify the impact of spirituality on your patient's life. These behaviors will give the patient strength for enduring or recovering from illness.

PAGE 190: CRITICAL THINKING ACTIVITY #3

- **African Americans:** Allow privacy when clergy visits, offer to read Bible passages for patient or pray with patient.
- **Amish:** Allow time for prayer before meals, rather than rushing patient to start eating. Allow for consultation with religious leader and family.
- **Appalachians:** Realize that patient may refuse some types of therapies, including preventive measures because of fatalistic viewpoint. Allow privacy for prayer. May need to reassure patient that it is all right to pray in hospital.
- **Arab Americans:** Provide for washing and hygiene before prayers (five times a day). Help patient to face East. Also realize that patient may prefer to endure pain rather than expect relief from it.
- **Chinese Americans:** Allow privacy for prayer.
- **Cuban Americans:** Realize that fatalism may interfere with planned medical and nursing therapies. Although many are Catholic, may rely on folk medicine as well. Allow use of folk remedies as long as they have been cleared as safe.
- **Egyptian Americans:** Allow practice of Christian or Muslim rituals as appropriate.
- **Filipino Americans:** Prayer is important; provide privacy. Usually Catholic. Use of religious symbols (rosary, crucifix) is source of comfort. Realize that patient may be fatalistic, making treatment difficult. Respect patient's decisions.
- **French Canadians:** Although prayer is important, there is little acceptance of spirituality as important to health. Because family is important source of strength, provide for adequate visiting.
- **Greek Americans:** Encourage display of icons of family saint or Virgin Mary. Allow for daily prayer.
- **Iranian Americans:** Allow for Muslim prayers (five times a day). Help patient to face East and provide for washing before prayers. Fatalistic, so respect patient's decisions not to pursue therapies.
- **Irish Americans:** Most are Catholic. Encourage symbols of faith (rosary, crucifix). Offer to call priest as needed. Provide privacy for prayer.
- **Jewish Americans:** Provide for observance of dietary laws if this is important to patient. Allow time for prayer. Orthodox men may wear yarmulke at all times or only when praying.
- **Mexican Americans:** Usually Catholic. Similar to Cuban Americans and Filipino Americans. Folk medicine is important and should be allowed if not contraindicated. Encourage use of cultural symbols of the spirit and discussion of the soul.
- **Navajo Native Americans:** Provide for healing ceremonies by tribal shaman as complementary to traditional medicine. Provide privacy. Arrange room assignments to permit visibility of outdoors.

- **Vietnamese Americans:** Allow modified altar (without candles lighted) in room, if possible. Arrange for family visits; family and prayer are major sources of strength.

PAGE 194: CRITICAL THINKING ACTIVITY #4

Strengths

- Close relationship with sister.
- Church affiliation (active in church before hospitalization).

Areas of Concern

- Not geographically close to daughter.
- Does not drive.
- Feels isolated from church community.

PAGE 194: CASE STUDY EVALUATION

Biographical Data

- 68 years old.
- Husband died 1 year ago.
- One grown daughter who lives in California with her husband and two sons, ages 4 and 6.
- Lives in two-story home; sister lives across the street in her own home.
- Born in Grady Falls, Pennsylvania.
- African American.
- Episcopal religion.
- MS in education, retired from teaching elementary school.
- Has Medicare with supplemental HMO from teacher's union.
- Hospital admission for CHF.
- Good historian; alert and seems reliable.

Current Health Status

- Difficulty breathing at rest, lower extremity edema, change in sleep patterns, uses two pillows.
- Currently hospitalized for exacerbation of CHF.
- Also has HTN.
- Current medications are lisinopril (Prinivil), 10 mg a day, for HTN.

Past Health History

- CHF for 3 to 5 years.
- HTN, treated for 15 years.

Family History

- Father died of heart failure at age 89.
- Mother died at age 70 from complications of a stroke.

Psychosocial Profile

- Typical day starts at 7:00 A.M., when Mrs. Mays arises, makes her bed, eats breakfast, and reads the paper.

Usually does light housework, such as dusting and washing breakfast dishes. She then rests because the activity leaves her short of breath. Generally calls her sister to come over for lunch and watch soap operas. Sister leaves by 4 P.M. to get dinner for her husband. Mrs. Mays usually eats a light supper and then does needlework until 10 P.M., when she watches the news. Goes to bed at 10:30 P.M.

- Eats three meals a day. Lunch is her main meal, and she enjoys it most because she usually has her sister for company. Doesn't like to cook just for herself, so eats a light supper.
- Until last week, walked to the corner store for her daily paper. Now has it delivered because the walk is too difficult. Also unable to walk to services at her church, which is three blocks away. Does not drive.
- Goes to bed at 10:30 P.M. and arises at 7 A.M. Not sleeping well because of breathing problems caused by CHF.
- Sees her doctor regularly for HTN and CHF.
- Sister lives across the street, and they spend most weekday afternoons together. Only child and two grandchildren live in California. They talk on the phone weekly, but she sees them rarely. Attended church and taught Sunday school and vacation Bible school until recently.
- Retired teacher.
- Hobbies are needlepoint, reading, watching TV.
- Lives in two-story house with bedroom on second floor.
- Has never smoked; doesn't drink alcohol.
- Says she doesn't like to make a "big fuss" over her health problems or "burden anyone" with them.

PAGE 201: LIST KEY HISTORY FINDINGS

- 68-year-old widow (1 year).
- Lives alone.
- Episcopal religion.
- Only child lives in distant state.
- Retired from teaching.
- CHF for 3 to 5 years; recent hospitalization for exacerbation.
- HTN for 15 years.
- Mother died of stroke-related causes at age 70.
- SOB makes walking difficult, so can't attend church.

PAGE 201: LIST KEY SPIRITUAL ASSESSMENT FINDINGS

- BP elevated despite medication.
- Lies in bed with face turned away from door, hands folded, and lips moving silently.
- Says, "I'm no use now."
- No visitors in 2 days; wonders why sister hasn't come.
- Wears cross, has Bible on nightstand.

PAGE 201: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Questions meaning of suffering and meaning of own existence; verbalizes concern about God's plan; unable to practice usual religious rituals; has alteration in behavior or mood as evidenced by withdrawal and anxiety.
- Hospitalizations necessary about every 6 weeks; statement about how her life has changed.
- Unable to get to church; lack of visitors; lack of relationship with pastor.

PAGE 201: ADDITIONAL NURSING DIAGNOSES

- Grieving, Anticipatory
- Coping, Ineffective

PAGE 201: NURSING INTERVENTIONS/RATIONALES

- Demonstrate respect for and acceptance of Mrs. Mays' values and beliefs. To foster a therapeutic nurse/patient relationship, the nurse must be accepting of the patient and her beliefs and values.
- Adapt nursing activities to accommodate values and religious beliefs (e.g., Mrs. Mays' desire for visit from clergyperson and privacy to receive communion).
- Maintain and respect patient's preferences during hospitalization.
- Allowance for rituals provides spiritual comfort. Encourage family (sister) to bring significant symbols to Mrs. Mays (Bible, religious jewelry).
- Promote comfort.

PAGE 202: REVIEW QUESTIONS

1. b. Certain religious faiths have dietary restrictions. The Jewish faith restriction with certain foods, such as dairy products and pork.
2. c. A person who is a Jehovah's Witness may refuse blood transfusion based upon religious beliefs.
3. b. Patients of Catholic faith may request a priest to administer the Sacrament of the Sick.
4. a. A person of Christian Scientist faith may refuse treatment based upon the belief that God will heal through prayer.
5. a. Islamic faith requires female patients to be cared for by female nurses.

CHAPTER 10

PAGE 222: CRITICAL THINKING ACTIVITY #1

- Fair skin.
- Family history of basal cell carcinoma and malignant melanoma.
- Only occasional use of sunscreen.

PAGE 222: CRITICAL THINKING ACTIVITY #2

- Usual state of health good.
- Nutritious diet.
- Motivation and health-seeking behaviors.

PAGE 222: CASE STUDY EVALUATION

Biographical Data

- 48-year-old Caucasian female, married, with two adult children (22 and 27 years old).
- Works part time in retail sales.
- Born in the United States, of English-Scottish descent.
- Protestant religion.
- Husband is accountant, health insurance provided by his employer.

Current Health Status

- Feels that she is in very good health. No current problems.
- Has recently started taking estrogen and progesterone (Prempro) for menopause symptoms. Is tolerating therapy well.
- Occasionally takes aspirin or acetaminophen for headache or back/shoulder discomfort, but not monthly.

Past Health History

- Usual childhood diseases, without sequelae.
- One miscarriage at 10 weeks of pregnancy, age 22.
- Hospitalized for two childbirths, with uncomplicated vaginal deliveries; cholecystectomy 2 years ago.
- No history of allergies or skin disorders.
- No recent exposures to people with infections.
- Unsure about immunization status. Received all childhood immunizations but cannot recall tetanus or other shots.

Family History

- Mother died of breast cancer at age 65, had previously had two basal cell lesions removed.
- Brother (age 45) has HTN and a history of childhood asthma.
- Father (age 73) has "borderline" BP and adult-onset DM, but is in generally good health. He has also had two basal cell cancers on his face and has several "sun spots."
- Two older siblings (ages 49 and 50) are in good health, although one had a small malignant melanoma between two toes, which was surgically removed 3 weeks ago, with no lesions or spread detected.
- Children are both in good health.

Review of Systems

- **General Health Survey:** Feels well generally.
- **Integumentary:** Skin is somewhat dry, itches occasionally. No sores or rashes. No changes in hair or nails.
- **HEENT:** No runny nose or itchy eyes.
- **Respiratory:** No SOB, congestion, or cough.
- **Cardiovascular:** No chest pain.
- **Gastrointestinal:** No abdominal discomfort, nausea, or change in bowel habits.
- **Genitourinary:** No change in bladder habits. Last menstrual period was about 1.5 years ago. Before that, periods were regular until the final year, when they fluctuated a little.
- **Musculoskeletal:** No joint pain except mild discomfort after extensive gardening.
- **Neurological:** No dizziness or headache.
- **Endocrine:** No problems.
- **Lymphatic/Hematologic:** No problems.

Psychosocial Profile

- Bathes or showers daily, usually with very warm water. Shampoos her hair every other day and rarely uses any hair spray, mousse, or other hair cosmetics. Nails closely filed with clear polish. Wears cosmetics daily and has never had problems tolerating them. Colored her hair in the past but does not do so now.
- Is often out in the Florida sun as she walks and gardens, but only occasionally wears sunscreen. Does not reapply sunscreen when she has been out in the sun for extended time. Worries about damage to skin now that her sister has had skin cancer.
- Typical day starts at 7:00 A.M., when her alarm wakes her. On days when she works (2 to 3 days a week), she showers, dresses, eats breakfast, and works from 9 A.M. to 5 P.M. On her nonworking days, she cleans her house, works in the yard/garden when weather permits, and just “keeps busy” until about 5 P.M. She then fixes dinner and watches TV or does needlework in the evening. She generally goes to bed at 10:30 to 11:00 P.M.
- Eats three meals a day with foods from all categories. Feels that she should lose 10 to 15 lb, which she has gradually gained over the past 10 years. Tries to remember to drink several glasses of water daily, but sometimes drinks only one or two 8-oz. glasses.
- Gardening is one of her main forms of exercise. Also walks 2 or 3 miles a day three or four times a week.
- Sleeps very well, with rare interruptions to void (two or three times a week).
- Only medications are those mentioned previously. Drinks 1 cup of coffee and 4 to 5 glasses of iced tea daily. Rarely drinks colas. Has never smoked but

was exposed to second-hand cigarette smoke until her husband quit smoking 5 years ago. Does not drink alcohol or use recreational drugs.

- Is not exposed to chemicals or unusual environmental situations at work, but uses various chemicals in gardening (fertilizers, pesticides, and weed killers). Does not always wear gloves, masks, or other protective equipment.
- Has no pets. Has lived in current ranch-type home for 12 years. House is adequate for needs and has central air and heat.
- Enjoys meeting customers at work and has several close friends with whom she often has lunch or shops. Has close and caring relationship with husband and good relationship with their children, who live nearby. Father and siblings live 600 miles away, but she sees them two or three times a year for an enjoyable visit. Is not currently involved in any specific community group and rarely attends church.

PAGE 256: LIST KEY HISTORY FINDINGS

- Takes estrogen and progesterone.
- Has no allergies or skin disorders.
- Mother and father both had two basal cell lesions removed.
- Brother had childhood asthma.
- Gardens and walks for exercise.
- Rarely wears sunscreen.
- Sometimes uses fertilizers, pesticides, and weed killers without protection.
- Exposed to husband's second-hand smoke until 5 years ago.

PAGE 256: LIST KEY PHYSICAL EXAMINATION FINDINGS

- Well-developed, 48-year-old, fair-skinned female, appearance consistent with stated age.
- No signs of distress, appears comfortable and provides history with no difficulty.
- Temperature, 97.4°F; pulse, 82 and regular; respirations, 14 and unlabored; BP, 134/82 left arm.
- Height 5'4", weight 138 lb.
 - Hair
 - Generally brown with scant amount of gray interspersed and distributed evenly over scalp.
 - Cut short and well groomed.
 - Shiny, with no signs of damage.
 - No areas of excess hair growth on body.
- Nails
 - Short, with smooth edges.
 - Coat of clear, glossy polish present.
 - Nailbed pink, firm to palpation.
 - No hemorrhages or discoloration of nails or surrounding tissues.
 - No clubbing.

- Skin
 - Warm, with good turgor indicated by brisk recoil.
 - Generally intact, with smooth texture.
 - Coloring symmetrical: Pale pink on unexposed areas, moderately tanned on exposed areas, with diffusely scattered freckles, consistently light brown and all 0.2 to 0.4 cm in diameter.
 - Approximately 30 moles, 0.2 to 0.4 cm in diameter, with well-circumscribed, regular borders, all dark brown and consistent in color.
 - Three patches of slightly pink, scaly, flat skin, approximately 0.8 cm each (two on right forearm, one on left forearm); no telangiectasia, ulcerations of the sites, or changes in surrounding skin.

PAGE 256: CRITICAL THINKING ACTIVITY #3

Strengths

- Patient proactive
- Health-seeking behaviors

Areas of Concern

- Sun exposure with inconsistent sunscreen use
- Chemical exposure
- Family history of skin cancer
- Fair skin
- Multiple lesions on sun exposed areas > 0.5 cm in size

PAGE 256: NURSING DIAGNOSIS CLUSTERED SUPPORTING DATA

- Expresses concern about skin cancer history among relatives and past exposures to environmental skin irritants.
- Frequent sun exposure with failure to use sunscreen consistently.
- Frequent use of gardening-related chemicals without protective equipment.
- Expresses fear regarding potential for skin cancer related to family history and sun exposure.

PAGE 256: ADDITIONAL NURSING DIAGNOSES

- Infection, risk for, related to impaired skin integrity
- Injury, risk for, related to environmental conditions, exposures, trauma
- Skin Integrity, impaired, related to various external environmental or internal factors
- Tissue Perfusion, ineffective, related to interrupted arterial or venous circulation
- Pain (itching), related to chemical or inflammatory stimulation of superficial nerves
- Body Image, disturbed, related to real or perceived disfigurement of skin, hair, or nails
- Social Isolation, related to alterations in physical appearance

PAGE 264: REVIEW QUESTIONS

1. d. Turgor is a reflection of skin elasticity. Since skin elasticity decreases with aging, older patients normally have decreased skin turgor.
2. d. Turgor is best assessed on unexposed areas, such as below the clavicle. Areas such as the face, hands, and neck may be affected by the sun and reflect turgor.
3. a. The dorsal aspect of the hand is best for assessing skin temperature.
4. d. The oral mucosa is best for assessing central cyanosis, which reflects hypoxia.
5. b. Color changes are difficult to detect in dark-skinned individuals. The oral mucosa is best for assessing color changes in dark-skinned individuals.
6. d. Pressure ulcers that are covered with eschar cannot be staged. There is no way to determine the depth of the ulcer until the eschar is removed.
7. c. The normal angle of attachment of the nail is 160 degrees.
8. a. A stage I pressure ulcer is a reddened area with skin intact.
9. d. Nits are usually found close to the scalp. Nits found 1/4 inch or more from the scalp probably have already hatched.
10. a, c, d. Many skin changes are seen during pregnancy due to hormonal changes associated with pregnancy. Striae, cholasma and palmar erythema are examples of skin changes associated with pregnancy.

CHAPTER 11

PAGE 285: CASE STUDY EVALUATION

Biographical Data

- 17-year-old unmarried male.
- Works part-time in fast-food restaurant.
- Born in the United States, was adopted immediately after birth.
- Protestant religion.
- Adoptive father is computer salesman and adoptive mother is radiology technician.
- Source: Self, reliable.

Past Health History

- Usual childhood illnesses, without complications.
- Tonsils and adenoids removed at age 4 after frequent throat infections.
- Tympanoplasty at age 3 after frequent bouts of ear infections.
- Immunizations are up to date. Tetanus updated last year when patient stepped on a piece of glass while wading on a shoreline.

- Seasonal hay fever since childhood.
- Takes OTC antihistamines when he has no prescribed antihistamines on hand, but usually takes Claritin (fexofenidine) during seasonal allergy attacks. Has always taken one multivitamin daily. Takes no other medications.
- No recent exposures to known infections, but notes that he works in a busy restaurant and attends public school, so could be exposed to “whatever’s going around.”

Family History

- Biological family history is unknown; would like to know his family health history, but has been told this is not available.
- Adoptive parents are both in good health.

Review of Systems

- **General Health Survey:** States that health is usually good. No changes in weight or energy level.
- **Integumentary:** No changes in skin, hair, and nails. No skin rashes.
- **HEENT:** No swollen nodes or masses in head or neck. “Allergy attack” causes ear fullness; nasal congestion and clear drainage; postnasal drip; sneezing; and scratchy/itchy throat, eyes, and nose. Vision 20/20, no problems reported.
- **Respiratory:** No cough, SOB, or wheezing.
- **Cardiovascular:** No chest pain or history of cardiovascular disease.
- **Gastrointestinal:** Appetite good, no dietary restrictions, no nausea or vomiting, has daily bowel movement.
- **Genitourinary:** No history of renal disease.
- **Reproductive:** Heterosexual, not currently sexually active, is aware of safe sex practices.
- **Musculoskeletal:** No weakness reported.
- **Neurological:** No headaches, dizziness, tremors, or paresthesia.
- **Endocrine:** No changes in weight or shoe size (shoe size 11).
- **Lymphatic/Hematologic:** Positive seasonal allergies, no bleeding disorders.

Psychosocial Profile

- Bathes daily; uses no cosmetics on face other than occasional OTC acne medication; sees doctor and dentist for checkup once a year.
- Typical day starts at 6:45 A.M., when his mother awakens him to get ready for school. Attends school from 8 A.M. until 3:15 P.M. on weekdays, and works 5 P.M. until 11 P.M. Monday and Friday nights and noon until 6 P.M. on Saturday. Goes to bed at 11:30 P.M. on school nights and 1 A.M. on weekends.

- Eats three meals plus one or two snacks daily. Breakfast consists of cereal, milk, and juice. Lunch is fast food, obtained between classes. Dinner is usually a cooked meal, shared with his parents. No specific dietary restrictions or food intolerances.
- Plays baseball and competes on swim team; practices 1 to 2 hours a day, depending on time of year. Wears protective gear while playing baseball. Tolerates swimming well, wearing goggles. Does some amount of physical activity daily, and this is well tolerated.
- Normally sleeps well and awakens rested. During “allergy season” sometimes has sleep difficulty. When he takes OTC antihistamines, they make it difficult to sleep and he awakens frequently during the night. If he takes no antihistamines or decongestants, he often awakens with congestion and/or sinus pressure.
- No medications other than seasonal use of antihistamines and daily vitamin. Uses no recreational drugs. Admits to rare use of alcohol (beer).
- Works 5 P.M. until 11 P.M. Monday and Friday nights and noon until 6 P.M. on Saturday. Exposed to second-hand smoke in the restaurant where he works; otherwise, not generally around smokers or exposed to any known chemicals. Tries to avoid exposure to smoke because he is concerned about the potential health hazards.
- Lives in a single family, two-story home described as “more than adequate” for his family.
- Has many friends and enjoys spending time with groups. Dates, but has no steady girlfriend.
- Does well in school and has been accepted to the state university. No major yet chosen.
- Not currently sexually active.

PAGE 285: CRITICAL THINKING ACTIVITY #1

John’s history identifies several conditions or behaviors that might place him at risk for injury to this system:

- Sports activity, even with protective gear, can result in major injury to the head, neck, and specific structures of the face.
- Work in a public setting and school attendance provide exposure to various infectious agents.
- Use of OTC and prescribed antihistamines and occasional alcohol use puts patient at risk for traumatic injury while driving.
- Recurrent allergic episodes increase the risk of injury to the upper airway structures and infection.
- Although his history indicates he is not currently sexually active, the potential for sexual activity in adolescent males places John at risk for exposures to STDs, which affect the head and neck.
- Plans to participate in diving activities, which increase risk for injury of decompression illness, as well as barotrauma to the sinus or middle ear.

- The absence of a family history makes a comprehensive risk assessment incomplete.

PAGE 285: CRITICAL THINKING ACTIVITY #2

John's history includes several areas of potential strength. These include:

- Wearing protective equipment during athletic activities.
- Awareness of sexual activity–related risks.
- Healthy dietary practices.
- Daily physical activity.
- Adequate rest habits.
- Obtaining necessary healthcare and immunizations in the past.
- Desire to maximize his health.
- Adequacy of family financial resources.

PAGE 314: KEY HISTORY FINDINGS

- Current “good health” in “health-directed” young man.
- Frequent exposure to potential injury, with protection, during sports participation.
- History of seasonal “hay fever,” treated with antihistamines.
- Rare allergy symptoms, including congestion, sneezing, sleep disturbance.
- Up-to-date immunization status.
- Unknown family history.
- Exposure to second-hand smoke.
- Occasional alcohol intake.
- Desires clearance for deep-sea diving class.

PAGE 314: KEY PHYSICAL ASSESSMENT FINDINGS

- Mildly pale nasal mucosa, with slightly enlarged turbinates

PAGE 314: CRITICAL THINKING ACTIVITY #3

Strengths

- Healthful outlook.
- Positive behaviors (e.g., healthy diet, rest, activity, protection).
- Health-seeking activities.
- Strong adoptive family.
- Ability to learn and understand instructions and guidance.

Areas of Concern

- Potential for sexual and casual exposures to infectious diseases.
- Chronic, recurrent allergic symptoms that cause intermittent distress.
- Risks associated with the various athletic and recreational activities.

- Absence of biological family history.
- Chronic changes to nasal mucosa and turbinates.
- Occasional use of medications and alcohol that may alter mental status and safety.

PAGE 314: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Verbalizes interest to maximize health; reports good health; practices health-directed behaviors with diet and participation in regular physical activity; responsibly maintains up-to-date immunizations; avoids exposures to tobacco smoke as able; takes daily vitamin; made appointment for proper clearance before new activity.
- Participates in team sports and recreation associated with some risks (baseball, future diving).
- 17-year-old driver; uses substances that can affect mental status (occasional alcohol, although no indications of drinking while driving, and antihistamines).
- Describes difficulty sleeping after OTC antihistamines and frequent awakening during episodes of congestion/sinus pressure.

PAGE 314: ADDITIONAL NURSING DIAGNOSES

- Activity Intolerance, related to discomfort, respiratory impairment
- Nutrition: imbalanced, less than body requirements, related to poor dentition, oral pain, or difficulty swallowing
- Fluid Volume, deficient, related to difficulty swallowing, oral pain
- Swallowing, impaired, related to inflammatory throat pain
- Body Image, disturbed, related to real or perceived disfigurement of the head, face, or neck
- Social Isolation, related to altered physical appearance
- Oral Mucous Membrane, impaired, related to infection, ulcerations, dehydration, effects of medications, or inflammation
- Pain, acute, related to inflammation of the mucous membranes
- Airway Clearance, ineffective, related to swelling of the mucous membranes and nasal congestion
- Sensory Perception, disturbed (olfactory), related to swelling of nasal mucosa

PAGE 319: REVIEW QUESTIONS

1. a. Only the ethmoid and maxillary sinuses are developed at birth; the other sinuses develop over the next 7 years.
2. a. There are 20 deciduous (baby) teeth and 32 permanent teeth.
3. a. The nasolabial fold, the crease between the corner of the nose and the edge of the lip (the smile crease) is an ideal place to assess for facial symmetry.

4. c. The trapezius and sternocleidomastoid muscles make up the anterior and posterior triangle.
5. d. The occipital lymph nodes are located at the base of the skull.
6. c. The submental nodes are located under the tip of the chin.
7. b. The tonsillar nodes are located at the angle of the jaw.
8. c. The posterior lymph nodes are located in the posterior triangle behind the trapezius muscle.
9. c. The isthmus of the thyroid is located below the cricoid cartilage.
10. c. To facilitate palpation of the thyroid gland have the patient swallow. As the patient swallows, the gland rises up and may be palpable if enlarged.
11. a. The parotid glands are located in front of the ears.

CHAPTER 12

Eye

PAGE 331: CRITICAL THINKING ACTIVITY #1

Considering Mrs. Scammel's age of 55, you might expect to observe greater laxity of the eyelids and periorbital tissues and pupils that are smaller at rest than a younger adult's and are slower to react to light and accommodation. Also, she may be less likely to focus clearly on near objects, and the general background of her eyes may be paler.

PAGE 333: CRITICAL THINKING ACTIVITY #2

In chronic HTN, the ophthalmic examination of the fundus may reveal papilledema or blurred, nondistinct margins of the optic disk, alteration in A/V ratio, and cotton-wool spots on the general background. In DM, eye changes may include diminished visual acuity, diminished peripheral vision, and microhemorrhages on the fundus.

PAGE 337: CASE STUDY EVALUATION (MRS. SCAMMEL)

Biographical Data

- 55-year-old Caucasian widow and mother of one child, age 30.
- Works full-time as a secretary at a large law firm.
- Has major medical insurance coverage through her employer. Does not have coverage for dental or vision examinations.
- Catholic, of Irish Italian descent.

Current Health Status

- Complains of blurred vision that has increased over the past month. Blurred vision is constant and worsened by fatigue. States that nothing seems to cause

the problem or has helped relieve it. Wonders if she needs new glasses.

- Typically works at a computer for at least 4 hours each day.
- Blurred vision has not prevented her from caring for herself at home; however, it is difficult to read the material she is typing on the computer screen, even with reading glasses.
- Denies tearing, ear pain, drainage, double vision, sensitivity to light, or changes in the appearance of the eyes. Currently takes Lasix 20 mg bid, Captopril 25 mg bid, Glucotrol 5 mg bid, and 400 to 600 mg of ibuprofen as needed for joint pain nearly every day.

Past Health History

- History of HTN for past 10 years and poorly controlled adult-onset DM diagnosed 3 years ago.
- Denies trauma or surgery affecting the eyes or hospitalizations except for childbirth. One child, no abortions.
- Denies allergies to foods, drugs, or environment.
- Last physical exam 10 months ago; last eye exam 5 years ago. Mammogram and Pap smear negative.

Family History

- Has one child, age 30, alive and well.
- Only sibling, a brother, died 2 years ago at age 50 of a MI.
- Father had history of colon cancer and died of heart failure at age 56.
- Mother, age 68, lives in nursing home and has history of non-insulin-dependent (type 2) DM, multiple sclerosis, and stroke 4 years ago.
- No family history of glaucoma or vision loss.

Review of Systems

- **General Health Survey:** States that she is usually in good health.
- **Integumentary:** No skin lesions or allergies.
- **HEENT:** No eye tearing or drainage. No history of headache or head injury. No symptoms of URI.
- **Respiratory:** No history of respiratory disease.
- **Cardiovascular:** No history of cardiac symptoms; positive history of HTN.
- **Gastrointestinal:** No history of liver or kidney disease.
- **Musculoskeletal:** No joint pain or deformity or rheumatoid arthritis.
- **Neurological:** No muscle weakness, parasthesia, multiple sclerosis, or myasthenia gravis.
- **Endocrine:** Positive history type 2 DM; no history of thyroid disease.
- **Lymphatic:** No history of HIV, chemotherapy, or immunotherapy.

Psychosocial Profile

- Last physical exam 10 months ago; last eye exam 5 years ago. Gets yearly mammograms and Pap tests.
- Typical day includes: Rises at 6:30 A.M., showers and dresses, has breakfast (bagel and coffee), leaves for work at 8 A.M. Works 8:30 A.M. to 5 P.M. Monday through Friday. Eats lunch (sandwich) at diner most days. Usually returns home by 6 P.M. unless she runs errands or shops for groceries. Rarely cooks a full dinner unless her son visits. Watches TV or reads after dinner. Goes to bed at 11 P.M. Spends weekends gardening in warm weather, visiting mother in nursing home, visiting son, running errands, doing housework. Goes to movies with friends occasionally.
- Admits difficulty following prescribed diabetic and low-sodium diets. Tries to eat regularly but has a limited appetite and finds it difficult to cook balanced meals for one person.
- Main source of exercise is gardening and yard work.
- Since loss of her husband, has tried to become more involved in her favorite hobbies: reading, watching TV, and gardening.
- Generally sleeps 8 hours a day without napping.
- Drinks socially three to four times a year. Smoked 1 pack of cigarettes a day since age 18 until 10 years ago when diagnosed with high BP.
- Was a full-time homemaker during her 22-year-marriage, but returned to workforce after husband died of a heart attack 2 years ago. Now works full time as a secretary, with occasional overtime. Reports using computers for an average of 4 hours a day.
- Lives in a rural community just outside of town, in her own home. Commutes 30 minutes to and from work.
- Sees son once a week (lives in a neighboring town). Visits mother in nursing home once a week.
- Is Catholic, but is not currently active in church. Denies any specific ethnic or cultural practices.
- Main social supports are friends from neighborhood and son and daughter-in-law.
- Stress and coping mechanisms include gardening and talking over problems with son and longtime friends from neighborhood.

PAGE 338: CRITICAL THINKING ACTIVITY #3

Mrs. Scammel's visual problems may affect her ability to function safely and independently at home and at work. Diminished visual acuity and peripheral vision may affect her ability to drive safely to and from work, so she might limit her driving to only essential trips, which decreases her sense of independence. Diminished visual acuity may also affect her ability to engage in her favorite pastimes, such as reading and watching TV, and may make performing her secretarial duties more difficult.

PAGE 338: CRITICAL THINKING ACTIVITY #4

Mrs. Scammel may need to make modifications at work that will enable her to function optimally. For example, she might use magnifying devices, print items to be proofed in large or bold print, and take frequent breaks from close-range computer work to reduce eyestrain. She will also need more frequent eye exams and close monitoring of blood sugar levels and BP.

PAGE 338: CRITICAL THINKING ACTIVITY #5

Mrs. Scammel's strengths include her desire to be active and independent, as evident in her returning to work after her husband's death. She has a variety of interests outside of work and generally engages in healthy habits of getting enough sleep, not smoking, and drinking in moderation only. When confronted with health problems that can benefit from a change in lifestyle, she is usually successful in changing. She was successful in quitting smoking and tries to avoid foods that would worsen her HTN and DM.

PAGE 338: CRITICAL THINKING ACTIVITY #6

Both of Mrs. Scammel's medical diagnoses of HTN and DM have effects on the eyes. Her last eye exam was 5 years ago. During the interim, she may have experienced some irreversible damage to the eyes. Also, her past smoking history may have narrowed or damaged the vasculature of the eyes.

PAGE 355: CRITICAL THINKING ACTIVITY #7

The bulbar conjunctiva is clear and covers the sclera, which is white. When the conjunctiva becomes inflamed during an infection such as conjunctivitis, the blood vessels that normally are not visible become irritated, more prominent, and more visible against the white sclera.

PAGE 360: KEY HISTORY FINDINGS

- Age 55.
- Overweight.
- Complaints of blurred vision.
- History of HTN and non-insulin-dependent DM.
- Insurance does not cover eye exams.
- Wears corrective lenses.
- Works at computer 4 hours a day.
- Positive family history of DM, MS, cardiovascular disease, and cerebrovascular disease.

PAGE 360: KEY PHYSICAL EXAMINATION FINDINGS

- Overweight.
- BP 170/108.
- Near vision blurred.
- Decreased peripheral vision in temporal fields.
- Hazy cornea OD.
- Arcus senilis.

- Abnormal fundoscopic findings (nicking, blurred disc margins, and scattered dot hemorrhages).

PAGE 360: NURSING DIAGNOSES SUPPORTING DATA

- Sensory Perception, disturbed (visual): Complaint of blurred vision; near vision blurred; decreased peripheral vision in temporal fields; hazy cornea; fundoscopic changes
- Injury, risk for: Complaint of blurred vision; near vision blurred; decreased peripheral vision in temporal fields; fundoscopic changes; drives to work 30 minutes each way
- Social Interaction, impaired: Visual changes; widowed; lives alone

PAGE 360: ADDITIONAL NURSING DIAGNOSES

- Knowledge, deficient (management of DM, HTN)
- Health Maintenance, ineffective

PAGE 364: REVIEW QUESTIONS (EYE)

- a. Presbyopia is an age-related vision change characterized by difficulty seeing near.
- d. A sign of digitoxin toxicity includes diplopia and altered color visions, yellow or green spots.
- a. 20/10 OU means that the patient can see at 20 feet what a person with normal vision can see at 10 feet. 20/10 vision.
- a. The corneal light reflex is a test of extraocular membrane. Shine a penlight directly on the eyes and look for the light reflection on the cornea in each eye.
- b. 20/20 vision is usually achieved by school-age. A 3-year-old child's far vision is normally about 20/40.
- b. The normal response of the cornea reflex is for the client to blink and there may be some tearing. The corneal reflex tests CNs V and VII.
- a. To test accommodation, have the patient follow your finger from afar as you move it closer. The normal response to accommodation is convergence and constriction of the pupils.
- b. The eyes work in conjugate movement, what one eye is doing, normally so should the other eye. If the eye receiving the light stimulus normally constricts, the opposite eye should also constrict. This is referred to as a *consensual reaction*.
- c. The nasolacrimal glands are located below the eyebrows and drain through the lacrimal ducts. The lacrimal ducts drain in the inner canthus of the eyes.
- a. When performing an ophthalmoscopic exam on a patient whose pupils are undilated, the smallest white light aperture should be used. The small white light aperture will cause minimal constriction of the pupil.

Ear

PAGE 375: CASE STUDY EVALUATION (MR. ARNEZ)

Biographical Data

- Married, 42-year-old father of two (ages 14 and 16).
- Full-time construction supervisor with bachelor's degree.
- Family's main source of financial support; wife is disabled.
- Carries HMO insurance through his employer that covers entire family.
- Native American and Spanish descent.
- Catholic religion.
- Source: Patient, who appears reliable.

Current Health Status

- Chief complaint is intermittent fullness and pressure in both ears for past 2 weeks.
- Denies precipitating factors, such as recent URI or recent altitude changes.
- Related symptoms include intermittent, bilateral hearing loss described as "like hearing in a tunnel" and intermittent dizziness.
- Rates severity of his ear fullness and pressure as 7 on a 10-point scale.
- Denies ear pain, drainage, or recent trauma to the ears.

Past Health History

- History of frequent ear infections as an infant and young child. Treated with antibiotics.
- Hospitalized once at age 3 with pneumonia and concurrent bilateral otitis media.
- Denies asthma, DM, or HTN.
- Received polio, MMR, DPT vaccinations.
- Allergic to sulfa, penicillin, dust, and pollen.
- Current medications include an OTC allergy medication.

Family History

- Father, age 68, has controlled HTN.
- Mother, age 60, diagnosed with multiple sclerosis at age 45, currently in remission.
- One sister, age 40, diagnosed with neurofibromatosis with acoustic neuroma.
- One brother, age 36, recently diagnosed with HTN.
- Patient has two teen-aged girls, ages 14 and 16, both alive and well.

Review of Systems

- **General Health Survey:** Denies fatigue or recent weight loss. Health is usually good.
- **Integumentary:** Denies rashes, lesions, or slow wound healing.

- **HEENT:** Denies ear pain, drainage, tinnitus, or ear trauma. Reports recent URI 4 weeks ago, with ear-ache, resolved with antibiotics. History of frequent childhood ear infections and occasional ear infections as an adult, occurring yearly or less often. Environmental allergy to dust and pollen.
- **Gastrointestinal:** Denies nausea, vomiting, diarrhea, or recent changes in bowel habits.
- **Genitourinary:** Denies changes in bladder habits.
- **Neurological/Musculoskeletal:** Intermittent problem with balance since onset of ear pressure. Denies vertigo, muscle weakness, or paresthesia.

Psychosocial Profile

- Last physical examination was 6 months ago. Last ear exam was 4 weeks ago during recent ear infection. Denies ever having a hearing test.
- Typical day: Rises at 5:30 A.M. on workdays (works 7–3 shift). Showers, has breakfast (cereal and coffee), leaves for work by 6:15 A.M. Eats quick lunch in hospital cafeteria. Returns home by 4 P.M. most days unless he has errands to run. Cooks dinner for wife and two daughters, then often drives daughters to various school or social activities. Spends evenings watching TV. Goes to bed by 10 P.M. Spends weekends attending sporting events, running errands, chauffeuring daughters. (Wife cannot drive because of disability.)
- Ear pressure and fullness have caused intermittent problems in work environment and socializing. States that he can usually compensate for the hearing loss at home.
- States that hearing loss causes difficulties at work.
- Denies chronic or recent exposure to loud noise at home or at work.

PAGE 375: CRITICAL THINKING ACTIVITY #9

Factors in Mr. Arnez's history that may explain his hearing problem include a history of recurrent ear infections in childhood and as an adult and a family history of acoustic neuroma and multiple sclerosis.

PAGE 376: CRITICAL THINKING ACTIVITY #10

Hearing loss may make Mr. Arnez less confident in work and in social situations in which verbal communication is emphasized. Intermittent dizziness may make him fearful of being injured and more reluctant to be active or drive a car.

PAGE 376: CRITICAL THINKING ACTIVITY #11

Strengths that Mr. Arnez possesses include independence and being the sole financial support for his family. His level of education will facilitate understanding of his medical condition and its treatment. He also has insurance coverage and family members nearby for emotional support.

PAGE 385: KEY HISTORY FINDINGS

- 42-year-old construction supervisor, sole support of family.
- Complains of intermittent fullness and pressure in ears for past 2 weeks, severity 7/10, dizziness; describes hearing as “hearing in a tunnel” and reports hearing loss.
- Recent URI.
- Frequent ear infections as a child.
- Positive family history of HTN, multiple sclerosis, and acoustic neuroma.
- Positive environmental allergies.
- Hearing problems have affected work and socialization.

PAGE 385: KEY PHYSICAL EXAMINATION FINDINGS

- Scarring of tympanic membrane.
- Rinne test AC < 2 x BC.
- Dizziness with head rotation.
- Positive Romberg's test.

PAGE 385: NURSING DIAGNOSES SUPPORTING DATA

- Reports changes in hearing like “hearing in a tunnel.” Difficulty hearing when auscultating.
- Rinne test AC < 2 x BC. Scarring on tympanic membrane.
- Complains of pressure and fullness in ears, severity 7/10.
- Recent URI; fullness and pressure in ears.

PAGE 385: ADDITIONAL NURSING DIAGNOSES

- Diversional Activity, deficient
- Communication, impaired verbal
- Social Interaction, impaired
- Injury, risk for

PAGE 390: REVIEW QUESTIONS (EAR)

1. a. Presbycusis is a hearing loss associated with aging. Presbycusis involves hearing loss of high-pitched sounds.
2. b. Tinnitus, ringing in the ears, is a sign of aspirin toxicity.
3. a. The ears should be in alignment with the outer canthus of the eyes. Low-set ears in a newborn may indicate Down syndrome.
4. a. The helix, tragus, and mastoid should be palpated for tenderness prior to insertion of the otoscope. If tenderness is present, proceed carefully.
5. a. The Weber test is a hearing test for lateralization of sound. Normally the sound should be heard equally in both ears.
6. b. The Rinne test compares air conduction with bone conduction. Normally, sound transmission is

- twice as long through air as through bone. If bone conduction is greater than air conduction, the patient may have a conductive hearing loss with a problem in the external or middle ear.
7. b. The shape of the ear canal assumes normal position by school age. When performing an otoscopic exam on a 2-year-old child, the ear is pulled down to straighten the canal. To straighten an adult ear canal, the helix is pulled up and back.
 8. b. The length of the ear canal of an adult is about 1 inch. The otoscope should be inserted about 1/2 inch. The inner two-thirds of the canal is over the temporal bone and is very sensitive. Care should be taken to avoid inserting the scope in too far for it would be uncomfortable for the patient.
 9. c. The color of a healthy tympanic membrane is pearly gray.
 10. b. Cerumen, ear wax, is the only normal drainage from the ear canal. Cerumen lubricates the ear canal and helps remove foreign objects.

CHAPTER 13

PAGE 408: CASE STUDY EVALUATION

Biographical Data

- 38-year-old Caucasian female.
- Married, mother of three children, ages 10, 8, and 4.
- Part-time accountant; BA degree in accounting.
- Born and raised in United States, Baptist religion.
- Healthcare insurance (HMO) through husband's work.
- Referral: Preadmission workup for surgery.
- Source: Self, seems reliable.

Current Health Status

- Brief morning cough for past 5 years, productive of small amount of thin, whitish-gray mucus, no odor or bad taste. Cough does not occur with activity or exercise. Positive history of smoking.
- No dyspnea or chest pain.
- No edema, loss of usual energy, change in sleeping patterns (except when gallbladder discomfort awakens her).

Past Medical History

- Frequent URIs as a child.
- No history of asthma, pneumonia, or TB.
- Never had surgery. Hospitalized only for childbirth: uncomplicated vaginal delivery of three children.
- No known exposure to people with respiratory infections.
- Denies allergies to food, drugs, or environmental factors.

- No medical problems except for gallbladder disease.
- Immunizations up to date. Last PPD a year ago was negative.
- Denies taking prescribed drugs. Takes acetaminophen about twice a month for headache relief.

Family History

- Grandfather smoked and died of lung cancer at age 63.
- Father, age 65, recently quit after smoking a pack a day for 40 years (40 pack-years). Diagnosed with lung cancer a year ago.
- All other known relatives alive and well.

Review of Systems

- **General Health Survey:** States that usual health is good. Has gained 15 lb in past 2 years.
- **Integumentary:** No changes in skin, hair, or nails.
- **HEENT:** *Head/Neck:* No swelling or masses.
Eyes/Ears: Vision 20/20; hearing good. *Nose/Throat:* Productive morning cough with whitish-gray mucus. Has occasional cold.
- **Cardiovascular:** No chest pain.
- **Gastrointestinal:** Appetite good, no nausea or vomiting. Daily bowel movement.
- **Urinary:** No changes reported.
- **Reproductive:** Is in a monogamous relationship; satisfied with sexual performance.
- **Musculoskeletal:** No weakness or muscle wasting reported.
- **Neurological:** Memory good. No changes noted. No tremors.
- **Immunologic/Hematologic:** No allergies or anemia.

Psychosocial Profile

- Has a physical exam and Pap smear every 2 years. Made a special appointment when she noticed RUQ colicky abdominal pain after eating high-fat meals. Scheduled surgery as soon as recommended by physician.
- Typical day consists of arising at 6 A.M., having breakfast with family, and getting children off to school. Walks 5 blocks to and from work at CPA office 4 days a week. Works 8 A.M. to 4 P.M. at desk with computer. Spends evenings playing with children, helping them with homework, shopping, fixing dinner, doing housework, visiting with husband, watching TV, or reading. Showers, then goes to bed at 11 P.M.
- Usually eats a healthy diet, but has gained about 15 lb over the past 2 years. Attributes this to sedentary job, "sweet tooth," and lack of exercise.
- Walks to and from work and likes to ride a bike, but does not have much time because of family, home, and work responsibilities.

- No pets; hobbies include reading and biking.
- Sleeps about 6 to 7 hours a night, uninterrupted, with one pillow. Denies problems falling asleep, but awakens with morning cough.
- Has smoked 1 pack a day of filtered cigarettes for past 10 years (10 pack-years). Lived with smokers all her life. Has tried to quit by cutting back without success. Drinks alcohol about three times a month at social events. Denies recreational drug use.
- Denies respiratory irritants at work.
- Lives in two-story, single-family home in suburbs with ample living space for five people; has hot water heating and air conditioning; home negative for radon gas; denies recent travel outside local community. Will stay in downstairs bedroom for the first week after surgery.
- Supportive, caring husband; good relationship with children. Concerned about father's prognosis and mother's ability to care for him. Aware of health risks of second-hand smoke for children, so tries not to smoke at home. Husband also smokes, both would like to "kick the habit." No community involvement at this time, but has several friends who will help out after surgery.
- Copes with stress by talking with husband or friends or by ignoring it; also by smoking.

PAGE 408: CRITICAL THINKING ACTIVITY #1

History of smoking and type of surgery place her at risk for altered respiratory function, hypoventilation, and a potential complication, atelectasis.

PAGE 408: CRITICAL THINKING ACTIVITY #2

Strengths are understanding risks associated with smoking, willingness to stop smoking, supportive family, health-seeking behaviors.

PAGE 408: CRITICAL THINKING ACTIVITY #3

Encourage coughing and deep breathing, control pain, early ambulation, and adequate hydration.

PAGE 410: CRITICAL THINKING ACTIVITY #4

Because the gallbladder is located in the RUQ. So postoperatively, Mrs. Kane may tend to hypoventilate because taking deep breaths may increase pain.

PAGE 431: KEY HISTORY FINDINGS

- Brief morning cough for past 5 years, productive of small amount of thin, whitish-gray mucus.
- Frequent URIs as a child.
- Family history of lung cancer (grandfather and father).
- Has smoked 1 pack a day of filtered cigarettes for past 10 years (10 pack-years).
- Family history of cigarette smoking and long-term exposure to second-hand smoke.

- Wants to stop smoking. Has tried to quit by cutting back without success. Accepts difficulty of quitting smoking.
- Understands potential health effects of smoking on self and family.
- Sedentary occupation, walks to and from work.
- 15-lb weight gain over 2 years.
- RUQ pain from gallbladder disease. Understands potential for postsurgical discomfort.

PAGE 431: KEY PHYSICAL EXAMINATION FINDINGS

- Skin pink, warm and dry, good turgor; mucous membranes pink and moist.
- Nails pink, firm, capillary refill < 3 seconds, nicotine stains on fingers and nails, no clubbing.
- Trachea midline.
- Chest: AP:lateral 1:2, costal angle < 90°, equal chest excursion, normal spinal curvatures, skin intact, no use of accessory muscles, no retraction.
- Chest nontender, symmetrical chest excursion, no masses, crepitus.
- Tactile fremitus equal anterior and posterior to mid thorax.
- Resonance noted throughout lung fields. Diaphragmatic excursion 4 cm bilaterally.
- Lungs CTA except for expiratory wheezing that clears with coughing; no crackles or rubs.

PAGE 432: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Scheduled for cholecystectomy. States that she is frightened. Asking questions about pre- and postoperative course.
- RUQ pain from gallbladder disease. Expiratory wheeze that clears with coughing. 10-pack-year smoking history.
- Brief morning cough for past 5 years productive of small amount of thin, whitish-gray mucus. Unsuccessful at attempting to stop smoking.
- Desires to stop smoking; previously unsuccessful attempts. Walks to and from work. Aware of effects of second-hand smoke on children.
- Frequent URIs as a child. Family history positive for lung cancer. Grandfather smoked and died of lung cancer at age 63. Father, age 65, recently quit after smoking 1 pack a day for 40 years (40 pack-years); diagnosed with lung cancer a year ago.
- Weight gain of 15 lb over past 2 years. Sedentary occupation.

PAGE 432: ADDITIONAL NURSING DIAGNOSES

- Infection, risk for
- Activity Intolerance
- Fatigue
- Breathing Pattern, ineffective

- Pain, self-management
- Anxiety

PAGE 432: COLLABORATIVE NURSING DIAGNOSES

- Potential complication: atelectasis.
- Scheduled for cholecystectomy with general anesthesia. Smoking history.

PAGE 437: REVIEW QUESTIONS

1. a. Vesicular breath sounds are heard in most of the lung fields. Vesicular sounds are low-pitched, soft sounds with a long inspiratory phase and shorter expiratory phase.
2. b. Bronchovesicular sounds are heard over the manubrium. Bronchovesicular sounds are medium in pitch and intensity with equal inspiratory and expiratory phases.
3. c. Bronchial breath sounds are normally heard over the trachea. Tracheal breath sounds are high-pitched, loud sounds with a short inspiratory phase and longer expiratory phase.
4. b. Rhonchi are abnormal, added sounds caused by a narrowing of the airways with mucus occurring predominantly on expiration in the larger airways.
5. d. Clubbing is a sign of long-standing cardiopulmonary problems. The normal angle of attachment of the nail is 160 degrees. Clubbing is present if the angle of attachment is 180 degrees or greater.
6. a. The normal shape of an adult chest is oval with the AP:lateral ratio is 1:2. Patients with COPD have a 1:1 AP: lateral ratio due to overinflation of the lung.
7. b. The normal percussion sound over healthy lung tissue is resonance, a low-pitched, loud, long sound. Hyperresonance is the extreme of resonance, very low-pitched, very loud, and very long in duration, and may be heard in patients with COPD.
8. b. Tactile fremitus is useful in assessing airflow. Since patients with COPD have problems moving air, fremitus is decreased.
9. b. Breath sounds are produced as a person moves air throughout his airways. Since patients with COPD have problems moving air, breath sounds are usually decreased at the bases.
10. d. Color changes in dark-skinned patients cannot readily be seen. Color changes are more readily seen in the mucous membranes.

CHAPTER 14

PAGE 459: CRITICAL THINKING ACTIVITY #1

Risk Factors

- Age
- Male gender
- Family history

- Newly diagnosed HTN
- Diet; 60 lb overweight
- Sedentary lifestyle
- Work schedule
- Lack of exercise
- Alcohol intake
- Stress

Modifiable

- HTN controllable
- Diet
- Weight
- Lack of exercise
- Sedentary lifestyle
- Alcohol intake
- Work schedule
- Stress

Unmodifiable

- Age
- Male gender
- Family history

PAGE 459: CASE STUDY EVALUATION

Biographical Data

- 68-year-old Caucasian male, married, father of seven grown children, self-employed entrepreneur. Born and raised in the United States, Italian descent.
- Catholic religion.
- BS in engineering.
- Blue Cross/Blue Shield medical insurance plan.
- Referral: Follow-up by PCP.
- Source: Self, reliable.

Current Health Status

- Denies chest pain, dyspnea, palpitations, or edema.
- Reports fatigue, loss of energy, and occasional dizzy spells.

Past Health History

- No history of rheumatic fever in childhood.
- No history of murmurs.
- No history of injuries.
- Inguinal hernia repair.
- ECG revealed LVH.
- Hospitalized 3 weeks ago for HTN.
- No known food, drug, or environmental allergies.
- Immunizations up to date.
- No prescribed medications except Vasotec (enalapril) 5 mg bid and weekly use of antacid for indigestion.

Family History

- Positive for HTN and stroke. Mother had HTN and died at age 78 from stroke. Paternal uncle died at age 79 from MI.

Review of Systems

- **General Health Survey:** Reports fatigue, weight gain of 60 lb over past 3 years.
- **Integumentary:** Reports feet cold, thick nails, tight shoes.
- **HEENT:** Reports two dizzy spells over past 6 months. *Eyes:* Wears glasses, no visual complaints, yearly eye exam.
- **Respiratory:** Reports being “short-winded” with activity.
- **Gastrointestinal:** Indigestion on weekly basis.
- **Genitourinary:** Awakes at least once a night to go to bathroom.
- **Male Reproductive:** Little sexual activity because he is “too tired;” also states that it is becoming difficult to have an erection.
- **Musculoskeletal/Neurological:** General weakness, cramps in legs on walking.

Psychosocial Profile

- Typical day starts at 7 A.M., when patient showers, has breakfast, then goes to work. Returns home by 6 P.M., eats dinner, watches TV until 11:30 P.M., but usually falls asleep before news is over. Usually in bed by 12:00 midnight. Usually works 7 days a week.
- 24-hour recall reveals a diet high in carbohydrates and fats and lacking in fruits and vegetables. Heavy-handed with salt shaker. Admits he has gained weight over the years and is 60 lb overweight.
- No exercise program; states that he is “too busy running my business.”
- Sleeps about 7 hours a night but usually feels he is not getting enough sleep, lately is more and more tired. Wife states that he snores.
- Does not have routine checkups. States, “I only go to the doctor’s when I’m sick.”
- Has never smoked. Drinks a bottle of wine every night with dinner.
- Hobbies include reading, crossword puzzles, and antique collecting.
- Has a large, close, caring family. Feels that running his own business can be very stressful, but feels he can handle it alone and doesn’t need anyone to help him.
- Lives with wife of 45 years in a two-story, single home in suburbs with ample living space.

PAGE 487: CRITICAL THINKING ACTIVITY #2

HTN increases the work of the heart so that it has to pump against greater resistance. This could lead to LVH.

PAGE 487: CRITICAL THINKING ACTIVITY #3

Left-sided CHF can result from LVH that developed secondary to HTN. Clinical signs of left CHF include breathing difficulties, such as SOB, DOE, or PND.

PAGE 487: CRITICAL THINKING ACTIVITY #4

The carotid pulsation is normally palpable because the arterial system is a high-pressure system. The venous system is a low-pressure system, so although the jugular venous pulsation may be visible, it is not palpable. The jugular venous pulse is obliterated with palpation.

PAGE 487: LIST KEY HISTORY FINDINGS

- 68-year-old Caucasian male.
- Positive fatigue and dizzy spells.
- Newly diagnosed with HTN.
- Vasotec (enalapril) 5 mg bid.
- Use of antacids weekly.
- Positive family history of HTN, stroke, MI.
- Diet high in fats, carbohydrates, and salt.
- Weight gain, 60 lb overweight.
- Sedentary lifestyle.
- Lack of routine activity or exercise program.
- SOB with activity.
- Changes in skin.
- Intermittent claudication.
- Indigestion weekly.
- Change in sexual activity.
- Lack of restful sleep, snoring, nocturia.
- Lack of health promotion and prevention activities.
- Alcohol intake.
- Large, close, caring family.

PAGE 487: LIST KEY PHYSICAL EXAMINATION FINDINGS

- BP 150/90.
- Arcus senilis, AV nicking, and cotton-wool spots.
- +1 pedal pulses; skin cool, pale, dry, thin, and hairless with trace of edema.
- +3 carotids.
- Sustained pulsation displaced lateral to apex, PMI 3 cm.
- Accentuated S₂, +S₄.

PAGE 487: CRITICAL THINKING ACTIVITY #5

Strengths

- Usual state of health good.
- BP is more stable.
- Supportive family.

Areas of Concern

- Newly diagnosed HTN.
- Family history of cardiovascular disease.
- Overweight.
- Unsatisfactory diet.
- Sedentary lifestyle.
- Stress.
- Inactivity, lack of routine exercise.
- Arcus senilis and fundoscopic changes.

- +1 pedal pulses with skin changes.
- 3-cm PMI with displacement.
- S₄ and accentuated S₂.

PAGE 488: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Positive pedal pulses; skin and nail changes in extremities; BP.
- 60 lb overweight; diet high in fats and carbohydrates; high salt intake.
- HTN, ECG with LVH; S₄; enlarged and displaced PMI.
- Newly diagnosed with HTN.

PAGE 488: ADDITIONAL NURSING DIAGNOSES

- Health Maintenance, ineffective, related to lack of knowledge regarding risk factors for heart disease
- Activity Intolerance, related to fatigue secondary to cardiovascular disease
- Coping, ineffective, related to stress of work
- Sleep pattern, disturbed, related to snoring (sleep apnea)
- Sexuality patterns, ineffective, related to activity intolerance

PAGE 490: REVIEW QUESTIONS

1. b. Chest pain of cardiac origin is not affected by palpation of the chest. Chest pain associated with movement or palpation is often musculoskeletal in nature.
2. c. There are many possible causes of chest pain. Since chest pain of cardiac nature may be life-threatening, it must always be considered first.
3. c. The PMI, point of maximum impulse, also referred to as the LVI, left ventricular impulse, is normally located at the apex, fifth intercostal space at the left mid-clavicular line.
4. a. A thrill is a vibratory sensation, created by turbulent blood flow. If a thrill is present on exam of the precordium, a murmur will be heard on auscultation.
5. d. An opening snap is an abnormal, high-pitched, early diastolic sound heard best at the apex with the diaphragm portion of the stethoscope. Opening of the valves is not usually heard, but if the valves are diseased, the sound opening of the mitral or tricuspid valves may be heard.
6. d. An S₄ is a late diastolic filling sound that occurs when the atria contracts against resistance. An S₄ is an extra diastolic, low-pitched sound heard best at the apex with the bell portion of the stethoscope in such conditions as HTN, CAD.
7. c. An S₃ is an early diastolic sound; a sound of a distressed heart, in such conditions as CHF. An S₃ is an extra diastolic, low-pitched sound heard best

at the apex with the bell portion of the stethoscope.

8. a. A carotid thrill is a palpable vibration caused by a narrowing of the carotid artery. If a carotid thrill is present, a bruit may be heard when auscultating the carotid.
9. a. The jugular venous system is a low-pressure system. Normally, the jugular veins are visible only when the patient is in the supine position.
10. b. Measurements of the jugular venous pressure should be obtained with the patient's head of the bed at a 30- to 45-degree angle.

CHAPTER 15

PAGE 504: CASE STUDY EVALUATION

Biographical Data

- 38-year-old African American female, married, one child.
- Works full-time as a secretary.
- Born and raised in United States, Baptist religion.
- High school graduate, interested in attending community college when her baby gets older.
- Healthcare insurance, PPO, through husband's work plan.
- Husband is firefighter.
- Referral: Private practice, internal medicine office.
- Source: Self, seems nervous but reliable.

Current Health Status

- 6-day history of dull, aching pain and tenderness in left leg that is progressively getting worse.
- Pain is worse when standing and walking and is minimally relieved by rest and elevation.
- Aspirin has not helped.
- Denies chest pain or SOB.

Past Medical History

- Delivered premature baby 1 week ago.
- Prolonged bedrest 2 months before delivery.
- No history of surgery. Hospitalized only for childbirth. Uncomplicated vaginal delivery; baby's gestational age is 31 weeks.
- No history of chronic illness.
- Denies prescribed drugs.
- Has smoked 1 pack of cigarettes a day for 22 years (22 pack-years).

Family History

- Mother, age 65, has HTN and is obese.
- Father, age 67, had MI at age 50; quit smoking 17 years ago.
- All other known relatives alive and well.

Review of Systems

- **General Health Survey:** General health good, feeling tired since delivery of baby.
- **Integumentary:** Inflammation and swelling of left calf.
- **HEENT:** *Head/neck:* No swelling or masses, no sore throat.
- **Respiratory:** No breathing problems.
- **Cardiovascular:** No chest pain.
- **Gastrointestinal:** No complaints, daily bowel movement.
- **Genitourinary:** No nocturia since delivery of baby.
- **Reproductive:** Has not been sexually active for about the past 3 months because of complications of pregnancy. No problems before this. Plans to bottle-feed baby.
- **Musculoskeletal:** Tenderness in left calf.
- **Neurological:** No numbness, tingling, or loss of sensation.
- **Endocrine:** No history of thyroid disease or DM.

Psychosocial Profile

- Has Pap smear by gynecologist every year. Last complete physical exam 5 years ago.
- Typical day consists of arising at 7 A.M., showering and dressing, and having breakfast with husband. Takes train to work. Works 8 A.M. to 5 P.M. 5 days a week as secretary, which involves mostly sitting at desk with computer. Spends evenings watching TV, reading, going out socially with husband. This routine was dramatically altered 2 months ago because of premature labor and bedrest.
- Usually eats lots of fast foods but has worked hard to maintain a healthy diet during her pregnancy. Gained 50 lb during pregnancy, which she attributes primarily to 2 months of bedrest.
- Likes to walk and continued this activity throughout pregnancy until she was put on bedrest. Before pregnancy, played tennis once a week with husband.
- Normally sleeps about 8 hours nightly without difficulty. During pregnancy, sleep was interrupted because of need to void.
- No pets. Hobbies include walking, tennis, sewing, and reading.
- Has smoked 1 pack a day of filtered cigarettes for past 22 years (22 pack-years). Has tried behavior modification, cold turkey, and nicotine patches without success.
- Drinks wine with dinner on weekends, but none since pregnancy. Denies recreational drug use.
- Sits for long hours in front of computer at work. Denies other occupational hazards.
- Lives in ranch home in suburbs with ample living space, air conditioning, and hot water heating. Denies travel outside of state.

- Has supportive, caring husband who worries about premature baby who will be coming home from hospital shortly. Visits baby daily. Concerned about patient's ability to care for baby with her leg problem. Would like her to quit smoking before baby comes home from hospital.
- Denies religious or cultural influences on health.
- Denies any sexual problems. Had sexual relations with husband until bedrest was ordered.
- Copes with stress by smoking and talking with husband and friends.

PAGE 504: CRITICAL THINKING ACTIVITY #1

- Circulatory problem: Thrombophlebitis.
- Contributing factors: Prolonged bedrest, pregnancy, smoking.

PAGE 504: CRITICAL THINKING ACTIVITY #2

Child care for newborn, parent-newborn relationship.

PAGE 504: KEY HISTORY FINDINGS

- 6-day history of dull, aching pain and tenderness in left leg that is progressively getting worse.
- Delivered premature baby 1 week ago. Uncomplicated vaginal delivery; baby's gestational age is 31 weeks.
- Prolonged bedrest 2 months before delivery.
- 50-lb weight gain with pregnancy.
- Sedentary occupation. Understands need to resume previous exercise regime.
- Concerned over new role as mother of premature child.
- 22-pack-year smoking history. Attempts to stop smoking unsuccessful in past. Understands potential health effects of smoking on self and family.

PAGE 518: KEY PHYSICAL EXAMINATION FINDINGS

- Positive Homans' sign.
- Left leg edematous and warm to touch, slightly cyanotic.
- Femoral and popliteal pulses strong; left posterior and pedal pulses difficult to locate.
- Left leg circumference, 41 cm; right leg circumference, 38 cm.
- Temperature, 99.8°F.

PAGE 518: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Describes pain as dull and achy with progressive worsening over 1 week. Pain gets worse with standing or walking and is only minimally relieved by rest and leg elevation. Positive Homan's sign.
- Appears anxious. States she is worried about having DVT and about how it will affect her ability to care

for her baby. Also worried about her new role as mother, especially with a premature baby. This is her first child.

- States that she is having difficulty walking and standing because of leg pain. Is aware that management of DVT requires bedrest and limb elevation.
- States that she wants to stop smoking despite previous unsuccessful attempts. Desire to stop smoking is enhanced by wanting a good environment for her child. Also wants to lose weight gained during pregnancy. Past behaviors included both exercise and balanced diet.
- Gained 50 lb with pregnancy. States that she wants to lose weight. Has sedentary occupation but exercised before complications of pregnancy. Is aware that medical management of DVT requires additional bedrest.

PAGE 518: ADDITIONAL NURSING DIAGNOSES

- Coping, ineffective, related to necessary lifestyle changes
- Family Processes, interrupted, related to loss of income and inability to work secondary to DVT and recent childbirth
- Mobility, impaired physical, related to intermittent claudication and fatigue
- Sleep Pattern, disturbed, related to rest pain
- Diversional Activity, deficient, related to prolonged immobility
- Injury, risk for, related to diminished arterial flow
- Skin Integrity, risk for impaired, related to decreased oxygenation of tissues secondary to vasospasm and arterial occlusion.
- Therapeutic regimen: ineffective management
- Noncompliance, related to failure to stop smoking cigarettes
- Tissue Integrity, impaired, caused by ineffective flow of fluid in the lymphatic vessels
- Body Image, disturbed, related to accumulation of fluid in the extremities
- Self-care deficit, related to limb enlargement secondary to lymphedema.

PAGE 518: COLLABORATIVE NURSING DIAGNOSES

- Potential complication: pulmonary embolus
- Risk for bleeding

PAGE 524: REVIEW QUESTIONS

1. b. The ulnar pulse is located on the lateral side of the wrist.
2. d. The temporal pulse is located on the lateral side of the forehead.
3. d. The popliteal pulse is located behind the knee.
4. b. The dorsalis pedis is located on the dorsum of the foot.
5. a. Normally leg pressures are higher than arm pressures. The ankle-brachial index is obtained by dividing the ankle systolic pressure by the brachial systolic pressure. If the ankle-brachial index is less than 1, it indicates vascular disease.
6. a. Prior to obtaining an arterial blood gas specimen from the wrist, the Allen test should be performed to ensure arterial patency of both the radial and the ulnar arteries.
7. d. The Color Change Test is a test for arterial insufficiency. Elevation pallor and dependent rubor are associated with arterial insufficiency.
8. a. The Homan's sign is a test for DVT. If dorsiflexion of the foot produces pain in the calf, the test is positive for DVT.
9. a. The epitrochlear nodes are located about one inch over the elbow,
10. b. The horizontal lymph nodes are located in the groin.

CHAPTER 16

PAGE 534: CRITICAL THINKING ACTIVITY #1

After ovulation, the breasts become engorged with blood and fluid because of the influence of progesterone. This causes them to become painful and/or tender.

PAGE 539: CASE STUDY EVALUATION

Biographical Data

- 46-year-old married woman.
- No pregnancies, no children.
- Caucasian, Catholic religion, Polish descent.
- College graduate; employed as high school teacher.
- Contact person: Husband, Edward Kobrynski, a steelworker.
- Insurance from full-time work.
- Source: Self, seems reliable.

Current Health Status

- Chief complaint is that her husband thinks she has a lump.
- Does not perform monthly BSE.

Past Health History

- Menarche at age 12; menstrual cycles have always been regular—every 28 days and lasting for 3 days.
- Had lumpectomy for benign breast lesion; otherwise, negative.
- No pregnancies; no history of infertility.
- No known food, drug, perfume, or deodorant allergies.
- Does not take any medications other than occasional Tylenol for headache. No history of BCP.

- No other medical problems. Does not examine breasts and has never had a CBE or mammogram.

Past Health History

- Positive for ductal breast cancer.
- Maternal grandmother deceased age 87; diagnosed with ductal breast cancer at age 85.
- Mother had benign breast lesion at age 57.
- Patient has two younger sisters, both alive and well.
- One maternal aunt, alive and well.
- No other cancer history in family.

Review of Systems

- **General Health Survey:** Usual state of health good.
- **Integumentary:** No dimpling, redness, or rashes on breasts or axillae.
- **Cardiovascular:** Reports no problems or medications.
- **Reproductive:** Menarche at age 12; menstrual cycle regular every 28 days, lasting 3 days; no history of BCP or HRT.

Psychosocial Profile

- Has not had physical exam in 5 years; has never had a mammogram; does not perform BSE; has not had CBE in 5 years.
- Typical day consists of awakening at 6 A.M., showering and dressing, eating breakfast with husband, and driving 30 minutes to school. Works 8 A.M. to 3:30 P.M. Monday through Friday. Returns home by 4 P.M., takes a walk or relaxes, fixes dinner. After dinner corrects papers, watches TV, or reads. Spends weekends catching up on housework, shopping and running errands, socializing with friends.
- Usually eats well, although diet premenstrually tends to be high in fats and salt. Temporarily gains 5 lb before she gets her period; otherwise, weight stable for past 10 years.
- Walks 3 miles, three to four times a week.
- Usually gets 8 hours sleep; no difficulty sleeping.
- Denies use of alcohol and caffeine.
- Employed full time, likes job, which does not require lifting.
- Satisfied with sex life.
- Married 15 years, husband and friends are supports.
- Goes to church weekly; religion is an important aspect of her life.

PAGE 539: CRITICAL THINKING ACTIVITY #2

- When did your husband first notice the lump?
- Do you have breast pain?
- Have you noticed any changes in your breasts?

PAGE 539: CRITICAL THINKING ACTIVITY #3

- Problem: Breast cancer.
- Risk factors: Family history of breast cancer; fact that she has never been pregnant; high-fat diet.

PAGE 548: LIST KEY HISTORY FINDINGS

- Age 46.
- No pregnancies.
- Husband thinks she has lump in breast.
- Does not perform BSE.
- Had lumpectomy for benign breast lesion.
- Has never had a CBE or mammogram.
- High-fat diet.
- Maternal grandmother died of ductal breast cancer at age 87.

PAGE 548: LIST KEY PHYSICAL EXAMINATION FINDINGS

- Healthy-appearing 46-year-old Caucasian female, appears stated age.
- Temperature, 98.4°F; pulse, 72 BPM; respirations, 18/min; BP, 110/68; height, 5'4"; weight, 116 lb.
- Day 7 of menstrual cycle.
- **Breasts:** Right breast slightly larger than left; uniform and symmetrical; ovoid in shape; no skin lesions noted; no redness or dimpling. Small, pea-sized (0.5 cm), easily movable, rubbery, smooth-edged lesion palpated in right breast at 2 o'clock; 4 cm from areola in the right upper outer quadrant; no lesions palpated in left breast.
- **Nipple:** No lesions observed or palpated; nipples erect and point in same direction; no inversion or eversion; patient denies discharge from nipple; unable to elicit discharge from either nipple; no specimens sent.
- **Axillae:** No lesions or thickening noted.

PAGE 549: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Does not perform BSE.
- Has not had physical for 5 years; never had a mammogram.
- Lump in left breast, slightly anxious.
- Possible need for breast surgery.

PAGE 553: REVIEW QUESTIONS

- d. Age is a major risk factor for breast cancer. The risk for breast cancer increases with age.
- c. A high-fat diet has been associated with increase risk of breast cancer, therefore, a low-fat diet is recommended to decrease the risk.
- d. For a premenopausal, it doesn't matter when she does a BSE. But, if the patient does the exam the same day every month, she is more apt to remember.

4. b. The areola is the most common site of breast cancer in men.
5. a. The central nodes are located high in the center of the axilla.
6. b. The lateral lymph nodes are located high in the axilla on the inner aspect of the humerus.
7. a. The epitrochlear nodes are located about 1 inch above the olecranon process.
8. a. Gynecomastia, male breast enlargement, is a side effect of digoxin.
9. c. Fibrocystic breast disease is characterized by tender masses.
10. d. Contraction of the pectoral muscles accentuates underlying masses. Sitting with hands pressed on hips contracts the pectoral muscles, accentuating dimpling or retraction of the breast tissue.

CHAPTER 17

PAGE 565: CRITICAL THINKING ACTIVITY #1

Appendicitis causes both visceral and parietal pain. The visceral pain results from distension of the appendix and results in umbilical pain; the parietal pain results from inflammation of the appendix and consequent inflammation of the peritoneum, which produces severe, localized pain that increases with movement.

PAGE 576: CASE STUDY EVALUATION

Biographical Data

- 56-year-old divorced woman, mother of two grown children.
- French Canadian descent, speaks English and French.
- Roman Catholic religion.
- High school graduate and mill worker.
- Source: Self, reliable.

Current Health Status

- No bowel movement in 4 days.
- Dull, intermittent lower abdominal pain rated 2 on a 0-to-10 point scale. Walking around makes it better and nothing makes it worse.
- Denies visible red blood in stool, black stool, or symptoms in other body systems.
- Says she has had problems with constipation for the past few years.

Past Health History

- Positive for varicella and measles.
- No history of GI problems as a child.
- Cholecystectomy 5 years ago; appendectomy 50 years ago.
- Denies having any GI assessment procedures or blood transfusions.

- Denies lactose intolerance or any allergies to substances, foods, or medications.
- Denies having any other health problems.
- Up to date on all immunizations; had hepatitis B series, pneumovax, and tetanus 5 years ago, yearly flu shot last fall.
- Takes no prescribed medications.
- Takes no OTC medications now. Has used laxatives to relieve constipation over past few years.
- Uses no herbal preparations.

Family History

- Mother and brother died of colon cancer.
- Father died of CHF.
- No family history of obesity or DM.

Review of Systems

- **General Health Survey:** No changes in energy level, just feels “bloated.”
- **Integumentary:** No changes.
- **HEENT:** No problems with head, neck, eyes, ears, nose, mouth, throat; no difficulty swallowing, last dental exam 6 months ago.
- **Respiratory:** No breathing difficulties.
- **Cardiovascular:** No problems.
- **Genitourinary:** No burning, frequency, or incontinence, no history of renal disease.
- **Neurological:** No loss of sensation, bowel and bladder control intact.
- **Musculoskeletal:** No changes, no history of arthritis.
- **Immune/Hematologic:** No allergies.
- **Endocrine:** No history of thyroid disease or DM.
- **Reproductive:** Postmenopausal.

Psychosocial Profile

- Typical day begins at 5:30 A.M., when she bathes, eats breakfast, and gets ready for work at the textile mill. Works 7 A.M. to 3 P.M. Goes grocery shopping after work, prepares supper, does housework, calls her sister, and then relaxes by watching TV. Goes to bed at 10 P.M. every night.
- 24-hour recall reveals that Mrs. Robichaud likes “junk food,” especially high-fat dishes traditional to the French culture. Breakfast usually consists of pancakes with butter and syrup; for lunch, she has chips and a sandwich; and for dinner, baked beans and brown bread. Says desserts are her weakness. She eats 1 serving of fruit; 1 serving of vegetables; 3 servings of meat; 1/2 serving of milk; 2 servings of white breads/pasta/rice/grains; and at least 5 fats a day. She reports that her usual weight is 175 lb and she is 5'3".
- No exercise program; uses car to get to work.

- Sleeps 7½ hours a night, longer on weekends.
- Goes for yearly physical exam.
- Does not drink alcohol or smoke cigarettes.
- Likes to make stained-glass objects in her spare time.
- Mother of two adult children, lives by herself and calls her sister every day. Has no pets.
- Lives alone in an apartment in a part of town where she doesn't feel safe walking alone.
- Mill where she works has been laying off employees, and she is worried that she will be next. Has been talking with her sister and other mill workers about this concern.
- Works in a textile mill, but does not know what chemicals she may be exposed to. Her apartment was built in 1985.

PAGE 576: CRITICAL THINKING ACTIVITY #2

- Age and weight.
- History of abdominal surgery.
- Family history of colon cancer.
- History of constipation.
- Use of laxatives.
- High-fat diet.
- Lack of routine exercise.
- Lead exposure from stained-glass hobby.
- Possible occupational exposure risk.

PAGE 576: CRITICAL THINKING ACTIVITY #3

- Abdominal discomfort as a result of no BM for past 4 days.
- Bloating feeling.
- Lack of exercise.
- History of constipation.
- Use of laxatives.
- History of abdominal surgery.
- Unsatisfactory diet.

PAGE 603: LIST KEY HISTORY FINDINGS

- Age 56, weight 175 lb.
- Lower abdominal discomfort, dull intermittent 2/10.
- No BM for 4 days.
- History of cholecystectomy, appendectomy.
- Use of laxatives.
- Positive family history of colon cancer.
- Junk food eater, high-fat diet.
- No routine exercise.
- Lead exposure from stained-glass hobby.
- Possible occupational exposure risk.

PAGE 603: LIST KEY PHYSICAL EXAMINATION FINDINGS

- Temperature, 98.8°F; pulse, 86 and regular; respirations, 22; BP, 168/98; height, 5'2", weight, 175 lb.

- Well-nourished, obese, 56-year-old woman who appears older than her years.
- In no acute distress, but mildly anxious; posture erect; pleasant and cooperative; provides history with ease.
- Abdomen uniformly light pink in color, protruberant and slightly distended in lower half, soft with complaints of slight tenderness in the lower quadrants.
- No striae, bruises, or hernias noted, umbilicus centered. Scar noted from previous cholecystectomy.
- No aortic pulsations, peristalsis noted.
- Hypoactive bowel sounds in the four quadrants.
- No bruits, friction rubs, or venous hums auscultated.
- Liver span 6.5 at MCL.
- Unable to percuss the bladder or spleen.
- Negative CVA tenderness.
- Liver, spleen, kidneys, bladder, and inguinal nodes not palpable.
- Abdominal aorta approximately 2 cm, palpable just left of the midline.
- Superficial abdominal reflexes intact.
- Negative fluid.

PAGE 603: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Abdominal discomfort rated 2/10; abdomen slightly distended and tender in lower quadrants.
- Dietary history of junk foods, fatty foods.
- High-fat, low-fiber diet.
- Family history of colon cancer; personal risk for colon cancer related to diet and lack of exercise.

PAGE 603: ADDITIONAL NURSING DIAGNOSES

- Activity Intolerance
- Nutrition: imbalanced, more than body requirements
- Pain
- Constipation
- Fatigue
- Knowledge, deficient, regarding diet and exercise

PAGE 603: COLLABORATIVE DIAGNOSES

Prevention of colon cancer through screening tests.

PAGE 603: CRITICAL THINKING ACTIVITY #5

- What is your daily fluid intake?
- Do you have trouble walking?
- How do you usually treat your constipation?
- Do you use laxatives?
- What medications are you on (may have a side effect of constipation)?

PAGE 603: CRITICAL THINKING ACTIVITY #6

- Increase fluid and fiber intake.
- Increase exercise.
- Respond promptly to defecation urge.

PAGE 610: REVIEW QUESTIONS

1. a. The ileopsoas test is a test for appendicitis. With the patient supine, the examiner applies pressure over the right thigh as the patient attempts to raise leg. Pain with this maneuver is a positive sign.
2. d. When the appendix ruptures, the contents of the intestines drain into the peritoneum causing peritonitis. Peritonitis usually paralyzes GI motility resulting in absent bowel sounds.
3. b. Bowel sounds normally occur every 5 to 15 seconds. Before you determine that bowel sounds are absent, you should auscultate for 5 minutes to be sure that the sounds are really absent rather than hypoactive.
4. c. The sequence for physical exam of the abdomen should be inspection, auscultation, percussion, then palpation. Auscultation should always occur prior to percussion and palpation so as not to alter the bowel sounds.
5. c. Underlying hernias can be accentuated by having the patient raise his head off the bed. Raising the head off the bed contracts the abdominal muscles, causing a hernia to become more pronounced.
6. a. The Scratch Test is a technique that is helpful in locating the lower border of the liver. The examiner places the diaphragm of the stethoscope in the right upper quadrant, then gently scratches the abdomen along the MCL moving up toward the RUQ. A scratching sound will be heard once the examiner is over the lower edge of the liver.
7. c. The normal liver span of an adult at the right MCL is 6 to 12 cm.
8. d. The color of stool may reflect an underlying GI problem. If the patient has a UGI bleed, the color of the stool will be black and tarry.
9. a. A patient with liver disease may develop venous congestion in the portal circulation. This may result in a venous hum, heard best in the RUQ.
10. a. Fluid in the abdominal cavity is called *ascites*. One technique for assessing ascites is shifting dullness. With the patient supine, percussion of the abdomen will reveal dullness laterally over the fluid and tympany over the umbilical area. The patient is then turned on his right side. As the patient turns, the fluid shifts. You will now note tympany on the left side of the abdomen and dullness on the right lateral aspects to the midline section.

CHAPTER 18

PAGE 628: CASE STUDY EVALUATION

Biographical Data

- 29-year-old African-American female.
- Married, mother of a 4-year-old girl.

- American by birth; Protestant religion.
- Attended 1 year of college after high school; homemaker at present. Husband is stockbroker for investment banking company.
- Healthcare insurance with husband's company.
- Source: Self, seems reliable.

Current Health Status

- Scant midcycle vaginal bleeding for the past 2 months with no other complaints at present.
- Seeking care for annual Pap smear, gynecological examination, and renewed prescription for oral contraceptives.

Past Health History

- Rheumatic fever at age 7.
- Usual childhood illnesses except for measles, mumps, and rubella.
- Denies any previous surgeries or serious injuries.
- Has taken oral contraceptives since birth of child 4 years ago. No other medications except occasional aspirin.
- No history of STDs.
- Gravida 1, para 1, no complications with pregnancy or vaginal delivery.
- Menarche at age 12, LMP 21 days ago. Menses last 5 days with moderate to light flow and occurs every 21 to 23 days. Denies dysmenorrhea.
- Denies exposure to DES.

Family History

- No gynecological carcinomas or maternal exposure to DES.
- Parents are alive and well.
- Has two healthy female siblings, ages 33 and 35.

Review of Systems

- **General Health Survey:** Usual state of health good, no changes in weight or energy level.
- **Integumentary:** No changes in skin, hair, or nails.
- **HEENT:** No sore throat, headaches, lumps, or masses in neck.
- **Respiratory:** No history of respiratory disease.
- **Cardiovascular:** No history of cardiovascular disease.
- **Gastrointestinal:** Appetite good, no GI problems.
- **Musculoskeletal:** No weakness, joint pain, or fractures.
- **Neurological:** No history of neurological problems.
- **Endocrine:** No history of DM or thyroid disease.
- **Lymphatic/Hematologic:** No lymphatic or hematologic disorders.

Psychosocial Profile

- Last gynecological examination and Pap smear 1 year ago with no abnormalities found. Had one cold in past year without seeking medical attention. Bathes daily and does not douche.
- Typical day includes waking at 6:00 A.M., preparing family breakfast, and performing household chores. Daughter goes to day care 2 days a week while patient volunteers at indigent healthcare facility.
- Eats three meals a day, drinks two cups of coffee. Does not eat at restaurants often and attempts to make healthy meals.
- Exercise and leisure activities include gardening, reading, and outside activities with family.
- Family pet is a 1-year-old beagle.
- Asleep by 10:30 P.M. (7 1/2 hours a night).
- Does not smoke, drinks one glass of wine occasionally with evening meals, denies recreational drug use.
- Lives in a suburban house of adequate size for a large family.
- No cultural or religious influences on healthcare. Family attends church most Sundays.
- Describes loving relationship with spouse and states that they intend to have three more children. Enjoys being a homemaker.
- Copes with stress by communicating with spouse.

Sexual History

- Sexually active in a monogamous, heterosexual relationship with husband.
- Satisfied with sexual performance.
- Uses oral contraceptives.

PAGE 628: CRITICAL THINKING ACTIVITY #1

Pelvic examination and Pap test.

PAGE 628: CRITICAL THINKING ACTIVITY #2

- Knowledge of oral contraceptives, side effects, and so forth.
- Compliance with taking oral contraceptives.

PAGE 643: KEY HISTORY FINDINGS

- Scant midcycle bleeding.
- Uses oral contraceptives.

PAGE 643: KEY PHYSICAL EXAMINATION FINDINGS

Normal gynecological examination without significant findings.

PAGE 643: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Waited until annual Pap smear was due to mention midcycle bleeding. Lack of concern over midcycle bleeding.

- Midcycle vaginal bleeding could indicate a uterine or vaginal infection, hormonal disturbance, or malignancy.

PAGE 643: ADDITIONAL NURSING DIAGNOSES

- Infection, risk for
- Sexuality Patterns, ineffective
- Growth and Development, delayed
- Sexual Dysfunction
- Role Performance, ineffective
- Self-Esteem, risk for situational low
- Pain, management
- Tissue Integrity, impaired, related to lesions

PAGE 643: COLLABORATIVE NURSING DIAGNOSES

Alteration in menstrual cycle patterns related to oral contraceptive use.

PAGE 647: REVIEW QUESTIONS

1. a. When a woman of child-bearing age presents with lower abdominal pain, one of the first questions asked should be “When was your last period?” You need to determine if the patient is pregnant. If the patient is pregnant, it will influence the direction of the exam and testing.
2. c. Excessive exercise may result in amenorrhea.
3. d. A cheesy, white vaginal discharge is characteristic of candidiasis, a fungal infection.
4. a. The Pap test is a screening test for cervical cancer.
5. c. A condylomata acuminatum is a painless, white, wartlike lesion caused by the human papillomavirus (HPV). It is sometimes referred to as *venereal warts*.
6. b. Water is the only lubricant that should be used on the speculum. Other lubricants may alter Pap test findings.
7. d. The cervix of a pregnant patient changes from pink in color to blue owing to an increase in venous congestion. This is referred to as *Chadwick’s sign*. Chadwick’s sign is usually seen by the 6th to 8th week of pregnancy and is a probable sign of pregnancy.
8. a. The color of a normal, nonpregnant, premenopausal cervix is pink.
9. c. The normal pubic hair distribution for an adult omen is an inverse triangle.
10. a. The KOH test is a quick test for *Candida*.

CHAPTER 19

PAGE 665: CASE STUDY EVALUATION

Biographical Data

- 35-year-old Caucasian male.
- American by birth; Protestant religion.
- Recently divorced, has a 10-year-old daughter.

- Employed as telephone lineman for 10 years; health-care insurance through work.
- High school education.
- Source: self, seems reliable.

Current Health Status

- Onset of dysuria 3 days ago. Clear urethral discharge occasionally at end of urine stream.
- Denies low back pain, blood in urine, or recent trauma.
- Denies suprapubic discomfort.
- Has had two different sex partners within the past 6 months.
- Uses condoms “sometimes.”

Past Health History

- Usual childhood illnesses except for measles, mumps, and rubella.
- Tonsillectomy in 1967, no other surgeries or hospitalizations.
- Had chlamydia approximately a year ago and received antibiotics.
- No history of renal calculi.
- Immunizations are current.
- Denies taking any prescription drugs; admits using aspirin occasionally for aches and pains.

Family History

- Father died at age 52 of heart attack.
- Mother, age 62, alive and well.
- Three siblings, all healthy.

Review of Systems

- **General Health Survey:** Usual state of health good.
- **Integumentary:** No changes in skin, hair, and nails; no lesions.
- **HEENT:** No swollen glands/nodes; denies sore throat; no eye drainage.
- **Respiratory:** No history of COPD.
- **Cardiovascular:** No history of HTN or cardiovascular disease.
- **Gastrointestinal:** No history of liver disease.
- **Musculoskeletal:** Denies joint pain or swelling.
- **Neurological:** No neurological problems, weakness or paralysis; denies personality changes or depression.
- **Endocrine:** No history of thyroid disease or DM.
- **Immunologic/Hematologic:** Has never been tested for HIV; no history of sickle cell anemia.

Psychosocial Profile

- Cannot remember when he last had a physical or went to a doctor for illness. Does not perform TSE.
- Typical day: Awakens at 6:30 A.M., has only coffee for

breakfast, gets to work at 7:30 A.M. Work is very physical and includes climbing up telephone poles and onto roofs of houses and crawling under houses. Usually arrives home by 5:30 P.M. Spends evenings watching TV or dating. Spends time with child every other weekend. Usually in bed about 10 P.M. on workdays.

- Skips breakfast, drinks four cups of coffee daily, eats out a lot, snacks on junk food, rarely has a nutritious meal.
- Besides work, hikes about every other weekend, plays tennis occasionally, does yard work.
- Sleeps about 7 hours a night on workdays, less on nights when he has dates. Naps on weekends to “catch up” on sleep.
- Has smoked a pack of cigarettes daily for the past 15 years. Drinks beer about three times per week and about 6 to 10 beers on weekends. Denies recreational drug use.
- Job is physically taxing and potentially dangerous because of climbing.
- Has good relationship with his one child and with his mother, who cleans, irons, and occasionally cooks for him.
- No community involvement, does not attend church.
- Lives alone since divorce 1 year ago, in single-family home with more space than he needs.
- Copes with stress by ignoring it.

Sexual History

- Currently sexually active.
- Heterosexual.
- Two sexual partners within past 6 months.
- Inconsistent use of condoms.
- Satisfied with sexual performance.
- No problems achieving erection or ejaculation.
- Does not perform TSE.

PAGE 665: CRITICAL THINKING ACTIVITY #1

- STDs.
- Safe sex practices.
- Nutritious diet.
- Smoking and alcohol.
- Health maintenance practices, such as TSE and yearly physicals.

PAGE 678: KEY HISTORY FINDINGS

- Onset of dysuria 3 days ago. Clear urethral discharge occasionally at end of urine stream.
- Had two different sex partners within the past 6 months.
- Uses condoms “sometimes.”
- Had chlamydia approximately a year ago and received antibiotics.
- Uses alcohol regularly.
- Has smoked 1 pack of cigarettes daily for 15 years.

- Does not perform TSE.
- Rarely has a nutritious meal.
- Cannot remember the last time he saw a doctor.

PAGE 678: KEY PHYSICAL EXAMINATION FINDINGS

- Reddened urethral meatus.
- Clear to white fluid discharge from penis.
- Right inguinal lymphadenopathy.

PAGE 678: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- States that he is concerned about STDs. Requests testing for all STDs including HIV.
- Is aware of effects of infections in reproductive system and of fact that some STDs are not curable. Verbalizes fears about being infected with an STD. States that he is concerned about telling sex partners. Scheduled for HIV antibody testing as well as tests for syphilis, gonorrhea, and chlamydia.
- Uses condoms only occasionally.
- Does not perform TSE. Uses tobacco and alcohol.

PAGE 678: ADDITIONAL NURSING DIAGNOSES

- Role Performance, ineffective
- Self-Esteem, risk for situational low
- Sexual Dysfunction
- Sexuality Patterns, ineffective
- Infection, risk for

PAGE 678: COLLABORATIVE NURSING DIAGNOSES

Potential for altered reproductive health: sexually transmitted disease.

PAGE 684: REVIEW QUESTIONS

1. b. Normally, the testes are about 5 cm by 2.5 cm, firm, rubbery in consistency.
2. b. Undescended testicles has been identified as a risk factor for testicular cancer.
3. d. Maternal use of DES is a risk factor for testicular cancer.
4. c. PSA is a screening test for prostate cancer.
5. a. Gonorrhea, caused by *Neisseria gonorrhoeae*, typically causes urinary symptoms and produces a thick, yellow penile discharge.
6. c. A condylomata acuminatum is a painless, white, wartlike lesion caused by HPV. It is sometimes referred to as *venereal warts*.
7. a. Male patient with a history of sickle cell anemia may develop priapism, a painful, sustained erection without sexual desire.
8. a. The position of the urinary meatus is assessed in male newborns. Normally, the meatus is located centrally on the tip of the penis. Epispadias is a urethral meatus opening on the dorsal (upper) side of the penis.
9. d. A hydrocele is a nontender accumulation of fluid between two layers of tunica vaginalis in the scrotum. The cause may be idiopathic or the result of trauma, inguinal surgery, epididymitis, or a tumor. In newborns, the cause is usually related to delivery.
10. d. The recommended age for prostate screening is age 50.

CHAPTER 20

PAGE 701: CASE STUDY EVALUATION

Biographical Data

- 68-year-old Caucasian female, widow.
- Works as housekeeper for parish priest.
- Born and raised in United States, Polish descent, Catholic religion.
- Current health insurance: Medicare part A and B, supplemented with Medicaid.
- Source: Self, seems reliable.

Current Health Status

- “Several-year” history of aching joints in the evening after working all day.
- Discomfort first occurred in back and neck.
- Takes OTC analgesic for pain with some relief.

Past Health History

- History of HTN.
- Wrist fracture at age 65, caused by a fall on the ice. No pinning was performed.
- Cholecystectomy at age 52, appendectomy at age 8, hysterectomy at age 35.
- Vaginal deliveries of two children with no complications.
- Denies allergies to food, drugs, or environmental factors.
- Medications include Zestril 10 mg OD for HTN and Aleve once daily for arthritic pain.

Family History

- Mother had osteoporosis and HTN; died of cardiac problems at age 82.
- Father had HTN and died of heart attack at age 65.
- Sister, age 72, positive for osteoporosis and HTN.
- One daughter and one son, alive and well.
- All other known relatives alive and well.

Review of Systems

- **General Health Survey:** Describes usual state of health as “fair.”
- **Integumentary:** Reports no changes in skin, hair, or nails.

- **HEENT:** Complains of pain and stiffness in neck; no swollen glands.
- **Respiratory:** No respiratory problems.
- **Cardiovascular:** Positive history of HTN, controlled with medication.
- **Gastrointestinal:** No abdominal problems.
- **Genitourinary:** No problems.
- **Reproductive:** Postmenopausal since age 35.
- **Neurological:** No changes in sensations.
- **Immunologic/Hematologic:** No infections or bleeding.
- **Endocrine:** No history of DM or thyroid disease.

Psychosocial Profile

- Has a yearly exam because of HTN. Has not seen a gynecologist since her hysterectomy. Visits family physician every 3 months for HTN check.
- Arises at 6 A.M. Does chores around own home and dresses for work. Workday is 9 A.M. to 5 P.M. Typical day consists of daily chores around employer's home, including cooking, cleaning, and running errands. After work, she relaxes and watches the news and some TV shows before going to bed at 9 P.M.
- Admits she is a "tea and toast lady." Lives alone, so if she cooks, it is something quick, unless she goes out to eat with friends or family. Drinks two to four cups of caffeinated coffee a day. Has never been overweight.
- Activities include housecleaning and running errands uptown. No formal exercise program.
- Has no pets. Knits for a hobby while watching TV. Occasionally has dinner out or sees a movie with friends or family.
- Sleeps 4 to 5 hours uninterrupted, but often gets up at night to void.
- Denies smoking, alcohol use, or illicit drug use.
- Job requires some lifting (e.g., carrying vacuum cleaner upstairs) as well as bending.
- Says her self-esteem is "pretty good." Says she feels proud she can still earn a living. Admits she worries about living alone and developing health problems as she gets older, especially those that limit her mobility.
- Lives alone in a ranch home in suburbs; no steps. Is not afraid to be alone because of good neighbors.
- Has lunch or dinner once a week with daughter or son. Talks on phone to sister several times a week. Has friends in neighborhood, including younger neighbors who check on her.
- Copes with stress by talking with children or priest.

PAGE 701: CRITICAL THINKING ACTIVITY #1

- **Musculoskeletal problem:** Osteoporosis.
- **Risk factors:** Age, sex, race, family history, history of fracture, being postmenopausal, drinking two to four cups coffee a day, not exercising.

PAGE 730: KEY HISTORY FINDINGS

- 68-year-old white female in for yearly routine examination.
- Concerned that she is "shrinking."
- Complains of back and neck discomfort at end of day for about past 5 years.
- Family history of osteoporosis in mother.
- Poor eating habits and sedentary lifestyle.
- Consumes two to four cups of coffee a day.
- History of hysterectomy because of dysfunctional bleeding at age 35. No use of estrogen replacement therapy.
- History of left wrist fracture age 65 caused by fall on ice.

PAGE 730: KEY PHYSICAL EXAMINATION FINDINGS

- Height 5'5"; weight 125 lb (1½-inch height loss since onset of menopause.)
- Gait steady, no support needed for ambulation.
- Mild muscle atrophy noted to quadriceps; all other muscles without atrophy. No involuntary movements.
- Shape and tone of muscles within normal limits.
- Contour smooth over joints. No tenderness, heat, redness, or pain palpated in any joint.
- Full ROM in all joints.
- Muscle strength 5/5.
- Limb length bilaterally: 80.5 cm leg; 40 cm thigh.
- Mild kyphosis noted when standing.

PAGE 731: CRITICAL THINKING ACTIVITY #2

- Strengths.
- General health fair.
- Health-seeking behavior.

Areas of Concern

Risk factors for osteoporosis: age, gender, ethnicity, postmenopausal, diet, caffeine intake, no exercise, history of fracture, family history of osteoporosis 1½ inch loss of height.

PAGE 731: COLLABORATIVE NURSING DIAGNOSES

- Complains of joint pain at end of day, back and neck pain.
- Pain, atrophy of quadriceps.
- Concerned about height loss; requesting information about osteoporosis.

PAGE 731: ADDITIONAL NURSING DIAGNOSES

- Activity Intolerance
- Body Image, disturbed
- Pain, chronic
- Self-Care Deficit
- Sleep Pattern, disturbed

PAGE 735: REVIEW QUESTIONS

1. b. An accentuated lumbar curve, lordosis, can be seen in women during the last trimester of pregnancy owing to the shift in the center of gravity.
2. a. The base of support is usually 2 to 4". If the person is having balance problems, the base of support is widened to maintain balance.
3. d. Scoliosis is a lateral "S" curve of the spine. To assess for scoliosis, have the patient bend at the waist and examine the spine. This helps you to differentiate a functional from structural problem. Unequal, accentuated scapula is a sign of scoliosis.
4. a. A side effect of steroids is osteoporosis.
5. a. A goniometer is an instrument useful in assessing the ROM of a joint.
6. d. The Romberg test is a test for balance. To perform the Romberg test have the patient stand with feet together, hands at side eyes, open then closed; note ability to maintain balance. A positive Romberg may indicate a cerebellar problem.
7. a, c, e. Several techniques can be used to assess balance. Tandem walk, heel-and-toe walking and the Romberg test can be used to assess balance.
8. b. Arm length measurements are taken from the acromion process to the tip of the middle finger.
9. a, b. The normal ROM of the knee are flexion and extension.
10. a, c, d. The normal ROM of the temporomandibular joint is retraction, elevation, and protraction.

CHAPTER 21

PAGE 761: CRITICAL THINKING ACTIVITY #1

- What does the pain feel like? How severe is it? Where does it hurt (point to spot)? Do you have any other symptoms, such as nausea or vision changes? Were you drinking or using drugs?

PAGE 761: CASE STUDY EVALUATION

Biographical Data

- 21-year-old African American male.
- Baptist religion.
- Senior in college; business major.
- Insurance under family's plan while in college.
- Lives on campus during school year; home is 1 hour from school.
- Contact person: Janet Jones, mother.
- Source: Self, seems reliable.

Current Health Status

- Complains of headache and neck pain from hitting head in motor vehicle accident.
- Had a few beers earlier in night.

- Felt a little confused after accident and is not sure what happened. Does not know whether he lost consciousness.

Past Health History

- No hospitalizations or surgeries.
- Concussion in high school while playing football, otherwise, negative for neurological problems.
- Fractured fibula freshman year in college while playing football; casted and healed without problem.
- Thinks immunizations are up to date but unsure of last tetanus shot.
- No known food, drug, or environmental allergies.
- No medications or other medical problems.

Family History

- Father, age 55, has HTN.
- Paternal uncle, age 60, has HTN.
- Paternal grandfather, died at 80 from stroke.
- Maternal aunt, age 50, has HTN.
- Maternal grandfather died at 75 from stroke.

Review of Systems

- **General Health Survey:** Usual state of health good until accident.
- **Integumentary:** No changes.
- **HEENT:** Headache and neck pain; a little "dizzy."
- **Respiratory:** No problems.
- **Cardiovascular:** No problems.
- **Gastrointestinal:** Appetite usually good, but felt "a little sick after accident." No vomiting.
- **Genitourinary:** No problems.
- **Reproductive:** Does not wish to discuss.
- **Endocrine:** Negative.
- **Lymphatic/Hematologic:** Negative.

Psychosocial Profile

- Gets yearly physical exam for sports.
- Typical day consists of awakening at 9 A.M., grabbing some breakfast, and heading to class. Goes to football practice from 3 P.M. to 6 P.M. 7 days a week, has dinner, relaxes, and does school work until about midnight. Usually goes to bed by 1 A.M.
- Usually eats well; diet high in carbohydrates and protein.
- Practices football with team 3 hours a day, 7 days a week. Wears protective gear.
- Usually gets at least 8 hours of sleep a night.
- Denies use of drugs or nicotine. Admits to drinking alcohol on weekends; amount varies.
- Does well in school (GPA 3.0).
- Does not wish to discuss sexuality patterns.
- Friends and family are supports.

- Copes with stress by exercising, partying with friends on weekends.

PAGE 761: CRITICAL THINKING ACTIVITY #2

- **Neurological problem:** Injury.
- **Contributing factors:** Nutritional deficits; alcohol abuse; family history of HTN and stroke.

PAGE 787: KEY HISTORY FINDINGS

- Drinks alcohol on weekends; had a few beers earlier in night.
- Felt a little confused after accident. Does not know whether he lost consciousness.
- Complains of headache, neck pain, and “a little dizziness.”
- No vomiting after accident.
- Concussion in high school while playing football; otherwise, no neurological problems.
- Family history of HTN/stroke.
- Wears protective gear while playing football.

PAGE 787: KEY PHYSICAL EXAMINATION FINDINGS

- Guarding and decreased ROM of neck.
- 2-inch laceration on forehead.
- Alcohol on breath.

PAGE 787: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

1. Reported head and neck injury from motor vehicle accident. Complains of headache and neck pain. Guarded movement of neck, increased neck ROM.
2. Reported head injury, headache, dizziness, recent memory loss.
3. Reported head injury, headache, dizziness, recent memory loss.

PAGE 787: ADDITIONAL NURSING DIAGNOSES

- Knowledge, deficient, related to substance abuse (alcohol)
- Potential for increased intracranial pressure related to head injury

PAGE 794: REVIEW QUESTIONS

1. b. The GCS is a universal tool that assess level of consciousness. The GCS evaluates best eye, motor, and verbal responses. Best score is 15, worst score 3.
2. a. A change in mental status is an early sign of increased intracranial pressure.
3. d. Level of orientation is assessed for time, place, and person. Usually disorientation to time, then place occur before a patient becomes disoriented to self. Since self-identity usually remains intact the longest, disorientation to self may be an ominous sign.

4. a. The muscles of mastication are controlled by the trigeminal nerve, CN V. To test the motor portion of CN V, have the patient bite down on a tongue blade and note strength of jaw.
5. b. The hypoglossal nerve, CN XII, controls the motor function of the tongue. The CN XII can be tested by having the patient say words that contain the letters d, l, n, and t. So, the phrase “You can’t teach an old dog new tricks” is commonly used.
6. b. Graphesthesia is a discriminatory sensation. With the patient’s eyes closed, test graphesthesia by writing a number or letter in the patient’s palm of his hand, and having him identify it.
7. b. Extinction is a discriminatory sensation. With the patient’s eyes closed, test extinction by simultaneously touching opposite sides of the patient’s body at the same time, and have him identify where he was touched.
8. b. The grading scale used for DTR is 0 to 4. 0 is absent, +1 diminished, +2 considered normal, +3 brisk, +4 hyperactive with clonus.
9. d. The Babinski reflex is an abnormal response in an adult. To test for a Babinski, stroke the sole of the patient’s foot in an arc from the lateral heel across the ball of the foot to the great toe. The normal response (plantar response) is flexion of the toes. Fanning of the toes with dorsiflexion of the great toe is a + Babinski.
10. b. Brudzinski sign is a test for meningitis. To test Brudzinski sign, have the patient flex head and touch chin to chest. If the hips flex, the test is positive for meningitis.

CHAPTER 22

PAGE 800: CRITICAL THINKING ACTIVITY #1

- Nutrition: imbalanced, more than body requirements
- Sexuality patterns, ineffective
- Health-seeking behavior, re: osteoporosis
- Health maintenance, ineffective, related to diet
- Knowledge, deficient, of breast self-examination

PAGE 800: CRITICAL THINKING ACTIVITY #2

- Breast exam.
- Cardiovascular exam.
- Musculoskeletal assessment for signs of osteoporosis such as loss in height.

PAGE 800: CASE STUDY EVALUATION

Biographical Data

- 40-year-old Caucasian female.
- Northern European extraction.
- Protestant religion.
- Married, with three children ages 12, 15, and 18.

- Has a full-time, sedentary job as an accountant.
- Source: Self; seems reliable.

Current Health Status

- No current health problems.
- Concerned about physiological changes of menopause and healthcare recommended for her age.
- Occasionally takes acetaminophen for tension headaches. No other medications.

Past Health History

- Had chickenpox, measles, mumps, and rubella as a child. Also had frequent bouts of strep throat as a child.
- Had an appendectomy at age 15. Was hospitalized for birth of her three children. Had a tubal ligation at age 35 for fertility control. Has never had a serious injury requiring hospitalization.
- No history of medical problems.
- No drug, environmental, or food allergies. Has taken penicillin without a reaction.
- Had all childhood immunizations. Last DPT shot 10 years ago. Negative PPD skin test 5 years ago.
- Has not traveled in a foreign country within past year. Has never served in the armed services.

Family History

- Husband, age 45, alive and well.
- Son, age 18, alive and well.
- Daughter, age 15, alive and well.
- Daughter, age 12, alive and well.
- Mother, age 76, had breast cancer at age 54; osteoporosis and fractured hip at age 63.
- Father, age 78, has history of HTN and had an MI at age 68.
- Younger brother died in car accident at age 35.
- Two sisters, ages 42 and 50, alive and well.
- Maternal grandmother died of breast cancer at age 78.
- Maternal grandfather died of MI at age 81.
- Maternal uncle, age 80, has history of HTN.
- Paternal grandmother died of stroke at age 82.
- Maternal aunt, age 78, has history of breast cancer and osteoporosis.
- Paternal grandfather died of MI at age 86.
- Paternal aunt, age 65, alive and well.
- Paternal aunt, age 69, history of breast cancer.
- Paternal uncle, age 71, history of HTN.

Review of Systems

- **General Health Survey:** Patient feels that her overall health is good. No fever, chills, fatigue, depression, or anxiety. She has occasional colds

and has gained 10 lb in past year without a change in diet.

- **Integumentary:** No rashes, lesions, mole changes, or bruising. Skin is dry.
- **HEENT:** No headaches, frequent sore throats, hoarseness, or problems with vision, hearing, or sinuses.
- **Respiratory:** No SOB, wheezing, or productive or nonproductive cough
- **Cardiovascular:** No history of HTN, heart problems or murmurs, blood clots/phlebitis, chest pain, palpitations, edema, orthopnea, or claudication.
- **Breast:** Performs BSE occasionally, when she remembers. Has never had a mammogram. No breast pain, tenderness, lumps, discharge, or change in nipples. Breastfed all 3 children.
- **Gastrointestinal:** No nausea/vomiting, abdominal pain, dysphasia, heartburn, jaundice, hemorrhoids, or blood in stool. Has daily bowel movement, brown in color.
- **Genitourinary:** No dysuria, frequency or urgency, although reports occasional stress incontinence. Satisfied with sexual life except for increasing vaginal dryness. No breast masses or nipple discharge. Has breast pain 7 days before menstrual cycle. Occasionally does BSE.
- **Reproductive:** LMP 21 days ago. Menarche at age 14. Gravida 3, para 3. Menses are irregular, ranging from 23 to 45 days (cycle used to be 28 to 30 days). Menses last 5 days with moderate to heavy flow. Contraception method is tubal ligation.
- **Musculoskeletal:** No joint pains, swelling, or muscle weakness.
- **Neurological:** No dizziness, vertigo, syncope, head injuries, seizures, numbness, or tingling of extremities. Reports forgetfulness and difficulty concentrating. Has never been treated for mental disorders, depression, or anxiety, although she did visit a counselor about a family problem 3 years ago.
- **Endocrine:** No DM, thyroid disease, polyuria/polydipsia, or heat or cold intolerance. Weight gain of 10 lb last year.
- **Immunologic/Hematologic:** No current infections, allergies, cancer, bleeding, anemia, or blood transfusions. Blood type A+.

Psychosocial Profile

- Does not usually seek routine healthcare or screenings. Feels that she is healthy and does not need healthcare unless she is sick. Does not know her cholesterol level, has never had a mammogram, has not had a Pap smear or physical examination in 3 years.
- Typical day consists of arising at 6:00 A.M., getting her children off to school and then going to work.

She works from 8:00 A.M. to 5:00 P.M. 5 days a week. After work, she makes dinner for her family and reads or watches TV. She goes to bed at 11:00 P.M. each night.

- Reports that she eats a balanced diet with foods from each of the major food groups each day, including 3 vegetables and 2 fruit servings. Admits to eating sweets and fried foods every day. Evaluation of daily calcium intake reveals that she usually ingests 500 mg a day. Has had an unexplained 10-lb weight gain in last year.
- 24-hour recall shows the following: Breakfast was two fried eggs, two slices of bacon, two slices of white toast with butter, one 4-oz. glass of orange juice, and a cinnamon bun with butter. Lunch was a chicken salad sandwich with mayonnaise on two slices of white toast, potato chips, two chocolate chip cookies, an 8-oz. cola, and a chocolate candy bar. Snack was one apple. Dinner was steak, baked potato with butter, string beans, tossed salad with Russian dressing, apple pie with ice cream, and 8 oz. of 2 percent low-fat milk.
- Admits that her only exercise is walking from her car to her office twice a day and walking up and down two flights of stairs several times a day. She walked on a daily basis in the past but stopped this year because of lack of time and inability to fit it into her schedule. For recreation she reads, does housework, and gardens on weekends.
- Hobbies include reading and gardening. Has a house cat.
- States that she usually sleeps well for 7 to 8 hours a night. Has occasional night sweats that awaken her.
- Has never smoked and is not exposed to smoke at home. Has smoke and carbon monoxide detectors at home. Drinks a glass of wine with dinner two to three times a week. Denies ever using illicit drugs.
- Identifies no occupational risks. Is not exposed to smoke at work. Drives 40 miles round trip to work each day. Wears seatbelt when driving.
- Attends church weekly with family. No religious or cultural influences that affect health practices.
- States that her marriage is supportive and comfortable. Describes good relationships with children, siblings, and parents.
- Uses prayer, friendships, and family support to deal with stress.

PAGE 801: CRITICAL THINKING ACTIVITY #3

Strengths

- Concerned about health and menopause.
- No history of medical problems.
- Supportive family.
- Normal physical examination.

Areas of Concern

- No regular exercise.
- Sedentary job.
- Weight gain of 10 lb.
- Diet high in fat and sugar, low in calcium.
- Infrequent BSE.
- Family history of cancer, osteoporosis, and cardiovascular disease.
- Does not seek routine healthcare and screening.

PAGE 808: LIST KEY HISTORY FINDINGS

- Concerned about menopause and healthcare recommended for her age group.
- Sedentary job; no regular exercise routine.
- Tubal ligation at age 35.
- History of breast cancer in family: Mother, grandmother, two aunts.
- History of osteoporosis in family: Mother and aunt.
- 10-lb weight gain in past year.
- Infrequent BSE; has never had mammogram.
- Does not go for routine health screenings.
- Has never had cholesterol checked.
- Has high-fat, high-calorie diet with inadequate calcium intake.

PAGE 808: LIST KEY PHYSICAL EXAMINATION FINDINGS

- Normal vital signs; height, 5'5"; weight, 145 lb.
- No abnormalities noted in any body system.

PAGE 808: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Weight gain of 10 lb over past year and high-fat, high-calorie diet.
- Evidenced by inadequate health maintenance, infrequent BSE, no pelvic examination for 3 years.

PAGE 808: COLLABORATIVE NURSING DIAGNOSES

- Potential complication: Cardiovascular disease.
- Potential complication: Osteoporosis.
- Potential complication: Breast cancer.

PAGE 808: REFERRALS

- Possible hormone replacement therapy: Perimenopausal.
- Pelvic examination and Pap test: 3 years since last examination, hormonal changes.
- Annual CBE: Infrequent BSE, family history of breast cancer, has never had a mammogram.
- Mammogram: Same as above.
- Cholesterol and thyroid screening: High-fat diet, family history of cardiovascular disease, perimenopausal.

- **DEXA scan for bone density:** Perimenopausal, family history of osteoporosis.

PAGE 809: REVIEW QUESTIONS

1. b. When performing a complete assessment, it is more efficient for both you and the patient to do the exam by region. It conserves the patient's energy by minimizing position changes.
2. a. When performing a focused assessment, a system approach is best allowing you to focus in on the area of concern.
3. a. When documenting your assessment findings, a systems approach is best for organizing the data.

CHAPTER 23

PAGE 823: CRITICAL THINKING ACTIVITY #1

During the antepartal period, encourage exercise and reassure your Irish American patient that stretching and other activities that involve lifting her arms over her head will not cause the umbilical cord to become wrapped around the baby's neck. With Mexican American patients, understand that because they view pregnancy as a "natural" and desirable condition, they may not seek early and ongoing prenatal care and instead rely on their family for advice and support. One approach to planning prenatal care would be to gain the support and acceptance of the family as you develop a timetable for the frequency of visits and discuss the goals of early and ongoing prenatal care.

In the intrapartum period, expect husbands of Arab American and Egyptian American patients to be voluntarily excluded from the birthing areas. For many cultures (Egyptian American, Filipino American, Greek American), the patient's mother will wish to be the primary labor support person.

Many cultures have certain beliefs regarding postpartum hygiene practices for the new mother. For example, your Egyptian American patient must avoid cold, which may result from bathing; your Filipino American patient may believe that showers can cause arthritis, so she will wish to take sponge baths only; and your Mexican American patient may wish to avoid showers altogether. Awareness of these different practices will be useful as you plan culturally appropriate care for your patients.

PAGE 823: CRITICAL THINKING ACTIVITY #2

- **African American:** Involve your patient's grandmother and maternal relatives in the plan for prenatal, intrapartum, and postpartum care.
- **Amish:** Make certain that the prenatal classes are scheduled at a time when fathers can attend.
- **Arab American:** Provide support, reassurance, and encouragement during labor, but do not emphasize relaxation and breathing techniques.

- **Chinese American:** Consult with the dietary department to plan healthy meals for the postpartum mother that do not include fruits or vegetables.
- **Cuban American, Egyptian American, Filipino American:** Involve the postpartum mother's family in the development of a plan of care for the new mother and her baby.
- **Greek American:** Plan for alternatives to showering for postpartum mothers.
- **Iranian American:** Soon after marriage, discuss preconception care and encourage healthy lifestyle practices to ensure a healthy pregnancy.
- **Irish American, Mexican American:** Educate about the rhythm method of birth control.
- **Jewish American:** Educate about birth control pills as one choice for contraception.
- **Navajo Native American:** Support and encourage the laboring woman who may wish to use special aids, such as a birthing necklace and a woven sash to facilitate the birth process.
- **Vietnamese American:** Avoid touching the antepartal woman's head or the newborn's head.

PAGE 823: CRITICAL THINKING ACTIVITY #3

- What do you and your family believe would be beneficial for you to do to remain healthy during your pregnancy?
- Can you identify some things that you could do, or not do, to improve your health and promote good health for your infant?
- Whom would you like to be present with you during labor?
- What activities would be important to you and to your family after your baby is born?
- What would you and your family like the nurses to do during your pregnancy, labor, and postpartum period?
- How would your family members like to participate in your birth experience?

PAGE 832: CRITICAL THINKING ACTIVITY #4

Mrs. Ramirez reports that her last normal period was April 10 to 17. Using Naegele's rule, $\text{April } 10 + 7 \text{ days} = \text{April } 17$; $3 \text{ months} = \text{January}$. Thus, her EDD is January 17.

PAGE 832: CRITICAL THINKING ACTIVITY #5

- Fatigue.
- Nausea and vomiting.
- Breast enlargement, tingling, and tenderness.
- Urinary frequency.
- Amenorrhea.

PAGE 832: CRITICAL THINKING ACTIVITY #6

- Avoid an empty or overfull stomach.
- Eat dry carbohydrates on awakening and remain in bed until nausea subsides.

- Eat five to six small meals a day.
- Avoid fried, odorous, greasy, spicy, or gas-producing foods.
- Maintain good posture and do not lie down immediately after eating.

PAGE 832: CRITICAL THINKING ACTIVITY #7

- Supportive and caring husband.
- Enjoys her work.
- Work is located near her home.
- Works in a safe environment.
- Obtains daily exercise.
- Obtains adequate rest.
- Maintains good nutrition.
- Enjoys daily recreation.
- Has yearly physical examination.
- Does not smoke, use drugs, or drink alcoholic beverages.
- Lives in a safe environment.
- Has family support persons nearby.
- Has a plan for infant care.

PAGE 832: CASE STUDY EVALUATION

Biographical Data

- 24 years old, married, first pregnancy.
- College graduate, full-time social worker.
- Born and reared in the United States, Hispanic descent, Catholic religion.
- Health insurance through own and husband's work plan.
- Referral: Routine prenatal care.
- Source: Self, seems reliable.

Current Health Status

- Fatigue.
- Nausea and vomiting for 3 weeks.
- Breast enlargement, tingling and tenderness.
- Urinary frequency.
- Amenorrhea.

Past Health History

- Usual childhood illnesses.
- No surgeries, hospitalizations, or allergies.
- Immunizations current.
- Takes 1 multivitamin with iron daily; no other medications.
- Physical examinations yearly for 18 years; normal Pap smears.
- Used oral contraceptives for 5 years without problems; discontinued them 8 months ago to attempt pregnancy.
- No previous pregnancies.

Family History

- Grandmother, age 64, takes medication to control high BP.
- All other relatives alive and well; two older sisters married with children and experienced no complications during pregnancies; one younger sister is single.

Review of Systems

- **General Health Survey:** Very tired for past couple of months, needs to take a nap every day when she gets home from work. Body weight 125 lb, on the low end of her reported range of 126 to 128 lb. Has lost 2 lb since becoming pregnant.
- **Integumentary:** Noticed nipples getting darker and line on abdomen. Hair very oily.
- **HEENT:** No changes in head or neck. Has nasal stuffiness but no colds, allergies, or flu. Gums bleed with brushing; last dental exam 6 months ago. No eye changes.
- **Respiratory:** No changes.
- **Cardiovascular:** No history of MVP, palpitations, or edema.
- **Breasts:** Enlargement, tenderness, and tingling sensation.
- **Gastrointestinal:** Nausea and vomiting in the morning, usually subside by noon; says certain foods "just turn my stomach"; reports constipation.
- **Genitourinary:** Reports frequency of urination, no burning
- **Reproductive:** LMP 4/10/01.
- **Musculoskeletal:** No changes.
- **Neurological:** Occasional "lightheadedness"; no history of depression; excited at possibility of being pregnant.
- **Endocrine:** No history of gestational DM; reports heat intolerance.
- **Immunologic/Hematologic:** No history of anemia.

Psychosocial Profile

- Patient is proactive regarding healthcare. Has yearly physical examination and Pap smear. Performs monthly BSE. Believes people are responsible for maintaining their own health.
- Typical day consists of arising at 6:30 A.M. and eating breakfast with husband. (Lately unable to enjoy usual breakfast of cold cereal and juice because of nausea.) Drives 4 miles to office 5 days a week and works from 8 P.M. to 5 P.M. at desk part of time and interviews patients in a clinic located on the ground floor of her office building for rest of time. Loves her work. In evenings, she and husband prepare and eat supper, clean up dishes, take a walk in neighborhood, then relax watching TV, playing

cards, or reading. Patient showers and goes to bed around 10 P.M.

- Usually eats a healthy diet, but lately has been skipping breakfast because of nausea. As day progresses, is gradually able to eat bland food such as baked chicken, potatoes, and steamed vegetables.
- Walks daily for 45 minutes with husband.
- No pets. Recreation includes daily walk with husband, dinner at restaurant, or movie once a week, family get-togethers. Hobbies include reading and needlework.
- Sleeps around 8 1/2 hours normally, but lately gets very sleepy at work, especially in midafternoon. Combats sleepiness with a walk outside her office and an afternoon snack of fruit and yogurt or crackers. Takes a short nap after work. No problems falling asleep.
- Does not smoke, use drugs, or drink alcoholic beverages.
- No chemical exposure at work.
- Lives in a one-story, two-bedroom single family home in suburbs; has hot water, heating, and air conditioning; negative for radon gas. Will convert guest room into baby's room. No recent travel outside local community.
- Catholic religion, Hispanic nationality (third generation). No religious or cultural influences that affect healthcare, pregnancy, or delivery.
- Supportive, caring husband (nonsmoker) who is computer analyst for national company; married 5 years; attempted pregnancy for 6 months.
- Mother, mother-in-law, and three sisters live in same community and will assist with infant care. Patient plans to breastfeed and stay at home for 6 weeks after birth and then return to job.

PAGE 842: KEY HISTORY FINDINGS

- Well-developed, well-nourished 24-year-old Hispanic woman in no acute distress.
- Oriented to time, place, situation, person; affect pleasant and appropriate.
- No complaints other than early pregnancy symptoms.
- Vital signs: Temperature, 99.0°F; pulse, 70 BPM, strong and regular; respirations, 16/min, unlabored; BP, 102/66.
- Height, 5'5"; weight, 125 lb; prepregnancy weight, 127 lb.
- Normal cyclic menstruator, certain of date of LMP; no contraception used during the past 8 months.
- Desires pregnancy, excited and asking appropriate questions.
- Taking multivitamins with iron and folic acid for past 8 months.
- Supportive social situation: Happily married, extended family nearby.
- No medical conditions, current or past.

- Exercises daily.
- Nutritional status good except for 2-lb weight loss related to morning sickness; urine ketones negative today.
- Desires to breastfeed.
- Wishes to enroll in childbirth preparation and infant care classes.

PAGE 842: KEY PHYSICAL EXAMINATION FINDINGS

- Urine pregnancy test positive today.
- Skin pink, warm, dry, good turgor, mucous membranes pink, moist.
- No palpable cervical chain lymph nodes; slight thyroid enlargement, no nodularity or tenderness.
- Lung sounds clear to auscultation bilaterally; heart, normal sinus rhythm.
- Breasts: Tenderness, venous congestion, prominence of Montgomery's tubercles, darkening of nipples and areolae, no nipple discharge.
- Abdomen: Linea nigra present, no masses, no enlargement, no FHTs auscultated with Doppler.
- External genitalia: Normal hair distribution, no varicosities, slight labial enlargement.
- Extremities normal color, no edema, no varicosities, no tenderness, no increased warmth, DTRs 1+.
- Pelvic examination:
 - Bartholin's and Skene's glands: Nontender, no discharge.
 - Cervix: Positive Chadwick's sign, slight leukorrhea.
 - Cervix long, thick, and closed. Cervical length, 3.0 cm; positive Hegar's sign.
 - Uterus approximately orange-sized; no adnexal masses.
 - Rectal mucosa pink, intact, no hemorrhoids.

PAGE 842: CRITICAL THINKING ACTIVITY #8

Strengths

- Healthy.
- No history of medical problems.
- Planned pregnancy.
- Supportive husband and family.
- Preconception critical use of multivitamins with folate, lifestyle conducive to health.

Areas of Concern

- None identified.

PAGE 842: CRITICAL THINKING ACTIVITY #9

- An overview of the plan for prenatal care.
- Self-care activities, including actions to prevent UTI (proper cleansing after urination, avoidance of bubble baths, adequate hydration, and frequent bladder emptying).

- Routine practice of Kegel's exercises to strengthen and maintain pelvic floor muscle tone.
- Maintaining adequate nutrition.
- Getting adequate rest and exercise.
- Involvement in childbirth preparation/parenthood education classes; development of a birth plan.
- Maintaining intimacy and making necessary sexual adjustments as pregnancy progresses.

PAGE 842: CRITICAL THINKING ACTIVITY #10

Unless problems develop, she will probably need to return to the clinic every 4 weeks until her 28th week of pregnancy, then every 2 weeks until her 36th week of pregnancy, then every week from the 37th week until the time of birth.

PAGE 842: CRITICAL THINKING ACTIVITY #11

The first prenatal visit is usually the most lengthy because you are obtaining the complete medical and obstetric history. On each subsequent visit, you should ask Mrs. Ramirez to describe any relevant events that have occurred since the last visit. Examples of questions you might ask are:

- How have you been doing since you were last here?
- Do you feel the baby move?
- Have you noticed any change in the baby's activity pattern?
- Have you experienced any pain?
- Have you experienced any nausea or vomiting?
- Have you experienced any bleeding?
- Have you noticed any unusual vaginal discharge, itching, or burning or experienced fluid leaking from your vagina?
- Have you experienced any severe back pain?
- Have you had any severe headaches or vision difficulties?
- Have you experienced a severe stomachache?
- Have you noticed any swelling in your hands, face, feet, or lower legs?
- Have you had any pain in your legs?
- Have you had any difficulty breathing?
- Have you had any abdominal pain or contractions?
- Do you have any problems when you urinate?
- Are you having regular bowel movements?
- Are you getting enough rest?
- Have you had any fever or chills?
- Are you getting daily exercise?
- How is your husband/significant other doing?
- Are you having any sexual difficulties?
- Have you made any changes to your birth plan?
- Have you enrolled in prenatal classes?
- Are you taking any medications?
- Is there anything you would like to share with me, or do you have concerns about something that I haven't covered?

Certainly, this list of questions is not exhaustive, but it serves as a guide for the types of questions that you should ask at each prenatal visit.

PAGE 842: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Pregnancy confirmed during this visit. Excited about this first pregnancy, asking about plan for prenatal care. Works full-time and wants to adjust work schedule to accommodate clinic visits.
- Weight loss of 2 lb since becoming pregnant. States that nausea and vomiting mostly occur in the morning on arising. Able to offset nausea and vomiting with diet modification. Urine negative for ketones today.
- States that she feels sleepy, especially in the midafternoon. Has begun going to bed earlier at night. Has a midafternoon snack and brief exercise in the midafternoon to offset fatigue.
- Has obtained a larger bra size; wears bra at all times when awake.
- Discontinued contraception 8 months ago to attempt conception. Began taking multivitamins with folic acid 8 months ago in anticipation of pregnancy. Maintains a healthy diet despite early morning nausea and occasional vomiting. Began prenatal care early in pregnancy. Exercises daily. Wishes to enroll in prenatal classes.

PAGE 842: POSSIBLE ADDITIONAL NURSING DIAGNOSES

- Body Image, disturbed
- Bowel Elimination, altered
- Sexuality Patterns, ineffective
- Adjustment, impaired
- Anxiety
- Infection, risk for
- Family Processes, interrupted
- Parenting, risk for impaired
- Comfort, altered
- Fear
- Coping, ineffective
- Self-Concept, altered

PAGE 847: REVIEW QUESTIONS

1. c. To calculate the EDB, add 7 days to the first day of the LMP, then subtract 3 months. 7 days to 11/30 = 12/7, minus 3 months = 9/7/07.
2. a. Presumptive signs of pregnancy are signs experienced by the patient. Presumptive signs include cessation of menses, nausea and vomiting, frequent urination, breast tenderness, quickening, skin changes, and fatigue.
3. d. Probable signs of pregnancy are signs observed by the examiner. Probable signs include abdominal enlargement, Piskacek's sign, Hegar's sign, Goodell's

- sign, Chadwick's sign, Braxton Hicks contractions, = pregnancy test and ballottement.
4. c. Positive signs of pregnancy attributed only to the presence of the fetus include FHB and visualization of the fetus.
 5. c. A bluish discoloration of the cervix is Chadwick's sign. This is seen at about the 6th to 8th week of pregnancy and is due to venous congestion.
 6. c. Since the patient had normal prepregnancy weight, her expected weight gain during pregnancy should be 25 to 35 lb.
 7. a. Lying supine may compress the aorta and vena cava, causing the patient to feel dizzy. Place the patient on her left side to relieve the pressure and improve flow.
 8. c. The normal FHB is 120 to 160 BPM. A rate less than 100, greater than 160, or a sudden change of plus or minus 20 BPM is cause for concern.
 9. b. A fetal Doppler can usually detect FHB by 10 to 12 weeks gestation.
 10. b. Leopold's maneuver is a technique used to determine the position of the fetus.

CHAPTER 24

PAGE 875: CRITICAL THINKING ACTIVITY #1

During your physical examination, invite the parents to watch as you assess their baby. Take this opportunity to describe and explain findings. Include infant care such as handling, bathing, diapering, and safety precautions in your discussion.

Contact the hospital's patient education services regarding parenting and breastfeeding classes. Offer audiovisual tapes and printed matter. Refer parents to a lactation consultant.

PAGE 875: CRITICAL THINKING ACTIVITY #2

Newborns should awaken and be alert for their feedings every 3 to 4 hours. Unarousable or lethargic behavior may indicate jaundice or infection, and medical advice should be sought.

PAGE 878: CASE STUDY EVALUATION

Health History

- Developmental.
- Sits.
- Bears weight on hands, stands with assistance.
- Rolls over.
- Controls head.
- Grasps objects.
- Is teething.
- Imitates sounds.
- Is attached to parents.
- Has separation anxiety.

- Safety
 - Smoke and carbon monoxide detectors.
 - Safety gates.
 - Cabinet locks.
 - Electrical outlet caps.
 - Car seats.
 - Medications and cleaning solutions out of reach.
- Immunizations
 - Hepatitis B 1 and 2.
 - DPT 1 and 2.
 - Polio 1 and 2.
 - HiB 1 and 2.
 - PCV 1 and 2.

Review of Systems

- **General Health Survey:** Very robust and active infant, healthy appearance.
- **Body Weight:** 16 lb, gained 2 lb since last month's visit for otitis media.
- **Integumentary:** No scaly skin.
- **Head and Neck:** Easily holds head upright and in alignment.
- **Eyes, Nose, and Throat:** Eyes: Normal response to visual stimuli, no drainage.
- **Ears:** Normal sound recognition and making sounds. No apparent abnormalities. Otolaryngology examination indicated because of recent otitis media.
- **Respiratory:** No breathing problems.
- **Cardiovascular:** No apparent distress. Active when awake.
- **Gastrointestinal:** Exclusively breastfed and nurses every 3 to 4 hours during the day, with daily bowel movements, yellow in color.
- **Genitourinary:** Normal urination pattern noted; eight diapers a day.
- **Musculoskeletal:** Attempts to crawl; rolled over and sat up on her own (normal findings).
- **Neurological:** No abnormal movements, nurses well, no problems.
- **Immunologic:** No infection aside from otitis media.
- **Developmental:** Laughs easily with mother but is fearful of examiner. Normal age-related behavior observed.
- **Relationships:** Warm, caring responses noted toward baby from parents. Both parents say they enjoy parenthood very much.

PAGE 884: CRITICAL THINKING ACTIVITY #3

Suggest that Mrs. Ramirez tap into other possible support systems, such as her other sisters, friends, neighbors, fellow workers, and members of her religious community. Community child care resources that may be of assistance to the Ramirez family include:

- On-site job child care center.
- Child care centers connected with places of worship.

- Accredited child care centers listed with the health department or local social service department.

PAGE 884: KEY HISTORY FINDINGS

- Otitis media—one episode last month.
- Family members who cared for infant during mother's working hours have moved to another state, so mother has lost child care/emotional supports.
- Mother concerned that infant is underweight.

PAGE 884: KEY PHYSICAL EXAMINATION FINDINGS

- Diaper rash on labia and buttocks.
- Infant appears well nourished and is normal weight.
- Developmentally on target.

PAGE 884: NURSING DIAGNOSES SUPPORTING DATA

- Irritating diaper rash noted on labia and buttocks. Mother eager to help condition.
- Otitis media—one episode last month.
- Mother's concern about Carmen's weight. Cultural interest toward large babies as healthy babies. Mother's lack of knowledge related to breastfeeding recommendation.
- Depended on family for emotional support and child care. Mother works full time. Husband supportive but works full time. No family or friends with children living locally.
- Mother eager to seek help and explore neighborhood resources for new friendships and child care.

PAGE 884: ADDITIONAL NURSING DIAGNOSES

- Aspiration, risk for
- Infection, risk for
- Injury, risk for
- Poisoning, risk for
- Suffocation, risk for
- Trauma, risk for
- Pain
- Parenting, impaired

PAGE 885: REVIEW QUESTIONS

- a. Acrocyanosis is peripheral cyanosis caused by thermoregulation adjustments and is normal for the first 12 hours after birth.
- c. An Apgar score assesses heart rate, respirations, color, muscle tone, and reflex irritability. This score is obtained at 1 minute and 5 minutes after birth. A score of 8 to 10 is excellent, 4 to 7 guarded, and 0 to 3 critical.
- d. Jaundice that is present within the first 24 hours of birth is usually pathological jaundice where as physiological jaundice may be seen after 24 hours of birth.

- a. The normal heart rate for a newborn in 120 to 160 BPM. A rate of 100 BPM warrants follow-up.
- a. Harlequin sign is a color change that is associated with position change.
- d. The anterior and posterior fontanelles are open at birth. The anterior fontanel usually closes between 12 and 18 months, and the posterior fontanel usually closes by 3 months.
- b. The glabellar reflex is assessed by gently tapping on the infant's forehead. The normal response to this reflex is blinking for the first few taps. Persistent blinking with repeated taps may indicate an extrapyramidal problem
- c. The magnet reflex is assessed by flexing the newborn's legs and applying pressure to the soles of the feet. The normal response should be extension of the legs. A breech birth or spinal cord problem may diminish the reflex.
- a. Ortolani's maneuver is a test for congenital hip dislocation. The infant's legs are flexed to the abdomen and then abducted and rotated. A clicking sound during the maneuver is a positive sign for hip dislocation.
- b. The normal breath sounds off a newborn are more bronchial owing to the size of the chest and airways.

CHAPTER 25

PAGE 897: CRITICAL THINKING ACTIVITY #1

The assessment is essentially the same as with other age groups except for the need to focus on growth and developmental changes that occur during the toddler/preschool years—for example, height and weight, body proportions (head), primary teeth, musculoskeletal and neurological development, and the struggle to develop a sense of autonomy.

PAGE 897: CRITICAL THINKING ACTIVITY #2

- Answer questions honestly.
- Use concrete terms.
- Explain what you are doing.
- Involve the parent or caregiver.
- Use toys or games (e.g., puppets).

PAGE 897: CRITICAL THINKING ACTIVITY #3

- Safety issues (injuries and poisonings).
- Nutrition.
- Toilet training.
- Sibling rivalry.
- Temper tantrums.
- Sleep problems.
- Communicable diseases.
- Abuse.

PAGE 897: CRITICAL THINKING ACTIVITY #4

For Child

- Anxiety, related to starting kindergarten.
- Anxiety, related to separation from parents
- Injury, risk for
- Nutrition: imbalanced, risk for
- Infection, risk for

For Parents

- Knowledge, deficient, related to dietary needs
- Knowledge, deficient, related to safety education

PAGE 898: REVIEW QUESTIONS

1. b. 20/20 vision is usually attained by school age. A 4-year-old's far vision is usually about 20/40.
2. a. Toddlers as they are learning to walk often have problems with balance. To maintain balance, the toddler will widen his base of support to prevent from himself from falling.
3. c. Toddlers with "potbellies" have a shift in their center of gravity. To compensate for this, you often see an accentuated lumbar curve, lordosis.
4. b. Bowel control is usually achieved by 3 years of age.
5. d. Full bladder control is usually achieved by 5 years of age.
6. a. When a child becomes ill or is hospitalized, it is not uncommon to see the child regress developmentally.
7. b. According to Erikson, the developmental task for the 1- to 3-year-old is autonomy vs. shame. The child should gain bowel and bladder control, tolerate separation from parents, differentiate self from others, develop socially acceptable behavior, develop verbal communication skills, become less egocentric, and tolerate delays in gratification.
8. c. According to Erikson, the developmental task for the 3- to 6-year-old is initiative vs. guilt. This is a time of playing and learning. The superego, or conscience, is developing.

CHAPTER 26

PAGE 911: CRITICAL THINKING ACTIVITY #1

The assessment is essentially the same as for other age groups, except for the need to focus on growth and developmental changes that occur during the school-age years—for example, height and weight, body proportions (head), secondary teeth, musculoskeletal and neurological development, and learning to work and play cooperatively.

PAGE 912: CRITICAL THINKING ACTIVITY #2

- Answer questions honestly,
- Be concrete and explain what you are doing.

PAGE 912: CRITICAL THINKING ACTIVITY #3

- Safety issues.
- Nutrition.
- Learning problems.
- Alcohol and drug abuse.
- Behavior problems (lying, cheating, and stealing).

PAGE 912: CRITICAL THINKING ACTIVITY #4

- Anxiety, related to starting school
- Injury, risk for
- Nutrition: imbalanced, risk for
- Knowledge, deficient, related to dietary needs
- Knowledge, deficient, related to sex education
- Knowledge, deficient, related to substance abuse

PAGE 924: CRITICAL THINKING ACTIVITY #5

The assessment is essentially the same as for other age groups, except for the need to focus on growth and developmental changes that occur during adolescence—for example, height and weight, sexual maturation, musculoskeletal development, and separating from parents and moving toward self-autonomy.

PAGE 924: CRITICAL THINKING ACTIVITY #6

- Listen actively.
- Answer questions honestly.
- Be nonjudgmental.
- Avoid scare tactics.
- Maintain confidentiality.
- Explain what you are doing.

PAGE 924: CRITICAL THINKING ACTIVITY #7

- Eating disorders.
- Pregnancy.
- STDs.
- Emotional problems.
- Substance abuse.
- Violence.
- Injuries and accidents.

PAGE 924: CRITICAL THINKING ACTIVITY #8

- Risk for identity conflict related to sexual maturation
- Nutrition: imbalanced, more or less than body requirements
- Sleep Pattern, disturbed, risk for
- Knowledge, deficient, related to safe sex practices
- Knowledge, deficient, related to sexual maturation

PAGE 925: REVIEW QUESTIONS

1. b. Myopia, nearsightedness, is often seen during adolescence.
2. c. Many hormonal changes occur during puberty. The thyroid gland may be palpable during this time owing to the increased activity.

3. a. According to Erikson, the developmental task for the 6- to 12-year-old is industry vs. inferiority. During this stage, the child becomes more independent, developing skills and competencies. Social skills develop as the child learns to work with others. Peer acceptance becomes very important. Sexual roles develop during this time.
4. d. According to Erikson, the developmental task for the 12- to 18-year-old is identity vs. role diffusion. During this time, the child establishes own identity and self-worth. The child's peers become very important in an attempt to belong and be accepted.
5. c. Scoliosis is a common musculoskeletal problem among adolescent girls. Screening for scoliosis is routinely done in middle school and high school.
6. d. Since the risk of meningitis is greater among college students, prior to college, the adolescent should receive the meningitis vaccine.
7. b. During puberty, it is not uncommon for boys to develop gynecomastia. This usually resolves on its own once the child is through puberty.

CHAPTER 27

PAGE 932: CRITICAL THINKING ACTIVITY #1

In cultures in which older people are viewed with respect and as valued members of the family and society, healthcare is readily available. If the older person lives with her or his family, the family can be an important factor in the healthcare planning for the person. However, in cultures in which older people are perceived as a burden, healthcare for them may not be a priority.

PAGE 948: CRITICAL THINKING ACTIVITY #2

- Liver: Decreased drug metabolism.
- GI tract: Decreased absorption of medications.
- Renal system: Decreased creatinine clearance may result in decreased drug clearance. Because of these changes, dosages of medications may need to be adjusted.

PAGE 948: CRITICAL THINKING ACTIVITY #3

- Decreased GI tract motility.
- Increased food intolerance.
- Decreased dentin (may cause loss of teeth).
- Decreased papillae on tongue (causes decreased taste).
- Delayed stomach emptying (causes decreased hunger sensation).

PAGE 948: CRITICAL THINKING ACTIVITY #4

Strengths

- Independent.
- Supportive family.

- Self-referral for healthcare.
- Socially active.
- Exercises routinely.

Areas of Concern

- Fatigue.
- Decreased appetite, activity tolerance, and muscle strength.
- History of breast cancer, HTN, DJD.
- Pain.
- History of falls.

PAGE 948: CASE STUDY EVALUATION

Biographical Data

- 80-year-old widow with four daughters.
- Widowed for 15 years.
- Resident of continuing care retirement community for 8 years.
- Catholic religion, Slavic ancestry.
- Not referred by doctor—came to outpatient office of her own accord.

Current Health Status

- Complains of vague fatigue all day, decreased appetite, and inability to engage in usual activities such as walking group. Fatigue has been going on for several weeks and has gotten worse in the past week, despite going to bed around 9 P.M. and getting up at 8 A.M. Is forcing herself to participate in social activities such as going to the dining room with friends for dinner. However, has no desire to eat.
- Takes Lasix 20 mg PO qd for HTN and Tamoxifen 20 mg bid to prevent recurrence of breast cancer.

Past Health History

- Breast cancer 20 years ago.
- HTN controlled by diuretics.
- DJD, mostly in hips.
- Cataract surgery 12 years ago (right eye) and 11 years ago (left eye).
- Lumpectomy, right breast, 23 years ago.
- Elective left total hip replacement 17 years ago.

Family History

- No known history of cognitive problems.
- Believes that both of her parents died of heart disease in their 70s and 80s.
- Does not know of any family history of cancer.

Review of Systems

- **General Health Survey:** Complains of fatigue and “not feeling ‘like myself.’” No changes in height or weight reported.

- **Integumentary:** Fair skin with no abnormal lesions. Complains of dry skin in winter.
- **HEENT:** No vision changes, dizziness, changes in smell or taste, or difficulty swallowing. Wears corrective lenses; has yearly eye exam. Reports being satisfied with quality of hearing.
- **Respiratory:** No cough or SOB.
- **Cardiovascular:** No chest pain or SOB during the day or at night. No SOB while walking to the dining room. No leg cramping or pain either at rest or walking. Some swelling in ankles on arising, usually disappears in a few hours.
- **Gastrointestinal:** No indigestion or burning. Has a regular bowel movement daily and takes MOM rarely, only if she has not had a bowel movement in several days. Has not felt much like eating, but has not had stomach pain, nausea, or vomiting.
- **Genitourinary:** No discharge. Loses urine when she coughs, sneezes, or laughs, so wears a sanitary pad, which she changes two or three times a day. This problem has not gotten any worse recently. No vaginal discharge or bleeding. Menopause at age 51; was never on hormone replacement therapy. Last Pap test about 14 years ago; last mammogram 3 years ago.
- **Neurological:** No dizziness, change in cognition, problems with balance, tremor, seizure, or difficulty with concentration. Is very happy at continuing care retirement community and feels strongly that this was the best move for her because she could no longer manage her home independently. Enjoys living near friends, participating in activities, and having her daughters visit. Is glad they do not have to take care of her or worry about her being home alone. No previous history of depression.
- **Musculoskeletal:** Mild morning stiffness, usually gone by the time she has breakfast. Takes Tylenol when pain is bothersome. Had two falls since moving into the facility, but no falls over past 6 months. No increase in joint pain or warmth or swelling of joints.
- **Immunologic/Hematologic:** Reports feeling more tired than usual.
- **Endocrine:** No weight changes, increased thirst or urination. Heart does not flutter, race, or skip beats.

Psychosocial Profile

- Typical day starts around 8 A.M. She fixes herself breakfast in her efficiency apartment, feeds her cat, then has coffee and watches TV. Does laundry, light housecleaning, and “putters around” apartment. Midmorning usually goes to the activity room to socialize or play cards with other residents. Fixes own lunch. Until recently, walked 20 minutes three times a week with walking group in afternoon. Goes on planned outings with other residents. Sees daughters every weekend for lunch or dinner, shop-

ping, and mass on Sunday. Previously went to bed at 11 P.M. but has been retiring at 9 P.M. for past few weeks because of fatigue.

- Does not smoke and rarely has a glass of wine socially.

Functional Assessment

- Performs all personal care independently. Does own laundry, shopping, finances, and cleaning aside from the vacuuming, which is done by the facility every 2 weeks.
- Walks independently with a rolling walker. Has been participating in walking group and until recently was walking for 20 minutes three times per week.

PAGE 958: LIST KEY HISTORY FINDINGS

- Age 80, widow.
- Lives in retirement community.
- Supportive family.
- Self-referral for healthcare.
- Independent.
- Fatigue for several weeks.
- Decreased appetite.
- Decreased activity tolerance.
- History of HTN, breast cancer, DJD with total hip replacement.
- Medications: Tylenol prn for joint pain, Lasix, Tamoxifen.
- No bowel movement for several days.
- No abdominal pain, nausea, or vomiting.
- Increase in sleep from 9 to 12 hours.
- Uses walker to ambulate.
- Usually walks 20 minutes three times a week.
- History of falls.

PAGE 958: LIST KEY PHYSICAL EXAMINATION FINDINGS

- Well groomed, pleasant, appropriate behavior.
- AAO X 3, memory intact.
- Vital signs within normal limits, afebrile.
- Wears glasses.
- Abdomen: Positive tenderness RUQ, positive palpable liver edge.
- Decreased muscle strength in lower extremities, decreased ROM.
- Gait steady with walker and wide base of support.
- Kyphosis.
- No evidence of depression.

PAGE 958: NURSING DIAGNOSES CLUSTERED SUPPORTING DATA

- Reports loss of appetite; does not feel like eating.
- DJD; history of falls; uses assistive device (walker).
- Complains of fatigue; unable to participate in walking program.

- Complains of having no energy; unable to participate in usual activities.


PAGE 958: ADDITIONAL NURSING DIAGNOSES

- Pain, Chronic, related to joint stiffness
- Incontinence, Stress Urinary, related to relaxed pelvic muscles
- Fluid Volume, Risk for Deficient, related to use of diuretics and loss of appetite
- Skin Integrity, Risk for Impaired, related to stress incontinence

PAGE 960: REVIEW QUESTIONS

1. c. The nails of an older adult may become thick and brittle. This is a normal variation in the nails associated with aging.
2. b. Arcus senilis, a white ring around the edge of the cornea, is a common, normal finding in older adult.

3. b. Decreased skin turgor due to a loss of elasticity is a common, normal finding in older adults.
4. c. The pupils normally decrease in size with age.
5. a. Kyphosis, an accentuated thoracic curve, is a common finding in older adults.
6. a. Difficulty in hearing high-pitched sounds is a common finding in older adults.
7. a. The vaginal walls and cervix are normally pale pink in postmenopausal women.
8. b. The ovaries are not normally palpable in postmenopausal women. Palpable ovaries in a postmenopausal women would warrant follow-up evaluation.
9. a. Stress incontinence is characterized by lack of bladder control during coughing, sneezing, or laughing.
10. c. A loss of height about 6 to 10 cm occurs with aging owing to a narrowing of the intervertebral discs.



APPENDIX


B

Growth Charts

Birth to 36 months: Boys
Length-for-age and Weight-for-age percentiles

NAME _____

RECORD # _____




Published May 30, 2000 (modified 4/20/01).
 SOURCE: Developed by the National Center for Health Statistics in collaboration with the National Center for Chronic Disease Prevention and Health Promotion (2000).
<http://www.cdc.gov/growthcharts>



Birth to 36 months: Boys
Head circumference-for-age and
Weight-for-length percentiles

NAME _____

RECORD # _____



Published May 30, 2000 (modified 10/16/00).


SOURCE: Developed by the National Center for Health Statistics in collaboration with the National Center for Chronic Disease Prevention and Health Promotion (2000).
<http://www.cdc.gov/growthcharts>



Birth to 36 months: Girls
Length-for-age and Weight-for-age percentiles

NAME _____

RECORD # _____




Published May 30, 2000 (modified 4/20/01).
 SOURCE: Developed by the National Center for Health Statistics in collaboration with the National Center for Chronic Disease Prevention and Health Promotion (2000).
<http://www.cdc.gov/growthcharts>



Birth to 36 months: Girls
Head circumference-for-age and
Weight-for-length percentiles

NAME _____

RECORD # _____




Published May 30, 2000 (modified 10/16/00).

SOURCE: Developed by the National Center for Health Statistics in collaboration with the National Center for Chronic Disease Prevention and Health Promotion (2000).
<http://www.cdc.gov/growthcharts>



SAFER • HEALTHIER • PEOPLE™



APPENDIX

C

Guidelines

Guidelines for Cultural Assessment*

According to Purnell and Paulanka (2003), culture is the “totality of socially transmitted behavioral patterns, arts, beliefs, values, customs, lifeways, and all other products of human work and thought characteristics of a population of people that guide their worldview and decision making.”

Because cultural background encompasses every aspect of a person’s life, you need to be aware of how it influences your patient’s health and wellness. Consider the following areas when determining the influence of culture on your patient’s health.

Overview

- Inhabited localities and topography
 - In what part of the world does this person’s cultural or ethnic group originate?
 - What are the climate and topography there?
- Heritage and residence
 - Where does this person’s cultural or ethnic group reside now?
 - What is the approximate population of the group here?
- Reason for migration and associated economic factors
 - What were the major factors that motivated this person’s cultural/ethnic group to emigrate?
 - What were the economic or political factors that influenced this group’s acculturation and professional development in America?
- Educational status and occupations
 - What value does the person’s cultural group place on education?
 - What are the predominant occupations of the group’s members?

Communication

- Dominant language and dialects
 - What is the dominant language of the group?
 - Does the person use that language or a dialect that may interfere with communication?
 - Are there specific contextual speech patterns for this group? If so, what are they?
 - What is the usual volume and tone of speech?
- Cultural communication patterns
 - Is the person willing to share thoughts, feelings, and ideas?
 - What is the practice and meaning of touch in the person’s society? With family, friends, strangers, same sex, opposite sex, and healthcare providers?

- What are the personal spatial and distancing characteristics when communicating one-to-one? With friends versus strangers?
- Does this group use eye contact? Does avoidance of eye contact have special meaning? Is eye contact influenced by socioeconomic status?
- Do various facial expressions have specific meanings? Are facial expressions used to express emotions?
- Are there acceptable ways of standing and greeting outsiders? If so, what are they?
- Temporal relationships
 - Are people primarily past, present, or future oriented? How do they see the context of past, present, and future?
 - Are there differences in interpretation of social time versus clock time? If so, what are they?
 - Are people expected to be punctual in terms of jobs, appointments, and social engagements?
- Format and names
 - What is the format for a person’s name?
 - How does one expect to be greeted by strangers and healthcare practitioners?

Family Roles and Organization

- Head of household and gender roles
 - Who is the perceived head of household?
 - How does this role change during different developmental aspects of life?
 - What are the gender-related roles of men and women in the family system?
- Prescriptive, restrictive, and taboo behaviors
 - What are the prescriptive, restrictive, and taboo behaviors for children?
 - What are the prescriptive, restrictive, and taboo behaviors for adults?
- Family roles and priorities
 - What family goals and priorities are emphasized by this culture?
 - What are the developmental tasks of this group?
 - What are the status and role of older people in the family?
 - What are the roles and importance of extended family members?
 - How does one gain social status in this cultural system? Is there a caste system?
- Alternative lifestyles
 - How are alternative lifestyles and nontraditional families viewed by the society?

Workforce Issues

- Culture in the workplace
 - Are workforce issues, such as education, affected by immigration? If so, how?

*Source: Adapted from Purnell & Paulanka, 2003.

- What are the specific multicultural considerations when working with this culturally diverse person or group?
- What factors influence patterns of acculturation in this cultural group?
- How does the person's or group's healthcare practices influence the workforce?
- Issues related to autonomy
 - What are the cultural issues related to professional autonomy, superior or subordinate control, religion, and gender in the workforce?
 - Are there language barriers? For example, do people sometimes misunderstand English expressions by interpreting them concretely?

Biocultural Ecology

- Skin color and biological variations
 - Are there skin color and physical variations for this group? If so, what are they?
 - What special problems or concerns might the skin color pose for healthcare practitioners?
 - What are the biological variations in body habitus or structure?
- Diseases and health conditions
 - What are the risk factors for people related to topography or climate?
 - Are there any hereditary or genetic diseases or conditions that are common with this group? If so, what are they?
 - Are there any endemic diseases specific to this cultural or ethnic group? If so, what are they?
 - Are there any diseases or health conditions for which this group has increased susceptibility? If so, what are they?
- Variations in drug metabolism
 - Does this group have any specific variations in drug metabolism, drug interactions, and related side effects? If so, what are they?

High-Risk Behaviors

- Are any high-risk behaviors common in this group? If so, what are they?
- What are the patterns of use of alcohol, tobacco, recreational drugs, and other substances in this group?

Healthcare Practices

- What are typical health-seeking behaviors for this group?
- What is this group's usual level of physical activity?
- Do people use safety measures, such as seat belts?

Nutrition

- Meaning of food
 - What does food mean to this group?

- Common foods and food rituals
 - What specific foods, preparation practices, and major ingredients are commonly used by this group?
 - Are there any specific food rituals for this group? If so, what are they?
- Dietary practices for health promotion
 - Are enzyme deficiencies or food intolerances commonly experienced by this group? If so, what are they?
 - Are large-scale or significant nutritional deficiencies experienced by this group? If so, what are they?
 - Are there native food limitations in America that may cause special health difficulties? If so, what are they?

Pregnancy and Childbearing Practices

- Fertility practices and views regarding pregnancy
 - What are the cultural views and practices related to fertility control?
 - What are the cultural views and practices regarding pregnancy?
- Prescriptive, restrictive, and taboo practices in the childbearing family
 - What are the prescriptive, restrictive, and taboo practices related to pregnancy, such as food, exercise, intercourse, and avoidance of weather-related conditions?
 - What are the prescriptive, restrictive, and taboo practices related to the birthing process, such as reactions during labor, presence of men, position of mother for delivery, preferred types of health practitioners, or place of delivery?
 - What are the prescriptive, restrictive, and taboo practices related to the postpartum period, such as bathing, cord care, exercise, food, and roles of men?

Death Rituals

- Death rituals and expectations
 - Are there culturally specific death rituals and expectations? If so, what are they?
 - What is the purpose of the death rituals and mourning practices?
 - What specific practices (e.g., cremation) are used for disposal of the body?
- Responses to death and grief
 - How are people expected to show grief and respond to the death of a family member?
 - What is the meaning of death, dying, and afterlife?

Spirituality

- Religious practices and use of prayer
 - How does the dominant religion of this group influence healthcare practices?

- Are there activities such as prayer, meditation, or symbols that help people reach fulfillment? If so, what are they?
- Meaning of life and individual sources of strength
 - What gives meaning to people's lives?
 - What is the person's source of strength?
- Spiritual beliefs and healthcare practices
 - What is the relationship between spiritual beliefs and healthcare practices?

Healthcare Practices

- Health-seeking beliefs and behaviors
 - What predominant beliefs influence healthcare practices?
 - What is the influence of health promotion and prevention practices?
- Responsibility for healthcare
 - Is the focus of acute care practice curative or fatalistic?
 - Who assumes responsibility for healthcare in this culture?
 - What is the role of health insurance in this culture?
 - What are the behaviors associated with the use of over-the-counter (OTC) medications?
- Folklore practices
 - How do magicoreligious beliefs, folklore, and traditional beliefs influence healthcare behaviors?
- Barriers to healthcare
 - Are there barriers to healthcare such as language, economics, and geography for this group? If so, what are they?
- Cultural responses to health and illness
 - Are there cultural beliefs and responses to pain that influence interventions? If so, what are they?
 - Does pain have special meaning?
 - What are the beliefs and views about mental and physical illness in this culture?
 - Does this culture view mental handicaps differently from physical handicaps?
 - What are the cultural beliefs and practices related to chronicity and rehabilitation?
 - Are there any restrictions to the acceptance of blood transfusions, organ donation, and organ transplantation for this group? If so, what are they?

Healthcare Practitioners

- Traditional versus biomedical care
 - What are the roles of traditional, folklore, and magicoreligious practitioners, and how do they influence health practitioners?
 - How does this culture feel about healthcare practitioners providing care to patients of the opposite sex?
 - Does the age of the practitioner make a difference? If so, what?

- Status of healthcare providers
 - How does this culture feel about healthcare providers?
 - What is the status of healthcare providers in this culture?
 - How do different healthcare practitioners in this culture view each other?

Example: Cuban Americans

Overview

- Cubans are of Hispanic origin.
- Culture influenced by Spain, United States, and Soviet Union.
- 860,000 in the United States, mainly in New York and Florida.
- Cuban climate tropical.
- Political and economic reasons for immigration.
- Educational attainment higher than that of other Hispanic groups.
- High proportion self-employed.
- High proportion in industry, such as wholesale/retail trade, banking/credit, insurance, and real estate.

Communication

- Dominant language is Spanish.
- Values *simpatico*—smooth interpersonal relationships, courtesy, respect without harsh criticism.
- *Personalismo*—interpersonal relationships more important than bureaucratic relationships.
- *Choteo*—light-hearted attitude with teasing and bantering.
- Animated facial expressions, eye contact, hand gestures, speech loud and fast.
- Touching in form of handshakes and hugs.
- More present oriented, deal with current issues, use a flexible time frame, which may make appointment setting difficult.
- Feel a sense of “specialness” about themselves and culture and convey this in communication.
- Use both mother's and father's surnames. For example, Regina Morales Colon, in which Morales is the patriarchal surname and Colon the matriarchal. If Regina marries Mr. Ordonez, she would drop the matriarchal name and add her husband's name, becoming Regina Morales de Ordonez.

Family Roles and Organization

- Family is most important unit.
- Patriarchal, male dominant, and aggressive; female dependent.
- Double standard applied to male and female behavior.
- Children are pampered and protected.
- Children are expected to respect parents, follow *el buen camino*—the straight and narrow.
- Expectations for boys and girls are different.

- Families are tightly knit.
- Family is most important source of emotional and physical support
- *Compaadrazgo* (godparents) are considered part of family.
- High proportion of older persons live with family members.
- High proportion of women are divorced.
- Gay lifestyle is contradictory to prevailing machismo image.

Workforce Issues

- Have been economically successful in United States, with 57 percent employed in professional and technical jobs.
- Many own businesses.
- Tend to be lineal in relationships at work, rather than collegial.
- Carry concept of *personalismo* through to work.
- May speak Spanish when talking with Cuban friends and family and speak English professionally.

Biocultural Ecology

- Skin and hair vary from light to dark.
- Slightly higher incidence of obesity than other Hispanic groups.
- Lower incidence of hypertension than whites and African Americans.
- High incidence of dental problems may be a result of diet high in sugars and starches.
- 42 percent of Cuban men smoke.
- Cuban American women in 30s have higher incidence of smoking than national average.
- High mortality rates among adolescents and young adult because of violence.
- Suicide rates higher than white non-Hispanics.
- A little heaviness seen as attractive. Diets are high in calories, starches, and saturated fats.
- Have high levels of preventive health behaviors.

Nutrition

- Food has powerful social meaning—promotes sense of community, perpetuates customs and heritage.
- Root crops, such as yams, yucca, malanga, boniato, plantains, and grains, are staples.
- Main course is meat, pork, or chicken.
- Diet high in calories, starches, and saturated fats.
- Leisurely noon meal and late dinner.
- Diet lacking in leafy green vegetables (fiber).

Pregnancy and Childbearing Practices

- Low rate of reproduction.
- Many folk beliefs about pregnancy. For example, believe you need to eat for two, which leads to excessive weight gain during pregnancy.

- Childbirth is time of celebration.
- Traditionally, men do not participate in birthing.
- Family members and friends often care for the mother and baby for 4 weeks postpartum.
- Breastfeeding considered better than bottle. Child is weaned early at 3 months and introduced to solid foods.
- Believe that fat child is healthy child.
- Believe that breastfeeding can lead to deformity or asymmetry of breasts.

Death Rituals

- Large gathering of family at deathbed.
- Death rite may include animal sacrifice or *santería*.
- Candles used after death to light path to afterlife.
- Wake (*velorio*) lasts 2 to 3 days until funeral.
- Bereavement is expressed openly.
- Relatives light candles, pray, and visit gravesite on anniversary of death

Spirituality

- 85 percent Catholic.
- Devoted to Blessed Virgin Mary, Jesus, and saints.
- Pictures of saints (*estampitas*) may be placed under pillow or beside bed of someone who is ill.
- Formal religious observance is an important source of support.
- Family is most important source of strength, identity, and emotional security.
- Take fatalistic view, feeling they have no control over circumstances influencing life.
- Practice folk remedies, most of which are harmless.

Healthcare Practices

- Family is primary source of health advice, usually older woman in the family.
- Poverty may be a barrier to healthcare.
- Herbal teas used to treat common ailments.
- Loneliness, depression, anger, and anxiety may occur from losses caused by migrating from homeland.
- Dependency acceptable with sick role.
- Blood transfusions acceptable.
- Use both traditional and biomedical healthcare.
- Underrepresented in healthcare occupations.

Guidelines for Environmental Assessment*

Shorter stays in acute care settings have increased the need for nursing care at home. Home care nursing can occur at both the primary and the tertiary levels. At the primary care level, you will make postpartum visits to new mothers and babies. At the tertiary level, you will make follow-up visits to patients discharged from the hospital.

Assessing the Home

In the home, your assessment is based on the health history and physical exam findings. You need to assess your patient's response to the treatment plan and also identify any risk factors in his or her environment that may affect his or her health and well-being. Remember, the treatment plan established in the hospital will be

effective only if your patient is able to follow it at home.

Begin by determining if your patient is able to perform activities of daily living (ADLs); then assess basic needs and environmental safety hazards. Also assess support systems and self-esteem or self-actualization. Keep in mind that financial status and religious and cultural beliefs will influence health beliefs and practices.

Assessing the Home

AREAS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

PHYSICAL NEEDS

Food

- What does the patient eat and drink?
- Who prepares food? Who buys food?
- Is food being stored properly?
- Is kitchen accessible? Clean? Are appliances in good operating condition?
- If you detect a nutritional deficit, does patient's illness have an effect on appetite?
- Is there an unexplained weight loss or gain?
- Does the patient drink alcohol?
- Does she or he have a dental problem?
- Are financial hardships limiting purchase of foods?
- Is the patient taking multiple medicines?
- Does he or she eat alone?
- Does she or he need assistance in eating?

Elimination

- Is the bathroom accessible?
- Does patient need a commode?
- Does he or she need a raised toilet seat? Grab bars?

Bathing

- Does patient take baths, showers, or sponge baths?
- Does she or he care for her or his hair and teeth?
- Does he or she need assistance in bathing? Does he or she have a shower chair?
- Does the bathtub have grab bars? Rubber mats? Nonskid tiles?

Dressing

- Does patient have clean clothes?
- Do clothes fit?
- Does she or he need assistance with dressing?
- Do shoes fit properly?
- Who does the laundry?

Sleep

- Where does the patient sleep?
- How much time does she or he spend in bed?
- Does the patient need a special mattress or bed?
- Are bedrails needed?

Identifies

- Need for referrals for assistance, such as financial support, Meals on Wheels, home health aides.
- Self-care deficits.
- Teaching needs.
- Nutritional deficits or problems affecting nutritional status.

Identifies

- Need for assistive devices.
- Self-care deficits.
- Risk for falls.
- Risk for incontinence problems

Identifies

- Need for assistive devices.
- Self-care deficits.
- Risk for falls.

Identifies

- Self-care deficits.
- Risks for falls or skin breakdown.
- Need for home health aide.

Identifies

- Risk for skin breakdown.
- Need for special equipment.
- Safety issues.

(continued)

Assessing the Home (continued)

AREAS/QUESTIONS TO ASK

RATIONALE/SIGNIFICANCE

- How far is the bed from the bathroom? From other family members?

- Is there privacy for members in household?

Medications

- Is patient able to take own medications as prescribed?
- Is he or she taking any OTC medications, vitamins, or herbal supplements?
- Does she or he need medications pre-poured?

- Does he or she have any impairments that would prevent him or her from self-administering medications safely (e.g., cognitive or visual impairments)?

- Are medications safely stored?

- If using syringes, how does patient dispose of them? Obtain supplies?

- What is the name of patient's pharmacy? Her or his medication insurance plan?

- Does patient have the finances to pay for medications and treatment?

- Does patient understand what medications he or she is taking and their purpose?

Shelter

- Is home maintained and clean?

- Who is responsible for managing and cleaning it?

- Are plumbing and sewage systems working properly?

- What type of heating is there (gas, electric, oil, wood)? Central heating or space heating? Is heating system working properly?

- Is there air conditioning? Fans?

- Is ventilation adequate? Do windows open and close easily and completely?

- What type of insulation is there?

- Is home in need of repair (e.g., peeling paint or cracks in foundation or windows)?

- Is there evidence of insect or rodent infestation?

- Has home been tested for radon?

ASSESSING ENVIRONMENTAL SAFETY

Mobility/Fall Prevention

- Is patient able to walk? Is her or his gait steady?

- Does patient use assistive devices correctly?

- Do devices fit through pathways and doorways?

- Is house one level or more? Are there elevators or stairs?

- Can patient enter and exit home without difficulty?

- Are pathways and stairs clear?

- Are there throw rugs?

- Are there sturdy handrails on the stairs? Are first and last steps clearly marked?

- Are floors slippery or uneven?

- Is there adequate lighting in hallways, stairs, and path to bathroom?

- Is there need for restraints? If yes, what type and when?

- Are carpets in good repair without tears?

Identifies

- Compliance with medical treatment plan and reasons for noncompliance.

- Problems associated with polypharmacy (e.g., side effects, drug interactions).

- Safety issue surrounding medication preparation and administration.

- Can patient open medication containers?

- Need for assistive devices.

- Need for referral or visiting nurse.

- Financial needs for obtaining prescriptions.

- Teaching needs.

Identifies

- Safety issues.

- Need for referrals.

- Need for assistance with home management and maintenance.

Identifies

- Risk for fall/injury.

- Need for assistive devices.

- Need for referrals.

- Need for teaching.

AREAS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Can patient walk on carpet? Does he or she have to hold onto furniture to maintain balance? If yes, is furniture sturdy and stable enough to provide support? ■ Are chairs sturdy and stable? ■ Are there any cords or wires that may present a tripping hazard? 	
<p>Fire/Burn Prevention</p> <ul style="list-style-type: none"> ■ Are there working smoke detectors on each floor? ■ Carbon monoxide detectors? Fire extinguisher? ■ Is there an escape plan in case of fire? ■ Are wires, plugs, and electrical equipment in good working condition? ■ Is patient using a heating pad or portable heaters safely? ■ Does she or he smoke? If yes, does she or he smoke safely? ■ Are there signs of cigarette burns? ■ Are there signs of burns in the kitchen? Is stove free of grease? ■ Does patient use oxygen? If yes, is tank stored safely away from heat or flame? 	<p>Identifies</p> <ul style="list-style-type: none"> ■ Risk for injury. ■ Safety issues. ■ Teaching needs. ■ Referrals (e.g., local fire company for free smoke detectors).
<p>Crime/Injury</p> <ul style="list-style-type: none"> ■ Are there working locks on doors and windows? ■ Is patient able to make emergency calls? Are emergency numbers readily available? Is phone readily accessible? ■ Are there firearms in home? If yes, are they safely secured with ammunition stored separately? ■ Is there any evidence of criminal activity? ■ Are poisonous or toxic substances properly stored? 	<p>Identifies Safety Issues</p> <ul style="list-style-type: none"> ■ Need for referral to local police. ■ Teaching needs.
<p>ASSESSING SUPPORT SYSTEMS, SELF-ESTEEM, AND SELF-ACTUALIZATION</p>	
<p>Roles</p> <ul style="list-style-type: none"> ■ What roles does person play because of illness? ■ How has this affected other family members? 	<p>Identifies</p> <ul style="list-style-type: none"> ■ Source of stress, depression, and anxiety. ■ Need for referrals.
<p>Caregivers</p> <ul style="list-style-type: none"> ■ Is there a caregiver? Is caregiver competent, willing, and supportive? ■ Does caregiver need support? ■ Can caregiver hear patient? Is there a need for an intercom, "baby monitor," or bell? 	<p>Identifies</p> <ul style="list-style-type: none"> ■ Supports. ■ Need for referrals (e.g., caregiver may need support, respite care, or home health aide). ■ Need for assistive devices. ■ Teaching needs.
<p>COMMUNICATION</p> <ul style="list-style-type: none"> ■ Is phone in easy reach? ■ Can patient dial phone and see numbers? Does he or she need oversized numbers, audio enhancer, or a memory feature? ■ Is there a daily safety check system? Should there be an alert system like Lifeline? ■ Are emergency numbers for police, fire, ambulance, nurse, doctor, relative, or neighbor clearly marked? ■ How does patient get mail? 	<p>Identifies</p> <ul style="list-style-type: none"> ■ Emergency supports. ■ Teaching needs. ■ Need for assistive devices. ■ Safety issues.
<p>FAMILY/FRIENDS/PETS</p> <ul style="list-style-type: none"> ■ Who visits patient? Family, friends, church members? ■ Who can drive patient to doctor appointments, church, and other places? 	<p>Identifies</p> <ul style="list-style-type: none"> ■ Supports. ■ Need for referrals. ■ Ability to maintain follow-up care.

(continued)

Assessing the Home (continued)

AREAS/QUESTIONS TO ASK	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> Are there any pets? Is patient able to take care of them properly? Are pets well behaved? <p>Self-Esteem and Self-Actualization</p> <ul style="list-style-type: none"> What does patient like to do? Are there creative ways to enable her or him to do activities she or hshe enjoys? Are there meaningful solitary activities patient can do, such as reading or listening to music? Interactive activities? 	<ul style="list-style-type: none"> Teaching needs. <p>Identifies</p> <ul style="list-style-type: none"> Sources of meaning in patient's life. Need for referral to community resources (e.g., library, senior citizen groups).

Source: Adapted with permission from Narayan, M. (1997). Environmental assessment. *Home Healthcare Nurse* 15(11):798. Philadelphia: Lippincott-Raven.

Assessing the Community

According to Smith and Maurer (2000), a community is an “open social system characterized by people in a place over time who have common goals.” Some examples of a community are a town, a school, or a nursing home.

Assessment involves assessing the people within the community, the environment of the community, and the interaction between the two to identify any actual or potential health problems.

Assessing the Community

QUESTIONS TO ASK YOURSELF	RATIONALE/SIGNIFICANCE
<p>BOUNDARIES</p> <ul style="list-style-type: none"> What is the geographic description of the community (e.g., a town) or the criteria for membership in the community (e.g., a school)? What are the neighboring areas (e.g., a city if the community is a suburb, or an archdiocese if the community is a private Catholic school)? Are boundaries open or closed? <ul style="list-style-type: none"> What is the purpose or goals of the community (e.g., a school's mission statement)? <p>PHYSICAL CHARACTERISTICS</p> <ul style="list-style-type: none"> How old is the community? <ul style="list-style-type: none"> What are community's demographics (e.g., age, race, sex, ethnicity, housing, density of population)? 	<ul style="list-style-type: none"> Can be real, concrete, or conceptual. Identify who and what are included in the community. Allows you to focus assessment on the community. Some boundaries may exclude certain groups (e.g., communities with only high-priced housing may exclude lower socioeconomic groups). Reflects purpose of community. For example, the goal of a Catholic school would be to educate children in Christian values. Older, well-established community may have more resources available than new, developing community. Or, older community's resources may be outdated, not meeting current needs. Can identify healthcare needs of community by identifying health problems associated with age, gender, or race. For example: <ul style="list-style-type: none"> Older adults have a higher incidence of health problems and need more services than younger adults. Hypertension (HTN) screening services are greater for African American males because they have a higher incidence of this problem.

QUESTIONS TO ASK YOURSELF	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ What are community's physical features? 	<ul style="list-style-type: none"> ■ Women need women's healthcare services such as BSE teaching and mammography. ■ Ethnic groups may have specific health concerns. Cultural beliefs, values, and customs influence what and how health issues are addressed. ■ Type and condition of housing influences health (e.g., overcrowding increases incidence of communicable diseases). Also reflects how community values individual members. ■ Can influence community's behavior and health (e.g., exposure to toxic substances).
<p>PSYCHOSOCIAL CHARACTERISTICS</p> <ul style="list-style-type: none"> ■ What is the community's predominant religion, socioeconomic class, educational level, type of occupation, and marital status? 	<ul style="list-style-type: none"> ■ Religion influences what and how health issues are addressed (e.g., abortion/birth control contradicts Catholic beliefs). ■ Socioeconomic class reflects affordability and accessibility of healthcare services. People in low socioeconomic areas may not have financial resources for healthcare or be able to practice preventive healthcare. Limited financial access to healthcare raises community's morbidity and mortality rates. ■ Educational level identifies health teaching needs and approaches. The higher their educational level, the more likely people are to practice preventive health behaviors. ■ Occupation can identify specific health issues for the group (e.g., blue-collar workers may have a higher incidence of musculoskeletal problems, whereas white-collar workers may have a higher incidence of stress-related diseases). ■ Marital status may identify stability and support sources within community.
<p>EXTERNAL INFLUENCES</p> <ul style="list-style-type: none"> ■ Does the community receive any external funding? ■ Are there facilities outside community that are available to community members? Is there access/transportation to these facilities? ■ Are adequate healthcare providers available to community? ■ Are volunteer groups available to the community? ■ How is health information communicated to community? ■ What laws affect the community? 	<ul style="list-style-type: none"> ■ Federal or state funding and grants may be available for health services. ■ Identifies healthcare facilities needed (e.g., rural areas may need to go outside community for health services). ■ Volunteer groups can be a valuable resource, especially if healthcare providers are limited. ■ Determines if there is a need for further and better means of communication. ■ Laws, such as zoning or pollution laws, can influence healthcare issues. Federal and state programs also can affect the health and well-being of the community, such as senior citizens.

(continued)

Assessing the Community (continued)

QUESTIONS TO ASK YOURSELF

- Are values of external influences consistent with those of community?

INTERNAL FUNCTIONS

Human Services

- What human services are available within community? Nurses, doctors, volunteers? Is access to services adequate?
- What is community's budget? How much is allocated for healthcare services?
- What and how many healthcare facilities (hospitals, nursing homes, day care) are available in community? Are there enough to meet people's needs? Are they well equipped? Accessible?
- Does facility have goods and supplies to produce its goods? What is the product? What is the facility's contribution to the community?

- Is education appropriate, accessible, and adequate for community? What types of schools are there? How much money is budgeted for education?

Politics

- What is the organizational structure of the community? Elected vs. appointed positions? Terms? Formal vs. informal leaders?
- How are decisions made? Majority rule or consensus? Is community independent or dependent?

- What are rules and laws of community?
- Formal and informal?

COMMUNICATION

- *Nonverbal*: What is personality of community? How do people respond to outsiders?
- *Verbal*: Who communicates with whom? What are means of communication? Is communication vertical or horizontal?

Values

- What does the community value? What is important?
- Does the community have any traditions?
- Are there subgroups within the community?
- What is the condition of the physical environment? Clean, dirty, in disrepair?
- How is health defined by the community?
- How much does community value health?
- What type of healthcare facilities are there? How frequently are they used?
- Is the community homogeneous or heterogenous?

RATIONALE/SIGNIFICANCE

- Inconsistent values can affect healthcare issues. For example, a rural community may depend on federal funding for free mammograms, but the amount of funding fails to adequately cover the service. Or a community may feel that cleaning up a polluted stream is urgent, but external influences may disagree.

- Determines if services meet community's needs.
- Identifies value of healthcare services and need for external funding sources.
- Determines availability and accessibility of healthcare facilities.
- Identifies both positive and negative effects to community (e.g., if facility is drug treatment center, drug dealers and abusers, as a community in itself, exert a negative effect on community at large).
- Identifies adequacy and availability of schools.

- Helps identify how decisions are made and who has power.
- Identifies approaches needed to bring about change. Identifies role of nurse within community (e.g., if community is dependent, nurse may need to take more active role).
- Identifies laws that regulate behaviors.
- Identifies expectations, peer pressures.

- Identifies approaches toward community.

- Identifies what is important to community.
- Can influence healthcare practices.
- Can have own values and norms that affect community and influence healthcare practices.
- Condition of environment reflects value place on it by community.
- Identifies value community places on health.

- Identifies approaches toward community.

QUESTIONS TO ASK YOURSELF	RATIONALE/SIGNIFICANCE
HEALTH BEHAVIOR/HEALTH STATUS	
People	
<ul style="list-style-type: none"> ■ What is the growth rate of the community? Relationship between birth and death rates? Relationship between immigration and emigration? Is population young or old? Is it mobile? 	<ul style="list-style-type: none"> ■ Identifies health status and needs of community (e.g., health needs for retirement community are different from those of a community with mostly young couples).
<ul style="list-style-type: none"> ■ What are morbidity and mortality rates? The prevalence and incidence of disease? ■ What types of risky behavior occur in the community? Are there at-risk groups? ■ What is the incidence of presymptomatic illnesses, such as HIV, HTN, or high cholesterol? 	<ul style="list-style-type: none"> ■ Indirectly identifies teaching needs. ■ Identifies factors within community that affect health. ■ Identifies focus and direction of healthcare needed by community. ■ Identifies health teaching needs for community. ■ Identifies at-risk groups (e.g., homeless, drug addicts).
<ul style="list-style-type: none"> ■ What is the level of functioning of community (e.g., dependent vs. independent)? ■ Are there people with disabilities? How many people and what types of disabilities? 	<ul style="list-style-type: none"> ■ Identifies need for intervention and teaching (e.g., binge drinking on college campuses). ■ Identifies need for screening programs. ■ Determines effectiveness of existing screening programs. ■ Identifies health teaching needs. ■ Identifies type and direction of intervention needed to maintain and promote health. ■ Identifies resources required to meet need of community members with disabilities.
Environment	
<ul style="list-style-type: none"> ■ What is the quality of the air? ■ What is the quality of the food supply? 	<ul style="list-style-type: none"> ■ Identifies possible exposure to pollutants. ■ Identifies possible contamination causing GI diseases (e.g., salmonella).
<ul style="list-style-type: none"> ■ What is the quality of the water supply? Is the water public or well? Is it fluoridated? ■ What is the quality of the soil? 	<ul style="list-style-type: none"> ■ Identifies possible contamination causing GI problems (e.g., <i>Giardia lamblia</i>, lead). ■ Identifies possible contamination with radioactive material, human or animal excreta, <i>Ascaris</i> worm.
<ul style="list-style-type: none"> ■ Is there adequate housing? 	<ul style="list-style-type: none"> ■ Identifies possible crowding, radon and lead exposure.
<ul style="list-style-type: none"> ■ What is the quality of home and work site? ■ What is the quality of solid waste disposal? ■ What is the quality of hazardous waste disposal? 	<ul style="list-style-type: none"> ■ Identifies possible occupational health risks. ■ Identifies possible source of contamination. ■ Identifies possible contamination from toxic substances.
<ul style="list-style-type: none"> ■ Is there infestation by insects, rodents, and animals? Use of pesticides? 	<ul style="list-style-type: none"> ■ Identifies disease carriers (e.g., mosquito carries malaria and West Nile virus; deer ticks carry Lyme disease; animals can transmit rabies). ■ Pesticide use increases risk for exposure to toxic chemicals.
<ul style="list-style-type: none"> ■ Are there natural disasters in the community? Incidences of violence or terrorism? 	<ul style="list-style-type: none"> ■ Identifies sources of stress (e.g., tornados, school shootings, terrorist attacks).

Source: Data from Smith, C., and Maurer, F. (2000). *Community health nursing theory and practice*, ed. 2. Philadelphia: W.B. Saunders.

Guidelines for Assessing the Homeless Person*

Homelessness can be defined as a state in which a person has no permanent night-time residence or resides in a temporary shelter or a public or private place not intended for sleeping.

Homelessness affects men, women, and children of all ages, races, religions, and walks of life. An estimated 760,000 people in the United States are homeless, and this estimate is probably low. Of the homeless, about 46 percent are single men, 36.5 percent are families with children, and 14 percent are single women. Fifty-six percent of the homeless are African American, 27 percent white, and 13 percent Hispanic. Approximately 20 percent of the urban homeless are veterans. In all, 40 percent of homeless men are veterans, 25 to 33 percent are mentally ill, and 45 percent suffer from substance abuse. Families with children (often single mothers) are the fastest-growing group among the homeless.

Stages of Homelessness

- *Stage 1:* Episodic—homeless from time to time.
- *Stage 2:* Temporary—homeless, but continue to identify with the mainstream of society.
- *Stage 3:* Chronic—homelessness becomes the norm.

Causes of Homelessness

Some causes of homelessness include unemployment, underemployment, poverty, domestic violence (50 percent of homeless women with children have experienced abuse), eviction, and deinstitutionalization.

Effects of Homelessness

- Lack of food, clothing, medical services, and social supports.
- Vulnerability to crimes such as robbery, assault, and rape.
- Impaired recovery from illness as a result of inadequate wound care, poor nutrition, and exposure.
- Increased risk for episodic and chronic health problems.
- Lack of follow-up care because there is no phone or address.

- Feelings of low self-esteem, hopelessness, and powerlessness.

Barriers to Healthcare for Homeless

- No health insurance.
- No primary care provider.
- Noncompliance with treatment; no follow-up.
- Negative attitudes of healthcare providers.

Performing the Assessment of the Homeless Person

The components of the health history and physical exam are the same as for other patients. However, when dealing with homeless patients, your approach and focus differ.

Approach and Focus

- Provide unconditional positive regard for the homeless patient.
- Be accepting and nonjudgmental.
- Listen.
- Be sensitive to the reality of the homeless situation.
- Be direct in your communication with the patient.
- Ask permission to touch the patient. (Physical privacy may be one of the few prerogatives the patient has left.)
- Focus more on addressing an immediate problem than on getting a complete history and physical exam.
- Use event markers for time periods; days, months, and years may not be meaningful to the homeless person.
- Schedule healthcare appointments early in the day because shelter management requires a person be present in the afternoon to secure a bed for the night. “Claiming a bed” may take priority over healthcare.
- Respond accurately and simply if the homeless person expresses concern about how her or his personal revelations are recorded and how the information will be used.
- Realize that the homeless person may be homeless by choice, be impatient, refuse to make eye contact, take a long time to respond, express minor complaints such as headache, have a mailing address or contact person, and see healthcare needs and priorities differently from you.

Health History

The table below outlines a thorough health history for a homeless patient.

Taking the Health History

AREA/TYPES OF QUESTIONS

RATIONALE/SIGNIFICANCE

Biographical Data

- Ask where patient is staying.

- Less threatening to ask than “What is your address?”
- Questions about family members, telephone numbers, or additional demographic data may not be appropriate and may be met with silence. May also be reluctant to identify government agencies.

AREA/TYPES OF QUESTIONS	RATIONALE/SIGNIFICANCE
Current Health Survey	
<ul style="list-style-type: none"> ■ Ask about injuries first. 	<ul style="list-style-type: none"> ■ Patient may synthesize life and environmental factors to explain symptoms in ways that may not have occurred to you. Begin with questions about injuries; this may be a marker for time in an unstructured life.
<ul style="list-style-type: none"> ■ Ask about symptoms of communicable diseases that may not have been reported. 	<ul style="list-style-type: none"> ■ Increased risk for communicable diseases as a result of crowded living quarters.
<ul style="list-style-type: none"> ■ Ask about respiratory diseases, such as tuberculosis (TB). 	<ul style="list-style-type: none"> ■ Risk is high for those who sleep in close quarters in shelters or crowded spaces.
<ul style="list-style-type: none"> ■ Ask about dermatological problems such as lesions, infestation. 	<ul style="list-style-type: none"> ■ No change of socks, no clean clothes, exposure to extremes, poor hygiene, living in close quarters increase risk.
Past Health History	
<ul style="list-style-type: none"> ■ Ask about past injuries, infections, communicable diseases, hospitalizations, substance abuse, psychiatric problems. 	<ul style="list-style-type: none"> ■ Determines compliance and/or completion of treatment and follow-up.
Family History	
<ul style="list-style-type: none"> ■ Ask about the patient's family history. 	<ul style="list-style-type: none"> ■ May not want to discuss family.
Review of Systems	
<ul style="list-style-type: none"> ■ Ask about problems in the following systems that you may have missed in the Current Health Survey. 	<ul style="list-style-type: none"> ■ Homeless patients are at increased risk for:
Integumentary	<ul style="list-style-type: none"> ■ Skin problems (see Current Health Status); scabies, lice.
HEENT	<ul style="list-style-type: none"> ■ Ear and eye problems with children; dental problems with all ages.
<i>Respiratory</i>	<ul style="list-style-type: none"> ■ Upper respiratory infections (URIs), TB.
<i>Cardiovascular</i>	<ul style="list-style-type: none"> ■ HTN.
<i>Gastrointestinal</i>	<ul style="list-style-type: none"> ■ Alcohol abuse increases risk for GI disorders and poor nutrition.
<i>Genitourinary</i>	<ul style="list-style-type: none"> ■ <i>Women:</i> Pregnancy, lack of prenatal care.
Musculoskeletal/Neurological	<ul style="list-style-type: none"> ■ <i>All:</i> Poor nutrition, complications, STDs.
Hematologic/Immune/Endocrine	<ul style="list-style-type: none"> ■ Neurological and psychiatric disorders.
Psychosocial Profile	<ul style="list-style-type: none"> ■ Anemia related to dietary deficiency, HIV/AIDS; diabetes.
Ask about:	<ul style="list-style-type: none"> ■ Gives patient opportunity to present positive aspects of life. Acknowledging positive behaviors may help establish bond between you and patient.
<ul style="list-style-type: none"> ■ Health Practices and Beliefs 	<ul style="list-style-type: none"> ■ Barriers to healthcare and limited resources limit preventive behaviors. Patient's healthcare practices are often crisis/problem oriented and treated symptomatically. Follow-up is difficult.
<ul style="list-style-type: none"> ■ Typical Day 	<ul style="list-style-type: none"> ■ May consist of being on feet all day, looking for food and a place to sleep.
<ul style="list-style-type: none"> ■ Nutritional Patterns 	<ul style="list-style-type: none"> ■ Nutrition is a problem. Tell patient about available resources (e.g., food stamps, soup kitchens).
<ul style="list-style-type: none"> ■ Activity and Exercise Patterns 	<ul style="list-style-type: none"> ■ Center on survival and meeting basic needs.
<ul style="list-style-type: none"> ■ Sleep/Rest Patterns 	<ul style="list-style-type: none"> ■ Finding a safe place to sleep is a problem.
<ul style="list-style-type: none"> ■ Personal Habits 	<ul style="list-style-type: none"> ■ Substance abuse and associated health problems are common among homeless people, but do not assume this is the case with your patient. Ask, "Do you smoke, drink alcohol, or use drugs?" If the answer is yes, ask "How much?"
<ul style="list-style-type: none"> ■ Occupational Health Patterns 	<ul style="list-style-type: none"> ■ Patient may be working, but income may not be sufficient to maintain housing.

(continued)

Taking the Health History (continued)

AREA/TYPES OF QUESTIONS	RATIONALE/SIGNIFICANCE
<ul style="list-style-type: none"> ■ Socioeconomic Status ■ Environmental Health Patterns ■ Roles/Relationships/Self-Concept ■ Cultural Influences ■ Religious Influences ■ Sexuality Patterns 	<ul style="list-style-type: none"> ■ Homelessness denotes low socioeconomic status, poverty. Ask if patient is a veteran. Identify available resources and make appropriate referrals. ■ Thermoregulatory problems (hypo/hyperthermia) resulting from exposure. Alcohol increases risk for hyperthermia. Using fire to keep warm increases risk for burns. Because the homeless person often wears everything he or she owns, risk for heat exhaustion, heat stroke, and dehydration increases in warm weather. ■ Increased risk for being a victim of crime, robbery, assault, and rape. ■ Low self-esteem, anxiety, and depression are common. ■ Can be seen in both legal and illegal immigrants. Many groups have high poverty rates (e.g., 30 percent of Hispanics live in poverty). In homeless, language barriers can also add to problem of accessing healthcare. ■ Faith-based organizations are often major service providers for basic needs. Spiritual support may be integrated. ■ Homeless patient may still be sexually active and in a committed relationship. Ask, "Are you sexually active? With men, women, or both? Do you have genital itching, burning, or other symptoms?" ■ Ask if patient is a victim of abuse—50 percent of homeless women with children have been victims of domestic abuse.

Physical Assessment

Conduct the physical assessment using tools and techniques appropriate for the age and gender of the patient. Use gloves if risk for exposure to bodily fluids exists; otherwise, direct touch can convey trust. With the homeless patient, the customary head-to-toe approach may not be practical. Close, face-to-face contact at the beginning of the exam may be seen as invasive or threatening by the patient. Simply reverse the order, working from toe to head. This also minimizes the power position of the nurse standing over the patient, giving directions while performing the exam.

Pay special attention to the patient's feet. The homeless are at risk for foot problems because they are often on their feet most of the day, wearing poorly fitted

shoes with no change of socks. Regardless of your findings, the act of touching and examining the feet demonstrates a thoroughness of approach and simple caring.

Also keep in mind that the homeless person may have all of his or her possessions on his or her person and may be reluctant to remove clothes. Assure the person that his or her clothing and belongings will be safe, and allow him or her to keep personal items.

Performing the Physical Examination outlines a thorough physical assessment of a homeless patient. Instead of performing a total assessment, you will usually focus on a specific area identified as a problem from the history data. Be as thorough as necessary to meet your patient's health needs and as thorough as your patient permits.

Performing the Physical Examination

AREA	RATIONALE/SIGNIFICANCE
General Health Survey <ul style="list-style-type: none"> ■ Inspect general appearance, signs of injury, dress and grooming. ■ Note any odors. ■ Take vital signs: temperature, respirations, BP. 	<ul style="list-style-type: none"> ■ May have signs of injury—homeless people are more vulnerable to crime. ■ Poor hygiene resulting from lack of resources may warrant referral. ■ Body odors may be caused by poor hygiene or alcohol abuse. ■ Hypo/hyperthermia from exposure often present. ■ Upper respiratory problems common. ■ HTN is chronic health problem.

AREA	RATIONALE/SIGNIFICANCE
<p>Integumentary</p> <ul style="list-style-type: none"> ■ Inspect skin for lesions, color changes; inspect hair for infestation. ■ Inspect feet for lesions and edema. <p>HEENT</p> <ul style="list-style-type: none"> ■ Inspect mouth and throat. <p><i>Respiratory</i></p> <ul style="list-style-type: none"> ■ Auscultate lungs. <p><i>Cardiovascular</i></p> <ul style="list-style-type: none"> ■ Palpate peripheral pulses ■ Auscultate heart for normal and extra sounds. <p><i>Gastrointestinal</i></p> <ul style="list-style-type: none"> ■ Auscultate bowel sounds. ■ Palpate abdomen. <p><i>Genitourinary</i></p> <ul style="list-style-type: none"> ■ Inspect genitalia. <p><i>Musculoskeletal</i></p> <ul style="list-style-type: none"> ■ Inspect and palpate muscles. <p><i>Neurological</i></p> <ul style="list-style-type: none"> ■ Assess mental status. 	<ul style="list-style-type: none"> ■ Lice or scabies caused by crowded living conditions. ■ High incidence of hepatitis related to intravenous drug use. ■ Cirrhosis related to alcohol abuse. ■ Minor skin problems common. ■ Poor wound healing because of poor nutrition. ■ Being on feet most of day increases risk for peripheral-vascular disease (PVD). ■ Diabetes is also a chronic problem. ■ Dental problems and URIs common among homeless. ■ Respiratory problems and TB common. ■ Peripheral-vascular disease common. ■ HTN common. ■ GI problems common. ■ Cirrhosis, pancreatitis associated with alcohol abuse common. ■ Homeless people, especially women, are susceptible to rape, increasing risk for STDs and pregnancy. ■ Homeless people are susceptible to trauma from beatings. ■ Head trauma may result from beatings. ■ Substance abuse (drugs and alcohol) can affect neurological status. ■ Seizures can result from trauma and substance abuse. ■ Psychiatric problems common, including depression, schizophrenia, and organic brain syndrome.

Guidelines for Assessing Abuse*

Abuse can take many forms, can affect any age group, and knows no socioeconomic boundaries. It can be physical, sexual, emotional, or a combination of two or all of these. It may also be directed at property. As a nurse, you need to be alert for signs and symptoms of abuse at all levels of health care for all of your patients. This section covers child abuse, spousal or partner (domestic) abuse, and elder abuse.

Types of Abuse

- **Physical Abuse:** Physical contact with intent to harm, ranging from slapping, hitting, and biting to murder.

Physical abuse includes **neglect**—depriving a person of basic needs such as food, water, or sleep.

- **Sexual Abuse:** Sexually oriented behavior without the consent of the other person or persons involved, ranging from sexually degrading remarks to rape.
- **Psychological (Emotional) Abuse:** Verbal abuse or actions that can be considered degrading, belittling, or threatening, used for control, often by evoking fear in the victim. This type of abuse is difficult to assess.
- **Property Abuse:** Deliberate destruction of a person's belongings.

Assessing Child Abuse

Child abuse and neglect are the leading cause of death in children under 3 years.

More than 2 million cases are reported every year. Child abuse can take the following forms:

- **Physical Abuse:** Includes actual physical trauma, such as bruising, breaking bones, intentional burns, and shaken-baby syndrome. It also includes Munchausen syndrome by proxy (MSBP), in which the parent or caretaker intentionally causes a child to be ill to gain sympathy and recognition.
- **Neglect:** Failure to meet the child's basic needs.
- **Sexual Abuse:** Includes actual intercourse with either vaginal or anal penetration, genital fondling and other inappropriate touching, or pornographic photography of the child.
- **Psychological Abuse:** Includes emotional detachment from the child, constant belittling, fostering and enabling substance abuse and delinquency, and failing to provide adequate supervision.

Assessing for Physical Abuse

As you go through the assessment, ask yourself: Are the injuries inconsistent with the child's age or developmental stage (e.g., 1-month-old baby falling out of crib)? Is evidence of old fractures or trauma seen on current x rays?

Taking the Health History

Obtain separate statements from parents or caregivers. Be alert for inconsistencies (e.g., parent saying that hand-mark bruising on the child was inflicted by another child). Ask the following questions:

- Ask yourself: What is your perception of the child (e.g., difficult or quiet)?
- Ask the parent or caregiver:
- Does the child have a history of past health problems? Where was the child treated? (Frequent emergency room visits signal abuse. Parents may use different healthcare centers to avoid suspicion.)
- What is the child's feeding history? (Compare the history given with the child's physical growth and development.)
- Ask the child (if old enough): What types of stress do you have in your life? Who are your supports?

Suspect MSBP if:

- Child does not respond to usual treatment.
- Laboratory results are inconsistent with history or physically impossible.
- Parent seems very knowledgeable about medical treatment.
- Signs and symptoms of illness do not occur when parent is not present.
- Similar findings or unexplained deaths have occurred in siblings.
- Parent craves adulation for her or his care of child.

Performing the Physical Examination

Obtain growth measurements and compare them with previous measurements. Nutritional neglect can cause failure to thrive.

- Look for distended abdomen, a sign of malnutrition.
- Observe child's grooming, dress, and hygiene.
- Examine all body surfaces for bruises, burns, and skin lesions, looking especially for the following signs of physical abuse:
 - Bruises that form the outline of a hand (from parent grabbing child).
 - Linear bruises (from belts).
 - Bruises on head, face, ears, buttocks, and lower back (consistent with abuse).
 - Obviously nonaccidental burns (e.g., burns on both lower legs).
 - Burn outline of an entire object, such as multiple cigarette burns on various parts of body.
 - Detached retina and hemorrhages and subdural hematoma (shaken-baby syndrome).

Assessing for Sexual Abuse

Taking the Health History

Be nonjudgmental when questioning caregivers about possible sexual abuse. Obtain the following information from patients or their caregivers:

- Ask children over 2½ years old directly about being touched in "private parts." With older children, use an approach such as: "Sometimes people you know may touch or kiss you in a way that you feel is wrong. Has this ever happened to you?" With adolescents, use an approach such as: "Sometimes people touch you in ways you feel are wrong. This can be frightening, and it is wrong for people to do that to you. Has this ever happened to you?"
- Determine whether there has been a sudden onset of bedwetting in a child who was previously not a bedwetter.
- Determine whether the child masturbates or sexually acts out with other children.
- Determine whether the child has genitourinary symptoms, such as burning, itching, or vaginal discharge.
- Determine whether the child has a history of running away from home, especially to unsafe situations as opposed to a friend's home.

Performing the Physical Assessment

Look for genital and anal irritation, discharge, swelling, redness, and bruising. In girls, carefully evaluate the vaginal introitus for evidence of penetration. During the exam, observe the child's behavior:

- Is the child overly solicitous in a sexual manner?
- Does he or she dress or act provocatively?

- Is she or he overly fearful of the exam, especially the genital exam?

Assessing for Psychological Abuse

Taking the Health History

Ask the parent or caregiver the following questions:

- Was this child the result of a planned pregnancy? What were the birth and postpartum period like? Were there many stressors at home at the time of your child's birth?
- How would you describe your child now?
- How is this child compared with others in your family?
- Is there a history of physical or developmental problems? If so, what kinds of supports do you receive?
- Has the child had many illnesses? If so, what are they? How has this affected your life?
- What are your expectations of the child?
- What form of discipline works best for the child?

Performing the Psychological Assessment

During the exam, observe the following:

- Compare parental description of the child to what you actually observe.
- Observe child-parent interaction. Is it distant or engaging? How does the parent comfort the child?
- If other siblings are present, does the parent respond to them differently from the child in question?
- Does the parent openly and repeatedly belittle the child (e.g., saying: "You're so dumb")?
- Does the child have developmental delays, poor social skills, speech problems, or regression?

Assessing Spousal or Partner Abuse (Domestic Violence)

Domestic violence is the leading cause of injury to women. One in every three women has been abused by a spouse or partner at least one time during the relationship. Here are more facts about domestic violence:

- Fifty percent of homeless women and children have been victims of abuse.
- The battering cycle consists of the tension-building phase, the battering phase, and the apologetic phase.
- Abuse occurs in same-sex relationships as well as in heterosexual relationships.
- Abusers are usually emotionally dependent and egocentric.
- The battered partner is usually unsure, economically and emotionally dependent, and exhibiting learned helplessness.
- Spousal or partner abuse can take the form of physical, sexual, or psychological abuse (or a combination of two or all of these), or property abuse.
- Emotional abuse is difficult to assess because the effects are not as visible as those of physical abuse.

- To gain power and control, the abuser may:
 - Use coercion and threats, such as threatening to leave the partner or commit suicide.
 - Use intimidation to evoke fear by destroying personal belongings or hurting pets.
 - Degrade and belittle, making the partner feel that it is his or her fault, make him or her feel guilty, or blame abuse on him or her.
 - Isolate the partner by limiting contacts with family and friends.
 - Control or limit finances.
 - Put children in the middle of the situation or threaten to take them away.
 - If male, use male privilege by enforcing dominant role.

Assessing for Physical Abuse

Taking the Health History

Ask the patient:

- What is your chief complaint? Complaints are often vague and nonspecific; for example, headaches, gastrointestinal (GI) complaints, asthma, fatigue, and chronic vague pain.
- Do you have trouble sleeping? Do you have nightmares?
- Have you ever been hospitalized? Frequent emergency room visits are common, with injuries becoming progressively more severe. Current x-rays may reveal old fractures.
- How did the injury happen? Injury may not match the explanation (e.g., black eye from walking into door).
- Do you have a history of depression or attempted suicide? Abuse victim with a sense of hopelessness and powerlessness may be depressed and attempt suicide.
- Are you taking any prescribed or OTC medications? Do you drink alcohol? If so, how much? Substance abuse may be used as an attempt to deal with physical abuse.
- Have you had any weight or appetite changes? Anorexia and bulimia are not uncommon.
- Have you ever been sexually assaulted? Are you satisfied with your sexual relationship with spouse or partner? Domestic violence can take the form of sexual abuse.

Performing the Physical Examination

Assess for injuries suggesting abuse, such as cigarette burns; black eyes; facial injuries, injuries to the chest, back, breast, or abdomen; bruising on genitalia; or bruises in the shape of a hand or belt.

During the exam, observe for the following:

- Patient's lethargic, passive behavior.
- Patient's poor eye contact and anxious or fearful behavior.
- Patient's visible fear when partner is in room; looking to partner before responding to questions.
- Partner answering for patient and being overly condescending.

Assessing Elder Abuse and Neglect

More than 800,000 cases of elder mistreatment are reported annually. Elder abuse can be physical, sexual, or psychological. It can also take the form of neglect or financial or property abuse. During your assessment, be alert if the caregiver speaks for the patient, refuses to let the patient be examined alone, or underreacts when confronted with findings suggesting abuse.

Assessing for Physical Abuse

Taking the Health History

Ask the following questions of the patient and caregiver:

- How did the injury happen? Suspect abuse if injury does not match explanation (e.g., caregiver says injury is from fall, but patient has no difficulty with balance or walking).
- When did the injury occur? Lengthy interval between injury and treatment suggests abuse.
- Who is the primary healthcare provider? “Doctor shopping” (using different healthcare providers) suggests abuse.
- To patient: Do you feel safe in your home?
- To patient: Are there any situations in which you feel afraid?
- To patient or caregiver: Have you ever given/taken the wrong dose of medication? Repeated medication errors by caregiver (e.g., oversedating patient) suggest abuse.

Performing the Physical Examination

During the exam, observe for the following:

- Whiplash burns from rope or cord.
- Cigarette burns on palms and soles of feet.
- Injuries with a pattern, such as that of a belt buckle or electrical cord.
- Oral ecchymosis or injury from forced oral sex.
- Whiplash injuries from shaking.
- Hyphema, subconjunctival hemorrhage, detached retina, ruptured tympanic membrane.
- Bruises on trunk, breast, abdomen, genitalia, and buttocks (bathing suit zone).
- Bleeding or bruising of genitalia, poor sphincter tone, and bruises on inner thighs.
- Bruising on wrists and ankles from being bound or restrained.

- Immersion burns (may follow a stocking-glove pattern).
- Evidence of old injuries.
- Anxiety and depression.

Assessing for Neglect

The following are signs of neglect in a patient:

- Never changing clothes or wearing clothes that are dirty or inappropriate for the weather.
- Poor hygiene or body odor.
- Ingrown nails.
- Decayed or missing teeth.
- Untreated sores or pressure sores.
- Matted hair.
- Hypo/hyperthermia.
- Untreated medical problems.
- Dehydration.
- Malnutrition, failure-to-thrive syndrome.
- Weight loss.
- Abnormal blood chemistry.
- If you are making a home visit: Clutter, disconnected utilities, neglected pets, animal or human excrement, spoiled food.


Assessing for Financial Abuse

The following are signs of financial abuse:

- Unusual activity in patient’s bank account.
- Bank statements diverted from patient’s home.
- Sudden disappearance of caregiver.
- Patient asked to sign documents such as power of attorney.
- Large withdrawals from patient’s account.
- Family that is reluctant to spend money on patient.
- Disappearance of personal belongings.
- Isolation of patient by caregiver.
- Forged signatures on checks and documents.
- Caregiver who is evasive about sources of income.
- Lack of solid, legal financial arrangements in place for older person.

Documentation

Physical, sexual, and financial abuse are crimes in all states. In most states, suspicion of abuse is grounds for reporting. Make sure that you document your findings clearly and objectively. Be specific and thorough because your assessment findings are crucial in early detection and prompt intervention.



APPENDIX

D

Genetics

Family History, the First Genetic Test

Introduction and Background

Collecting a family history has been part of the health assessment interview at least since Hippocrates and probably for generations before he was born. Although health-care practitioners instinctively understood the link between family history and health problems, until recently, little could be done except to suggest lifestyle changes (increasing weight-bearing exercise and dietary calcium for a woman at risk for osteoporosis), or increasing surveillance (encouraging regular breast exams and annual mammograms for a woman at risk for breast cancer). Over the past 50 years, however, our understanding of the risk factors for and causes of many diseases has exploded, in large part because we have a better understanding of biochemistry and genetics. Fifty years after Watson and Crick first described the structure of DNA in the April 1953 edition of the journal *Nature*, the National Human Genome Research Institute (NHGRI) announced the completion of the Human Genome. The entire sequence is available online at no charge at the Online Mendelian Inheritance in Man (OMIM) website (<http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM>; see Genetic Resources) to anyone interested in exploring it. We are rapidly beginning to uncover the specific genetic causes for many diseases, and it is proposed that virtually all human health conditions (including accidental injury) have at least some genetic component. Collecting a comprehensive family history—the first genetic test—has never been more important. Nurses are perfectly positioned to assist patients and families move healthcare into the 21st century.

The process for gathering a family history and the rationale for doing so are explained very well in Chapter 2, The Health History. If, however, after collecting the family history, you suspect that a patient may be at increased risk for inheriting a genetic condition, you should ask a series of detailed questions to get more specific information. The main reasons to collect more information are to clarify social and biological relationships, establish patterns of inheritance, identify at-risk individuals, reach a correct diagnosis, make a presymptomatic diagnosis, or determine a prognosis. When collecting the genetic history, be on the lookout for genetic “red flags” (see Genetic Red Flags). If any of these appear in the history, your patient may be at increased risk for inheriting a genetic disorder.

Pedigree Collection

If at all possible, you should record the family history using standard pedigree symbols (Bennett et al., 1995),

Genetic Red Flags

- Multiple family members affected with the same or related conditions
 - Evidence of autosomal dominant inheritance (vertical transmission on a pedigree)
 - Early age at onset than expected
 - Alzheimer's disease at age 50
 - Breast cancer age 36
 - Occurrence of condition in the less-often-affected sex
 - Breast cancer in male
 - Multifocal or bilateral occurrence
 - Multisystem
 - Deafness and nephritis “Alport’s syndrome”
 - Individual or couple with three or more pregnancy losses
 - Medical problems in children or parents who are closely related
 - Within an individual look for:
 - A medical condition and dysmorphic features
 - Developmental delay with dysmorphic features and/or physical birth anomalies
 - Learning disabilities or behavioral problems
 - Unexplained movement disorders, hypotonia, ataxia
 - Congenital/juvenile deafness, blindness, or cataracts
- Unexplained infertility**

because it visually represents the family, revealing relationships between individuals and exposing inheritance patterns. There are a few basic steps to pedigree collection: Begin with the “proband” and identify this individual with an arrow. The proband is defined by Genetics Home Reference as “The affected individual through whom a family with a genetic disorder is ascertained; may or may not be the consultand (the individual presenting for genetic counseling).”

Ask about ethnic origin if you suspect an autosomal recessive condition (like Tay Sachs, sickle cell disease, or cystic fibrosis).

Be sure to include information from both the maternal and the paternal relatives. For example, a woman carrying a mutation for the BRCA1 gene (a breast cancer mutation) is at increased risk for developing breast cancer at a young age. The BRCA1 gene is located on chromosome 17, which can be inherited from either the woman’s mother or her father. If a woman inherits the gene from her father, it may not appear, or it may appear to have skipped a generation, because men rarely get breast cancer. But if you do not ask about his family history, you may never find out about the aunts and female cousins who have had early-onset breast or ovarian cancer, and you may miss an opportunity to make a presymptomatic diagnosis in this proband.

Ideally, you should collect a three-generational pedigree—one generation below and two above the proband, but how far up and down the family tree you can go depends upon the proband. If the proband is an infant, three generations might include siblings, parents, aunts/uncles, nieces/nephews, and grandparents. If the proband is an adult, the pedigree may include siblings, parents, grandparents, aunts/uncles, nieces/nephews, cousins, children, grandchildren, and possibly, even great-grandchildren. Be sure to record all pregnancy losses, adopted children (in or out), twins (whether they were mono- or dizygotic), and any known consanguinity (a common ancestor). Other important information to record are the current health status of family members, the age and cause of death if deceased, the age of disease onset (disease of interest), and the primary site for a cancer (if cancer is the genetic disease of concern).

Be sure to document the date the history was collected and who recorded it in the medical record. Remember that family history is dynamic; pedigrees should be reviewed at least annually and updated as more information emerges. You will quickly discover that family histories can be quite complex and disorders may be layered or intertwined. One individual, for example, may have a strong family for cancer, diabetes, hypertension, and mental illness. Where do you start? In this situation, it may take time to build a complete “picture,” and you may choose to focus the history on the most “urgent” problem and ask the patient to go home and collect more information.

Resources

Several groups, including the Centers for Disease Control and Prevention (CDC) and the U.S. Surgeon General, have been hard at work developing electronic tools to assist in the collection of family history information. Some tools are designed to be completed by individuals at home (U.S. Surgeon General), whereas others are an integral part of an electronic health record and data are collected and entered into the system by a healthcare professional (often a nurse). Information and a free download of the U.S. Surgeon General’s Family History tool are available at <http://www.hhs.gov/familyhistory/>.

This tool was widely advertised on Thanksgiving weekend in 2004 and 2005 in the hopes that people would discuss and record their family medical histories while gathered together to celebrate the holiday. The goal is to have everyone complete the tool and bring a paper copy of the pedigree to their next office visit. Then it’s up to healthcare providers to verify the information, evaluate the history, and make recommendations based on any familial risk factors.

Evaluating the history: Risk Stratification and Referral

When evaluating a family history, clinicians should ask themselves three questions:

1. Is there evidence of a genetic disease or syndrome?

The “red flags” in the box provide clues to the presence of a genetic disorder or syndrome, so all of these items should be carefully reviewed with the family member.
2. What might make a family history difficult to interpret?
 - The disorder may be new and the proband the first person to have the disease.
 - Nonpaternity. The person identified as the “father” on the pedigree may not be the biological father.
 - The family may be very small, or the person providing the history is not aware of all the details of the family history, has inaccurate information (“ovarian cancer” was actually “cervical cancer”), or is deliberately withholding information (perhaps from fear of discrimination).
 - The early death of a family member from other causes (for example, in a war or an accident).
 - Disorders that are masked because they are inherited from an individual who does not typically express the disorder. For example, colorblindness is not usually seen in women, because the mutation is on the X chromosome, and women have 2 Xs (basically a back-up system). If you are looking for colorblindness patterns in a family in which there are many women/girls, but few boys and the men died early of heart disease or never talked about having any vision problems, colorblindness can appear to be a new condition, when in fact, it may have been present for many generations.
 - Incomplete penetrance. Some disorders have a “range” of expression from mild to severe. For example, Treacher Collins syndrome is caused by a mutation on the fifth chromosome. This gene directs the growth of facial bones and structures in human embryos. If this gene is abnormal, the face does not form normally. This mutation is not completely “penetrant,” however, and in some people, the facial abnormality is so mild that the individual may never realize they have a genetic mutation until they have a child with a severe facial anomaly.
 - Multifactorial diseases. The environment or other genes can regulate the expression of a mutation. Phenylketonuria (PKU) is one example of a multifactorial genetic condition. Individuals with PKU lack an enzyme to break down phenylalanine, a byproduct of protein metabolism. The normal American diet contains large amounts of protein, and in infants with PKU, phenylalanine levels

climb quickly, causing irreversible brain damage. If a low-protein diet is adhered to (the environment is modified), brain damage does not develop.

3. What is the most likely mode of inheritance?

Key features to look for:

- Autosomal dominant. The disorder shows up in every generation (vertical transmission), every affected person has an affected parent, the disorder occurs equally in both sexes, and approximately 50 percent of offspring have the disorder.
- Autosomal recessive. Nonaffected parents with affected offspring, approximately 25 percent of offspring have the disorder, the disorder occurs equally in both sexes, and consanguinity increases the risk.
- X-linked disorders. More affected males (pseudodominance); nonaffected carrier females; all females of affected males are obligate carriers.
- Mitochondrial disorders. Variable age of onset, disease severity, and prognosis; universal weakness and fatigue; unique maternal inheritance pattern—all offspring (male and female) of women are affected; men may have the disorder, but they do not pass it on.

Genetic Counseling: Who, When, How, and Why

Once you've decided that a family history is significant, referral to an appropriate provider is typically the next step. There are several healthcare disciplines to choose from. Here's a brief explanation of who is working in the genetic field to help you decide who might be the most appropriate referral resource.

Medical Geneticist

The American Society of Human Genetics (ASHG) was founded in 1948 and is the oldest human genetic organization in the United States. The ASHG's mission is to provide leadership in research, education, and service in human genetics. At the time of its founding, virtually everyone interested in genetics belonged to this one organization. Over time as genetics diversified, smaller societies began separating from the parent organization. In 1979, the genetic counseling community separated to form The National Society of Genetic Counselors (NSGC). In 1991, The American College of Medical Genetics (ACMG) and the American College of Medical Genetics Foundation (ACMGF) were established to represent clinical practitioners, and since 1993, ACMG has been recognized as the body that represents doctoral-

prepared researchers and physicians with advanced education in genetics.

In 1981, ASHG formed a separate certification organization, the American Board of Medical Genetics (ABMG). The ABMG writes and administers certification exams for medical and clinical laboratory professionals. When referring a patient to a medical geneticist, look for credentials like FACMG (Fellow in the American College of Medical Genetics) or ASHG/ABMG. More information on medical geneticists is available on the ACMG (<http://www.acmg.net/>) and ASHG (<http://genetics.faseb.org/genetics/ashg/ashgmenu.htm>) websites.

Genetic Nurse (BSN and APN)

Genetic nurses perform risk assessments, analyze genetic contributions to disease risk, help individuals and families explore the meaning and impact of the disease, and develop management plans to assist families in coping with genetic disorders. They also provide genetic education and nursing care to patients with genetic conditions and conduct research in genetics. Two credentials nurses can work toward in the area of genetics are The Genetic Clinical Nurse (GCN), developed for baccalaureate-prepared nurses, and the Advanced Practice Nurse in Genetics (APNG), offering a masters-prepared nurse specialty credentialing in the area of genetics. Obtaining either credential is rigorous and time consuming. These nurses are well prepared to care for patients with genetic conditions.

When referring a patient to a nurse geneticist, look for credentials GCN or APNG. More information about both of these credentials is available on the International Society of Nurses in Genetics (ISONG) website: www.isong.org.

Genetic Counselor (CGC)

Genetic counselors are masters-prepared health professionals who have specialized in the areas of medical genetics. Most have undergraduate degrees in a science or healthcare field (biology, genetics, nursing, psychology, public health, social work). Genetic counselors provide information and support to individuals and families who are at risk for or who have genetic disorders. They provide supportive counseling to families, serve as patient advocates, and refer individuals and families to community or state support services. Genetic counselors also serve as an important source of education and information about genetic conditions for other health professionals and for the general public. When referring a patient to a genetic counselor, look for the CGC credentials. More information about genetic counselors is available on the National Society of Genetic Counselors website at www.nsgc.org.

Genetic Resources

Recommended Textbooks

Everyone should have at least one genetic textbook in their library—one that they find interesting and are willing to read. Here are two suggestions:

- *Nursing Care in the Genomic Era: A Case Based Approach* by Jean Jenkins and Dale Lea. This book, published by Jones & Bartlett in 2005, is a great resource that every nurse should have in his or her library. It is unique because it is both very readable and very detailed. Jenkins and Lea do a wonderful job making sure all the essential areas are covered; biology, genetic testing, pharmacogenetics, results disclosure, genetic research, and resources. (Chapter 14 on resources stands out.) However, what makes it truly different from other genetic textbooks are the stories. Throughout the book, you hear the voices of people with genetic conditions or the voices of people who have family members with genetic conditions; they make genetics come alive.
- *Clinical Genetics in Nursing Practice*, ed. 3, by Felissa Lashley. Lashley is the Dean and Professor of the College of Nursing at Rutgers in New Jersey. She is a nurse who received her Ph.D. in human genetics (along with a minor in biochemistry). She was also the first nurse to be credentialed as a Ph.D. Medical Geneticist by the ABMG and was a founding fellow of the ACMG. This might seem like quite enough for one person to do, but she's perhaps best known as the author of this textbook, which was published by Springer in 2005 and is now in its third edition. This text is another one of those resources that all nurses should have on their bookshelf. This book is similar to *Nursing Care in the Genomic Era*, in that it addresses biology, genetic testing, family history collection, and so on, but it goes into more detail and includes chapters on selected genetic disorders. If you want to learn more about a particular a genetic condition, this is a great resource to turn to.


Recommended Online Resources

Genetics “lives” on the web, and just by entering a few words into a search engine like Google, you can find links to literally hundreds of websites. How do you decide

which ones are “good”? Here are a few suggestions to get you started.

- **DNAi:** (www.dnai.org)
This site is an amazing, interactive, multimedia resource developed for high school and introductory college biology students and teachers. That said, it is also a wonderful place for nurses to learn about genetics. The site was developed to commemorate the 50th anniversary (2003) of James Watson and Francis Crick's discovery of the structure of DNA. The Dolan DNA Learning Center at Cold Spring Harbor was funded by the Howard Hughes Medical Institute to develop it, and it has won a number of awards including the British Academy of Film and Television Arts award in 2004 and the Exploratorium's Ten Cool Sites Award for educational excellence in 2003. It contains more than 200 video clips and animations; over 50 interviews with scientists, clinicians, and patients; and 3D animations that really show how DNA is copied, transported, and turned into proteins. There is a module on mitochondrial inheritance called “Uncovering the Romanov's” in which you can explore the mystery of the death and discovery of the Romanov graves.
- **Online Mendelian Inheritance in Man (OMIM):** <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM>
This is where you should go if you want detailed, specific information about a particular gene or genetic condition. It's an online catalog of human genes and genetic disorders. The content is edited by Dr. Victor McKusick at Johns Hopkins along with medical colleagues from around the globe. The website is supported by the National Center for Biotechnology Information (NCBI). The intended audience is physicians, genetic counselors, genetic researchers, and advanced students in science and medicine. Although the database is free and open to the public, it's not intended for patient education and it can be tough going for people who aren't totally “up to speed” in genetics.
- **Infogenetics:** <http://www.infogenetics.org/>
This is the place to go if you're wondering where to start looking—for a genetic clinical trial, for information about a genetic test, for a link to OMIM, for answers to a newborn screening question, and so on. Infogenetics is described an “Internet Manager with Clinical Practice Tools.” This can take the place of your “favorites” folder.

Handwritten text in a cursive script, likely a Latin manuscript, located at the top of the page. The text is partially obscured by the top edge of the drawing's circle.



Bibliography

Chapter 1

- American Nurses' Association (1980). *Nursing: A Social Policy Statement*. (Publication Code: NP-63, 35M.) Kansas City, Mo.:ANA.
- American Nurses' Association (1998). *Standards of Clinical Nursing Practice*, ed. 2. Washington, D.C.: American Nurses' Publishing, American Nurses' Foundation/ANA.
- American Nurses' Association (2004). Code for Nurses.
- Benner, P. (1984). *From Novice to Expert: Excellence and Power in Clinical Nursing Practice*. Menlo Park, Calif.: Addison-Wesley.
- Benner, P., Tanner, C., and Chesla, C. (1996). *Expertise in Nursing Practice, Caring, Clinical Judgment and Ethics*. New York: Springer Publishing.
- Carpenito, L. (1997). *Nursing Diagnosis Application to Clinical Practice*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- Doenges, M., Moorhouse, M., and Geissler, A. (1997). *Nursing Care Plans: Guidelines for Individualizing Patient Care*, ed. 5. Philadelphia: F.A. Davis.
- Ennis, R. (1985). *A Taxonomy of Critical Thinking Dispositions and Abilities*. Illinois Critical Thinking Project. Champaign, Ill.: University of Illinois.
- Green, C. (2000). *Critical Thinking in Nursing*. Upper Saddle River, N.J.: Prentice Hall.
- Gordon, M. (2002). *Manual of Nursing Diagnosis*, ed. 10. St. Louis: Mosby.
- HIPAA. www.hhs.gov/ocr/hipaa
- Kelly-Heidenthal, P., and Marthaler, M. (2005). *Delegation of Nursing Care*. Clifton Park, N.Y.: Thomson Delmar Learning.
- Maslow, A. (1970). *Motivation and Personality*, ed. 2. New York: Harper & Row.
- Morton, P. (1993). *Health Assessment in Nursing*, ed. 2. Philadelphia: F.A. Davis.
- NANDA (2005). <http://www.nanda.org/html/nursing>
- Nightingale, F. (1859). *Notes on Nursing: What It Is, and What It Is Not*. Philadelphia: J.B. Lippincott.
- NJPC. See Morton, P. (1993).
- Paul, R. (1990). *Critical Thinking*. Rohnert Park, Calif.: The Center for Critical Thinking and Moral Critique, Sonoma State University.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. . Philadelphia: F.A. Davis.
- Standards of Practice (1992). Kansas City, Mo.: American Nurses Association.
- Stuart, G., and Laraia, M. (2001). *Principles and Practice of Psychiatric Nursing*, ed. 7. Philadelphia: C.V. Mosby.
- Taylor, C., Lillis, C., and LeMone, P. (2001). *Fundamentals of Nursing: The Art and Science of Nursing Care*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.
- Weeks, C. (1891). *Textbook on Nursing*. New York: Appleton.
- Wilkinson, J. (1991). *Nursing Process in Action*. Redwood City, Calif.: Addison-Wesley Nursing.

Chapter 2

- Carpenito, L. (1997). *Nursing Diagnosis Application to Clinical Practice*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- Craven, R., and Hirnle, C. (2000). *Fundamentals of Nursing Human Health and Function*, ed. 3. Philadelphia: Lippincott Williams & Wilkins.
- Doenges, M., Moorehouse, M., and Geissler, A. (1997). *Nursing Care Plans: Guidelines for Individualizing Patient Care*, ed. 4. Philadelphia: F.A. Davis.
- Ecomap website. www.uic.edu/nursing/genetics/Lecture/Family/ecomap.htm
- Morton, P. (1993). *Health Assessment in Nursing*, ed. 2. Philadelphia: F.A. Davis.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Stuart, G., and Laraia, M. (2001). *Principles and Practice of Psychiatric Nursing*, ed. 7. Philadelphia: C.V. Mosby.

- Taylor, C., Lillis, C., and LeMone, P. (2001). *Fundamentals of Nursing: The Art and Science of Nursing Care*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.
- U.S. Department of Health and Human Services (1998). *Put Prevention Into Practice: Clinician's Handbook of Preventative Services*, ed. 2. Washington, DC: AHCPR.
- Wong, D., and Hockenberry-Eaton, M. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: C.V. Mosby.

Chapter 3

- Lewis, S., Heitkemper, M., and Dirksen, S. (2000). *Medical Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Lowdermilk, D., Perry, S., and Bobak, I. (1999). *Maternity Nursing*, ed. 5. Philadelphia: C.V. Mosby.
- Morton, P. (1992). *Health Assessment in Nursing*, ed. 2. Philadelphia: F.A. Davis.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Taylor, C., Lillis, C., and LeMone, P. (2001). *Fundamentals of Nursing: The art and Science of Caring*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.
- Wong, D., and Hockenberry-Eaton, M. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: C.V. Mosby.

Chapter 4

- Attia, J., et al. (1987). Measurement of postoperative pain and narcotic administration in infants using a new clinical scoring system. *Anesthesiology* 67(3A):A532.
- Beal, J., and Freda, M. (2004). Nursing assessment of infant pain. *The American Journal of Maternal/Child Nursing* 29(5):330.
- Beck, C. (1995). Screening methods for postpartum depression. *Journal of Obstetrics and Gynecological Nursing* 4(4): 308.
- Beyer, J.E., Denyes, M.J., and Villarruel, A.M. (1992). The creation, validation and continuing development of the Oucher: A measure of pain intensity in children. *Journal of Pediatric Nursing* 7(5):335-346.
- Clark, M. (2004). *Pain: Current understanding of assessment, management and treatments*. American Pain Society www.ampainsoc.org
- Cline, M.E., et al. (1992). Standardization of the visual analogue scale. *Nursing Research* 41(6):378-380.
- Eland, J.A., and Banner, W. (1999). Analgesia, sedation, and neuromuscular blockage in pediatric critical care. In Hazinski, M.E. (ed.): *Manual of Pediatric Critical Care*. St. Louis: Mosby.
- Hannallah, R.S., et al. (1987). Comparison of caudal and ilioinguinal/iliohypogastric nerve blocks for control of post-orchiopey pain in pediatric ambulatory surgery. *Anesthesiology* 66:832-834.
- Hester, N.O., Foster, R. L., and Kristensen, K. (1998). Putting pain measurement into clinical practice. In Finley G.A., and McGrath, P.J. (eds.): *Measurement of Pain in Infants and Children*, vol. 10. Seattle, Wash.: International Association for the Study of Pain Press.
- Hodgkinson, K., et al. (1994). Measuring pain in neonates: Evaluating an instrument and developing a common language. *Australian Journal of Advanced Nursing* 12(1):17-22.
- International Association for the Study of Pain. (1994).
- Joyce, B.A., et al. (1994). Reliability and validity of preverbal pain assessment tool. *Issues Compr. Pediatric Nursing* 17:121-135.
- Kozier, B., Erb, G., Berman, A., and Snyder, S. (2004). *Fundamentals of Nursing*, ed. 7. Upper Saddle River, N.J.: Pearson/Prentice Hall.
- Kretchel and Bildner. (1995). CRIES Neonatal Pain Scale developed at University of Missouri-Columbia.
- LaDuke, S. (2002). Undertreated pain. *Nursing* 32:18.
- Lane, P. (2004). Assessing pain in patients with advanced dementia. *Nursing* 34(8):17.
- Lawrence, J., et al. (1993). The development of a tool to assess neonatal pain. *Neonatal Network* 12(6):59-66.

- Mayer, D., Torma, L., Byock, I., and Norris, K. (2001). Speaking the language of pain. *American Journal of Nursing* 101(2):44–49.
- McCaffery, M., and Passero, C. (1999). *Pain: Clinical Manual*, ed. 2. Philadelphia: Mosby.
- McGrath, P.J. et al. (1985). The Children's Hospital of Eastern Ontario Pain Scale (CHEOPS): A behavioral scale for rating post-operative pain in children. In Fields, H.L., Dubner, R., and Cervero, F. (eds.): *Advances in Pain Research and Therapy*. New York: Raven Press.
- National Pharmaceutical Council. (2001).
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Reyes, S. (2003). Nursing assessment of infant pain. *Journal of Perinatal and Neonatal Nursing* 17(4):291–303.
- Robieux, I., et al. (1991). Assessing pain and analgesia with a lidocaine-prilocaine emulsion in infants and toddlers during venipuncture. *Journal of Pediatrics* 118(6):971–973.
- Schade, J.G., et al. (1996). Comparison of three preverbal scales for post-operative pain assessment in a diverse pediatric sample. *Journal of Pain Symptom Management* 12(6):348–359.
- Stevens, B. (1990). Development and testing of a pediatric pain management sheet. *Pediatric Nursing* 16(6):543–548.
- Stevens, B., et al. (1996). Premature Infant Pain Profile: Development and initial validation. *Clinical Journal of Pain* 12:13–22.
- Stuart, M.C., and Lieberman, J.A. (1993). *The Fifteen-Minute Hour: Applied Psychotherapy for the Primary Care Physician*, ed. 3. Westport, CT: Praeger, 101–183.
- Taddio, A., et al. (1995). A revised measure of acute pain in infants. *Journal of Pain and Symptom Management* 10(6):456–463.
- Tesler, M., et al. (1991). The word-graphic rating scale as a measure of children's and adolescents' pain intensity. *Research Nurse Health* 14:361–371.
- Victor, K. (2001). Properly assessing pain in the elderly. *RN* 64(5):45–49.
- Wong, D., and Hockenberry-Eaton, M. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: Mosby.

Chapter 5

- APA. (2000).
- Beck, (1995).
- Boyd, M.A. (2005). *Psychiatric Nursing Contemporary Practice*. Philadelphia: Lippincott, Williams & Wilkins.
- Brigham and Women's Hospital. (2001). *Depression: A Guide to Diagnosis and Treatment*. Boston: Brigham and Women's Hospital.
- Cox, Holden, and Sagovsky. (1989).
- Folstein, M., Folstein, S.G., and McHugh, P. (1975). Mini-Mental State, a practical method for grading the cognitive state of patients for the clinician. *Journal of Psychiatric Research* 12:189–198.
- Gorman, L. Raines, M., and Sultan, D. (2005). *Psychosocial Nursing for General Patient Care*, ed. 2. Philadelphia: F.A. Davis.
- Haddad, M., Plummer, S., Taverner, A., Gray, R., Lee, S., Payne, E., and Knight, D. (2005). District nurses' involvement and attitudes to mental health problems: A three area cross-sectional study. *Journal of Clinical Nursing* 14(8):976–985.
- Healthy People 2010.
- Jahoda, M. (1958). *Current Concepts of Positive Mental Health*. New York: Basic Books.
- Lowdermilk, D., Perry, S., and Bobak, I. (1999). *Maternity Nursing*, ed. 5. Philadelphia: Mosby.
- Pedersen, D. (2005) *PsychNotes: Clinical Pocket Guide*. Philadelphia: F.A. Davis.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Seyle, H. (1976). *The Stress of Life*. New York: McGraw-Hill.
- Stuart, G., and Laraia, M. (2005). *Principles and Practice of Psychiatric Nursing*, ed. 7. Philadelphia: Mosby.

- Stuart, and Lieberman. (1993).
- Townsend M. (2006). *Psychiatric Mental Health Nursing*, ed. 5. Philadelphia: F.A. Davis.
- U.S. Public Health Service (USPHS). (1999). *The Surgeon General's Call to Action to Prevent Suicide*. Washington, D.C.: U.S. Department of Health and Human Services. <http://www.surgeongeneral.gov/library/calltoaction.htm>.
- WHO. (2005). www.healthypeople.gov/Document/HTML/volume2/18Mental.htm
- www.stopthechokinggame.com
- Yesavage, J.A., Brink, T.L., Rose, T.L., Lum, D., Huang, V., Adey, M.B., and Leirer, V.O. (1983). Development and validation of a geriatric depression screening scale: A preliminary report. *Journal of Psychiatric Research* 17:37–49.

Chapter 6

- American Red Cross (March 1997). *Health and Safety Services Instructional System: Manual for the Instructor Trainer Candidate*. American Red Cross.
- Eiss, A.E., and Harbeck, M.B. (1969). *Behavioral Objectives in the Affective Domain*. Washington, D.C.: National Education Association.
- Felder, R.M., and Solomon, B.A. (retrieved 17 March 2002). *Learning styles and strategies*. Published online at ww2.ncsu.edu/unity/lockers/users/f/felder/public/ILSdir/styles.htm
- Houle, C.O. (1974). *The Design of Education*. San Francisco: Jossey-Bass.
- Martin, B.L. (1989). A checklist for designing instruction in the affective domain. *Educational Technology* 29:7–15.
- Taylor, C., Lillis, C., and LeMone (2001). *Fundamentals of Nursing: The Art and Science of Nursing Care*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.

Chapter 7

- American Academy of Family Physicians (1999). *Summary of Policy Recommendations for Periodic Health Examination*. Washington, D.C.: American Academy of Family Physicians.
- Brazelton, T. (1990). Saving the bathwater. *Child Development* 61:1661–1671.
- Burggraf, V., and Barry, R.J. (2000). Healthy People 2010—Protecting the health of older individuals. *Journal of Gerontological Nursing* 26(12):16–22.
- Child Health Guide* (2000). Washington, D.C.: U.S. Department of Health and Human Services, Public Health Service, Agency for Healthcare Research and Quality.
- Erikson, E. (1963). *Childhood and Society*, ed. 2. New York: Norton.
- Lindberg, J.B., Hunter, M.L., and Kruszewski, A.Z. (1998). *Introduction to Nursing: Concepts, Issues, and Opportunities*, ed. 3. New York: Lippincott Williams & Wilkins.
- Pender, N.J. (1996). *Health Promotion in Nursing Practice*, ed. 3. Stamford, Conn.: Appleton & Lange.
- Richardson, G.E., and Berry, N.E. (1987). Strength intervention: An approach to lifestyle modification. *Health Education* 18(3):42–46.
- Staying Healthy at 50+* (2000). Washington, D.C.: U.S. Department of Health and Human Services, Public Health Service, Agency for Healthcare Research and Quality.
- Stolte, K.M. (1996). *Wellness Nursing Diagnosis for Health Promotion*. New York: Lippincott Raven.
- U.S. Department of Health and Human Services. (November 2000). *Healthy People 2010*, ed. 2. Washington, D.C.: U.S. Government Printing Office.
- U.S. Preventative Task Force (2005). *The Guide to Clinical Preventative Services*. Washington, D.C.: U.S. Department of Health and Human Services, Public Health Service, Agency for Healthcare Research and Quality.

Chapter 8

- American Diabetes Association (1993). *Medical Management of Pregnancy Complicated by Diabetes*. Alexandria, Va.: American Diabetes Association.
- American Diabetes Association (1994). *Maximizing the Role of Nutrition in Diabetes Management*. Alexandria, Va.: American Diabetes Association.
- American Heart Association (1999). *Heart and Stroke Guide*. www-medlib.med.utah.edu/infofair/slides/hile/tsld016.htm
- Centers for Disease Control and Prevention (1999). *Overweight Prevalence Among Adults 20–74 Years Old, According to Age, Race, Sex, and Hispanic Origin, Selected Periods, 1960–1994*. <http://webapp.cdc.gov/aging/obesez.html>
- Dudek, S. (1997). *Nutrition Handbook for Nursing Practice*, ed. 3. Philadelphia: Lippincott Williams & Wilkins.
- Estes, M.E.Z. (1998). *Health Assessment and Physical Examination*. Albany, N.Y.: Delmar Publishers.
- Frisancho, A.R. (1981). New norms of upper limb fat and muscle areas for assessment of nutritional status. *American Journal of Clinical Nutrition Association* 30:2540–2548.
- Glenn, F.B., Glenn, W.D., and Duncan, R.C. (1982). Fluoride tablet supplementation during pregnancy for caries immunity: A study of the offspring produced. *American Journal of Obstetrics and Gynecology* 143(5):560–564.
- Havala, S., and Dwyer, J. (1993). Position of the American Dietetic Association: Vegetarian diets. *Journal of the American Dietetic Association* 93(11):1317–1319.
- Jarvis, C. (1996). *Physical Examination and Health Assessment*, ed. 2. Philadelphia: W.B. Saunders.
- Lee, R., and Nieman, D. (1996). *Nutritional Assessment*, ed. 2. St. Louis: C.V. Mosby.
- Leitch, I. (1942). The evolution of dietary standards. *Nutrition Abstracts and Reviews* 11:109–117.
- Looker, A.C., Dallman, P.R., Carroll, M.D., Gunter, E.W., and Johnson, C.L. (1997). Prevalence of iron deficiency in the United States. *Journal of the American Medical Association* 277(12): 973–976.
- Lutz, C., and Przytulski, K. (2001). *Nutrition and Diet Therapy*, ed. 3. Philadelphia: F.A. Davis.
- Messina, V.K., and Burke, K.I. (1997). Position of the American Dietetic Association: Vegetarian diets. *Journal of the American Dietetic Association* 97(11):1317–1321.
- Metropolitan Life Insurance Company (1999). *Height and weight tables (online)*: <http://www.metlife.com/Lifeadvice/Tools/Heightweight/Docs/frameable.html>
- Mitchell, M.K. (1994). *Nutrition Across the Lifespan*. Philadelphia: W.B. Saunders.
- National Center for Health Statistics. (1976). *NCHS Growth Charts. Monthly Vital Statistics Report* 25(3). Supplement (HRA)76–1120. Rockville, Md.: Health Resources Administration.
- National Center for Health Statistics (1998). *Health, United States, 1998, With Socioeconomic Status and Health Chartbook*. Hyattsville, Md.: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics. DHHS Publication number (PHS) 98–1232.
- National Cholesterol Education Program. (1993). *Second Report of the Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults*. Bethesda, Md.: U.S. Dept. of Health and Human Services, Public Health Service; National Institutes of Health; National Heart, Lung, and Blood Institute.
- Pipes, P.L., and Trahms, C.M. (1993). *Nutrition in Infancy and Childhood*. St. Louis: C.V. Mosby.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. F.A. Davis, Philadelphia.
- Ritchev, A.K. (1987). Iron deficiency in children. Update on an old problem. *Postgraduate Medicine* 82(2):59–63, 67–69.
- Rexrode, K.M., Carey, V.J., Hennekens, C.H., Walters, E.E., Colditz, G.A., Stampfer, M.J., Willett, W.C., and Manson, J.E. (1998). Abdominal adiposity and coronary heart disease in women. *Journal of the American Medical Association* 280(21):1843–1848.
- Rose, E.A., Porcerelli, J.H., and Neale, A.V. (2000). Pica: Common but commonly missed. *Journal of the American Board of Family Practice* 13(5):353–358.
- Rowland, M.L. (1989). A nomogram for computing body mass index; using the nomogram. *Dietetic Currents* 16(2):6–7. Columbus, Ohio: Ross Laboratories.
- U.S. Department of Agriculture and Department of Health and Human Services: *Food guide pyramid*. <http://www.usda.gov/cnpp/pyramid.html>
- U.S. Department of Agriculture (1995). *Nutrition and Your Health: Dietary Guidelines for Americans*. <http://www.hoptechno.com/dietary.htm>
- U.S. Department of Agriculture and Department of Health and Human Services (1990). *Nutrition and Your Health: Dietary Guidelines for Americans*. Home and Garden Bulletin No. 232, Washington, D.C.: U.S. Government Printing Office.
- Walker, S.P., Rimm, E.B., Ascherio, A., Kawachi, I., Stampfer, M.J., and Willett, W.C. (1996). Body size and fat distribution as predictors of stroke among U.S. men. *American Journal of Epidemiology* 144(12):1143–1150.
- Whaley, L.E., & Wong, D.L. (1991) *Nursing Care of Infants and Children*, ed. 4. St. Louis: Mosby–Year Book.
- Williams, C.L., Bollela, M., and Wynder, E.L. (1995). A new recommendation for dietary fiber in childhood. *Pediatrics* 96:985–988.
- World Health Organization (2000). Nutrition: Aims and Objectives. <http://www.who.int/nut/aim.htm>

Chapter 9

- Berger, K.J., and Williams, M.B. (1999). *Fundamentals of Nursing: Collaborating for Optimal Health*, ed. 2. Stamford, Conn.: Appleton & Lange.
- Burgess, W.A. (1997). *Psychiatric Nursing*. Stamford, Conn.: Appleton & Lange.
- Carson, V.B. (1989). *Spiritual Dimensions of Nursing Practice*. Philadelphia: W.B. Saunders.
- Carson, V.B., and Green, H. (1992). Spiritual well-being: A predictor of hardness in persons with acquired immunodeficiency syndrome. *Journal of Professional Nursing* 8(4):209–220.
- Dossey, B., and Dossey, L. (1998). Attending to holistic care. *American Journal of Nursing* 98(8):35–38.
- Ellis, J.R., and Nowlin, E.A. (1994). *Nursing: A Human Needs Approach*, ed. 5. Philadelphia: Lippincott Raven.
- Ellison, C., and Paloutzian, R. (1983). Spiritual well-being: Conceptualization and measurement. *Journal of Psychology and Theology* 11(4):330–340.
- Fehring, R.J., Brennan, P.F., and Keller, M.L. (1987). Psychological and spiritual well-being in college students. *Research in Nursing and Health* 10:391–398.
- Fehring, R.J., Miller, J.E., and Shaw, C. (1997). Spiritual well-being, religiosity, hope, depression, and other mood states in elderly people coping with cancer. *Oncology Nursing Forum* 24(4):663–671.
- Highfield, M.F. (1997). Spiritual assessment across the cancer trajectory: Methods and reflections. *Seminars in Oncology Nursing* 13(4):237–241.
- Hungelman, J., Kenkel-Rossi, E., Klassen, L., & Stollenwerk, R. (1987). Milwaukee, Wis.: Marquette University College of Nursing.
- Hungelmann, J., Kenkel-Rossi, E., Klassen, L., and Stollenwerk, R. (1996). Focus on spiritual well-being: Harmonious interconnectedness of mind-body-spirit-use of the JAREL Spiritual Well-Being Scale. *Geriatric Nursing: American Journal of Care for the Aging* 17(6):262–266.

- Kozier, B., Erb, G., Blais, K., and Wilkinson, J.M. (1998). *Fundamentals of Nursing: Concepts, Process, and Practice*, ed. 5. New York: Addison-Wesley.
- Mickley, J.R., Soeken, K., and Belcher, A. (1992). Spiritual well-being, Religiousness and hope among women with breast cancer. *IMAGE: Journal of Nursing Scholarship* 24(4):267–272.
- Moberg, D.O. (1984). Subjective measures of spiritual well-being. *Review of Religious Research* 25:351–364.
- Nagai-Jacobson, M.G., and Burkhardt, M.A. (1989) Spirituality: Cornerstone of holistic nursing practice. *Holistic Nursing Practice* 3(3):18–26.
- Potter, P.A., and Perry, A.G. (2001). *Fundamentals of Nursing*, ed. 5. St. Louis: C.V. Mosby.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Reed, P. (1986). Developmental resources and depression in the elderly. *Nursing Research* 35:368–374.
- Ross, L. (1995). The spiritual dimension: Its importance to client's health, well-being and quality of life and its implications for nursing practice. *International Journal of Nursing Studies* 32(5):457–468.
- Shelly, J.A., and Fish, S. (1988). *Spiritual Care: The Nurse's Role*, ed. 3. Downers Grove, Ill.: Intervarsity.
- Stoll, R.I. (1989). The essence of spirituality. In Carson V.B. (ed.): *Spiritual Dimensions of Nursing Practice*. Philadelphia: W.B. Saunders.

Chapter 10

- American Cancer Society. (2005). *All About Skin Cancer-Melanoma*. Available at: www.cancer.org. Last accessed January 23, 2005.
- Carpenter, D. (ed.) (1998). *Handbook of Signs and Symptoms*. Springhouse, Pa.: Springhouse Corporation.
- Epstein, O., Perkin, G., deBono, D., and Cookson, J. (1997). *Clinical Examination*, ed. 2. Philadelphia: C.V. Mosby.
- Goldsmith, L., Lazarus, G., and Tharp, M. (1997). *Adult and Pediatric Dermatology: A Color Guide to Diagnosis and Treatment*. Philadelphia: F.A. Davis.
- Goroll, A., and Mulley, A. (2000). *Primary Care Medicine*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.
- Lewis, S., Heitkemper, M., and Dirksen, S. (2000). *Medical-Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Mangione, S. (2000). *Physical Diagnosis Secrets*. Philadelphia: Hanley & Belfus, Inc.
- McPhee, S., Lingappa, V., Ganong, W., and Lange, J. (1997). *Pathophysiology of Disease: An Introduction to Clinical Medicine*. Stamford, Conn.: Appleton & Lange.
- Morton, P. (1993). *Health Assessment in Nursing*, ed. 2. Philadelphia: F.A. Davis.
- National Institutes of Health (NIH). (2000). Of Mice and Skin: Custom Rodents. COX-1 Show Value Against Skin Cancers. National Institute of Arthritis and Musculoskeletal and Skin Diseases. www.nih.gov/niams/news/spotlight/ofmice.htm
- National Institutes of Health. (2000). Scientists Scrutinize Sun and Skin. National Institute of Arthritis and Musculoskeletal and Skin Diseases. www.nih.gov/niams/news/spotlight/ofmice.htm
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Ries, L., Eisner, M., Kosary, C., Hanbey, B., Miller, B., Clegg, L., and Edwards, B. (eds.). (2000). SEER Cancer Statistics Review, 1973–1997. Bethesda, Md.: National Cancer Institute.
- Sellers, R. (1996). *Differential Diagnosis of Common Complaints*, ed. 3. Philadelphia: W.B. Saunders.
- Sommers, M., and Johnson, S. (1997). *Davis's Manual of Nursing Therapeutics for Diseases and Disorders*. Philadelphia: F.A. Davis.
- Swartz, M. (1998). *Textbook of Physical Diagnosis: History and Examination*. Philadelphia: W.B. Saunders.

- Thompson, J., McFarland, G., Hirsch, J., and Tucker, S. (1997). *Mosby's Clinical Nursing*. St. Louis: C.V. Mosby.
- Van Wynsberghe, D., Noback, C., & Carola, R. (1995). *Human Anatomy and Physiology*, ed. 3. New York: McGraw-Hill.
- Venes, D. (ed.). (2001). *Taber's Cyclopedic Medical Dictionary*, ed. 20. Philadelphia: F.A. Davis.
- Weber, J., and Kelley, J. (1998). *Health Assessment in Nursing*. New York: Lippincott Raven.
- Whitmore, S.E., Morison, W.L., Potten, C.S., and Chadwick, C. (2001). Tanning salon exposure and molecular alterations. *Journal of the American Academy of Dermatology* 44(5):755–780.

Chapter 11

- Greenlee, R., Hill-Harmon, M.B., Murray, T., and Then, M. (2001). Cancer statistics. *CA: A Cancer Journal for Clinicians* 51(1):15–36.
- Hurst, J. (1996). *Medicine for the Practicing Physician*. Stamford, Conn.: Appleton & Lange.
- Koonings, P., and Gaylis, B. (2002). Mouth cancer. *Clinical Obstetrics and Gynecology* 45(3):952–954.
- Mengel, M., and Schwieber, L. (1996). *Ambulatory Medicine: The Primary Care of Families*. Stamford, Conn.: Appleton & Lange.
- National Institutes of Health. (2000). *CancerNet: Screening for Oral Cancer*. <http://www.meb.uni-bonn.de/cancernet/304725.html>
- National Institutes of Health. (2000). *Cancer of the Larynx*. <http://www.cancer.gov/cancerinfo/wyntk/larynx>
- National Institutes of Health. (2000). *Oral Cancer*. <http://webmd.lycos.com/encyclopedia/article/2730.308>
- Pai, S., Ghezzi, E., and Ship, J. (2000). Development of a visual analogue scale questionnaire for subjective assessment of salivary function. *Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics* 91(3):311–316.
- Porth, C. (1998). *Pathophysiology: Concepts of Altered Health States*. Philadelphia: Lippincott Williams & Wilkins.
- Springhouse (2000). *Professional Guide to Signs and Symptoms*. Springhouse, Pa.: Springhouse Corporation.
- Stahl, S., Meskin, L.H., and Brown, L.J. (2004). The American Dental Association's Oral Cancer Campaign: The impact on consumers and dentists. *Journal of the American Dental Association* 135(9):1261–1267, 1303–1306.
- Swartz, M. (1998). *Textbook of Physical Diagnosis: History and Examination*. Philadelphia: W.B. Saunders.
- Thompson, J., McFarland, G., Hirsch, J., and Tucker, S. (1997). *Mosby's Clinical Nursing*. St. Louis: C.V. Mosby.

Chapter 12: The Eye

- American Academy of Ophthalmology (2005).
- Barkauskas, V., Stoltenberg-Allen, K., Baumann, L., and Darling-Fisher, C. (1998). *Health and Physical Assessment*, ed. 2. St. Louis: C.V. Mosby.
- Coleman, A.L. (1999). Glaucoma. *The Lancet* 354:1803–1810.
- Committee on Practice and Ambulatory Medicine, Section on Ophthalmology (1996). Examination and vision screening in infants, children and young adults. *Pediatrics* 98(1):153–158.
- Downey, D., and Leigh, J. (1998). Eye movements: Pathophysiology, examination, and clinical importance. *Journal of Neuroscience Nursing* 30(1):15–22.
- Fong, D.S. (2000). Age-related macular degeneration: Update for primary care. *American Family Physician* 61:3035–3042.
- Gilman, S., and Newman, S. (1996). *Essentials of Clinical Neuroanatomy and Neurophysiology*, ed. 9. Philadelphia: F.A. Davis.
- Greenstein, B., and Greenstein, A. (2000). *Color Atlas of Neuroscience*. New York: Thieme.
- Hoglund, T. (2000). *Food and Drug Administration Approves Photodynamic Therapy in the Treatment of Wet AMD*. The Foundation Fighting Blindness, <http://www.blindness.org>

- Jarvis, C. (2000). *Physical Examination and Health Assessment*, ed. 3. Philadelphia: W.B. Saunders.
- Knoop, K.J., Stack, L.B., and Storrow, A.B. (1997). *Atlas of Emergency Medicine*. New York: McGraw-Hill.
- Mir, A.M. (1995). *Atlas of Clinical Diagnosis*. Philadelphia: W.B. Saunders.
- Morton, P.G. (ed.). (1993). *Health Assessment in Nursing*, ed. 2. Springhouse, Pa.: Springhouse Corporation.
- National Eye Institute, NEI Statement, "Gene found to increase risk of the most common cause of blindness, March 10, 2005, http://w.nei.nih.gov/news/statements_amd.asp The Foundation Fighting Blindness (available online at <http://www.blindness.org>).
- Shingleton, B., and O'Donoghue, M. (2000). Blurred vision. *New England Journal of Medicine* 343(8):556–562.
- Snell, R., and Lemp, M. (1998). *Clinical Anatomy of the Eye*, ed. 2. Malden, Mass.: Blackwell Science, Inc.
- Skorin, L. (2000). Corneal and eyelid anomalies. *Consultant* 265–272.
- U.S. Department of Health and Human Services (1998). *Put Prevention Into Practice: Clinician's Handbook of Preventative Services*, ed. 2. Washington, D.C.: U.S. Government Printing Office.
- Weinstock, E.J., and Weinstock, M.B. (1996). Common eye disorders: Six clients to refer. *Postgraduate Medicine* 99(4):107–116.
- Wong, D., Wilson, D., Hockenberry-Eaton, M., Winkelstein, M., and Schwartz, P. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: C.V. Mosby.

Chapter 12: The Ear

- Barkauskas, V., Stoltenberg-Allen, K., Baumann, L., and Darling-Fisher, C. (1998). *Health and Physical Assessment*, ed. 2. St. Louis: C.V. Mosby.
- Gilman, S., and Newman, S. (1996). *Essentials of Clinical Neuroanatomy and Neurophysiology*, ed. 9. Philadelphia: F.A. Davis.
- Greenstein, B., and Greenstein, A. (2000). *Color Atlas of Neuroscience*. New York: Thieme.
- Jarvis, C. (2000). *Physical Examination and Health Assessment*, ed. 3. Philadelphia: W.B. Saunders.
- Juhn, S., Li, W., Kim, J., Javel, E., Levine, S., and Odland, R. (1999). Effects of stress-related hormones on inner ear fluid homeostasis and function. *The American Journal of Otolaryngology* 20:800–806.
- Knoop, K.J., Stack, L.B., & Storrow, A.B. (1997). *Atlas of Emergency Medicine*. New York: McGraw-Hill.
- Lang, L. (1994). Environmental impact on hearing: Is anyone listening? *Environmental Health Perspectives Supplements* 102(11):924–930.
- Meyer, C., Witte, J., Hildmann, A., Hennecke, K.H., Schunck, K.U., Maul, K., Franke, U., Fahnenstich, H., Rabe, H., Rossie, H., Hartmann, S., and Gortner, L. (1999). Neonatal screening for hearing disorders in infants at risk: Incidence, risk factors, and follow-up. *Pediatrics* 104(4):900–904.
- Mir, A.M. (1995). *Atlas of Clinical Diagnosis*. Philadelphia: W.B. Saunders.
- Moore, K.L., and Persaud, T.V. (1998). *The Developing Human: Clinically Oriented Embryology*, ed. 6. Philadelphia: W.B. Saunders.
- Prasher, D. (1998). New strategies for prevention and treatment of noise-induced hearing loss. *The Lancet* 352:1240–1242.
- Schlauch, R.S., and Samuel, L. (1995). Evaluating hearing threshold differences between ears as a screen for acoustic neuroma. *Journal of Speech and Hearing Research* 38(5):1168–1174.
- Silverthorn, D. (1998). *Human Physiology: An Integrated Approach*. Upper Saddle River, NJ: Prentice Hall.
- U.S. Department of Health and Human Services. (1998). *Put Prevention Into Practice: Clinician's Handbook of Preventative Services*, ed. 2. Washington, D.C.: U.S. Government Printing Office.
- Watts, R.L., and Koller, W.C. (1997). *Movement Disorders: Neurological Principles and Practice*. New York: McGraw-Hill.
- Yueh, B., Shapiro, N., MacLean, C., and Shellelle, P. (2003). Screening and management of adult hearing loss in primary care: Scientific Review. *Journal of the American Medical Association* 289(15):1976–1985.

Chapter 13

- Ambrose, M. (1998). Chronic dyspnea. *Nursing* 98 28(5):41–46.
- Brooks-Brunn, J. (1995). Postoperative atelectasis and pneumonia. *Heart and Lung* 24(2):94–111.
- American Cancer Society. (2005).
- Carpenito, L. (1997). *Nursing Diagnosis Application to Clinical Practice*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- Carpenter, D. (ed.). (1998). *Handbook of Signs and Symptoms*. Springhouse, Pa.: Springhouse Corporation.
- Considine, J. (2005). The role of nurses in preventing adverse events related to respiratory dysfunction: Literature review. *Journal of Advanced Nursing* 49(6):624–633.
- DeGowin, R.L. (1994). *DeGowin and DeGowin's Diagnostic Examination*, ed. 6. New York: McGraw-Hill.
- Deshpande, V., Pibeam, S., and Dixon, R. (1988). *A Comprehensive Review of Respiratory Care*. Norwalk, Conn.: Appleton & Lange.
- Doenges, M.E., and Moorhouse, M.F. (1998). *Nurse's Pocket Guide: Diagnoses, Interventions, and Rationales*. Philadelphia: F.A. Davis.
- Epstein, O., Perkin, G., deBono, D., and Cookson, J. (1997). *Clinical Examination*, ed. 2. Philadelphia: C.V. Mosby.
- Goroll, A., May, L., and Mulley, A. (1995). *Primary Care Medicine Office Evaluation and Management of the Adult Patient*, ed. 3. Philadelphia: Lippincott-Raven.
- Guyton, A.C. (1991). *Textbook of Medical Physiology*, ed. 8. Philadelphia: W.B. Saunders.
- Hanson, C. (1996). *Instant Nursing Assessment: Gerontologic*. New York: Delmar.
- Isselbacher, K., Braunwald, E., Wilson, J., Martin, J., Fauci, A., and Kasper, D. (eds.). (1995). *Harrison's Principles of Internal Medicine*, ed. 13. *Companion Handbook*. New York: McGraw-Hill.
- Marieb, E.N. (1992). *Human Anatomy and Physiology*, ed. 2. Menlo Park, Calif.: Benjamin/Cummings.
- Mezey, M., Rauckhorst, L., and Stokes, S. (1993). *Health Assessment of the Older Individual*, ed. 2. New York: Springer.
- Morton, P. (1993). *Health Assessment in Nursing*, ed. 2. Springhouse, Pa.: Springhouse.
- Polaski, A., and Tatro, S. (1996). *Luckmann's Core Principles and Practice of Medical-Surgical Nursing*. Philadelphia: W.B. Saunders.
- Scanlon, V., and Sanders, T. (1995). *Essentials of Anatomy and Physiology*, ed. 2. Philadelphia: F.A. Davis.
- Sellers, R. (1996). *Differential Diagnosis of Common Complaints*, ed. 3. Philadelphia: W.B. Saunders.
- Thomas, C.L. (ed.). (1997). *Taber's Cyclopedic Medical Dictionary*, ed. 20. Philadelphia: F.A. Davis.
- Wilkins, R., Hodgkins, J., and Lopez, B. (1996). *Lung Sounds: A Practical Guide*, ed. 2. Philadelphia: C.V. Mosby.

Chapter 14

- American Heart Association. (2005).
- Carpenter, D. (ed.). (1998). *Handbook of Signs and Symptoms*. Springhouse, Pa.: Springhouse Corporation.
- Cray, J., Kothare, V., and Weinstock, D. (eds.). (1999). *Auscultation Skills: Breath and Heart Sounds*. Springhouse, Pa.: Springhouse Corporation.
- Doenges, M., Moorehouse, M., and Geissler, A. (1997). *Nursing Care Plans: Guidelines for Individualizing Patient Care*, ed. 4. Philadelphia: F.A. Davis.
- Epstein, O., Perkin, G., de Bono, D., and Cookson, J. (1997). *Clinical Examination*, ed. 2. Philadelphia: C.V. Mosby.
- Erickson, B., Littman, D., Liebson, P., Stovall, W., and Tavel, M. (1997). *3M Education Program Cardiac Auscultation-Heart Sounds Study Guide*. St. Paul, Minn.: 3M.

- Goroll, A., and Mulley, A. (2000). *Primary Care Medicine*. Philadelphia: Lippincott Williams & Wilkins.
- Isselbacher, K., Braunwald, E., Wilson, J., Martin, J., Fauci, A., and Kasper, D. (1995). *Harrison's Principles of Internal Medicine*, ed. 15. New York: McGraw-Hill.
- Koenig, W., Loel, H., Baumert, J., and Meisinger, C. (2004). C-reactive protein modulates risk prediction based on the Framingham Score: Implications for future risk assessment: Results from a large cohort study in southern Germany. *Circulation* 109(11):1349–1353.
- Lewis, S., Heitkemper, M., and Dirksen, S. (2000). *Medical-Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Mangione, S. (2000). *Physical Diagnosis Secrets*. Philadelphia: Hanley & Belfus, Inc.
- Mason, D. (2000). *Listening to the Heart*. Philadelphia: FA. Davis.
- Morton, P. (1993). *Health Assessment in Nursing*, ed. 2. Philadelphia: FA. Davis.
- Ridker, P., Stampfer, M., and Rifai, N. (2001). Novel factors for systemic Atherosclerosis: A comparison of C-reactive protein, fibrinogen, homocysteine, lipoprotein (a), and standard cholesterol screening as predictors of peripheral arterial disease. *Journal of the American Medical Association* 285(19):2481–2485.
- Scanlon, V., and Sanders, T. (1995). *Essentials of Anatomy and Physiology*, ed. 2. Philadelphia: FA. Davis.
- Sellers, R. (1996). *Differential Diagnosis of Common Complaints*, ed. 3. Philadelphia: W.B. Saunders.
- Sommers, M., and Johnson, S. (1997). *Davis's Manual of Nursing Therapeutics for Diseases and Disorders*. Philadelphia: FA. Davis.
- Stampfer, M., Hu, F., Manson, J., Rimm, E., and Willett, W. (2000). Primary prevention of coronary heart disease in women through diet and lifestyle. *The New England Journal of Medicine* 343(1):16–22.
- Van Wynsberghe, D., Noback, C., and Carola, R. (1995). *Human Anatomy and Physiology*, ed. 3. New York: McGraw-Hill.
- Venes, D. (ed.). (2001). *Taber's Cyclopedic Medical Dictionary*, ed. 20. Philadelphia: FA. Davis.

Chapter 15

- Anand, S.S., et al. (1998). Does this patient have deep vein thrombosis? *Journal of the American Medical Association* 279:1094.
- Barkauskas, V.H., Stoltenberg-Allen, K., Bauman, L.C., and Darling-Fisher, C. (1994). *Health and Physical Assessment*. St. Louis: C.V. Mosby.
- Bichler, L.M. (1999). Foot ulcers in diabetes. *Advance for Nurse Practitioners* 7:50.
- Bickley, L.S., Hoekelman, R.A., and Bates, B. (1998). *Bates' Guide to Physical Examination and History Taking*. Philadelphia: Lippincott Williams & Wilkins.
- Blank, C.A., and Irwin, G.H. (1990). PV disorders—Assessment and intervention 7:777.
- Carman, T.L., and Fernandez, B.B. (1999). Issues and controversies in venous thromboembolism. *66*:113, 1999.
- Carpenito, L. (1997). *Nursing Diagnosis Application to Clinical Practice*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- Criqui, M.H., and Denenberg, J.O. (1999). American Heart Association's Hypertension Primer. Council on High Blood Pressure Research. Dallas: American Heart Association.
- Dains, J.E., Baumann, L.C., and Scheibel, P. (1998). *Advanced Health Assessment and Clinical Diagnosis in Primary Care*. St. Louis: C.V. Mosby.
- Ginsberg, J.S. (1996). Management of venous thromboembolism. *New England Journal of Medicine* 335:1816.
- Gray, B.H., and Sullivan, T.M. (1997). Vascular claudication: How to individualize treatment. *Cleveland Clinic Journal of Medicine* 64:429.
- Guyton, A.C. (1991). *Textbook of Medical Physiology*, ed. 8. Philadelphia: W.B. Saunders.
- Howes, P.S. (1998). Treating intermittent claudication. *RN* 98:16A.

- LeMone, P., and Burke, K.M. (1996). *Medical Surgical Nursing—Critical Thinking in Client Care*. Menlo Park, Calif.: Addison-Wesley.
- Lymphoedema Foundation at Post Office Box 834, San Diego, CA 92014–0823 or Toll-free at 1-800-LYMPH-DX.
- Mohler, E. (2003). Peripheral arterial disease: Identification and implications. *Archives of Internal Medicine* 163(19):2306–2314.
- Morton, P.G. (1995). *Health Assessment*, ed. 2. Philadelphia: Springhouse Corporation.
- Petrella, R.J. (1999). Exercise for older patients with chronic disease: *The Physician and Sportsmedicine* 27:79.
- Powers, K.B., Vacek, J.L., and Lee, S. (1999). Noninvasive approaches to PV disease. *Postgraduate Medicine* 106:52.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: FA. Davis.
- Rice, K.L. (1998). Peripheral arterial occlusive disease, part 1: Navigating a bottleneck. *Nursing98* 28:33.
- Ridker, P.M. (1997). Age-specific incidence rates of venous thromboembolism among heterozygous carriers of factor V Leiden mutation. *Annals of Internal Medicine* 126:528.
- Seidel, H.M., Ball, J.W., Dains, J.E., and Benedict, G.W. (1995). *Mosby's Guide to Physical Examination*, ed. 3. St. Louis: C.V. Mosby.
- Seller, R.H. (1996). *Differential Diagnosis of Common Complaints*. Philadelphia: W.B. Saunders.
- Taber's Cyclopedic Medical Dictionary*. Philadelphia: FA. Davis.
- UKCC Professional Development Categories. (1997). The assessment and treatment of patients with lymphoedema, *Nursing Standard* 11:49.
- Zafar, M.U., and Farkouh, M.E. (1999). Diagnosing and managing lower extremity arterial disease. *Internal Medicine* 20:31.

Chapter 16

- American Cancer Society (2002). *CA: A Cancer Journal for Clinicians Cancer Statistics 2002*. 52(1). Atlanta: American Cancer Society.
- American Cancer Society. (2005). <http://www.cancer.org>
- Furberg, A.S., Veierod, M.B., Wilsgaard, T., Bernstein, L., and Thune, I. (2004). Serum high-density lipoprotein cholesterol, metabolic profile, and breast cancer risk. *Journal of the National Cancer Institute* 96(15):1152–1160.
- Lewis, S., Heitkemper, M., & Dirksen, S. (2000). *Medical-Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Molyneux, D. (1995). "Predicting the practice and teaching of breast self-examination of student nurses." Unpublished doctoral dissertation, Widener University, Chester, Pa.
- Morton, P. (1993). *Health Assessment in Nursing*, ed. 2. Philadelphia: FA. Davis.
- Tanner, J. (1962). *Growth at Adolescence*, ed. 2. Oxford: Blackwell Scientific.

Chapter 17

- Alberts, D.S., Martinez, M.E., Roe, D.J., Guillen-Rodriguez, J.M., Marshall, J.R., Van Leeuwen, J.B., Reid, M.E., Ritenbaugh, C., Vargus, P.A., Battacharyya, A.B., Earnest, D.L., Sampliner, R.E., and the Phoenix Colon Cancer Prevention Physician's Network. (2000). Lack of effect of a high-fiber cereal supplement on the recurrence of colorectal adenomas. *The New England Journal of Medicine* 342(16):1156–1162.
- American Cancer Society (2000). The colon and rectum cancer resource center. Available at: www3.cancer.org/cancerinfo/load_cont.asp?st=ds&ct=10 and www3.cancer.org/cancerinfo/load_cont.asp?st=pr&ct=10
- American Cancer Society. (2005). <http://www.cancer.org>
- American Institute for Cancer Research. (2000). Cancer researchers agree: New colon cancer studies send wrong message. Located on the Internet at: <http://www.aicr.org/r042000.htm>

- Anderson, K. (1996). *Mosby's Medical, Nursing and Allied Health Dictionary*, ed. 4. St. Louis: C.V. Mosby.
- Barkauskas, V.H., Stoltenberg-Allen, K., Baumann, L.C., and Darling-Fisher, C. (1998). *Health and Physical Assessment*, ed. 2. St. Louis: Mosby—Year Book.
- Bates, B., Bickley, L.S., and Hoekelman, R.A. (1995). *Physical Examination and History Taking*, ed. 6. Philadelphia: Lippincott-Raven.
- Becker, K.L., and Stevens, S.A. (1988). Performing in-depth abdominal assessment. *Nursing88* 18(6):59–63.
- Black, J.M., & Matassarin-Jacobs, E. (1993). *Luckmann and Sorenson's Medical-Surgical Nursing: A Psychophysiological Approach*, ed. 4. Philadelphia: W.B. Saunders.
- Burkhart, C. (1992). Guidelines for rapid assessment of abdominal pain indicative of acute surgical abdomen. *Nurse Practitioner: American Journal of Primary Healthcare* 17(6):39, 43–46.
- Carpenito, L.J. (1997). *Nursing Diagnosis: Application to Clinical Practice*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- Centers for Disease Control and Prevention. Trends in screening for colorectal cancer—United States, 1997 and 1999. www.cdc.gov/mmwr/preview/mmwrhtml/mm5009a2.htm
- Giger, J.N., and Davidhizar, R.E. (1995). *Transcultural Nursing Assessment and Intervention*, ed. 3. St. Louis: C.V. Mosby.
- Harris, J. (1986). Pediatric abdominal assessment. *Pediatric Nursing* 12(5):355–362.
- Holmgren, C. (1992). Perfecting the art of abdominal assessment. *RN* 53(3):28–34.
- Ignatavicius, D.D., Workman, M.L., and Mishler, M.A. (1995). *Medical-Surgical Nursing: A Nursing Process Approach*, ed. 2. Philadelphia: W.B. Saunders.
- Jarvis, C. (2000). *Physical Examination and Health Assessment*, ed. 3. Philadelphia: W.B. Saunders.
- Langan, J.C. (1998). Abdominal assessment in the home: From A–Z. *Home Health-Care Nurs*, 16(1):50–58.
- Lawrence, D.M. (1993). Gastrointestinal trauma. *Critical Care Nursing Clinics of North America* 5(1):127–140.
- Metheny, N.A., Stewart, B.J., Smith, L., Yan, H., Diebold, M., and Clouse, R.E. (1999). pH and concentration of bilirubin in feeding tube aspirates as predictors of tube placement. *Nursing Research* 48(4):189–197.
- Munn, N.E. (1988). Diagnosis: Acute abdomen. *Nursing88* 18(9):34–42.
- Niles, M.A., Buffington, C., Cowan, G., and Hepworth, J.T. (1998). Comparison of lifestyles among obese and nonobese African American and European American women in the community. *Nursing Research* 47(4):251–257.
- O'Hanlon-Nichols, T. (1998). Basic assessment series: Gastrointestinal System. *American Journal of Nursing* 98(4):48–53.
- O'Toole, M.T. (1990). Advanced assessment of the abdomen and gastrointestinal problems. *Nursing Clinics of North America* 25(4):771–775.
- Piscara, V.H. (1999). Recognizing the various presentations of appendicitis. *The Nurse Practitioner* 24(8):48–55.
- Porth, C.M. (1994). *Pathophysiology Concepts of Altered Health States*, ed. 4. Philadelphia: Lippincott-Raven.
- Santilli, J., and Santilli, S. (1997). Diagnosis and treatment of abdominal aortic aneurysms. *American Family Physician [on line]* 56(4):1–12.
- Schatzin, A., Lanza, E., Corle, D., Lance, P., Iber, E., Caan, B., Shike, M., Weissfeld, J., Burt, R., Cooper, M.R., Kikendall, J.W., Cahill, J., and the Polyp Prevention Trial Study Group. (2000). Lack of effect of a low-fat, high-fiber diet on the recurrence of colorectal adenomas. *The New England Journal of Medicine* 342(16):1149–1155.
- Seidel, H.M., Ball, J.W., Dains, J.E., and Benedict, G.W. (1999). *Mosby's Guide to Physical Examination*, ed. 4. St. Louis: C.V. Mosby.
- Shaw, B. (1995). Primary care for women: Comprehensive assessment of gastrointestinal disorders. *Journal of Nurse-Midwifery* 40(2):216–230, 259–264.
- Shaw, B. (1996). Management and treatment of gastrointestinal disorders. *Journal of Nurse-Midwifery* 41(2):155–172.
- Slota, M. (1982). Abdominal assessment. *Critical Care Nurse* 2:78–81.
- Smith, C.E. (1988). Assessing bowel sounds: More than just listening. *Nursing88* 18(2):42–43.
- Stone, R. (1996). Primary care diagnosis of acute abdominal pain. *The Nurse Practitioner* 21(12):19–41.
- Thibodeau, G.A., and Patton, K.T. (1993). *Anatomy and Physiology*, ed. 2. St. Louis: Mosby—Year Book.
- Thibodeau, G.A., and Patton, K.T. (1999). *Anthony's Textbook of Anatomy and Physiology*, ed. 16. St. Louis: C.V. Mosby.
- Thompson, J.M., and Wilson, S.F. (1996). *Health Assessment for Nursing Practice*. St. Louis: Mosby—Year Book.
- Town, J. (1997). Bringing acute abdomen into focus. *Nursing97* 27(5):52–58.
- Walsh, J., Posner, S., and Perez-Stable, E., (2002). Colon cancer screening in the ambulatory setting. *Preventive Medicine* 35(3):209–218.
- Wong, D. (1999). *Nursing Care of Infants and Children*, ed. 6. St. Louis: Mosby.
- Wright, J. (1997). Seven abdominal assessment signs every emergency nurse should know. *Journal of Emergency Nursing* 23:446–450.
- Zator Estes, M.E. (1998). *Health Assessment and Physical Examination*. Albany, N.Y.: Delmar Publishers.

Chapter 18

- Allen, K., and Phillips, J. (1997). *Women's Health Across a Lifespan*. Philadelphia: Lippincott Williams & Wilkins.
- Barkauskas, V., Stoltenberg-Allen, K., Baumann, L., and Darling-Fisher, C. (1998). *Health and Physical Assessment*, ed. 2. St. Louis: C.V. Mosby.
- Bates, B. (1999). *A Guide to Physical Examination*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- Centers for Disease Control and Prevention (1998). *Guidelines for Treatment of Sexually Transmitted Diseases. MMWR Morbidity and Mortality Weekly Report* Rep. 47(RR-1):1–118.
- Harlan, L.C., Bernstein, A.M., and Kessler, L.F. (1991). Cervical cancer screening. *Journal of Public Health* 81(7):855–890.
- Hawkins, J., and Roberto-Nichols, D. (1997). *Protocols for Nurse Practitioners in Gynecologic Settings*. Tiresias Press.
- Morton, P.G. (1993). *Health Assessment in Nursing*, ed. 2. Springhouse, Pa.: Springhouse Corporation.
- Noble, J. (1996). *Textbook of Primary Care Medicine*. St. Louis: C.V. Mosby.
- Sawaya, G., McConnell, K., Kulasingam, S., Lawson, H., Kerlikowske, K., Melnikow, J., Lee, N., Gildengorin, G., Meyers, E., and Washington, E. (2003). Risk of cervical cancer associated with extending the interval between cervical cancer screening. *The New England Journal of Medicine* 349(16):1501–1509.
- Seidel, H., et al. (1999). *Mosby's Guide to Physical Exam*. St. Louis: C.V. Mosby.

Chapter 19

- American Academy of Pediatrics (1999). *Periodic survey of fellows #37, pediatrician attitudes, experiences mixed on circumcision counseling*. <http://www.aap.org/research/p537a.htm>
- American Cancer Society. (1994). *Cancer Facts and Figures*. [On-line]. Internet Address: bcic/acsz/gifs/fact.map
- Bates, B. (1999). *A Guide to Physical Examination*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- Centers for Disease Control and Prevention (1998). Guidelines for the treatment of sexually transmitted diseases. *MMWR Morbidity and Mortality Weekly Report*, Rep 47(RR-1):1–118.
- Gilligan, T., Wang, P., Levin, R., Kantoff, P., and Avorn, J. (2004). Racial differences in screening for prostate cancer in the elderly. *Archives of Internal Medicine* 164(17):1858–1864.

- Holmes, K.K., March, P.A., and Sparling, P.F. (1990). *Sexually Transmitted Diseases*, ed. 2. New York: McGraw-Hill.
- Jaffe, M., and McVan, B. (1997). *Davis's Laboratory and Diagnostic Test Handbook*. Philadelphia: F.A. Davis.
- Lerman, S.E., & Liao, J.C. (2001). Neonatal circumcision. *Pediatric Clinics of North America* 48(6):1539–1557.
- Lewis, S., McLean Heitkemper, M., and Ruff Dirksen, S. (2000). *Medical Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Morton, P.G. (1993). *Health Assessment in Nursing*, ed. 2. Philadelphia: F.A. Davis.
- Smith, R., Malkoicz, S., Whittington, R., Van Arsdalen, K., Tochner, Z., and Wein, A. (2004). Identification of clinically significant prostate cancer by prostate-specific antigen screening. *Archives of Internal Medicine* 164(11):1227–1230.
- Tanner, J.M. (1962). *Growth at Adolescence*, ed. 2. Oxford: Blackwell Scientific Publications.
- Walsh, P.C., Retik, A.P., Stamey, T.A., and Vaughn, E.D. (1992). *Campbell's Urology*, ed. 6. Philadelphia: W.B. Saunders.
- Wynsberghe, D., Noback, C., and Carola, R. (1995). *Human Anatomy and Physiology*, ed. 3. New York: McGraw-Hill.

Chapter 20

- Alexander, M., and Kuo, K.N. (1997). Musculoskeletal assessment of the newborn. *Orthopaedic Nursing* 16(1):21–31.
- Allison, M., and Keller, C. (1997). Physical activity in the elderly: Benefits and intervention strategies. *Nurse Practitioner* 22(8):53–69.
- Behrman, R.E., and Vaughan, V.C. (eds). (1996). *Nelson Textbook of Pediatrics*, ed. 15. Philadelphia: W.B. Saunders.
- Benetti, M.C., and Marchese, T. (1996). Management of common musculoskeletal disorders. *Journal of Nurse-Midwifery* 41(2):173–187.
- Birnbaum, J.S. (1976). *The Musculoskeletal Manual*. Orlando, Fla.: Grune & Stratton.
- Chestnut, C. (2001). Osteoporosis, an underdiagnosed disease. *Journal of the American Medical Association* 286(22):2865–2866.
- DeGowin, E., and DeGowin, R. (1987). *Bedside Diagnostic Examination*, ed. 5. New York: Macmillan.
- Gordon, M. (1979). Assessing activity tolerance. *American Journal of Nursing* 76:72–75.
- <http://www.arthritis.org>
- http://www.cdc.gov/arthritis/arthritis/risk_factors.htm
- Hunt, A.H. (1996). The relationship between height change and bone mineral density. *Orthopaedic Nursing* 15(3):57–64.
- Ignatavicius, D., and Workman, M. (2002). *Medical-Surgical Nursing: Critical Thinking for Collaborative Care*, ed. 4. Philadelphia: Saunders.
- Jones, A.K. (1997). Primary care management of acute low back pain. *Nurse Practitioner* 22(7):50–68.
- Jones-Walton, P. (1990). Orthopaedic nursing assessment. *Advanced Clinical Care* 5(3):22.
- Kemmler, W., Engelke, K., Lauber, D., Weineck, J., Hensen, J. and Kalender, W. (2002). Exercise effects on fitness and bone mineral density in early postmenopausal women: 1-year EFOPS results. *Medicine and Science in Sports & Exercise* 34(12):2115–2123.
- Killam, P. (1989). Orthopaedics assessment of young children: Developmental variations. *Nurse Practitioner* 14(7):27–32.
- Lewis, S., Heitkemper, M., and Dirksen, S. (2001). *Medical Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Maldonado, A. (1995). Comprehensive assessment of common musculoskeletal disorders. *Journal of Nurse-Midwifery* 40(2):202–215.
- Mason, K. (1989). Pediatric orthopaedics: Developmental norms. *Orthopedic Nursing* 8(4):45–50.
- McIntosh, E. (1997). Low back pain in adults: Guidelines for the history and physical exam. *Advanced Nursing Practice* 5(8):16–25.

- Milde, F. (1988). Impaired physical mobility. *Journal of Gerontological Nursing* 14(3):20–24, 38–40.
- Muscari, M.E. (1998). Preventing sports injuries. *American Journal of Nursing* 98(7):58–60.
- O'Hanlon-Nichols, T. (1998). A review of the adult musculoskeletal system. *American Journal of Nursing* 98(6):48–52.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Riggs, B., and Melton, L. (1995). The worldwide problem of osteoporosis: Insights afforded by epidemiology. *Bone* 17(Suppl. 5):505S–511S.
- Ross, C.A. (1997). A comparison of osteoarthritis and rheumatoid arthritis: Diagnosis and treatment. *Nurse Practitioner* 22(9):20–41.
- Siris, E.S., Miller, P.D., Barrett-Connor, E., Abbot, T., Sherwood, L., and Berge, M. (2001). Identification and fracture outcomes of undiagnosed low bone mineral density in postmenopausal women: Results from the National Osteoporosis Risk Assessment. *Journal of the American Medical Association* 286:2815–2822.
- Trembl, L.A. (1996). Assessing patient mobility: Mobility screening a part of a community-based geriatric assessment. *Home Care Provider* 1(1):26–29.
- Shaefer, K.M. (1997). Health patterns of women with fibromyalgia. *Journal of Advanced Nursing* 26:565.
- Spence, A., and Mason, E. (1987). *Human Anatomy and Physiology*, ed. 3. Menlo Park, Calif.: Benjamin-Cummings.
- Wallace, M. (1994). Assessment and management of pain in the elderly. *MEDSURG Nursing* 3:293–298.
- Williams, M.A., Oberst, M.T., Bjorklund, B.C., and Hugher, S.H. (1996). Family caregiving in cases of hip fracture. *Rehabilitation Nursing* 21(3):124–131, 138.
- Woodhead, G.A. and Moss, N.M. (1998). Osteoporosis: Diagnosis and prevention. *Nurse Practitioner* 23(11):18–37.

Chapter 21

- Carrozzella, J., and Jauch, E. (2002). Emergency stroke management: A new era. *Nursing Clinics of North America* 37(1):35–58.
- Folstein. (1975).
- Goroll, A., and Mulley, A. (2000). *Primary Care Medicine*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.
- Hock, N. (1999). Brain attack: The stroke continuum. *Nursing Clinics of North America* 34(3):689–724.
- Huff, J. Farace, E., Brady, W., Kheir, J., and Shaver, G. (2001). The Quick Confusion Scale in the ED: Comparison with the Mini-Mental State Examination. *The American Journal of Emergency Medicine* 19(6).
- Iron, M.J., Farace, E., Brady, W.J., and Huff, J.S. (2002). Mental status screening of emergency department patients: Normative study of the Quick Confusion Scale. *Academic Emergency Medicine* 9(10):989–994.
- Lewis, S., Heitkemper, M., and Dirksen, S. (2001). *Medical Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Lower, J. (2002). Facing neuro assessment fearlessly. *Nursing* 2002 32(2):58–64.
- Morton, P. (1995). *Health Assessment in Nursing*, ed. 2. Philadelphia: F.A. Davis.
- National Institute of Neurological Diseases, U.S. Department of Health and Human Services, National Institutes of Health: Stroke scales.
- Neatherlin, J. (1999). Foundation for practice: Neuroassessment for neuroscience nurse. *Nursing Clinics of North America* 34(3):573–592.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: F.A. Davis.
- Ringleb, P., Schellinger, P., Schranz, C., and Hacke, W. (2002). Thrombolytic therapy within 3 to 6 hours after the onset of ischemic stroke: Useful or harmful? *Stroke* 33(5):1437–1441.
- Scanlon, V., and Sanders, T. (1995). *Essentials of Anatomy and Physiology*, ed. 2. Philadelphia: F.A. Davis.

- Thomas, C. (ed.). (1997). *Taber's Cyclopedic Medical Dictionary*, ed. 18. Philadelphia: FA. Davis.
- U.S. National Institutes of Health, NINDS (1999–2000). *Brain basics: Preventing stroke*. Bethesda, Md.: NIH Neurological Institute.
- Van Wynsberghe, D., Noback, C., and Carola, R. (1995). *Human Anatomy and Physiology*, ed. 3. New York: McGraw-Hill.
- Wong, D., and Hockenberry-Eaton, M. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: Mosby.

Chapter 22

- AWHONN. (1999). *Monograph I: Postmenopausal health and the importance of prevention*. Chicago: Pragmaton.
- AWHONN (1999). *Monograph II: Disease prevention strategies among postmenopausal women*. Chicago: Pragmaton.
- Carpenito, L. (1997). *Nursing Diagnosis: Application to Clinical Practice*, ed. 7. Philadelphia: Lippincott Williams & Wilkins.
- DeGowin, R.L. (1994). *DeGowin and De Gowin's Diagnostic Examination*, ed. 6. New York: McGraw-Hill.
- Doenges, M.E., Moorhouse, M.F., and Geissler, A.C. (1997). *Nursing Care Plans: Guidelines for Individualizing Patient Care*, ed. 4. Philadelphia: FA. Davis.
- Goroll, A., May, L., and Mulley, A. (1995). *Primary Care Medicine: Office Evaluation and Management of the Adult Patient*, ed. 3. Philadelphia: Lippincott-Raven.
- Morton, P. (1993). *Health Assessment in Nursing*, ed. 2. Springhouse, Pa.: Springhouse Corporation.
- North American Nursing Diagnosis Association (2001). *NANDA Nursing Diagnoses: Definitions and Classification, 2001–2002*. Philadelphia: NANDA.
- Polanski, A., and Tatro, S. (1996). *Luckmann's Core Principles and Practice of Medical-Surgical Nursing*. Philadelphia: W.B. Saunders.
- Scanlon, V., and Saunders, T. (1999). *Essentials of Anatomy and Physiology*, ed. 3. Philadelphia: FA. Davis.
- Taylor, C., Lillis, C., and LeMone, P. (2001). *Fundamentals of Nursing: The Art and Science of Nursing Care*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.
- U.S. Preventive Services Task Force. (1996). *Guide to Clinical Preventive Services: An Assessment of 169 Interventions*. Baltimore, Md.: Williams & Wilkins.
- U.S. Preventive Services Task Force. (2005). <http://www.ahrq.gov/clinic/uspstfix.htm>.
- Venes, D. (ed.). (2001). *Taber's Cyclopedic Medical Dictionary*, ed. 19. Philadelphia: FA. Davis.

Chapter 23

- Carcio, H.A. (1999). *Advanced Health Assessment of Women*. Philadelphia: Lippincott Williams & Wilkins.
- Cunningham, F.G., MacDonald, P.C., Gant, N.F., Leveno, K.J., Gilstrap L. C., III, Hankins, G.D.V., and Clark, S.L. (1997). *Williams' Obstetrics*, ed. 20. Stamford, Conn.: Appleton & Lange.
- Darcus, J.V., Meyer, N L., and Sibai, B.M. (1996). How preconception counseling improves pregnancy outcome. *Contemporary OB/GYN* 40(6):11–115.
- Ferguson, H.W., & Hisley, S.M. (2000). *Nurse Test: Maternal-Newborn Nursing*, ed. 2. Springhouse, Pa.: Springhouse Corporation.
- Gorrie, T.M., McKinney, E S., and Murray, S.S. (1998). *Foundations of Maternal-Newborn Nursing*, ed. 2. Philadelphia: W.B. Saunders.
- Hoebel, C.J. (1998). Routine antenatal laboratory tests and specific screening tests. *Contemporary OB/GYN* 56(2):25–34.
- Jarvis, C. (1996). *Physical Examination and Health Assessment*, ed. 2. Philadelphia: W.B. Saunders.
- Lowdermilk, D.L., Perry, S.E., and Bobak, I.M. (1997). *Maternity and Women's Health Care*, ed. 6. St. Louis: C.V. Mosby.
- Niebyl, J. (1995). Folic acid supplementation to prevent birth defects. *Contemporary OB/GYN* 40(6):43–39.

- O'Brien, B., and Zhou, Q. (1995). Variables related to nausea and vomiting during pregnancy. *Birth* 22(2):93–96.
- Pillitteri, A. (1995). *Maternal & Child Health Nursing*. Philadelphia: Lippincott-Raven.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: FA. Davis.
- Rogers, J., and Davis, B. (1995). How risky are hot tubs and saunas for pregnant women? *MCN American Journal of Maternal Child Nursing* 20(3):137–139.
- Simpson, K.R., and Creehan, P.A. (1996). *AWHONN Perinatal Nursing*. Philadelphia: Lippincott-Raven.
- Spector, R.E. (1991). *Cultural Diversity in Health and Illness*, ed. 3. Norwalk, Conn.: Appleton & Lange.
- Weber, J., and Kelley, J. (1998). *Health Assessment in Nursing*. Philadelphia: Lippincott Williams & Wilkins.
- Wheeler, L. (1997). *Nurse Mid-wifery Handbook: A Practical Guide to Prenatal and Postpartum Care*. Philadelphia: Lippincott-Raven.
- Youngkin, E.Q., and Davis, M.S. (1998). *Women's Health: A Primary Care Clinical Guide*, ed. 2. Stamford, Conn.: Appleton & Lange.

Chapter 24

- Ballard, J., et al. (1991). New Ballard score, expanded to include extremely premature infants. *Journal of Pediatrics* 119(3):417–423.
- Barkauskas, V., Stoltenberg-Allen, K., Baumann, L., and Darling-Fisher, C. (1998). *Health and Physical Assessment*, ed. 2. St. Louis: C.V. Mosby.
- Creehan, P. (2001/2002). Sending baby home safely. *Lifeline* 5(6):60–70.
- Dickason, E., Lang Silverman, B., and Kaplan, J. (1998). *Maternal-Infant Nursing Care*, ed. 3. Philadelphia: C.V. Mosby.
- Fuller, J., and Schaller-Ayers, J. (2000). *Health Assessment: A Nursing Approach*. Philadelphia: Lippincott Williams & Wilkins.
- Gardner, P., and Georges, P. (2001). Recommended schedules for routine immunization of children and adults. *Infectious Disease Clinics of North America* 15(1):1–8.
- Holcomb, S. (2005). Managing jaundice in full-term infants. *The Nurse Practitioner The American Journal of Primary Health Care* 30(1):6–12.
- Lowdermilk, D.L., Perry, S E., and Bobak, I.M. (2000). *Maternity and Women's Health Care*, ed. 7. St. Louis: C.V. Mosby.
- National Highway Traffic Safety Administration. (2000). Traffic Safety Facts for 1999. Washington, D.C.: U.S. Department of Transportation. <http://www.nsc.org/lrs/statinfo/99report.htm>
- Nichols, F.H., & Zwelling, E. (1997). *Maternal-newborn Nursing: Theory and Practice*. Philadelphia: W.B. Saunders.
- Olds, S.B., London, M.L., and Weiland Ladewig, P.A. (2000). *Maternal Newborn Nursing: A Family- and Community-Based Approach*, ed. 6. Upper Saddle River, N.J.: Prentice Hall Health.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: FA. Davis.
- Weber, J.R. (2001). *Nurses' Handbook of Health Assessment*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.
- Wilson, D.D. (1999). *Nurses' Guide to Understanding Laboratory and Diagnostic Tests*. Philadelphia: Lippincott Williams & Wilkins.
- Wilson, S.F., and Giddens, J.F. (2001). *Health Assessment for Nursing Practice*, ed. 2. St. Louis: C.V. Mosby.
- Wong, D., and Hockenberry-Eaton, M. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: C.V. Mosby.

Chapter 25

- American Academy of Pediatrics (1991). *Caring for Your Adolescent: Ages 12–21*. New York: Bantam Books.
- American Medical Association (1999). *Implementation Guide for School-Based Health Centers*. American Medical Association.

- Berti, L., Zylbert, S., and Rolnitzky, L. (2001). Comparison of health status of children using a school-based health center for comprehensive care. *Journal of Pediatric Health Care* 15(5):244–250.
- Dennison, B., Russo, T., Burdick, P., and Jenkins, P. (2004). An intervention to reduce television viewing by preschool children. *Archives Pediatric and Adolescent Medicine* 158(2):170–176.
- Erikson, E. (1968). *Identity Youth and Crisis*. New York: W.W. Norton.
- Green, M. (ed.). (1994). *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. Arlington, Va.: National Center for Education in Maternal and Child Health.
- Mobley, C., and Evanshevski, J. (2000). Evaluating health and safety knowledge of preschoolers: Assessing their early start to begin health smart. *Journal of Pediatric Health Care* 4(4):160–165.
- Morton, P. (1995). *Health Assessment in Nursing*. Philadelphia: E.A. Davis.
- Seidel, H.M., Ball, J.W., Dains, J.E., and Benedict, G.W. (1999). *Mosby's Guide to Physical Examination*, ed. 4. St. Louis: C.V. Mosby.
- Thomas, C. (ed.). (1997). *Taber's Cyclopedic Medical Dictionary*, ed. 18. Philadelphia: E.A. Davis.
- U.S. Department of Health and Human Services. (1997). *Put Prevention Into Practice*. Washington, D C : USDHHS.
- Wong, D., Hockenberry-Eaton, M., Wilson, D., Winkelstein, M., and Schwartz, P. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: C.V. Mosby.

Chapter 26

- American Academy of Pediatrics. (1991). *Caring for Your Adolescent: Ages 12–21*. New York: Bantam Books.
- American Medical Association. (1999). *Implementation Guide for School-Based Health Centers*. American Medical Association.
- Green, M. (ed.). (1994). *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. Arlington, Va.: National Center for Education in Maternal and Child Health.
- Hardy, L., Harrell, J., and Bell, R. (2004). Overweight children: Definitions, measurements, confounding factors and health consequences. *Journal of Pediatric Nursing* 19(6):376–384.
- Mobley, C., and Evanshevski, J. (2000). Evaluating health and safety knowledge of preschoolers: Assessing their early start to begin health smart. *Journal of Pediatric Health Care* 14(4):160–165.
- Morton, P. (1995). *Health Assessment in Nursing*. Philadelphia: E.A. Davis.
- Seidel, H.M., Ball, J.W., Dains, J.E., and Benedict, G.W. (1999). *Mosby's Guide to Physical Examination*, ed. 4. St. Louis: C.V. Mosby.
- Thomas, C. (ed.). (1997). *Taber's Cyclopedic Medical Dictionary*, ed. 18. Philadelphia: E.A. Davis.
- U.S. Department of Health and Human Services. (1997). *Put Prevention Into Practice*. Washington, D.C.: USDHHS.
- Using a school-based health center for comprehensive care. *Journal of Pediatric Health Care* 15(5):244–249.
- Wong, D., Hockenberry-Eaton, M., Wilson, D., Winkelstein, M., and Schwartz, P. (2001). *Wong's Essentials of Pediatric Nursing*, ed. 6. Philadelphia: C.V. Mosby.

Chapter 27

- Abdul, A., and Mourad, F. (1998). Constipation: Common-sense care of the older client. *Geriatrics* 51:28–36.
- Advisory Committee on Immunization Practices, Centers for Disease Control and Prevention. <http://www.americangeriatrics.org/imm-sched.html>
- Agachan, F., Chen, T., Pfeifer, J., Reissman, P., and Wexner, S. (1996). A constipation scoring system to simplify evaluation and management of constipated clients. *Diseases of the Colon and Rectum* 39:681–685.
- Alessi, C., Yoon, E., Schnelle, J., Al-Samarrai, N., and Cruise, P. (1999). A randomized trial of a combined physical activity and environmental intervention in nursing home residents: Do sleep and agitation improve? *Journal of the American Geriatric Society* 47:784–792.

- American Cancer Society. (1983). Mammography guidelines. *California Cancer Journal Clinicals* 33:255–259.
- American Heart Association. (2005).
- American Medical Association, Council on Scientific Affairs. (1997). Mammographic screening in asymptomatic women aged 40 years and older. *Journal of the American Medical Association* 261:2535–2542.
- Barthel DW, et al (1965) Functional Evaluation: The Barthel Index. *Maryland State Medical Journal* 14: 61–65.
- Bartkiw, T., and Pynn, B. (1994). Burning mouth syndrome: An overlooked condition in the geriatric population. *Geriatric Nursing* 15:241–245.
- Benson, S. (1975). Simple chronic constipation: Pathophysiology and management. *Post-graduate Medicine* 5:55–60.
- Bravo, G., Gauthier, P., Roy, P., Payette, H., Gaulin, P., Harvey, M., Peloquin, L., and Dubois, M. (1996). Impact of a 12-month exercise program on the physical and psychological health of osteopenic women. *Journal of the American Geriatrics Society* 44:756–762.
- Buchner, D., Cress, M., Wagner, E., de Lateur, B., Price, R., and Abrass, I. (1993). The Seattle FICSIT/MoveIt study: The effect of exercise on gait and balance in older adults. *Journal of the American Geriatrics Society* 41:321–325.
- Cassell, C., Cohen, H., Larson, E., Resnick, N., Rubenstein, L., and Sorensen, L. (1996). *Geriatric Medicine*. New York: Springer.
- Chandler, J. (1996). Understanding the relationship between strength and mobility in frail older persons: A review of the literature. *Topics of Geriatric Rehabilitation* 11:20–37.
- Chaouloff, F. (1997). Effects of acute physical exercise on central serotonergic systems. *Medicine and Science in Sports and Exercise* 29:58–62.
- Clark, D. (1999). Physical activity and its correlates among urban primary care clients aged 55 years or older. *The Journal of Gerontology* 54B:S41–S48.
- Costanza, M. (1992). Breast cancer screening in older women: Synopsis of a forum. *Cancer* 69:1925–1931.
- Daly, M. (2000). Nutritional status and involuntary weight loss. In Adelman, A., and Daly, M. (eds): *20 Common Problems in Geriatrics*. New York: McGraw-Hill, pp. 337–367.
- Dishman, R. (1994). Motivating older adults to exercise. *Southern Medical Journal* 87:S79–S82.
- Dobbs, R., Chalett, A., and Bowes, S. (1993). Is this walk normal? *Age and Ageing* 22:25–30.
- Dungan, J., Brown, K., and Ramsey, B. (1996). Health maintenance for independent frail older adults: Can it improve physical and mental health? *Journal of Advanced Nursing* 23:1185–1193.
- Dunlap, R. (1997). Teaching advance directives: The why, when and how. *Journal of Gerontological Nursing* 23:11–16.
- Duke University Center for the Study of Aging and Human Development, The multidimensional Functional Assessment Questionnaire, ed. 2. Durham, N.C.: Duke University, pp. 169–170; 154–156; 157–162.
- Elliott, J. (1988). Swallowing disorders in the elderly: A guide to diagnosis and treatment. *Geriatrics* 43:95–99.
- Elward, K., and Larson, E. (1992). Benefits of exercise of older adults. *Clinics of Geriatric Medicine* 8:35–50.
- Emanuel, E., and Emanuel, L. (1992). Proxy decision making. *Journal of the American Medical Association* 267:2221–2226.
- Erikson, E. (1982). *The Life Cycle Completed: A Review*. New York: Morton.
- Ettinger, W. (1996). Physical activity and older people: A walk a day keeps the doctor away. *Journal of the American Geriatrics Society* 44:207–208.
- Folstein, M., Folstein, S., and McHugh, P. (1975). Mini-Mental State: A practical method for grading the cognitive state of clients for the clinician. *Journal of Psychiatric Research* 12:189–198.
- Goldberg, T., and Chavin, S. (1997). Preventive medicine and screening in older adults. *Journal of the American Geriatrics Society* 45:341–354.
- Goroll, A., & Mulley, A. (2000). *Primary care medicine*, ed. 4. Philadelphia: Lippincott Williams & Wilkins.

- Gregg, E., Cauley, J., Seeley, D., Ensrud, K., and Bauer, D. (1998). Physical activity and osteoporotic fracture risk in older women. *Annals of Internal Medicine* 129:81-88.
- Gruber, M., and Lance, P. (1998). Colorectal cancer detection and screening. *Lippincott's Primary Care Practice* 2:369-376.
- Gudas, S. (1996). Implications of oncology in the aged. In Lewis, C. (ed.): *Aging: The Health Care Challenge*. Philadelphia: FA. Davis, pp. 225-260.
- Hanson, C. (1996). *Gerontologic Instant Nursing Assessment*. New York: Delmar.
- Heuvelen, M., Kemper, G., Orem, H., and Rispen, P. (1998). Physical fitness related to age and physical activity in older persons. *Medicine and Science in Sports and Exercise* 30:434-441.
- Hogan, T. (1996). Geriatric emergencies. In Cassel C., et al. (eds.): *Geriatric Medicine*. New York: Springer-Verlag, 1996.
- Jacoby, S. (1999). Great sex: What's age got to do with it? *Modern Maturity* 9:43-45.
- Johanson, J. E. (1998). Geographic distribution of constipation in the United States. *The American Journal of Gastroenterology* 93:188-191.
- Kane, R. (1996). The defeat of aging versus the importance of death. *Journal of the American Geriatrics Society* 44:321-325.
- Kaplan, G., Strawbridge, W., Camacho, T., and Cohen, R. (1993). Factors associated with change in physical functioning in the elderly. *Journal of Aging and Health* 5:140-153.
- Katz, S., Ford, A., and Moskowitz, R. (1963). Studies of illness in the aged: The index of ADL. *Journal of the American Medical Association* 185:914-919.
- King, A., Oman, R., Brassington, G., Bliwise, D., and Haskell, W. (1997). Moderate-intensity exercise and self-rated quality of sleep in older adults. *Journal of the American Medical Association* 257:32-37.
- Lachs, M., Feinstein, A., and Cooney, L. A. (1993). A simple procedure for general screening of functional disability in elderly clients. *Annals of Internal Medicine* 112:699-702.
- Lavker, R. (1995). Cutaneous aging: Chronologic versus photoaging. In Gilchrist B. (ed.): *Photodamage*. Boston: Blackwell Sciences, pp. 125-133.
- Lewis, S., Heitkemper, M., and Dirksen, S. (2000). *Medical Surgical Nursing Assessment and Management of Clinical Problems*, ed. 5. Philadelphia: C.V. Mosby.
- Lichtenstein, M., Bess, E., and Logan, S. (1988). Validation of screening tools for identifying hearing-impaired elderly in primary care. *Journal of the American Medical Association* 259:2875-2878.
- Mahoney, E., and Barthel, D. (1965). Functional evaluation: The Barthel Index. *Maryland State Medical Journal* 14(2):61-65.
- Mahoney, J., Drinka, T., Ablner, R., Gunter-Hunt, G., Matthews, C., Gravenstein, S., and Carnes, M. (1994). Screening for depression: Single question versus GDS. *Journal of The American Geriatrics Society* 42:1006-1007.
- Maslow, A. (1954). *Motivation and Personality*. New York: Harper & Row.
- Mayfield, D., McLeod, G., and Hall, P. (1974). The CAGE questionnaire: Validation of a new alcoholism screening instrument. *American Journal of Psychology* 131:1121-1123.
- Meador, J. (1995). Cerumen impaction in the elderly. *Journal of Gerontological Nursing* 25:43-45.
- Metter, E., Lynch, N., Conwit, R., Lindle, R., Tobin, J., and Hurley, B. (1997). Muscle quality and age: Cross-sectional and longitudinal comparisons. *International Journal of Sports Medicine* 3S225-S231.
- Mezey, M., Rauchhorst, L., and Stokes, S. (1993). *Health Assessment of the Older Individual*, ed. 2. New York: Springer.
- Miller, C. (1990). *Geriatric Nursing* 11(6):301-302.
- Morey, M., Pieper, C., and Cornoni-Huntley, J. (1998). Physical fitness and functional limitations in community-dwelling older adults. *Medicine & Science in Sports & Exercise* 30:715-723.
- Morrell, C. H., Gordon-Salant, S., Pearson, J. D., Brant, L. J., and Fozard, J. L. (1996). Age- and gender-specific reference ranges for hearing level and longitudinal changes in hearing level. *Journal of Acoustic Society of America* 100(4 Pt 1):1949-1967.
- Morris, B., Van Niman, S., Perlin, T., Lucie, K., Vieth, J., Agricola, K., and McMurray, M. (1995). Healthcare professionals? Accuracy in predicting clients? Preferred code status. *The Journal of Family Practice* 40:41-44.
- Nahm, E., and Resnick, B. (In press). End of life care preferences of older adults. *Nursing Ethics*.
- National Institutes of Health Consensus Development Conference Statement. (1995). Physical activity and cardiovascular health, pp 1-12.
- Naylor, M., Stephens, C., Bowles, K., and Bixby, B. (2005). Cognitively impaired older adults: From hospital to home: An exploratory study of these patients and their caregivers. *American Journal of Nursing* 105(2):52-61.
- Nesher, G., and Moore, T. (1994). Clinical presentation and treatment of arthritis in the elderly. *Clinics in Geriatric Medicine* 10:659-675.
- Odenheimer, G., Funkenstein, H., and Beckett, L. (1994). Comparison of neurologic changes in successfully aging persons vs. the total aging population. *Archives of Neurology* 51:573-580.
- Okumiya, K., Matsubayashi, K., Wada, T., Kimura, S., Doi, Y., and Ozawa, T. (1996). Effects of exercise on neurobehavioral function in community-dwelling older people more than 75 years of age. *Journal of the American Geriatrics Society* 44:569-572.
- Parkatti, T., Deeg, D., Bosscher, R., and Launer, L. (1998). Physical activity and self-rated health among 55- to 89-year-old Dutch people. *Journal of Aging and Health* 10:311-326.
- Parker, J., Vukov, L., and Wollan, P. (1996). Abdominal pain in the elderly: Use of temperature and laboratory testing to screen for surgical disease. *Family Medicine* 28:193-197.
- Pate, R., Pratt, M., and Blair, S. (1995). Physical activity and public health: A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *Journal of the American Medical Association* 253:402-407.
- Pearson, J. D., Morrell, C. H., Gordon-Salant, S., Brant, L. J., Metter, E. J., Klein, L. L., and Fozard, J. L. (1995). Gender differences in a longitudinal study of age-associated hearing loss. *Journal of Acoustic Society of America* 97(2):1196-1205.
- Pescatello, L., DiPietro, L., Fargo, A., Ostfeld, A., and Nadel, E. (1994). The impact of physical activity and physical fitness on health indicators among older adults. *Journal of Aging and Physical Activity* 2:2-13.
- Potter, P., and Perry, A. (1999). *Basic Nursing*, ed. 4. Philadelphia: C.V. Mosby.
- Province, M., Hadley, E., Hornbrook, M., Lipsitz, L., Miller, J., Mulrow, C., Ory, M., Sattin, R., Tinetti, M., and Wolf, S. (1995). The effects of exercise on falls in elderly clients. *Journal of the American Medical Association* 253:1341-1347.
- Purnell, L. D., and Paulanka, B. J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: FA. Davis.
- Resnick, B. (1998). Functional performance of older adults in a long-term care setting. *Clinical Nursing Research* 7:230-246.
- Resnick, B. (1998). Health care practices of the old-old. *American Academy Journal of Nurse Practitioners* 10:147-155.
- Resnick, B. (1999). Atrial fibrillation in the older adult: Presentation and Management. *Geriatric Nursing* 20:1-6.
- Resnick, B. (2000). Health promotion practices of the older adult. *Public Health Nursing* 17(3):160-168.
- Resnick, B. (2001). Geriatric health promotion. *Medscape Nursing* 1(1) @2001 Medscape, Inc.
- Rodeheffer, R., Gerstenblith, G., and Becker, L. (1994). Exercise cardiac output is maintained with advancing age in healthy subjects: Cardiac dilation and increased volume compensate for a diminished heart rate. *Circulation* 69:203-213.
- Royall, D., Mulroy, A., Chiodo, L., and Polk, M. (1999). Clock drawing is sensitive to executive control: A comparison of six methods. *Journal of Gerontology* 54B:P328-P333.
- Saltzstein, S., Behling, C., and Baergen, R. (1998). Features of cancer in nonagenarians and centenarians. *Journal of the American Geriatrics Society* 46:994-998.
- Scanlon, V., and Sanders, T. (1995). *Essentials of Anatomy and Physiology*, ed. 2. Philadelphia: FA. Davis.

- Schiffman, S. (1997). Taste and smell losses in normal aging and disease. *Journal of the American Medical Association* 256:1357–1360.
- Seeman, T., Charpentier, P., Berkman, L., Tinetti, M., Guralnik, J., Albert, M., Blazer, D., and Rowe, J. (1993). Predicting changes in physical performance in a high-functioning elderly cohort: MacArthur studies of successful aging. *Journal of Gerontology* 49(3):M97–M107.
- Sherman, J., Abel, E., and Tavakoli, A. (1996). Demographic predictors of clinical breast examination, mammography, and Pap test screening among older women. *Journal of the American Academy of Nurse Practitioners* 8(5):231–236.
- Ship, J.A., Pearson, J.D., Cruise, L.J., Brant, L.J., and Metter, E.J. (1996). Longitudinal changes in smell identification. *Journal of Gerontology, Series A, Biological Sciences and Medicine and Medical Sciences* 51(2):M86–M91.
- Sonnenberg, A., and Koch, T. (1989). Physician visits in the United States for constipation. *Digestive Diseases and Science* 34:606–611.
- Talley, N., Flemin, K., Evans, J., O'Keefe, E., Weaver, A., Sinsmeister, A., and Melton, L. Constipation in an elderly community: A study of prevalence and potential risk factors. *The American Journal of Gastroenterology* 91:19–25.
- Teno, J., Branco, K., Mor, V., Phillips, C., Hawes, C., Morris, J., and Fries, B. (1997). Changes in advance care planning in nursing homes before and after the client self-determination act: Report of a 10-state survey. *Journal of the American Geriatrics Society* 45:939–944.
- Tinetti, M., and Ginter, S. (1988). Identifying mobility dysfunctions in elderly clients: Standard neuromuscular examination or direct assessment? *Journal of the American Medical Association* 259:1190–1193.
- Van Wynsberghe, D., Noback, C., and Carola, R. (1995). *Human Anatomy and Physiology*, ed. 3. New York: McGraw-Hill.
- Venes, D. (ed.). (2001). *Taber's Cyclopedic Medical Dictionary*, ed. 19. Philadelphia: FA. Davis.
- Watson, Y., Arfken, C., and Birge, S. (1993). Clock completion: An objective screening test for dementia. *Journal of the American Geriatrics Society* 41:1235–1240.
- Weinberg, A., Pals, J., McGlinchey-Berroth, R., and Minaker, K. (1994). Indices of dehydration among frail nursing home clients: Highly variable but stable over time. *Journal of the American Geriatrics Society* 42:1070–1073.
- Weinstein, B. (1994). Age-related hearing loss: How to screen for it and when to intervene. *Geriatrics* 49:40–46.
- Whitehead, W., Chaussade, S., Corazziari, E., and Kumar, D. (1991). Report of an international workshop on management of constipation. *Gastroenterology International* 4:99–113.
- Whitehead, W., Drinkwater, W., and Cheskin, L. (1989). Constipation in the elderly living at home: Definition, prevalence, and relationship to lifestyle and health status. *Journal of the American Geriatrics Society* 37:423–429.
- Yesavage, J., Brink, T., Rose, T., Lum, O., Huang, V., Adey, M., and Leirer, O. (1983). Development and validation of a geriatric depression screening scale: A preliminary report. *Journal of Psychiatric Research* 17:37–49.
- Young, D., Appel, L., Jee, S., and Miller, E. (1999). The effects of aerobic exercise and tai chi on BP in older people: Results of a randomized trial. *Journal of the American Geriatrics Society* 47:257–284.

Appendix C Guidelines for Cultural Assessment

- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: FA. Davis.

Appendix C Guidelines for Environmental Assessment

- Narayan, M. (1997). Environmental assessment. *Home Healthcare Nurse* 15(11):798.
- Purnell, L.D., and Paulanka, B.J. (2003). *Transcultural Healthcare: A Culturally Competent Approach*, ed. 2. Philadelphia: FA. Davis.
- Smith, C., and Maurer, E. (2000). *Community Health Nursing Theory and Practice*, ed. 2. Philadelphia: W.B. Saunders.

Appendix C Guidelines for Assessing the Homeless Person

- Hitchcock, J., Schubert, P., and Thomas, S. (1999). *Community Health Nursing Caring in Action*. Albany, N.Y.: Delmar.
- O'Connell, J.J., and Groth, J. (1991). *The Manual of Common Communicable Diseases in Shelters*. Boston: Boston Health Care for the Homeless Program.
- Smith, C., and Maurer, E. (2000). *Community Health Nursing Theory and Practice*, ed. 2. Philadelphia: W.B. Saunders.
- The Stewart B. McKinney Homeless Assistance Act. (1987). Public Law 100–107.
- Web Sites:
 Healthcare for the Homeless Information Resource Center: General information, research, reports, bibliographies, history, links.
 The National Coalition for the Homeless: information about the homeless condition, legislation, and police.

Appendix D Genetics

- American College of Medical Geneticists (ACMG). [Online]. <http://www.acmg.net/>
- American Society of Human Geneticists (ASHG). [Online]. <http://genetics.faseb.org/genetics/ashg/ashgmenu.htm>
- Bennett, R.L., Steinhaus, K.A., Uhrich, S.B., O'Sullivan, C.K., Resta, R.G., Lochner-Doyle, D., Markel, D.S., Vincent, V., and Hamanishi, J. (1995). Recommendations for standardized human pedigree nomenclature. Pedigree Standardization Task Force of the National Society of Genetic Counselors. *American Journal of Human Genetics* 56(3):745–752.
- DNai. [Online]. www.dnai.org
- Genetics Home Reference (<http://ghr.nlm.nih.gov/ghr/glossary/proband>) retrieved 13 January 2006.
- Infogenetics. [Online]. <http://www.infogenetics.org>
- International Society of Nurses in Genetics (ISONG). [Online]. www.isong.org
- Jenkins, J., and Lea, D. (2005). *Nursing Care in the Genomic Era: A Case Based Approach*. Boston: Jones & Bartlett.
- Lashley, E.R. (2005). *Clinical Genetics in Nursing Practice*, ed. 3. New York: Springer.
- National Society of Genetic Counselors (NSGC). [Online]. www.nsgc.org
- Online Mendelian Inheritance in Man (OMIM). <http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=OMIM>
- Snustad, D.P., and Simmons, M.J. (2003). *Principles of Genetics*. New York: Wiley & Sons.
- U.S. Surgeon General's Family History tool [Online]. <http://www.hhs.gov/familyhistory/>

Illustration Credits

Chapter 1

Figure 1-1 p. 4: From Doenges, M., Moorehouse, M., & Geissler, A. (1997). *Nursing Care Plans: Guidelines for Individualizing Patient Care*, ed. 5. Philadelphia: FA. Davis, p. 7, with permission.

Chapter 4

Figure 4-1 p. 88: From Wilkinson, J., Van Leuven, K. (2007). *Fundamentals of Nursing*. Philadelphia: FA. Davis, with permission.

Figure 4-2 p. 90: From Wilkinson, J., Van Leuven, K. (2007). *Fundamentals of Nursing*. Philadelphia: FA. Davis, with permission.

Figure 4-3 p. 92: From Wilkinson, J., Van Leuven, K. (2007). *Fundamentals of Nursing*. Philadelphia: FA. Davis, with permission.

Figure 4-4 p. 100: *Wong's Essentials of Pediatric Nursing*, ed. 6. St. Louis: Mosby, p. 1301. Copyright by Mosby, Inc. Reprinted with permission.

Unnumbered Figure 4-2 p. 100: Tesler, M. et al. (1991) The word-graphic rating scale as a measure of children's and adolescents' pain intensity. *Research in Nursing and Health* 14:361-371.

Chapter 5

Figure 5-1 p. 113: From Scanlon, V.C. (1997). *Understanding Human Structure and Function*. Philadelphia: FA. Davis.

Figure 5-2 p. 113: Adapted from Scanlon, V.C., & Sanders, T. (2003). *Essentials of Anatomy and Physiology*, ed. 4. Philadelphia: FA. Davis.

Chapter 8

Figure 8-1 p. 166: Courtesy of United States Department of Agriculture and Department of Health and Human Services.

Chapter 10

Figure 10-4 p. 230: Courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA.

Figure 10-5 p. 230: Goldsmith, L.A., Lazarus, G.S., & Tharp, M.D. (1997): *Adult and Pediatric Dermatology: A Color Guide to Diagnosis and Treatment*, Philadelphia: FA. Davis, with permission.

Chapter 11

Art for Inspection of the Head and Face, pp. 289–290: Acromegaly, Cushing's syndrome courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA. Bell's palsy courtesy of Wills Eye Hospital, Philadelphia, PA.

Art for Inspection of the Nose, pp. 294–295: Polyps and Deviated septum courtesy of J. Barabe, CMSP Clearing House.

Art for Inspection of the Mouth and Throat, pp. 297–304: Cheilitis, Allergic response, Cancer on lip, Malocclusion, Tetracycline staining, Fluorosis, Leukemia, Early HIV periodontitis, Advanced HIV periodontitis, Lichen planus, Leukoplakia, Cancer on oral mucosa, Fordyce granules, Torus palatinus, Torus mandible, Cocaine use, Geographic tongue, Parotitis, Glossitis, Cancer of the tongue, Herpangioma courtesy of Tina S. Liang, DMD, Bala Cynwyd, PA. Chancre, Aphthous ulcer from Goldsmith, L.A., Lazarus, G.S., & Tharp, M.D. (1997). *Adult and Pediatric Dermatology: A Color*

Guide to Diagnosis and Treatment. Philadelphia, FA. Davis, with permission. Dental caries, Gum hyperplasia, Gingival recession, Chronic gingivitis, HIV palatal candidiasis courtesy of Robert A. Levine, DDS, PC, Philadelphia, PA, and Sheryl Radin, DDS, Pediatric Dentistry, Red, "beefy" tongue courtesy of SPL, CMSP Clearing House. Black, hairy tongue courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA. Enlarged tonsils with exudates courtesy of SIU BIOMED, CMSP Clearing House.

Art for Inspection of the Neck, p. 307: Cervical adenitis courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA.

Art for Common Abnormalities, pp. 315–318: Acute cervical adenitis courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA. Polyps courtesy of J. Barabe, CMSP Clearing House. Tonsillitis, Enlarged tonsils with exudates courtesy of SIU BIOMED, CMSP Clearing House.

Chapter 12

Figure 12-6 p. 326: From Scanlon, V.C., & Sanders, T. (1999). *Essentials of Anatomy and Physiology*, ed. 3. Philadelphia: FA. Davis.

Figure 12-11 p. 341: From Scanlon, V.C., & Sanders, T. (1999). *Essentials of Anatomy and Physiology*, ed. 3. Philadelphia: FA. Davis.

Figure 12-13 p. 355: Courtesy of Wills Eye Hospital, Philadelphia, PA. **Art of Assessing the Extraocular Muscles, pp. 358:** Exotropia, Congenital, exotropia courtesy of Wills Eye Hospital, Philadelphia, PA.

Art for Inspecting the External Structures, pp. 347–354: Soft contact lenses, Hard contact lenses, Lice, Blepharconjunctivitis, Entropion, Espophthalmos, Ptosis, Squamous cell carcinoma, Xanthelasma, Chalazion, Hordeolum, Dacryocystitis, Acute allergic conjunctivitis, Pterygium, Pinguecula, Subconjunctival hemorrhage, Nevus, Papilloma, Diffuse episcleritis, Arcus senilis, Corneal abrasion, Healing corneal ulcer, Mature cataract, Anisocoria, Adie's pupil, Hypopyon, Hyphema, Acute angle closure courtesy of Wills Eye Hospital, Philadelphia, PA.

Art for Performing an Ophthalmic Examination, pp. 356–358: Normal fundus, Acute papilledema, Chronic papilledema, Glaucomatous optic nerve, Optic neuritis, Optic nerve pallor, Hypertensive changes, Diabetic retinopathy, Malignant hypertension, Age-related macular degeneration, Advance macular degeneration courtesy Wills Eye Hospital, Philadelphia, PA.

Art for Common Abnormalities (eyes) pp. 360–362: All figures courtesy Wills Eye Hospital, Philadelphia, PA.

Art for Focused Assessment of the External Ear: Congenital ear anomaly and low-set ears from *Variations and Minor Departures in Infants*, Mead Johnson Nutritional Division, Evansville, IN, with permission.

Art for Performing an Otoscope Examination, p. 379–381: Exostosis, Foreign body, External otitis, Normal tympanic membrane, Serous otitis, Otitis media, Perforated TM, Hemotympanum, Adhesive otitis media, Cholesteatoma from *A Guide to the Use of Diagnostic Instruments in Eye and Ear Examinations*, Welch Allyn, Inc., Skaneateles Falls, NY, with permission.

Art for Common Abnormalities (ears) p. 388: All figures courtesy Ann Marie Ramsey, RN, MSN, CPNP, Section of Pediatric otolaryngology, University of Michigan Medical Center.

Chapter 15

Art for Inspecting the Veins and Lymph Nodes, pp. 508–509: Lymphedema courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA. Venous stasis ulcer and Arterial insufficiency from © 3M Health Care, used with permission.

Chapter 18

Art for Inspecting the External Genitalia, pp. 631–632: Pubic lice courtesy of Wills Eye Hospital, Philadelphia, PA. Herpes vulvovaginitis courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA.

Art for Speculum Examination of the Internal Genitalia, pp. 634–635: Chadwick's sign courtesy of Women's Health Care Group, Philadelphia, PA.

Chapter 19

Art for Inspecting the Genitalia, pp. 668–671: Chancere, Genital herpes from Goldsmith, L., Lazarus, G., & Tharp, M. (1997). *Adult and Pediatric Dermatology: A Color Guide to Diagnosis and Treatment*. Philadelphia: FA. Davis, p. 464. Chancroid, Genital warts on penis, Genital warts on scrotum, Tinea cruris courtesy of Hahnemann University Department of Dermatology, Philadelphia, PA.

Art for Common Abnormalities, pp. 680–682: Chancroid, Genital warts courtesy of Hahnemann University Department of Dermatology, Philadelphia, PA. Chancere, Genital herpes from Goldsmith, L., Lazarus, G., & Tharp, M. (1997). *Adult and Pediatric Dermatology: A Color Guide to Diagnosis and Treatment*. Philadelphia: FA. Davis, p. 464.

Chapter 20

Art for Assessing Joints and Muscles, pp. 717–728: rheumatoid arthritis, Heberden's nodes, Gouty arthritis, Callus courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA.

Chapter 23

Art for Head-to-Toe Physical Assessment, pp. 834–841:

Chadwick's sign, Circular cervical opening: nulliparous, Slit-like cervical opening: multiparous courtesy of Women's Health Care Group, Philadelphia, PA.


Chapter 24

Figure 24-1, p. 852: From Venes, D. (ed.) (2001): *Taber's Cyclopedic Medical Dictionary*, ed. 19. Philadelphia: FA. Davis, p. 425.

Chapter 27

Art for Head-to-Toe Physical Assessment, pp. 949–953:

Acrocordon, senile purpura, Heberden's nodes courtesy of MCP-Hahnemann University Department of Dermatology, Philadelphia, PA. Arcus senilis courtesy of Wills Eye Hospital, Philadelphia, PA.



INDEX

Note: Page numbers followed by f refer to figures; page numbers followed by b refer to boxes; and page numbers followed by t refer to assessment tables.

- A**
- A-delta fibers, 89
- Abdomen, 551–610. *See also* Gastrointestinal system
- acute, 564
 - in African Americans, 561t
 - anatomy of, 556–558, 556f, 556t–557t
 - mapping for, 576–577, 576f, 577f
 - in Asian Americans, 562t
 - assessment of, 562–602
 - abdominal reflexes in, 596, 597t–598t
 - anatomical landmarks in, 576, 576f
 - anatomical mapping for, 576–577, 576f, 577f
 - approach to, 577–578
 - auscultation in, 581, 585, 585f, 586t–587t
 - Ballance's sign in, 597, 598t–599t
 - ballottement in, 596–597, 598t
 - biographical data in, 562–563
 - bowel movements in, 565–566, 565b
 - cardiovascular system in, 572t, 580t
 - case study of, 562, 575–576, 602–603
 - checklist for, 608–609
 - cultural considerations in, 561, 561t–562t
 - current health status in, 563
 - developmental considerations in, 559, 561
 - drugs in, 568t–569t
 - endocrine system in, 572t, 582
 - family history in, 569, 570t
 - in female genitourinary disorders, 630t
 - focused history in, 563b
 - gastrointestinal system in, 572t, 582
 - general survey in, 571t, 578–579, 579t
 - genitourinary system in, 582
 - head-to-toe, 579, 579t–581t
 - health history in, 562–574
 - HEENT system in, 571t, 580t, 582
 - hematologic system in, 573t, 582
 - immunologic system in, 573t
 - indigestion in, 566
 - in infants, 881t–882t
 - inspection in, 581, 583t–586t
 - in integumentary disorders, 225
 - integumentary system in, 571t, 579t–580t, 582
 - Kehr's sign in, 597, 598t
 - lymphatic system in, 582
 - McBurney's sign in, 599t
 - for menopause, 805t
 - in mental health disorders, 127t
 - Murphy's sign in, 597–598, 599t
 - musculoskeletal system in, 572t, 581t, 582
 - nausea in, 566
 - neurologic system in, 572t, 581t, 582
 - in newborns, 864t
 - pain in, 563–564, 564b, 565b
 - palpation in, 591–595, 592t–595t
 - past health history in, 566, 567t–568t
 - in patient teaching, 151t
 - percussion in, 588, 588f, 589t–590t
 - in peripheral-vascular disorders, 508t
 - physical examination in, 577–603
 - in postpartal women, 844t
 - in pregnancy, 836t–839t
 - psychosocial profile in, 573, 573t–574t
 - rebound tenderness in, 601t
 - reproductive system in, 572t, 581t, 582
 - respiratory system in, 572t, 580t
 - review of systems in, 571, 571t–573t
 - symptom analysis in, 563–566
 - vomiting in, 566
 - weight in, 565, 565b
 - body systems interaction with, 558–559, 560
 - in children, 559, 561
 - circumference of, in newborns, 857t
 - contour of, 584t–585t
 - distension of, 584t–585t
 - dullness of, 595–596, 596f, 597, 598t
 - in infant, 559
 - in Jewish Americans, 562t
 - in Mediterranean Americans, 562t
 - in Native Americans, 562t
 - nine regions of, 577, 577f, 578b
 - in older adults, 561
 - pain in, 563–564, 564b, 565b
 - in pregnant women, 561
 - quadrants of, 576, 576f, 577b
 - pain in, 564, 564b
 - shapes of, 584t–585t
 - surface movements of, 585t
 - symmetry of, 584t
 - vascular sounds of, 585, 585f, 586t–587t
 - Abdominal aorta, auscultation of, 516t, 587t
 - Abdominojugular (hepatojugular) reflux, 473t, 476
 - Abducens nerve (CN VI), 328, 746t, 747f, 772t–773t
 - Abduction, 715f, 716t
 - Abscess, breast, 552t
 - Abstract thinking, in neurologic system
 - assessment, 767t, 771
 - Abuse, 1033–1036
 - child, 115, 889, 889b, 1033–1035
 - elder, 117, 1036
 - sexual, 1034–1035
 - spousal (partner), 1035
 - Accessory muscles, 394, 396, 398f
 - Accessory nerve (CN XI), 746t, 747f, 776t
 - Accommodation, in vision, 328, 353t
 - Acetic acid wash, 637, 639t
 - Acetylcholine, 112b
 - Achilles tendon, 687f
 - Achilles tendon squeeze, 769
 - Acid-base balance, 395–396
 - Acidosis, metabolic, 395–396
 - Acne, 211
 - Acne vulgaris, 235t, 257t
 - Acoustic nerve (CN VIII), 366f, 746t, 747f, 775t
 - Acoustic neuroma, 388t
 - Acrochordon, 949t, 956t
 - Acrocyanosis, in newborns, 850t, 859t, 860t
 - Acromegaly, 289t
 - Acromion process, 687f
 - Actinic keratosis, 211, 956t
 - Action cues, 8
 - Active listening, in patient interview, 10
 - Activities of daily living, 944, 944b–946b, 945t
 - Acute stress disorder, 139t
 - Adaptation, in vision, 328
 - Addiction, 133. *See also* Substance abuse
 - Addison's disease, 228t
 - Adduction, 715f, 716t
 - Adenitis, cervical, 307t, 315t
 - Adie's pupil, 352t
 - Adipose tissue, 206f, 207t
 - Adolescents, 913–925
 - assessment of, 915–923
 - biographical data in, 915
 - breast in, 917t, 922t
 - cardiovascular system in, 917t, 922t
 - case study of, 915, 924
 - communication in, 916b
 - cultural considerations in, 914, 915t
 - current health status in, 916
 - endocrine system in, 918t
 - family history in, 916
 - gastrointestinal system in, 918t, 922t
 - general survey in, 917t, 920f, 920t
 - genitourinary system in, 918t, 922t–923t
 - health history in, 915–916
 - HEENT system in, 917t, 921t
 - hematologic system in, 918t
 - integumentary system in, 917t, 921t
 - lymphatic system in, 918t
 - musculoskeletal system in, 918t, 923t
 - neurological system in, 918t, 923t
 - past health history in, 916
 - physical, 920, 920t–923t
 - psychosocial profile in, 916, 919–920
 - reproductive system in, 918t, 922t–923t
 - respiratory system in, 917t, 921t
 - review of systems in, 916, 917t–918t
 - blood pressure in, 494
 - breast in, 532, 532b
 - developmental changes in, 914
 - eating disorders in, 913
 - emotional changes in, 914
 - emotional disorders in, 913
 - exercise for, 160, 919
 - genitalia of, 654, 656t
 - health concerns in, 924–925
 - immunizations for, 924–925
 - injury in, 162, 913
 - integumentary system of, 211
 - neurologic system of, 749, 751
 - nutrition for, 167, 168t–169t, 919
 - obesity in, 913, 924
 - physical changes in, 914
 - pregnancy in, 818–819, 913
 - psychosocial changes in, 914
 - puberty in, 615, 617b
 - risk-taking behavior by, 115, 115b, 919
 - scoliosis in, 691
 - self-concept of, 919
 - sexuality of, 919–920
 - sexually transmitted diseases in, 913
 - sleep for, 159, 919

- spirituality in, 189
- stages of, 914
- stress in, 161, 920
- substance abuse in, 913
- suicide risk in, 115b
- violence by, 913–914
- wellness assessment in, 157
- Adrenal gland, 558f
 - hyperfunction of, 209
 - hypofunction of, 209
 - in pregnancy, 818t
- Adrenergic agents
 - neurologic system effects of, 755t
 - respiratory system effects of, 402t
- Adrenergic blockers, neurologic system effects of, 755t
- Adrenocorticotropic hormone,
 - musculoskeletal system effects of, 696t
- Advance directive, 946
- Affective skills, for assessment, 13
- Affirmation, in patient interview, 10
- African Americans
 - cardiovascular disease in, 454
 - childbearing in, 819t
 - eye examination in, 330, 360
 - facial skin of, 275
 - gastrointestinal symptoms in, 561t
 - integumentary system of, 211
 - musculoskeletal health problems in, 694t
 - neurologic disorders in, 751
 - newborn, 853t
 - nutrition and, 171t
 - older adult, 932t
 - pain response in, 96t
 - spirituality and, 191t
- Afterload, 444, 475–476t
- Age spots, 211
- Aging. *See* Older adults
- Agnosia, 766b
- Agnosticism, 198t
- AIDS (acquired immunodeficiency syndrome), 644
 - in newborns, 874t
 - palatal candidiasis in, 301t
 - periodontitis in, 299t
- Airway, 394, 394f. *See also* Respiratory system
- Akinesia, 766b
- Ala, 268f
- Alaskans, otitis media in, 370
- Albinism, 229t
- Alcohol use/abuse, 135t, 461t
 - nutrition and, 172t
 - screening for, 156t
 - sleep and, 158
- Alimentary canal, 556
- Alkalosis, metabolic, 396
- Allen cards, 58t
- Allen test, 329, 517b
- Allergic rhinitis, 316t
- Alopecia, 214, 252, 253t
 - adrenal hypofunction and, 209
 - cicatricial, 214
 - noncicatricial, 214
- Alopecia areata, 253t
- Alopecia universalis, 347t
- Alpha-fetoprotein, 833b
- Alveolus, 394f, 395t
 - fibrosis of, 398
- Alzheimer's disease, 789t–790t
- Amblyopia, 895t
- Amenorrhea, 619
- Amiloride, nutrition and, 172t
- Amino acids, 89
- Aminoglycosides
 - ear effects of, 372t
 - nutrition and, 172t
- Amish
 - adolescent, 915t
 - childbearing in, 819t
 - mental disorders and, 118t
 - musculoskeletal health problems in, 694t
 - newborn, 853t
 - nutrition and, 171t
 - school-age, 903t
 - spirituality and, 191t
 - views of women and, 618t
- Amphetamines
 - abuse of, 135t
 - bruxism with, 279t
 - sleep and, 159b
- Amphiarthrosis, 690t
- Amphotericin B, nutrition and, 172t
- Anagen, 214
- Anal culture, 637, 639t
- Analgesics
 - ear effects of, 372t
 - gastrointestinal system effects of, 568t
 - respiratory system effects of, 402t
- Androgens
 - breast effects of, 535t
 - female genitourinary system effects of, 621t
- Anemia, iron deficiency, 184t–185t
- Aneurysm, 521t, 605t
- Angina, 469t
 - Ludwig's, 317t
 - Prinzmetal's, 461t
- Angina pectoris, 487t
- Angioedema, 297t
- Angioma
 - cherry, 233t
 - during pregnancy, 814t
- Angle of Louis, 464f, 471t
- Anglican Church, 195t–196t
- Aniridia, 351t
- Anisocoria, 351t
- Ankle, 725t–726t
- Ankle-brachial index, 518b
- Ankle clonus, 785t
- Ankylosing spondylitis, 732, 732t–733t
- Anorexia, physiological, in toddlers, 892
- Anorexia nervosa, 141t–142t, 183t–184t, 580t, 608t
 - in adolescent, 913
- Anosmia, 772t
- Antacids
 - gastrointestinal system effects of, 568t
 - nutrition and, 172t
- Anterior chamber, 323f, 325t
 - inspection of, 353t
- Anterior cruciate ligament, 729
- Anterior spinothalamic tract, 742, 742f, 742t
- Anterior triangle, 286, 286f
- Anthropometry, 83–84, 84f
 - in adults, 180–182, 180f, 181t
 - in children, 179
 - in nutrition assessment, 177, 179–182
- Anti-infectives
 - ear effects of, 372t
 - eye effects of, 333t
 - gastrointestinal system effects of, 569t
 - integumentary system effects of, 216t
 - lymphatic system effects of, 500t
 - neurologic system effects of, 755t
 - nutrition and, 172t
 - respiratory system effects of, 402t
- Antiandrogens, male sexual function effects of, 659t
- Antianginals, neurologic system effects of, 755t
- Antianxiety agents
 - abuse of, 135t
 - cardiovascular system effects of, 457t
 - male sexual function effects of, 659t
- Antiarrhythmics
 - eye effects of, 333t
 - lymphatic system effects of, 499t
 - neurologic system effects of, 755t
 - respiratory system effects of, 402t
- Antiasthmatics, cardiovascular system effects of, 457t
- Anticholinergics
 - eye effects of, 333t
 - gastrointestinal system effects of, 568t
 - head, face, and neck effects of, 278t
 - male sexual function effects of, 659t
 - side effects of, 123t
- Anticonvulsants
 - head, face, and neck effects of, 278t
 - integumentary system effects of, 215t
 - lymphatic system effects of, 499t
 - male sexual function effects of, 660t
 - musculoskeletal system effects of, 696t
 - neurologic system effects of, 755t
 - side effects of, 123t
- Antidepressants
 - breast effects of, 535t
 - cardiovascular system effects of, 457t
 - female genitourinary system effects of, 621t
 - gastrointestinal system effects of, 568t
 - head, face, and neck effects of, 278t
 - male sexual function effects of, 660t
 - musculoskeletal system effects of, 696t
 - neurologic system effects of, 755t
 - sleep and, 159b
- Antidiabetics
 - gastrointestinal system effects of, 568t
 - integumentary system effects of, 216t
 - lymphatic system effects of, 499t
- Antifungals, cardiovascular system effects of, 457t
- Antigout agents, musculoskeletal system effects of, 697t
- Antihelix, 366f
- Antihistamines, head, face, and neck effects of, 278t
- Antihypertensives
 - female genitourinary system effects of, 621t
 - gastrointestinal system effects of, 569t
 - head, face, and neck effects of, 278t
 - lymphatic system effects of, 500t

- Antihypertensives (*continued*)
 male sexual function effects of, 660t
 neurologic system effects of, 755t
 respiratory system effects of, 402t
- Antitilepemics
 musculoskeletal system effects of, 697t
 nutrition and, 172t
- Antimalarials
 ear effects of, 372t
 integumentary system effects of, 215t
- Antineoplastics
 cardiovascular system effects of, 457t
 eye effects of, 333t
 female genitourinary system effects of, 621t
 gastrointestinal system effects of, 569t
 head, face, and neck effects of, 278t–279t
 integumentary system effects of, 215t
 lymphatic system effects of, 500t
 neurologic system effects of, 755t
 nutrition and, 172t
 respiratory system effects of, 402t
- Antiparkinsonian agents, neurologic system effects of, 755t
- Antipsychotics
 breast effects of, 535t
 cardiovascular system effects of, 457t
 eye effects of, 333t
 female genitourinary system effects of, 621t
 lymphatic system effects of, 500t
 male sexual function effects of, 660t
 neurologic system effects of, 755t
 side effects of, 123t
- Antipyretics, ear effects of, 372t
- Antisocial personality disorder, 140t
- Antitragus, 366f
- Antitubercular agents
 eye effects of, 333t
 gastrointestinal system effects of, 569t
- Anus, 612f, 631t, 651f
 in infants, 882t
- Anxiety, 139t
- Anxiolytics
 abuse of, 135t
 cardiovascular system effects of, 457t
 male sexual function effects of, 659t
- Aorta, 559t
 abdominal
 aneurysm of, 521t, 605t
 auscultation of, 516t, 587t
 palpation of, 591, 591t, 593, 593t
 dissection of, 521t
 thoracic, aneurysm of, 521t
- Aortic stenosis, 480t
- Aortic valve, 438f, 439t
- Apgar score, 854, 854t
- Aphasia, 766b
- Apical impulse, 474–475
- Apley's test, 730, 730f
- Apnea, 81, 415b
- Apneustic center, 396
- Apneustic respiration, 415b
- Apocrine glands, 206f, 207t, 208, 211
- Appalachians
 adolescent, 915t
 childbearing in, 819t–820t
 mental disorders and, 118t
 newborn, 853t
 nutrition and, 171t
 older adult, 932t
 pain response in, 96t
 respiratory disease in, 398
 school-age, 903t
 spirituality and, 191t
 views of women and, 618t
- Appendicitis, 599, 599t–601t, 605t
- Appendix, 556f, 557t
 in pregnant women, 561
- Apraxia, 766b
- Aqueous humor, 325t
- Arab Americans
 adolescent, 915t
 childbearing in, 820t
 mental disorders and, 118t
 newborn, 853t
 nutrition and, 171t
 pain response in, 96t
 school-age, 903t
 spirituality and, 191t
 views of women and, 618t
- Arcus senilis, 330, 351t, 469t, 950t
- Areolae, 528f, 530t, 545t
- Argyll Robertson pupil, 353t
- Arm
 circumference of, 705, 714t
 inspection of, 511t–512t
 length of, 713t–714t
 lymph nodes of, 512t
 midarm circumference of, 181, 181f
 midarm muscle circumference of, 181
 pronator drift test for, 728, 728f
 range of motion of, 721t
 triceps skin fatfold measurement in, 180, 180f
- Arrector pili muscle, 206f, 208t
- Arsenic poisoning, 229t
- Arterial embolism, 522t
- Arterial insufficiency, 509t
- Arterial pulses. *See* Pulse(s)
- Arterial thrombosis, 521t
- Arterioles, 441b, 441f
 cutaneous, 207t
- Arteriosclerosis obliterans, 522t
- Artery
 anatomy of, 441b, 441f
 pressure in, 450, 451f
- Arthritis, 732
 gouty, 727t
 juvenile, 732
 rheumatoid, 727t, 732
- Articulation, 686, 690t. *See also* Musculoskeletal system
- Arytenoid cartilage, 395t
- Ascites, 595–596, 595f, 596f
- Asian Americans. *See also* Chinese Americans; Vietnamese Americans
 cerumen in, 370
 gastrointestinal symptoms in, 562t
 integumentary system of, 211
- Assessment, 3–24, 4f. *See also at specific systems and patient populations*
 affective/interpersonal skills for, 13
 clinical decision making for, 12
 cognitive skills for, 11–12, 12b
 comprehensive, 17
 critical thinking for, 11, 11b–12b
 data collection for, 17–20
 ethical skills for, 13
 focused, 17
 medical vs. nursing, 14, 16
 nurse role in, 14–16, 15b
 physical. *See* Physical assessment
 problem-solving skills for, 12–13
 psychomotor skills for, 13
 purpose of, 11
 skills for, 11–13
 tips for, 20b
- Asthma, 432t
- Astigmatism, 326, 326f, 360t
- Ataxia, 709t
- Atheism, 198t
- Athlete, heart rate in, 80
- Athlete's foot, 263t
- Atopy, 209, 258t
- Atrial fibrillation, 472f
- Atrial kick, 442
- Atrial septal defect, 446
- Atrioventricular (AV) node, 444f, 445
- Atrium
 left, 438f, 439t
 right, 438f, 439t
- Audamina, 859t
- Audioscopy, 387
- Auditory stimuli, in neurologic system assessment, 764, 769
- Auricle, 365t, 366f
- Auscultation, 20, 71–72, 72b
 abdominal, 581, 585, 585f, 586t–587t
 abdominal aorta, 516t, 587t
 apical, 80–81, 81f, 482t
 bowel sounds, 586t
 in newborns, 864t
 cardiac, 478–485, 480t, 481t–482t, 485b–486b. *See also* Heart sound(s)
 in newborns, 864t
 carotid artery, 478, 478t, 516t
 chest, 425–430, 425t–426t
 direct, 71
 Erb's point, 482t
 indirect, 71, 72b
 LLSB (tricuspid), 481t
 lungs, 425–430, 425t–426t
 in newborns, 863t
 murmurs, 480t, 484, 483t–484t, 482f, 485t. *See also* Murmurs
 neck, 311, 312t, 478, 479t, 516t
 precordium, 478–485, 482b–483b, 483t–485t
 scrotal, 677, 678t
 thyroid gland, 311, 312t
- Auscultatory gap, 82b
- Automaticity, 444
- Autonomic nervous system, 453, 746, 749f
- Avoidant personality disorder, 140t
- Axilla
 inspection of, 545t
 palpation of, 546, 547t
- Axillary vein, 493f, 493t
- Axon, 738, 739f
- Azygos vein, 493f, 493t

- B**
- Babinski reflex, 784, 786t
in infant, 869t–870t
- Baby blues, 116, 116b
- Baby bottle caries, 895t
- Bachmann's bundle, 444f
- Back pain, 688
in pregnancy, 693
- Bacterial vaginosis, 644t
- Bacteriuria, screening for, 156t
- Balance, 695, 705, 705b. *See also* Gait
- Ballance's sign, 597, 598t–599t
- Ballottement, 69, 70f
abdominal, 596–597, 598t
patellar, 729, 729f
- Barbiturates
abuse of, 135t
integumentary system effects of, 216t
sleep and, 159b
- Barlow-Ortolani maneuver, 855, 866t
- Baroreceptors, 453
- Barthel Index, 945t
- Bartholin's gland, 612f, 614t
- Basal cell carcinoma, 211, 212–213, 258t, 348t, 957t
- Basilic vein, 493f, 493t
- BATHE technique, 131
- Beau's lines, 251t
- Bedwetting, 158
- Behavior(s)
health, 154–155
nonverbal, 7–9
problem
in adolescents, 115, 115b, 913–914, 919
in school-age children, 902
- Behavior Assessment System for Children, 115t
- Behavior Pain Score, 106
- Behavioral theory, 114
- Bell's palsy, 290t
- Benign prostatic hypertrophy, 653, 680t
- Benzodiazepines
abuse of, 135t
musculoskeletal system effects of, 697t
- Beta-blockers
female genitourinary system effects of, 621t
respiratory system effects of, 402t
sleep and, 159b
- Biases, personal, 19
- Biceps brachii, 687f
- Biceps femoris, 687f
- Bidigital examination, in male genitourinary system assessment, 677, 677f
- Biographical data, 28t, 31, 32t
in abdominal assessment, 562–563
in adolescent assessment, 915
in breast assessment, 533
in female genitourinary system assessment, 619
in integumentary system assessment, 212
in lymphatic system assessment, 497
in male genitourinary system assessment, 657–658
in mental health assessment, 120, 120t–121t
in neurologic system assessment, 752
in nutrition assessment, 170
in older adult assessment, 934
in pain assessment, 97
in peripheral-vascular system assessment, 497
in school-age child assessment, 904
in teaching assessment, 147
in toddler and preschooler assessment, 890
- Biological clock, 158
- Biotinidase deficiency, 874t
- Biot's respiration, 412, 415b
- Bipolar disorders, 141t
- Bisphosphonates, gastrointestinal system effects of, 569t
- Bladder, 556f, 558f, 559t, 612f, 615t, 651f, 653t
age-related changes in, 930t
continence and, 614
palpation of, 595, 596t
percussion of, 590t–591t
size of, 588
- Bleeding. *See also* Hemorrhage
vaginal, 619
- Blepharitis, 347t
- Blepharoconjunctivitis, 347t
- Blindness, 329f, 330
- Blink reflex, 329, 350t
- Blood pressure, 80t, 81
abnormal, 516t
in adolescents, 494
arm for, 83b
in cardiovascular system assessment, 465
in infants, 494
Korotkoff's sounds with, 83b, 453b
measurement of, 82b–83b, 83b, 517b
in newborns, 858t–859t
in older adults, 453–454
in pregnancy, 495
in school-age children, 908, 908f
sphygmomanometer for, 58t
in toddlers and preschoolers, 893, 893f
- Blood type incompatibility, 874t
- Body language, 8
culture and, 10
- Body mass index, 180, 181t
- Body systems interaction
in abdominal assessment, 558–559, 560
in breast assessment, 528, 530, 531
in cardiovascular system assessment, 452, 453
in ear assessment, 366–369
in eye assessment, 326–328
in facial assessment, 269–270, 273, 274
in female genitourinary system assessment, 614–615, 616
in gastrointestinal system assessment, 558–559, 560
in head assessment, 269–270, 273, 274
in integumentary system assessment, 208–209, 210
in lymphatic system assessment, 494, 496
in male genitourinary system assessment, 651, 653, 655
in musculoskeletal system assessment, 686, 688, 692
in neck assessment, 269–270, 273, 274
in neurologic system assessment, 746, 748, 750
in peripheral-vascular system assessment, 494
in respiratory system assessment, 396–397
- Body temperature, 78–80
axillary, 79t
in cardiovascular system assessment, 466
in children, 79, 80t
circadian rhythms of, 80
environmental factors and, 80
exercise and, 80
forehead, 79t
gender and, 80
measurement of, 79–80, 79f, 79t
in newborns, 79, 80t, 851t, 855, 858t
in older adults, 79, 80t
oral, 79t
rectal, 79t
regulation of, 78–79
in respiratory system assessment, 412
in school-age children, 908, 908f
temporal artery, 79t
in toddlers and preschoolers, 893, 893f
tympanic, 79f, 79t
- Body type, in physical assessment, 74
- Bone(s), 686, 687f, 690t. *See also* Musculoskeletal system
cancellous, 688f
compact, 686, 688f
cross-section of, 688f
spongy, 686
- Bone marrow, 686
- Borborygmi, 586t
- Borderline personality disorder, 140t
- Bouchard's nodes, 727t
- Boutonnière deformity, 727t
- Bowel movements, 565–566, 565b
- Bowel sounds, 586t
in newborns, 864t
in pregnant women, 561
- Bowlegs, 691, 708t
- Brachial pulse, 492f, 493t, 511t
- Brachial vein, 493f, 493t
- Brachioradialis, 687f
- Bradycardia, 80
- Bradypnea, 81, 412, 415b
- Brain, 739, 740f, 740t–741t. *See also* Neurologic system
age-related changes in, 931t
- Brain tumor, 790t
- Brainstem, 741t
- Brazilian Americans, older adult, 932t
- Breakthrough bleeding, 619
- Breast(s), 527–553
abnormalities of, 551t–552t
abscess of, 552t
in adolescents, 532, 532b
anatomy of, 528, 528f, 529f
assessment of, 38t, 533–548
in adolescents, 917t, 922t
anatomical landmarks in, 539, 540f
approach to, 539
biographical data in, 533
in breast reconstruction, 540
cardiovascular system in, 537t, 541t, 542
case study of, 533, 538–539, 548
checklist for, 552–553
concentric circle palpation in, 544, 544f
cultural considerations in, 532–533
current health status in, 533–534

- Breast(s) (*continued*)
- developmental considerations in, 530, 532, 532b
 - documentation for, 548, 548f
 - drugs in, 535t–536t
 - endocrine system in, 537t, 542
 - family history in, 536, 536t
 - focused history in, 533b
 - gastrointestinal system in, 537t, 541t, 542
 - general survey in, 536t, 540, 540t, 542
 - in genitourinary disorders, 623t, 630t
 - genitourinary system in, 541t, 542
 - head-to-toe, 540, 540t–541t
 - health history in, 533–538
 - HEENT system in, 541t, 542
 - hematologic system in, 542
 - inspection in, 541, 543f, 544t–545t
 - integumentary system in, 536t–537t, 540t, 542
 - lump in, 534
 - lymphatic system in, 542
 - in menopause, 804t
 - musculoskeletal system in, 541t, 542
 - neurological system in, 541t
 - nipple discharge in, 534
 - nursing diagnoses in, 549
 - pain in, 534
 - palpation in, 541, 543–544, 543f, 544f, 546, 546t–548t
 - past health history in, 534, 534t–535t
 - physical, 539–548
 - pie wedge palpation in, 543, 543f
 - positions for, 539, 544t–545t
 - in pregnancy, 828t, 835t–836t
 - psychosocial profile in, 537, 537t–538t
 - reproductive system in, 537t, 541t, 542
 - respiratory system in, 537t, 541t, 542
 - review of systems in, 536, 536t–537t
 - symptom analysis in, 534
 - vertical strip palpation in, 543, 543f
 - body systems interaction with, 528, 530, 531
 - cancer of, 546t, 551t
 - diet and, 549
 - incidence of, 532–533
 - risk factors for, 551
 - screening for, 156t
 - in children, 532, 532b
 - development of, 532, 532b
 - drug effects on, 535t–536t
 - fibroadenoma of, 551t
 - fibrocystic disease of, 551t
 - in infants, 530, 532
 - interductal papilloma of, 551t
 - lump in, 534, 546t
 - lymph nodes of, 529f, 530t
 - male, 528, 667t
 - mammary duct ectasia of, 552t
 - mammography of, 549, 959t
 - mastitis of, 552t
 - neuroendocrine system interaction with, 530
 - in older adults, 532
 - pain in, 534
 - in pregnancy, 532, 816t
 - quadrants of, 548f
 - reconstruction of, 540
 - reproductive system interaction with, 530
 - self-examination of, 533, 549, 550
- Breath sounds, 425–429
- abnormal, 426, 427
 - absent, 425t
 - adventitious, 425t–426t, 427–429, 428t–429t
 - bronchial, 428t
 - bronchovesicular, 428t
 - misplaced, 426
 - normal, 425t–426t, 426, 427f, 428t
 - vesicular, 428t
- Breathing. *See also* Respiration; Respiratory system
- pursed-lip, 413t
- Brief Pain Inventory, 99
- Broad ligament, 613f
- Broca's area, 740f
- Bronchi, 394f, 395t
- Bronchioles, 394, 394f, 395t
- Bronchitis, 432t
- Bronchodilators, cardiovascular system effects of, 457t
- Bronchophony, 426t, 429–430
- Bronzing, 228t
- Brudzinski's sign, 784, 784f
- Bruits
- abdominal, 516t, 586t–587t
 - carotid, 478, 516t
- Brushfield spots, 329, 351t
- Buerger's disease, 521t
- Bulbourethral gland, 650, 651f, 652t
- assessment of, 676–677, 677f
- Bulge test, 729, 729f
- Bulimia nervosa, 142t, 184t, 608t
- in adolescent, 913
- Bulla, 235t, 244t
- Bull's-eye lesion, 241t
- Bundle of His, 444f, 445
- Bunions, 727t, 955t
- Bursae, 686, 690t
- Bursitis, elbow, 726t
- C**
- C-fibers, 89
- C-reactive protein, 487
- Cachexia, cardiac, 465
- Café-au-lait spots, 860t
- Caffeine, sleep and, 158
- CAGE questionnaire, 134b
- Calcaneus, 687f
- Calcium
- nutrition and, 172t
 - during pregnancy, 169t
- Calculative skills, in neurologic system assessment, 767t, 771
- Calf
- circumference of, 514t
 - pain in, 514t
- Callus, 728t
- Cancer
- breast, 156t, 532–533, 546t, 549, 551, 551t
 - colorectal, 603–604, 605t
 - endometrial, 644, 645t
 - lip, 297t
 - lung, 431–432, 433t
 - oral, 296b, 300t, 303t, 314–315
 - pain in, 91t, 92
 - prostate gland, 679, 682t
 - stomach, 605t
 - tongue, 303t
- Candidiasis, 237t
- genital, 645t, 669t
 - oral, 300t, 301t, 316t, 862t
- Canthus
- lateral, 322f
 - medial, 322f, 324t
- Capillary, 441b, 441f
- Capillary refill test, 252t, 508t
- Caput medusa, 583t
- Carbohydrates, 164
- Cardiac agents
- breast effects of, 535t
 - eye effects of, 333t
 - female genitourinary system effects of, 621t
 - gastrointestinal system effects of, 569t
 - head, face, and neck effects of, 279t
- Cardiac cycle, 442
- Cardiac output, 80, 443
- Cardiac veins, 442b
- Cardinal fields of gaze test, 346t
- Cardiovascular system, 437–490. *See also* Peripheral-vascular system
- age-related changes in, 929t
 - anatomy of, 438–442, 438f, 439t
 - assessment of, 37t, 454–489
 - in adolescents, 917t, 922t
 - anatomical landmarks in, 463, 463f, 464f
 - approach to, 465–466
 - auscultation in, 463b, 478–485, 481t–482t, 485b
 - biographical data in, 454
 - in breast disorders, 537t, 541t, 542
 - carotid artery in, 463, 464f, 485f
 - case study of, 454, 462–463, 470, 485
 - checklist for, 488–489
 - cultural considerations in, 454
 - current health status in, 455–456
 - developmental considerations in, 453–454
 - drugs in, 457t–458t
 - in ear disorders, 377t, 386
 - endocrine system in, 460t, 467
 - in eye disorders, 334t, 339t, 340
 - family history in, 456, 458t
 - in female genitourinary disorders, 623t, 630t, 635
 - focused history in, 454b
 - in gastrointestinal disorders, 572t, 580t, 582
 - gastrointestinal system in, 459t, 467, 469t
 - general survey in, 458t, 465–466, 467, 468t
 - genitourinary system in, 459t, 467
 - in head, face, and neck disorders, 281t, 288t, 313
 - head-to-toe, 466, 468t–470t
 - health history in, 456–462
 - HEENT system in, 459t, 467, 469t
 - hematologic system in, 460t, 467
 - in infants, 877t, 881t
 - inspection in, 470, 470b, 470t–471t, 472f
 - in integumentary disorders, 218t, 224t, 225

- integumentary system in, 459t, 467, 468t–469t
- jugular veins in, 463, 463f
- in lymphatic disorders, 501t, 506t, 507
- lymphatic system in, 460t, 467
- in male genitourinary disorders, 661t, 666
- for menopause, 803t, 804t
- in mental health disorders, 123t, 127t
- in musculoskeletal disorders, 698t, 703t, 704
- musculoskeletal system in, 460t, 467, 470t
- in neurologic disorders, 757t, 764t, 765
- neurologic system in, 460t, 467, 470t
- in newborns, 855
- nursing diagnoses in, 486
- in nutritional disorders, 173t, 179t
- in older adults, 938t, 952t
- in pain disorders, 104t
- palpation in, 472, 473t–475t
- past health history in, 456, 456t–457t
- patient positions for, 466, 466f
- in patient teaching, 148t, 150t
- percussion in, 476t, 478
- in peripheral-vascular disorders, 501t, 506t
- physical, 464–488
- in postpartal women, 843t, 844t
- in pregnancy, 828t, 835t
- psychosocial profile in, 460, 460t–462t
- reproductive system in, 459t–460t
- in respiratory disorders, 404t, 414t, 419
- respiratory system in, 459t, 467, 469t
- review of systems in, 458t–461t
- in school-age children, 906t, 910t
- symptom analysis in, 455–456
- in toddlers and preschoolers, 891t, 895t
- vital signs in, 465–466
- body systems interaction with, 452, 453
- in children, 453
- drug effects on, 457t–458t
- ear interaction with, 368
- endocrine system interaction with, 452, 453
- eye interaction with, 327, 328
- female genitourinary system interaction with, 614, 616
- gastrointestinal system interaction with, 559, 560
- head, face, and neck interaction with, 270, 273
- in infants, 453
- integumentary system interaction with, 209, 210
- lymphatic system interaction with, 494
- male genitourinary system interaction with, 655
- male reproductive system interaction with, 651, 655
- musculoskeletal system interaction with, 692
- neurologic system interaction with, 452, 453, 748, 750
- in newborns, 851t
- in older adults, 453–454
- during pregnancy, 815t–816t
- in pregnant women, 453
- respiratory system interaction with, 396, 397, 452, 453
- Caries, 298t
- baby bottle, 895t
- Carotemia, 229t
- Carotid artery, 463, 464f
- auscultation of, 478, 478t, 516t
- inspection of, 470, 470b, 470t
- palpation of, 472, 473t
- pressure in, 472
- Carotid pulse, 492f, 493t, 510t
- Carpal tunnel syndrome, 733t
- Cartilage, 686, 690t
- Cartilaginous joint, 688f, 690t
- Caruncle, 322f
- Cataract, 330, 351t, 361t
- Categorical scales, for pain, 99
- Central nervous system, 739–745. *See also* Neurologic system
- Central venous pressure, 470
- Cephalic vein, 493f, 493t
- Cephalosporins, integumentary system effects of, 216t
- Cerebellar function, in musculoskeletal system
- assessment, 703, 705, 709t–711t
- Cerebellum, 740f, 740t, 741t
- Cerebral cortex, 740f
- Cerebral function, in neurologic system
- assessment, 764, 766t–768t
- Cerebrum, 453, 740f
- Cerumen, 370, 378t, 379t
- Cervical adenitis, 307t, 315t
- Cervical cancer, 643, 644
- screening for, 156t
- Cervical lymph nodes
- anatomy of, 270f
- in cancer, 308t
- in children, 275, 308
- deep, 270f, 309t
- enlargement of, 307t, 315t
- functions of, 272t
- in infection, 308t
- infraclavicular, 310t, 547t
- normal, 308t
- occipital, 270f, 309t
- palpation of, 306–308, 308t–310t, 511t, 515
- postauricular, 270f, 309t
- posterior, 270f, 309t
- preauricular, 270f, 309t
- submandibular, 270f, 309t
- submental, 270f, 309t
- superficial, 270f, 309t
- supraclavicular, 270f, 310t, 547t
- tonsillar, 270f, 309t
- Cervical smear, 65t, 634, 637, 638t, 959t
- Cervical spine, range of motion of, 718t–719t
- Cervical vertebrae, 687f
- Cervix, 612f, 613f, 615t
- bimanual examination of, 640t
- cancer of, 156t, 643, 644
- during pregnancy, 817t
- in pregnancy, 817t, 840t
- speculum examination of, 634, 636t–637t
- Chadwick's sign, 617, 636t, 817t, 840t
- Chalazion, 348t
- Chancre, 297t, 669t
- Chancroid, 632t, 669t, 680t
- Chandleier's sign, 640t
- Charcot-Marie-Tooth disease, hearing in, 372t
- Cheilitis, 238t, 297t
- Cheilosis, 297t
- Chemical digestion, 558
- Chemoreceptors, 453
- Chemotherapeutic agents. *See* Antineoplastics
- Chest. *See also* Respiratory system
- anatomical landmarks of, 409, 409f, 410f, 411f, 412f
- anterior, 416t–417t
- auscultation of, 425–430, 425t–426t
- barrel, 416f, 417t
- circumference of, in newborns, 856t
- excursion of, 418, 421t
- funnel, 416f, 863t
- in infants, 881t
- inspection of, 416, 416f, 416t–418t
- lateral, 417t
- pain in, 400, 420t, 455, 487t
- palpation of, 418, 420t–422t, 474–475
- percussion of, 422f, 423, 423t–424t
- pigeon, 416f, 863t
- posterior, 417t–418t
- size of, 398
- tenderness of, 420t
- Cheyne-Stokes respiration, 412, 415b
- Chief complaint, 31
- Child abuse, 115, 889, 889b, 1033–1035
- Child-bearing practices, 819, 819t–823t. *See also* Pregnancy
- Child Behavior Checklist, 115t
- Children. *See* Adolescents; Infants; Newborns; Preschoolers; School-age children; Toddlers
- Children's Depression Inventory, 115t
- Children's Hospital of Eastern Ontario Pain Scale, 106
- Chinese Americans
- adolescent, 915t
- chest size in, 398
- childbearing in, 820t
- facial skin of, 275
- integumentary system of, 211
- mental disorders and, 118t
- musculoskeletal health problems in, 694t
- newborn, 853t
- nutrition and, 171t
- older adult, 932t
- pain response in, 96t
- school-age, 903t
- spirituality and, 191t
- views of women and, 618t
- Chlamydial infection, 644t, 680t
- screening for, 156t
- Chloasma, 211, 228t, 275, 814t, 834t
- Chlorpropamide, nutrition and, 172t
- Choking game, 115, 115b
- Cholecystitis, 605t
- Cholesteatoma, 381t, 388t
- Cholesterol, 165
- breast cancer and, 549
- cutaneous, 207t
- in older adults, 959t
- Cholestyramine resin, nutrition and, 172t
- Cholinergic agents, respiratory system effects of, 402t

- Cholinergic blockers, neurologic system
effects of, 755t
- Chordee, 670t
- Choroid, 323f, 325t
- Christian Scientist, 196t
- Christianity, 195t–196t
- Chronic arterial insufficiency, 521t–522t
- Chronic obstructive pulmonary
disease, 400b
- Chronic pain syndrome, 91t, 92
- Chronic venous insufficiency, 522t
- Cigarette smoking, 137t, 461t
screening for, 157t
sleep and, 158
- Cilia, nasal, 271t
- Ciliary body, 323f, 325t, 328
- Cimetidine, breast effects of, 536t
- Circadian rhythms, 158
- Circulation, 440, 442
bronchial, 396, 398
cardiac, 440, 442
pulmonary, 396, 398, 440, 440f
systemic, 440, 440f
- Circumcision
female, 632t
male, 657
- Circumduction, 715f, 716t
- Circumlocution, 766b
- Cirrhosis, 605t
- Cisplatin, nutrition and, 172t
- Clarification, 12b
- Clarifying, in patient interview, 10
- Clavicle, 687f
- Cleft lip/palate, 862t
- Clichés, 19
- Clicking, joint, 716
- Clitoris, 612f, 614t, 631t
hypertrophy of, 632t
- Clock scoring, 957t
- Clonus, 783, 785t
- Clubbing, 208, 208f, 414b
- Cocaine
abuse of, 135t, 461t
palatal lesions and, 301t
- Coccidioidomycosis, 407b
- Cochlea, 365t, 366f
- Cochlear nerve, 366f
- Code for Nurses, 13b–14b
- Codeine, abuse of, 136t
- Coffee, 63t
- Cognitive function, in neurologic system
assessment, 766t–768t, 770
- Cognitive inhibition, 783
- Cognitive skills, for assessment, 11–12, 12b
- Cognitive theory, 114
- Cold caloric test, 775t
- Colestipol, nutrition and, 172t
- Collaboration, 16, 16b
- Collaborative problem, 5, 5b
- Collateral ligament, 729–730
- Colon, 556f
- Color change test, 518b
- Color Tool, for pain, 100
- Colorectal cancer, 603–604, 605t
screening for, 156t
- Colostomy, 602t
- Colostrum, 532
- Communication, 5, 7–11
culture and, 9–10
interpreter for, 9b
for interview, 10–11, 11b, 18–19
nonverbal, 5, 7–9
- Communication evaluation, in neurologic
system assessment, 768t, 771
- Community, assessment of, 1026, 1026t–1029t
- Concentric circles method, of breast
palpation, 544, 544f
- Conchae, 269f, 271t
- Conductivity, cardiac, 444
- Condyloma acuminatum, 669t, 680t
- Cones, 326
- Congenital adrenal hyperplasia, 874t
- Congestive heart failure, 400b, 486t
- Conjunctiva, 323f, 324t
inspection of, 349t–350t
- Conjunctivitis, 361t
allergic, 349t
- Connective tissue, 206f, 207t
- Constipation, 565, 565b, 605t–606t
in older adults, 938t
- Contact lenses, 347t
- Contraceptives, 624t
- Contractility, cardiac, 444–445
- Cooper's ligament, 529f, 530t
- Coordination, 695, 705, 710t–711t.
See also Gait
- COPD, 400b
- Cor pulmonale, 400b
- Cornea, 323f, 324t
abrasion of, 351t
cloudy, 351t
inspection of, 350t–351t
ulcer of, 351t
- Corneal light reflex, 329–330, 345t
- Corneal reflex, 350t, 773t–774t
- Corniculate cartilage, 395t
- Corns, 728t
- Coronal sutures, 268f, 271t
- Coronary arteries, 442b
- Coronary artery disease, 488t
C-reactive protein in, 487
epidemiology of, 487
risk factors for, 455b
screening for, 156t
- Corpus cavernosum, 651f
- Corpus spongiosum, 651f
- Corrigan's (water-hammer) pulse, 477f
- Corti, organ of, 365t, 367
- Corticobulbar tract, 743–744, 744t
- Corticospinal tract, 743–744, 743f, 744t
- Corticosteroids
female genitourinary system effects of,
621t
integumentary system effects of, 215t
musculoskeletal system effects of,
697t
neurologic system effects of, 756t
nutrition and, 172t
- Costovertebral angle, 71f, 588, 591t
- Cotton balls, 63t
- Cough, 400, 456
- Cover/uncover test, 345t
- Crack cocaine, abuse of, 135t
- Crackles, 425t, 427, 428t, 429
- Cranial nerve(s), 745, 746t, 747f, 771,
772t–777t. *See also* Neurologic
system
in older adults, 955t
- Cranial nerve palsy, 361t
- Cranium, 687f
- Crawling reflex, 872t
- Creativity, 12b
- Crepitus, 420t, 716
- Cricoid cartilage, 270f, 395t
- CRILES, for infant pain assessment, 102, 105t
- Crisis, 131, 132t
- Critical thinking, for assessment, 11, 11b–12b,
12b
- Crohn's disease, 606t–607t
- Crossed extension reflex, 872t
- Crust, 236t
- Cruveilhier-Baumgarten murmur, 587t
- Cryptorchidism, 673t, 680t
- Cuban Americans
adolescent, 915t
childbearing in, 820t–821t
mental disorders and, 119t
newborn, 853t
nutrition and, 171t
older adult, 932t
pain response in, 96t
school-age, 903t
spirituality and, 191t
views of women and, 618t
- Cubital vein, 493f, 493t
- Cullen's sign, 583t
- Culture, 78
abdomen assessment and, 561, 561t–562t
adolescent assessment and, 914, 915t
assessment of, 1019–1022
breast assessment and, 532–533
cardiovascular assessment and, 454
communication and, 9–10
eye assessment and, 330
female genitourinary system assessment
and, 617, 618t
head, face, neck assessment and, 275
integumentary system assessment and, 211
male genitourinary system assessment and,
657
mental health assessment and, 118t–119t
musculoskeletal assessment and, 693, 694t
neurologic system assessment and, 751
newborn assessment and, 850, 852, 853t
nutrition assessment and, 170, 171t
older adult assessment and, 932, 932t
pain assessment and, 95, 96t–97t
peripheral-vascular assessment and, 495,
497
pregnancy assessment and, 819, 819t–823t
respiratory system assessment and, 398–399
school-age child assessment and, 903,
903t–904t
spirituality assessment and, 190, 191t–192t
teaching effectiveness and, 147
views of women and, 618t
- Cuneiform cartilage, 395t
- Current health status, 28t, 31. *See also* Drugs
in abdomen assessment, 563
in adolescent assessment, 916
in breast assessment, 533–534

- in cardiovascular system assessment, 455–456
 - in ear assessment, 370
 - in eye assessment, 331
 - in face assessment, 275–276
 - in female genitourinary system assessment, 619
 - in head assessment, 275–276
 - in integumentary system assessment, 212–214
 - in lymphatic system assessment, 497–498
 - in male genitourinary system assessment, 658
 - in menopause assessment, 797
 - in musculoskeletal system assessment, 694
 - in neck assessment, 275–276
 - in neurologic system assessment, 752
 - in nutrition assessment, 170–171
 - in older adult assessment, 934–935
 - in pain assessment, 97–98
 - in peripheral-vascular system assessment, 497–498, 499t–500t
 - in pregnancy assessment, 824–825
 - in preschooler assessment, 890
 - in respiratory system assessment, 399
 - in school-age child assessment, 904
 - in spirituality assessment, 192
 - in teaching assessment, 147
 - in toddler assessment, 890
 - Cushing's syndrome, 290t
 - Cushing's triad, 762
 - Cutis marmorata, 850t
 - Cyanosis, 208, 226t, 413t
 - central, 208, 226t, 229t
 - clubbing in, 208, 208f
 - peripheral, 208, 226t, 229t
 - Cyclophosphamide, nutrition and, 172t
 - Cyclothymic disorder, 141t
 - Cyst, 234t
 - keratogenous, 234t
 - sebaceous, 670t
 - Cystocele, 632t
 - Cystourethrocele, 632t
 - Cytology brush, 65t
- D**
- Dacryocystitis, 348t
 - Dancing reflex, 870t
 - DAR documentation method, 23–24
 - Darwinian tubercle, 378t
 - Data, 17–24
 - collection of, 17–20, 22b
 - documentation of, 23–24
 - objective, 17
 - organization of, 21
 - primary, 17
 - prioritization of, 21–22
 - secondary, 17
 - validation of, 20
 - Decibels, 366
 - Decision making, 16b
 - Deep tendon reflexes, 744, 745t, 782–783, 783f
 - in pregnancy assessment, 841t
 - Deep vein thrombosis, 514t–515t, 522t
 - Defense mechanisms, 114
 - Delirium, 116, 117t
 - Deltoid, 687f
 - Dementia, 117t
 - pain assessment in, 106t, 107
 - Dendrites, 738, 739f
 - Dependent personality disorder, 140t
 - Depression, 138, 141t
 - assessment of, 132
 - in older adults, 117, 117b, 117t, 118b
 - postpartum, 116, 116b, 845b
 - screening for, 156t
 - Depression (body movement), 715f, 716t
 - Depression Attitude Questionnaire, 138
 - Dermatitis
 - atopic, 209, 258t
 - contact, 235t, 258t
 - stasis, 262t
 - Dermatomes, 745, 748f
 - Dermis, 206, 206f, 207t
 - Desert fever, 407b
 - Detumescence, 650
 - Development, 28t, 39, 39b–40b, 40t, 78
 - in abdomen assessment, 559, 561
 - in adolescent assessment, 914
 - in breast assessment, 530, 532, 532b
 - in cardiovascular system assessment, 453–454
 - in ear assessment, 369–370
 - in eye assessment, 328–330
 - in face assessment, 274–275
 - in female genitourinary system assessment, 615, 617, 617b
 - in head assessment, 274–275
 - in infant assessment, 850–852, 876, 878t
 - in lymphatic system assessment, 494–495
 - in menopause assessment, 796
 - in mental disorder assessment, 114–115
 - in musculoskeletal system assessment, 688–689, 691, 693
 - in neck assessment, 274–275
 - in neurologic system assessment, 749, 751
 - in newborn assessment, 850–852
 - in peripheral-vascular system assessment, 494–495
 - in preschooler assessment, 888
 - in respiratory system assessment, 398
 - in school-age child assessment, 902–903
 - in spirituality assessment, 188–190
 - teaching effectiveness and, 146–147
 - theories of, 39b–40b, 40t, 41t–42t, 114
 - in toddler assessment, 888
 - in wellness assessment, 155, 157–158
 - Dexamethasone, nutrition and, 172t
 - Diabetes mellitus, 184t
 - eye examination in, 360
 - in pregnancy, 826
 - retinopathy in, 358t
 - screening for, 156t
 - Diaphragm (musculoskeletal), 394, 396, 398f
 - excursion of, 423, 424t
 - Diaphysis, 688f
 - Diarrhea, 566
 - Diarthrosis, 690t
 - Diastasis recti, 583t
 - Diastole, ventricular, 442
 - Diencephalon, 741t
 - Diet. *See also* Nutrition
 - breast cancer and, 549
 - sleep and, 158
 - vegetarian, 164b
 - Digestion, 558
 - in older adults, 561
 - Digestive system, 556, 556f, 556t–557t. *See also* Gastrointestinal system
 - Digital rectal examination, 677, 677f, 679
 - Diplopia, 331
 - Disabled patient, 73–74
 - Discharge
 - nipple, 534
 - penile, 658
 - vaginal, 619, 632t, 633t
 - Distension, abdominal, 584t–585t
 - Distress signs, in physical assessment, 74
 - Diuretics
 - ear effects of, 372t
 - male sexual function effects of, 660t
 - musculoskeletal system effects of, 697t
 - nutrition and, 172t
 - sleep and, 159b
 - Divergent thinking, 12b
 - Diverticulitis, 606t
 - Dizziness, 456
 - in neurologic system assessment, 753
 - Documentation, 23–24
 - in breast assessment, 548, 548f
 - DAR, 23–24
 - of health history, 47, 47b–51b
 - in menopause assessment, 806t–807t
 - narrative, 24
 - in neurologic assessment, 769, 783f
 - of physical assessment, 85
 - PIE, 24
 - in pregnancy assessment, 824–825
 - problem-oriented, 23
 - SOAPIE, 23
 - source-oriented, 23
 - tips for, 24b
 - of vital signs, 83
 - Doll's eyes, 772t–773t
 - Dopamine, 112b
 - Doppler ultrasound, 57t
 - Dorsalis pedis pulse, 492f, 493t, 513t
 - Double vision, 331
 - Dowager's hump, 693, 707t
 - Down syndrome, 749, 862t
 - Drain, abdominal, 599, 601t–602t
 - Drainage
 - ear, 371, 378t
 - eye, 332
 - Drawer test, 729
 - Dress, 8
 - in physical assessment, 77–78
 - Drugs. *See also specific drug classes*
 - breast effects of, 535t–536t
 - cardiovascular system effects of, 457t–458t
 - ear effects of, 372t
 - female genitourinary system effects of, 621t–622t
 - gastrointestinal system effects of, 568t–569t
 - integumentary system effects of, 215t–216t
 - lymphatic system effects of, 499t–500t
 - mental health effects of, 122b, 123t

- Drugs (*continued*)
 musculoskeletal system effects of, 696t–697t
 neurologic system effects of, 755t–756t
 nutritional effects of, 172t, 943t
 in older adult assessment, 935
 sleep and, 159, 159b
- Ductus arteriosus, closure of, 453
- Ductus deferens, 651f
- Dullness, on percussion, 70t
- Duodenum, 557t, 558f
- Dysarthria, 766b
- Dysmenorrhea, 619
- Dyspepsia, 566
- Dysphasia, 766b
- Dysphonia, 766b
- Dyspnea, 81, 400, 456
- Dysthymia, 132
- Dysuria, 619, 658
- E**
- Ear(s), 365–389. *See also* HEENT system
 abnormalities of, 388t–389t
 age-related changes in, 929t
 anatomic landmarks of, 376, 376f
 anatomy of, 365, 365t, 366f
 assessment of, 37t, 370–385
 in adolescents, 917t, 921t
 biographical data in, 370
 cardiovascular system in, 377t
 case study of, 370, 374–375, 385
 checklist for, 389–390
 in children, 369
 current health status in, 370
 developmental considerations in, 369–370
 drugs in, 372t
 family history in, 371, 372t
 focused history in, 370b
 gastrointestinal system in, 373t
 general survey in, 373t, 376–377
 genitourinary system in, 373t
 in head, face, and neck disorders, 281t, 288t, 313
 head-to-toe, 377, 377t
 health history in, 370–374
 hearing tests in, 382–383, 383t–385t, 384t–385t
 HEENT system in, 373t, 377t, 386
 in infants, 369, 369f, 880t
 inspection in, 378t
 integumentary system in, 373t, 377t, 386
 lymphatic system in, 373t
 musculoskeletal system in, 373t, 377t, 386
 neurologic system in, 373t, 377t, 386
 in newborns, 861t
 nursing diagnosis in, 385
 in older adults, 369–370, 951t
 otoscopy in, 379t–381t
 palpation in, 379t
 past health history in, 371, 371t
 physical, 376–385
 psychosocial profile in, 373, 373t–374t
 reproductive system in, 373t
 respiratory system in, 373t, 377t, 386
 review of systems in, 373, 373t
 in school-age children, 905t, 909t
 in toddlers and preschoolers, 890t, 894t
 in young adults, 369
 attachment of, 376, 376f, 378t
 body systems interaction with, 366–369
 congenital anomalies of, 378t
 deviated, 376, 376f
 drainage from, 371, 378t
 drug effects on, 372t
 external, 365t, 366f
 foreign object in, 380t
 functions of, 365t
 inner, 365t, 366f
 low-set, 376, 376f, 378t
 middle, 365t, 366f
 musculoskeletal system interaction with, 366–367, 368
 neurologic system interaction with, 367–369
 pain in, 371
 surfer's, 380t
 swimmer's, 380t
- Ear wax, 370, 378t
- Earache, 371
- Eastern Orthodox Church, 195t
- Eating disorders, 141t–142t, 565b. *See also*
 Anorexia nervosa; Bulimia
 in adolescents, 913
 in school-age children, 902
- Ecchymosis, 209, 231t
 genital, 632t
- Eccrine glands, 206f, 207t, 208, 211
- Ecomapping, 36b
- Ectasia, mammary duct, 552t
- Ectasy, 136t
- Ectropion, 330, 348t
- Eczema, 258t
 nummular, 240t
- Edema, 209, 400, 456, 498. *See also*
 Lymphedema
 assessment of, 510b
 genital, 632t, 658
 in pregnancy, 839t
- Effusion, pleural, 433t
- Egophony, 426t, 430
- Egyptian Americans
 adolescent, 915t
 childbearing in, 821t
 mental disorders and, 119t
 musculoskeletal health problems in, 694t
 newborn, 853t
 nutrition and, 171t
 older adult, 932t
 pain response in, 96t
 school-age, 903t
 spirituality and, 191t
 touch and, 9
 views of women and, 618t
- Ejection click, 479t
- Elastic cartilage, 686
- Elbow
 golf, 726t
 range of motion of, 721t
 tennis, 726t
- Elderly patient. *See* Older adults
- Elevation (body movement), 715f, 716t
- Embolism, arterial, 521t
- Empathy, 11b
- Emphysema, 432t–433t
- End-diastolic pressure, 443
- End-diastolic volume, 442
- End-of-life care, 946
- End-systolic pressure, 443
- End-systolic volume, 443
- Endocarditis, temperature in, 466
- Endocardium, 438f, 439t
- Endocervical smear, 65t, 634, 637, 638t, 959t
- Endocrine system
 age-related changes in, 931t
 assessment of, 39t
 in adolescents, 918t
 in breast disorders, 537t, 542
 in cardiovascular disorders, 460t, 467
 in eye disorders, 334t–335t
 in female genitourinary disorders, 625t, 635
 in gastrointestinal disorders, 572t, 582
 in head, face, and neck disorders, 282t, 313
 in integumentary disorders, 219t, 225
 in lymphatic disorders, 501t
 in male genitourinary disorders, 662t, 666
 in musculoskeletal disorders, 698t, 704
 in neurologic disorders, 758t, 765
 in newborns, 855
 in older adults, 940t
 in patient teaching, 148t
 in peripheral-vascular disorders, 501t
 in pregnancy, 830t
 in respiratory disorders, 419
 breast interaction with, 530, 531
 cardiovascular system interaction with, 452, 453
 eye interaction with, 327
 female genitourinary system interaction with, 616
 gastrointestinal system interaction with, 559, 560
 head, face, and neck interaction with, 273, 274
 integumentary system interaction with, 209, 210
 male genitourinary system interaction with, 653, 655
 musculoskeletal system interaction with, 692
 neurologic system interaction with, 748, 750
 in pregnancy, 818t
 respiratory system interaction with, 397
- Endometrial cancer, 644, 645t
- Endometriosis, 645t
- Endosteum, 688f
- Entropion, 330, 347t
- Enuresis, nocturnal, 158
- Environment
 assessment of, 1022–1029
 for physical assessment, 72–73
- Epicardium, 438f, 439t
- Epidermis, 206, 206f, 207t
- Epididymis, 650, 651f, 652t
- Epididymitis, 680t
- Epigastric area, palpation of, 475t
- Epiglottis, 395t
- Epiphysis, 688f

- Episcleritis, 350t
 Episcopalian Church, 195t
 Epispadias, 671t, 681t
 Epistaxis, 276, 316t
 Epstein's pearls, 862t
 Epulis, 815t, 834t
 Equilibrium, disorders of, 369
 Equipment, for physical assessment, 55, 56t–66t
 Erb's palsy, 862t
 Erb's point, 464f
 auscultation of, 482t
 Erectile dysfunction, 657, 664
 Erosion, 237t
 Erythema, 229t, 242t
 oropharyngeal, 304t
 palmar, 495, 814t, 834t
 Erythema toxicum neonatorum, 211
 Erythroplakia, 300t
 Esophagus, 394f, 556t
 age-related changes in, 930t
 Estimated date of birth, 825
 Estrogens
 breast effects of, 530, 535t
 female genitourinary system effects of, 622t
 male sexual function effects of, 660t
 Ethical skills, for assessment, 13
 Ethmoid sinus, 268f
 Eupnea, 81, 412, 415b
 European Americans, older adult, 932t
 Eustachian tube, 269f, 365t, 366f
 Evaluation, 4f, 5
 Event sequencing, in patient interview, 10
 Eversion (body movement), 715f, 716t
 Excitability, 444
 Excitatory amino acids, 89
 Excoriation, 209, 237t
 Exercise
 for adolescents, 160, 919
 assessment of, 159–160
 for children, 892
 heart rate for, 160
 for older adults, 942
 for school-age children, 907
 sleep and, 158
 Exhalation, 396
 Exophthalmos, 348t
 Exostosis, in external auditory canal, 380t
 Exotropia, 345t
 Expiration, 81. *See also* Respiration
 Expiratory center, 396
 Extensor carpi radialis, 687f
 External acoustic meatus, 268f
 External auditory canal, 365t, 366f, 378t–379t
 Extraocular muscles, 328
 assessment of, 344, 345t–346t
 function of, 324t
 Extrusion reflex, in newborns, 871t
 Eye(s), 322–364. *See also* HEENT system
 abnormalities of, 360t–362t
 in African Americans, 330
 age-related changes in, 928t
 anatomy of, 322, 322f, 323f, 324f, 324t–325t
 appearance of, 332
 assessment of, 37t, 330–359
 in adolescents, 917t, 921t
 anatomical landmarks in, 338, 338f
 biographical data in, 331
 cardiovascular system in, 334t, 339t, 340
 case study of, 330, 336–337, 359–360
 checklist for, 363
 cultural considerations in, 330
 current health status in, 331
 developmental considerations in, 328–330
 drugs in, 333t
 endocrine system in, 334t–335t
 external structure examination in, 346, 347t–354t
 extraocular muscle testing in, 344, 345t–346t
 family history in, 333, 333t
 focused history in, 331b
 gastrointestinal system in, 334t, 339t, 340
 general survey in, 334t, 338–339
 genitourinary system in, 334t
 in head, face, and neck disorders, 281t, 288t, 313
 head-to-toe, 339, 339t
 health history in, 331–336
 HEENT system in, 334t, 339t, 340
 in infants, 880t
 integumentary system in, 334t, 339t, 340
 internal structures examination in, 355, 356t–359t, 555f
 lymphatic system in, 335t
 musculoskeletal system in, 334t, 339t, 340
 neurologic system in, 334t, 339t, 340
 in newborns, 861t
 nursing diagnoses in, 360
 in older adults, 936t–937t, 950t
 past health history in, 332, 332t
 physical, 341–359
 psychosocial profile in, 335, 335t–336t
 reproductive system in, 334t, 339t, 340
 respiratory system in, 334t, 339t, 340
 review of systems in, 334, 334t–335t
 in school-age children, 905t, 909t
 symptom analysis in, 331–332
 in toddlers and preschoolers, 890t, 894t
 visual acuity testing in, 341–342, 342t–343t
 body systems interaction with, 326–328
 cardiovascular system interaction with, 327, 328
 cross-section of, 323f
 drainage from, 332
 drug effects on, 333t
 dry, 331–332
 of infants, 328–329
 musculoskeletal system interaction with, 327, 328
 neurologic system interaction with, 327, 328, 329f
 pain in, 331
 Eye contact, 8
 culture and, 9
 Eyeball, 323f
 inspection of, 349t
 palpation of, 354t
 Eyelashes, 324t
 inspection of, 347t
 Eyelids, 322f, 324t
 inspection of, 348t
 movement of, 328
- F**
 Face. *See also* HEENT system
 anatomy of, 268, 268f, 269f, 270f, 271t
 appearance of, 290t
 assessment of, 275–314
 anatomical landmarks in, 286, 286f
 biographical data in, 275
 cardiovascular system in, 281t, 288t, 313
 case study of, 275, 284–285, 312, 314
 checklist for, 318–319
 cultural considerations in, 275
 current health status in, 275–276
 developmental considerations in, 274–275
 drugs in, 278t–279t, 279
 endocrine system in, 282t, 313
 in eye disorders, 334t
 family history in, 279, 279t–280t
 focused history in, 276b
 gastrointestinal system in, 281t, 288t, 313
 general survey in, 281t, 287, 287t, 313
 genitourinary system in, 282t, 288t, 313
 head-to-toe, 287, 287t–288t
 health history in, 275–284, 276b
 HEENT system in, 281t, 288t, 313
 hematologic system in, 282t, 313
 inspection in, 289, 289t–290t
 integumentary system in, 281t, 287t, 313
 lymphatic system in, 282t, 313
 musculoskeletal system in, 282t, 288t, 313
 neurologic system in, 282t, 288t, 313
 nursing diagnoses in, 314
 palpation in, 290, 291t
 past health history in, 276, 277t–278t
 physical, 286–318
 psychosocial profile in, 283, 283t–284t
 reproductive system in, 282t, 313
 respiratory system in, 288t, 313
 review of systems in, 280, 281t–282t
 symptom analysis in, 276
 body systems interaction with, 269–270, 273, 274
 cardiovascular system interaction with, 270, 273
 cultural variations in, 275
 drug effects on, 278t–279t, 279
 endocrine system interaction with, 273, 274
 hematologic system interaction with, 273, 274
 lymphatic system interaction with, 273, 274
 muscles of, 270, 274f
 musculoskeletal system interaction with, 270, 273
 neurologic system interaction with, 270, 273
 in physical assessment, 74
 in pregnant women, 275
 respiratory system interaction with, 269–270, 273
 skin color of, 275
 FACES Pain Rating Scale, 99, 100f
 Facial expression, 8
 culture and, 9
 Facial muscles, 687f
 Facial nerve (CN VII), 746t, 747f, 774t
 Facilitation, in patient interview, 10
 Fallopian tube, 612f, 613f, 615t

- Falls, in older adults, 941b
- Family history, 28t, 35, 35b, 35f, 43b
- in abdominal assessment, 569, 570t
 - in adolescent assessment, 916
 - in breast assessment, 536, 536t
 - in cardiovascular system assessment, 458, 458t
 - in ear assessment, 371, 372t
 - in eye assessment, 333, 333t
 - in facial assessment, 279, 279t–280t
 - in female genitourinary system assessment, 622, 622t
 - in head assessment, 279, 279t–280t
 - in integumentary system assessment, 216, 217t
 - in lymphatic system assessment, 500, 501t
 - in male genitourinary system assessment, 660, 660t
 - in menopause assessment, 798
 - in musculoskeletal system assessment, 695, 697t
 - in neck assessment, 279, 279t–280t
 - in neurologic system assessment, 756, 756t
 - in nutrition assessment, 172
 - in older adult assessment, 935
 - in pain assessment, 100
 - in peripheral-vascular system assessment, 500, 501t
 - in pregnancy assessment, 826–827
 - in respiratory system assessment, 400, 402t
 - in school-age child assessment, 904
 - in spirituality assessment, 193
 - in teaching assessment, 148
 - in toddler and preschooler assessment, 890
- Fasciculation, 766b
- Fatigue, 456, 498
- Fats, 164–165
- saturated, 165
- Fear
- vs. anxiety, 139t
 - in children, 889
- Fecal occult blood test, 959t
- Feet, flat, 728t
- Femoral hernia, 675t
- inspection for, 671t
 - palpation for, 674t–675t, 676, 677f
- Femoral pulse, 492f, 493t, 512t, 587t
- in newborns, 864t
- Femur, 687f
- Fertilization, 613
- Fetoscope, 57t
- Fetus. *See also* Pregnancy
- circulation of, 850, 852f
 - heart tones of, 837t
 - position of, 837t–838t
 - presentation of, 838t
- Fever, 79
- Fibroadenoma, of breast, 551t
- Fibrocystic disease, of breast, 551t
- Fibromyalgia, 732
- Fibrosis, pulmonary, 433t–434t
- Fibrous cartilage, 686
- Fibrous joint, 688f, 690t
- Fibula, 687f
- Fields of gaze test, 346t
- Fight or flight response, 559
- Filipino Americans
- adolescent, 915t
 - childbearing in, 821t
 - face of, 275
 - mental disorders and, 119t
 - musculoskeletal health problems in, 694t
 - newborn, 853t
 - nutrition and, 171t
 - older adult, 932t
 - pain response in, 96t
 - school-age, 903t
 - spirituality and, 191t
 - views of women and, 618t
- Finger(s)
- range of motion of, 722t–723t
 - rheumatoid arthritis of, 727t
 - strength of, 723t
 - swan-neck deformity of, 734t
- Finger-thumb opposition, 710t
- Finger-to-finger test, 710t
- Finger-to-nose test, 705, 710t
- Fingernails, 208, 208f
- age-related changes in, 928t
 - Beau's lines of, 251t
 - blue, 249t
 - brittle, 251t
 - capillary refill test on, 252t
 - in cardiovascular disorders, 459t, 469t
 - changes in, 214
 - clubbing of, 208, 208f, 249, 252
 - color of, 249t
 - concave, 250t
 - cyanotic, 413t
 - in gastrointestinal disorders, 580t
 - half-and-half, 249t
 - in head, face, and neck disorders, 287t
 - hemorrhages of, 250t
 - infection of, 250t
 - inspection of, 249, 249t–250t, 252
 - in older adults, 950t
 - palpation of, 251t, 252
 - pitting of, 250t, 251t
 - shape of, 250t
 - in thyroid disease, 209
 - white, 249t
 - yellow, 249t, 413t
- Fissure, 238t
- Fixative, 65t
- FLACC Postoperative Pain Tool, 106–107, 106t
- Flaccidity, 709t
- Flat feet, 728t
- Flatness, on percussion, 70t
- Flexion, 715f, 716t
- Fluent aphasia, 766b
- Fluid wave test, 595–596, 596f
- Fluoride, during pregnancy, 169t
- Fluorosis, 298t
- Focused history, 28, 28t, 29
- in abdomen assessment, 563b
 - in breast assessment, 533b
 - in cardiovascular system assessment, 454b
 - in ear assessment, 370b
 - in eye assessment, 331b
 - in face assessment, 276b
 - in female genitourinary system assessment, 618b
 - in head assessment, 276b
 - in integumentary system assessment, 212b
 - in lymphatic system assessment, 497b
 - in male genitourinary system assessment, 657b
 - in menopause assessment, 801b
 - in musculoskeletal system assessment, 694b
 - in neck assessment, 276b
 - in neurologic system assessment, 752b
 - in nutrition assessment, 175b
 - in peripheral-vascular system assessment, 497b
 - in pregnancy assessment, 824b
 - in respiratory system assessment, 399b
- Focusing, in patient interview, 10
- Folic acid, 169t, 751
- Fontanel, 274, 860t, 862t
- Food, digestion of, 558
- Food Guide Pyramid, 165–167, 166f
- Food intake records, in nutrition assessment, 176
- Foot drop, 709t
- Foramen ovale, closure of, 453
- Fordyce granules, 300t
- Foreign object, in ear, 380t
- Fornices, bimanual examination of, 641t
- Fovea centralis, 324f, 325t, 355f
- assessment of, 359t
- Framework, for data organization, 21, 21f, 21t
- Fremitus, 418, 421t–422t
- French Americans, school-age, 904t
- French Canadians
- childbearing in, 821t
 - mental disorders and, 119t
 - newborn, 853t
 - pain response in, 96t
 - spirituality and, 191t
- Frenulum
- labial, 271t
 - lingual, 271t, 296
- Frequency, of sound, 365–366
- Friction rub, 451f, 587t
- pericardial, 449–450, 479t
 - pleural, 426t, 429, 429t
- Frontal bone, 268f
- Frontal cortex, 89
- Frontal lobe, 740f, 740t
- Frontal sinus, 268f, 269f
- inspection of, 292t
 - palpation of, 293t
 - percussion of, 293t
 - transillumination, 292t
- Frontalis, 274f
- Fundoscopic examination, 355, 355b, 355f, 356t–359t
- Fungal infection, of nails, 249t
- G**
- Gait, 695, 702–703, 705, 705b, 705f, 708t–709t, 711t–713t
- in older adults, 693, 954t
 - in physical assessment, 74
 - in pregnancy, 693
 - propulsive, 711t
 - scissors, 709t, 712t
 - spastic, 712t
 - steppage, 713t
 - waddling, 712t

- Galactosemia, 874t
- Gallbladder, 556f, 558t
inflammation of, 605t
- Gallop rhythms, 448–449
- Gamma-aminobutyric acid, 112b
- Gastrocnemius, 687f
- Gastroenteritis, 579t
- Gastrointestinal agents, neurologic system effects of, 756t
- Gastrointestinal system. *See also* Abdomen
abnormalities of, 605t–608t
age-related changes in, 929t–930t
assessment of, 38t
in adolescents, 918t, 922t
in breast disorders, 537t, 541t, 542
in cardiovascular disorders, 459t, 467, 469t
in ear disorders, 373t
in eye disorders, 334t, 339t, 340
in female genitourinary disorders, 623t–624t, 635
in head, face, and neck disorders, 281t, 288t, 313
in infants, 877t, 881t–882t
in integumentary disorders, 218t, 224t, 225
in lymphatic disorders, 501t, 506t, 507
in male genitourinary disorders, 661t, 666
in mental health disorders, 124t
in musculoskeletal disorders, 698t, 703t, 704
in neurologic disorders, 757t–758t, 764t, 765
in newborns, 855
in nutritional disorders, 173t, 179t
in older adults, 938t, 953t
in pain disorders, 104t
in patient teaching, 148t
in peripheral-vascular disorders, 501t, 506t
in pregnancy, 829t, 836t–839t
in respiratory disorders, 404t, 414t, 419
in school-age children, 906t, 911t
in toddlers and preschoolers, 891t, 896t
- body systems interaction with, 558–559, 560
- cardiovascular system interaction with, 452, 559, 560
- drug effects on, 568t–569t
- endocrine system interaction with, 559, 560
- eye interaction with, 327
- female genitourinary system interaction with, 616
- integumentary system interaction with, 209, 210, 558, 560
- male genitourinary system interaction with, 655
- musculoskeletal system interaction with, 559, 560, 692
- neurologic system interaction with, 559, 560, 750
- in newborns, 851t
during pregnancy, 816t
respiratory system interaction with, 397, 559, 560
- Gastrostomy tube, 602t
- Gate-control theory, of pain, 90, 90f
- General knowledge, in neurologic system assessment, 767t, 771
- General survey, 74, 77–78
- Genetics, 1038–1041, 1038b
- Genitalia. *See also* Genitourinary system;
Reproductive system
female, 612–613, 612f, 614t–615t
bimanual examination of, 637, 640, 640t–641t
in children and adolescents, 615, 617b
in infants, 615
inspection of, 631t–632t, 634, 636t–637t, 637
in newborns, 865t
palpation of, 630, 633t, 634, 637, 640, 640t–641t
speculum examination of, 634, 636t–637t, 638t–639t
male, 650, 651f, 652t–653t
in adolescents, 654, 656t
bidigital examination of, 677, 677f
in children, 653–654
in infants, 653
inspection of, 667, 668t–671t
in newborns, 865t–866t
in older adults, 654, 657
palpation of, 672, 672t–675t, 676, 677f
- Genitourinary system. *See also* Reproductive system
assessment of, 38t
in adolescents, 918t, 922t–923t
in cardiovascular disorders, 459t, 467
in ear disorders, 373t
in eye disorders, 334t, 340
in gastrointestinal disorders, 572t, 582
in head, face, and neck disorders, 282t, 288t, 313
in infants, 877t
in integumentary disorders, 218t, 225
in lymphatic disorders, 501t, 506t, 507
in mental health disorders, 124t
in musculoskeletal disorders, 698t, 703t, 704
in neurologic disorders, 764t, 765
in newborns, 855
in older adults, 938t–939t
in pain disorders, 104t
in patient teaching, 148t
in peripheral-vascular disorders, 501t, 506t
in respiratory disorders, 404t, 419
in school-age children, 906t, 911t
in toddlers and preschoolers, 891t, 896t
- female, 611–648
abnormalities of, 644t–646t
in adolescents, 615, 617b
anatomy of, 612–613, 612f, 613f, 614t–615t
assessment of, 617–642
abdomen in, 630t
amenorrhea in, 619
approach to, 628
biographical data in, 619
bleeding in, 619
in breast disorders, 541t, 542
breast in, 623t, 630t
cardiovascular system in, 623t, 630t, 635
case study of, 617, 627–628, 643
checklist for, 646–647
cultural considerations in, 617, 618t
current health status in, 619
developmental considerations in, 615, 617, 617b
drugs in, 621t–622t
endocrine system in, 625t, 635
external genitalia in, 630, 631t–632t, 633t, 634
family history in, 622, 622t
focused history in, 618b
gastrointestinal system in, 623t, 635
general survey in, 622t–623t, 629, 629t
genitourinary system in, 635
head-to-toe, 629, 629t–630t
health history in, 617–627
HEENT system in, 623t, 629t, 635
hematologic system in, 625t, 630t, 635
in infants, 882t
inspection in, 630, 631t–632t, 634, 636t–637t, 637
integumentary system in, 623t, 629t, 635
internal genitalia in, 634, 636t–637t, 637, 638t–639t, 640, 640t–641t
lesions in, 619
lymphatic system in, 625t, 630t, 635
in menopause, 805t
musculoskeletal system in, 624t, 630t, 635
neurologic system in, 624t, 630t, 635
nursing diagnoses in, 643
pain in, 619
palpation in, 630, 633t, 634, 637, 640, 640t–641t
past health history in, 619, 620t–621t
physical, 628–642
position for, 629
in postpartal women, 844t
in pregnancy, 829t, 840t
psychosocial profile in, 625, 625t–626t
rectovaginal examination in, 642, 642t
reproductive system in, 624t
respiratory system in, 623t, 630t, 635
review of systems in, 622, 622t–625t, 635
sexual history in, 627
specimen collection in, 637, 638t–639t
symptom analysis in, 619
urinary symptoms in, 619
vaginal discharge in, 619, 632t, 633t
vaginal wall inspection in, 637, 639t
- body systems interaction with, 614–615, 616
- cardiovascular system interaction with, 452, 614, 616
- in children, 615
drug effects on, 621t–622t
ear interaction with, 368
endocrine system interaction with, 616
eye interaction with, 327
functions of, 613
gastrointestinal system interaction with, 560, 616
in infants, 615

Genitourinary system (*continued*)

- integumentary system interaction with, 209, 210, 616
 - lymphatic system interaction with, 616
 - musculoskeletal system interaction with, 614, 616, 692
 - neurologic system interaction with, 750
 - neurological system interaction with, 615, 616
 - in older adults, 617
 - in pregnant women, 615, 617, 817t
 - respiratory system interaction with, 397, 616
 - male, 649–684
 - abnormalities of, 680t–683t
 - in adolescents, 654, 656t
 - anatomy of, 650, 651f, 652t–653t
 - assessment of, 657–678
 - auscultation in, 677, 678t
 - bidigital examination in, 677, 677f
 - biographical data in, 657–658
 - bulbourethral gland examination in, 676–677, 677f
 - cardiovascular system in, 661t, 666
 - case study of, 657, 664–665, 678
 - checklist in, 683
 - cultural considerations in, 657
 - current health status in, 658
 - drugs in, 659t–660t
 - endocrine system in, 662t, 666
 - erectile dysfunction in, 664
 - family history in, 660, 660t
 - focused history in, 657b
 - gastrointestinal system in, 661t, 666, 667t
 - general survey in, 661t, 665, 667t
 - head-to-toe, 667, 667t–668t
 - health history in, 657–664, 657b
 - HEENT system in, 661t, 666, 667t
 - hematologic system in, 662t, 666
 - immunologic system in, 662t
 - in infants, 882t
 - inspection in, 667, 668t–671t
 - integumentary system in, 661t, 666, 667t
 - lesions in, 658
 - lymphatic system in, 666
 - musculoskeletal system in, 661t, 666, 668t
 - neurologic system in, 662t, 666, 668t
 - nursing diagnoses in, 678
 - pain in, 658
 - palpation in, 672, 672t–675t, 676, 677f
 - past health history in, 658–659, 659t
 - penile discharge in, 658
 - physical, 665–678
 - prostate gland examination in, 675t, 676–677, 677f
 - psychosocial profile in, 662, 662t–663t
 - rectal examination in, 675t, 676–677, 677f
 - respiratory system in, 661t, 666
 - review of systems in, 660, 661t–662t
 - sexual history in, 663–664
 - sexually transmitted diseases in, 658b
 - swelling in, 658
 - symptom analysis in, 658
 - urethral specimen in, 676b
 - urinary problems in, 658
 - body systems interaction with, 651, 653, 655
 - cardiovascular system interaction with, 452, 651, 655
 - in children, 653–654
 - drug effects on, 659t–660t
 - ear interaction with, 368
 - endocrine system interaction with, 653, 655
 - eye interaction with, 327
 - gastrointestinal system interaction with, 560, 655
 - in infants, 653
 - integumentary system interaction with, 209, 210, 655
 - lymphatic system interaction with, 655
 - musculoskeletal system interaction with, 653, 655, 692
 - neurologic system interaction with, 750
 - neurological system interaction with, 653, 655
 - in older adults, 654, 657
 - respiratory system interaction with, 397, 655
 - self-examination teaching about, 676b
 - Genogram, 35f
 - Genu valgum, 691, 708t
 - Genu varum, 691, 708t
 - Genuineness, for patient interview, 11b
 - German Americans, integumentary system of, 211
 - Gestational age, 874t
 - Gestures, 8
 - Gingivae (gums), 269f, 271t
 - hyperplasia of, 299t
 - hypertrophy of, 834t
 - inspection of, 299t–300t
 - in pregnant women, 275
 - recession of, 299t
 - Gingivitis, 299t
 - Glabellar reflex, 872t
 - Glans penis, 651f, 653t
 - Glasgow Coma Scale, 769, 770t
 - Glaucoma, 353t, 361t
 - Globe, 323f
 - Glossitis, 303t
 - Glossopalatine arch, 269f
 - Glossopharyngeal nerve (CN IX), 746t, 747f, 775t–776t
 - Gloves, 64t
 - Glucocorticoids, eye effects of, 334t
 - Gluteus maximus, 687f
 - Goiter, 274
 - Gold compounds
 - head, face, and neck effects of, 279t
 - integumentary system effects of, 216t
 - lymphatic system effects of, 500t
 - respiratory system effects of, 402t
 - Golf elbow, 726t
 - Goniometer, 61t
 - Gonococcal smear, 637, 638t
 - Gonorrhea, 681t
 - Goodell's sign, 640t, 817t
 - Gordon's functional health patterns, 21, 21t
 - Gout, 732
 - Gouty arthritis, 727t
 - Graafian follicle, 613
 - Graphesthesia, 777, 781t
 - Grasp reflex, 783, 869t
 - Greek Americans
 - adolescent, 915t
 - childbearing in, 821t–822t
 - gastrointestinal symptoms in, 562t
 - mental disorders and, 119t
 - newborn, 853t
 - older adult, 932t
 - pain response in, 96t
 - school-age, 904t
 - spirituality and, 191t
 - views of women and, 618t
 - Grey-Turner's sign, 583t
 - Grooming, 8
 - Group B streptococcal pharyngitis, 317t
 - Growth charts, 179, 1008–1017
 - Grunting, 429
 - Guillain-Barré syndrome, 790t–791t
- ## H
- Hair, 206, 206f, 208, 208t
 - age-related changes in, 928t
 - body, 254t
 - in cardiovascular disorders, 468t
 - color of, 254t
 - growth of, 214
 - in head, face, and neck disorders, 287t
 - inspection of, 252, 253t–254t, 254
 - loss of, 253t
 - in older adults, 950t
 - palpation of, 254, 255t
 - pubic
 - female, 617b
 - male, 654, 656t, 668t
 - scalp, 253t
 - in thyroid disease, 209
 - Hair follicle, 206f
 - Halitosis, 297t
 - Hallucinations, 141t
 - Hallucinogens, abuse of, 136t
 - Hallus valgus, 955t
 - Hallux valgus, 727t
 - Hammer, percussion, 71f
 - Hammer toe, 727t
 - Hand(s)
 - palpation of, 722t–723t
 - range of motion of, 722t–723t
 - strength of, 723t
 - Hand(s), for palpation, 68f
 - Hand gestures, 8
 - Handshake, 9
 - Hard palate, 269f, 272t, 301t
 - Harlequin sign, 850t, 859t
 - Hay fever, 316t
 - Head. *See also* HEENT system
 - anatomy of, 268, 268f, 269f, 270f
 - assessment of, 36t, 275–314
 - anatomical landmarks in, 286, 286f
 - biographical data in, 275
 - cardiovascular system in, 281t, 288t, 313
 - case study of, 275, 284–285, 312, 314
 - checklist for, 318–319
 - cultural considerations in, 275
 - current health status in, 275–276

- developmental considerations in, 274–275
- drugs in, 278t–279t, 279
- endocrine system in, 282t, 313
- in eye disorders, 334t, 339t
- family history in, 279, 279t–280t
- focused history in, 276b
- gastrointestinal system in, 281t, 288t, 313
- general survey in, 281t, 287, 287t, 313
- genitourinary system in, 282t, 288t, 313
- head-to-toe, 287, 287t–288t
- health history in, 275–284, 276b
- HEENT system in, 281t, 288t, 313
- hematologic system in, 282t, 313
- inspection in, 289, 289t–290t
- integumentary system in, 281t, 287t, 313
- lymphatic system in, 282t, 313
- musculoskeletal system in, 282t, 288t, 313
- neurologic system in, 282t, 288t, 313
- in newborns, 860t
- nursing diagnoses in, 314
- palpation in, 290, 291t
- past health history in, 276, 277t–278t
- physical, 286–318
- psychosocial profile in, 283t–284t
- reproductive system in, 282t, 313
- respiratory system in, 288t, 313
- review of systems in, 280, 281t–282t
- symptom analysis in, 276
- bobbing of, 465
- body systems interaction with, 269–270, 273, 274
- cardiovascular system interaction with, 270, 273
- circumference of, 274
 - in newborns, 856t
- contour of, 291t
- drug effects on, 278t–279t, 279
- endocrine system interaction with, 273, 274
- functions of, 271t–272t
- hematologic system interaction with, 273, 274
- in infants, 274
- lymphatic system interaction with, 273, 274
- musculoskeletal system interaction with, 270, 273
- neurologic system interaction with, 270, 273
- in newborns, 850t, 856t
- pain in, 276
- respiratory system interaction with, 269–270, 273
- shape of, 290t
- size of, 289t
- Head-to-toe physical assessment, 84–85
 - in abdominal assessment, 579, 579t–581t
 - in breast assessment, 540, 540t–541t
 - in cardiovascular system assessment, 466, 468t–470t
 - in ear assessment, 377, 377t
 - in eye assessment, 339, 339t
 - in female genitourinary system assessment, 629, 629t–630t
 - in head, face, and neck assessment, 287, 287t–288t
 - in infant assessment, 879, 879t–883t
 - in integumentary system assessment, 223, 223t–224t
 - in lymphatic system assessment, 505, 505t–506t
 - in male genitourinary system assessment, 667, 667t–668t
 - in menopause assessment of, 801, 802t–805t
 - in mental health assessment, 127t
 - in musculoskeletal system assessment, 702, 703t
 - in neurologic system assessment, 762, 763t–764t
 - in newborn assessment, 855, 856t–867t
 - in nutrition assessment, 178t–179t
 - in older adult assessment, 948, 949t–957t
 - in pain assessment, 104t
 - in peripheral-vascular system assessment, 505, 505t–506t
 - in pregnancy assessment, 833, 834t–841t
 - in respiratory system assessment, 412, 413t–414t
 - in school-age child assessment, 908, 908t–911t
- Headache, 752
 - migraine, 791t–792t
- Health behaviors, 154–155
- Health care access, 154–155
- Health history, 27–52
 - in abdominal assessment, 562–574
 - in adolescent assessment, 915–916
 - in breast assessment, 533–538
 - in cardiovascular system assessment, 456–462
 - communication for, 30, 30b
 - complete, 28–29, 28t
 - components of, 28t, 30–40
 - biographical data, 28t, 31, 32t
 - current health status, 28t, 31
 - developmental considerations, 28t, 39, 39b–40b, 40t
 - family history, 28t, 35, 35b, 35f, 43b
 - past health history, 28t, 33, 33t–34t
 - psychosocial profile, 28t, 29, 44, 44t–47t
 - reason for seeking health care, 28t, 31
 - review of systems, 28t, 36, 36t–39t
 - documentation of, 47, 47b–51b
 - in ear assessment, 370–374
 - environment for, 30, 30b
 - in eye assessment, 331–336
 - in face assessment, 275–284
 - family, 28t, 35, 35b, 35f, 43b
 - in female genitourinary system assessment, 617–627, 618b
 - focused, 28, 28t, 29. *See also* Focused history
 - forms for, 29
 - in head assessment, 275–284
 - in infant assessment, 876
 - in integumentary system assessment, 212–220
 - in lymphatic system assessment, 497–503
 - in male genitourinary system assessment, 657–664, 657b
 - medical, 30
 - in menopause assessment, 797–799
 - in mental health assessment, 114, 120, 120t–121t
 - in musculoskeletal system assessment, 693–700
 - in neck assessment, 275–284
 - in neurologic system assessment, 751–760
 - in newborn assessment, 854, 854t
 - nursing, 30
 - in nutrition assessment, 170–172
 - in older adult assessment, 933–934
 - in pain assessment, 97–98
 - past, 28–29, 28t, 33, 33t–34t. *See also* Past health history
 - patient's condition and, 29
 - in peripheral-vascular system assessment, 497–503
 - in pregnancy assessment, 823–831
 - purpose of, 28
 - in school-age child assessment, 904
 - in spirituality assessment, 192–193
 - in teaching assessment, 147–149
 - time for, 29
 - in toddler and preschooler assessment, 889–890
- Health Insurance Portability and Accountability Act, 14b
- Healthy People 2010, mental health objectives of, 138
- Hearing, 365–366, 367f
 - assessment of, 382–383, 383t–385t, 384t–385t, 387
 - case study of, 370
 - in children, 369
 - in infants, 369, 369f, 388
 - in middle-aged adults, 369
 - in older adults, 269–270
 - in young adults, 369
 - loss of, 371, 382–383, 387, 388
 - musculoskeletal system and, 366–367
 - neurological system and, 367, 369
- Hearing Handicap Inventory for the Elderly–Screening, 387
- Heart. *See also* Heart sound(s); Murmurs
 - anatomy of, 438f, 439, 439t–440t
 - electrical conduction system of, 444–445, 444f
- Heart failure, 400b, 488t
- Heart rate
 - abnormal, 473t
 - in athlete, 80
 - for exercise, 160
 - heart sounds and, 483
- Heart sound(s)
 - auscultation for, 478–485, 479t, 481t–482t, 485b–486b
 - in children, 453
 - duration of, 479
 - early ejection click, 447
 - first (S₁), 443f, 445–446, 445f, 479t, 483–484f
 - fourth (S₄), 448–449, 448f, 479t, 484f
 - frequency of, 478
 - grading of, 479
 - intensity of, 479
 - location of, 478
 - mechanisms of, 442, 443f, 445–449
 - midsystolic ejection click, 447
 - murmurs, 449–450, 449f, 450f

- Heart sound(s) (*continued*)
 in newborns, 864t
 opening snap, 447–448, 448f, 479t, 484f
 quality of, 479
 second (S₂), 443f, 446, 447f, 479t, 484f
 third (S₃), 448, 448f, 479t, 484f, 488
 timing of, 479
- Heberden's nodes, 727t, 955t
- Heel-and-toe walking, 709t
- Heel walking, 709t
- HEENT system, 268. *See also* Ear(s); Eye(s);
 Head; Nose; Throat
- assessment of
 in adolescents, 917t, 921t
 in breast disorders, 541t, 542
 in cardiovascular disorders, 459t, 467,
 469t
 in ear disorders, 373t, 377t, 386
 in eye disorders, 339t, 340
 in female genitourinary disorders, 623t,
 629t, 635
 in gastrointestinal disorders, 571t, 580t,
 582
 in head, face, and neck disorders, 281t,
 288t, 313
 in infants, 877t, 880t
 in integumentary disorders, 217t,
 223t–224t, 225
 in lymphatic disorders, 501t, 505t, 507
 in male genitourinary disorders, 661t,
 666, 667t
 in menopause evaluation, 802t–803t
 in mental health disorders, 123t, 127t
 in musculoskeletal disorders, 698t, 703t,
 704
 in neurologic disorders, 757t, 763t–764t,
 765
 in newborns, 855
 in nutritional disorders, 173t, 178t–179t
 in older adults, 936t–937t, 950t–951t
 in pain disorders, 104t
 in patient teaching, 148t, 150t
 in peripheral-vascular disorders, 501t,
 505t
 in postpartal women, 843t
 in pregnancy, 827t–828t, 834t–835t
 in respiratory disorders, 404t, 413t–414t,
 419
 in school-age children, 905t, 909t–910t
 in toddlers and preschoolers, 890t–891t,
 894t–895t
 during pregnancy, 815t
- Hegar's sign, 817t, 840t
- Height, 83–84, 84f
 in cardiovascular system assessment, 466
 in infants, 879t
 measurement of, 180, 180f
 in older adults, 693
- Helix, 366f
 Darwinian tubercle of, 378t
- Hemangioma
 capillary, 211, 232t
 in newborns, 859t, 860t
- Hemangioma simplex, 211
- Hematologic system
 age-related changes in, 931t
 assessment of, 39t
 in adolescents, 918t
 in breast disorders, 542
 in cardiovascular disorders, 460t, 467
 in female genitourinary disorders, 625t,
 630t, 635
 in gastrointestinal disorders, 573t, 582
 in head, face, and neck disorders, 282t,
 313
 in integumentary disorders, 218t, 225
 in male genitourinary disorders, 662t, 666
 in musculoskeletal disorders, 698t, 704
 in neurologic disorders, 758t, 765
 in older adults, 940t
 in patient teaching, 148t
 in pregnancy, 830t
 in respiratory disorders, 404t, 419
 in school-age children, 906t
 in toddlers and preschoolers, 892t
 eye interaction with, 327
 head, face, and neck interaction with, 273,
 274
 musculoskeletal system interaction with,
 692
 in pregnancy, 818t
- Hematoma, 209
 subdural, 790t
- Hemianopsia, 329f
- Hemocult test, 66t
- Hemochromatosis, 228t
- Hemorrhage
 nail, 250t
 splinter, 250t
 subconjunctival, 350t
- Hemotympanum, 381t, 388t
- Hepatitis, 606t
- Hepatitis B, screening for, 156t
- Hepatomegaly, 590t
- Hernia, 583t, 675t, 681t
 hiatal, 606t
 incisional, 606t
 inguinal, 606t
 inspection for, 671t
 palpation for, 674t–675t, 676, 677f
 umbilical, 583t–584t, 606t
- Heroin, abuse of, 136t
- Herpangioma, 304t
- Herpes simplex virus infection
 cutaneous, 234t, 236t, 240t, 259t
 genital, 246t, 632t, 645t, 669t, 681t
 of lips, 297t, 317t
- Herpes zoster, 241t, 245t, 259t
- Hesselbach's triangle, 676
- Heterochromia iridis, 351t
- Hinduism, 198t
- Hip, 729, 729f
 Barlow-Ortolani maneuver for,
 855, 866t
 range of motion of, 723t–724t
 strength of, 723t–724t
 tests for, 729, 729f
- Hippus phenomenon, 773t
- Hirsutism, 254t
- Hispanic Americans
 cardiovascular disease in, 454
 otitis media in, 370
- Histoplasmosis, 407b
- Histrionic personality disorder, 140t
- HIV infection, 644
 in newborns, 874t
 palatal candidiasis in, 301t
 periodontitis in, 299t
- Hive, 234t
- Hoarseness, 276
- Homans' sign, 414t, 514t–515t, 839t
- Home
 assessment of, 1023, 1023t–1026t
 in infant assessment, 876
- Homeless persons
 assessment of, 1030–1033
 school-age, 912
- Homeothermia, 79
- Hopelessness, 133t
- Hopping, 709t
- Hordeolum, 348t
- Hormones. *See also* Androgens; Estrogens
 cardiovascular system effects of, 457t
- Horner's syndrome, 352t
- Horns, keratotic, 211
- Human papilloma virus infection, 632t, 680t
- Humerus, 687f
- Humor, in patient interview, 10
- Hunter's syndrome, hearing in, 372t
- Hyaline cartilage, 686
- Hyaloid canal, 323f
- Hydrocele, 670t, 681t
- Hydrocephalus, 289t
- Hydromorphone, abuse of, 136t
- Hygiene, in physical assessment, 77–78
- Hymen, 612f, 614t
- Hyoid bone, 270f
- Hyperbilirubinemia, in infants, 884–885
- Hyperlipidemia, 469t
- Hyperopia, 326, 326f
- Hyperpigmentation, adrenal hypofunction
 and, 209
- Hyperresonance, on percussion, 70t
- Hypersensitivity, cutaneous, 600t
- Hypertension, 495, 497, 516t, 520
 optic disc in, 361t
 in pregnancy, 495, 826
 screening for, 156t
 stage 1, 81
 stage 2, 81
- Hyperthermia, 79
- Hyperthyroidism, 311, 312b, 312t
- Hyperventilation, 415b
- Hyphema, 354t
- Hypnotics
 abuse of, 135t
 neurologic system effects of, 756t
 respiratory system effects of, 403t
- Hypodermis, 206f, 207t
- Hypoglossal nerve (CN XII), 746t, 747f,
 776t–777t
- Hypopyon, 354t
- Hypospadias, 671t, 681t
- Hypotension, 81, 516t, 579
 orthostatic (postural), 83, 453, 516t
 during pregnancy, 453, 816t
- Hypothalamus, 453, 741t
- Hypothermia, 79
- Hypothyroidism, 311, 312b, 312t
 integumentary system in, 209
- Hypoxia, 400

I

- Ichthyosis, 228t, 956t
 Ileostomy, 602t
 Ileum, 557t
 Ileus, paralytic, 607t
 Iliac artery, 558f
 auscultation of, 587t
 Iliac crest, 687f
 Iliac vein, 493f, 493t, 558f
 Iliopsoas muscle test, 601t
 Immunizations
 for adolescents, 924–925
 for infants, 876
 for older adults, 959t
 for school-age children, 912
 for toddlers and preschoolers, 898
 Immunologic system
 age-related changes in, 931t
 assessment of, 39t
 in gastrointestinal disorders, 573t
 in integumentary disorders, 218t, 225
 in male genitourinary disorders, 662t
 in musculoskeletal disorders, 698t
 in newborns, 855
 in older adults, 940t
 in pain disorders, 104t
 in patient teaching, 148t
 in pregnancy, 830t
 integumentary system interaction with,
 209, 210
 in pregnancy, 494–495, 818t
 Implementation (intervention), 4f, 5
 Incontinence, in older adults, 938t–939t, 941t
 Incus, 365t, 366, 366f, 367f
 Indigestion, 566
 Infants, 849–886. *See also* Newborns
 abdomen of, 559
 assessment of, 876–884
 anorectal system in, 882t
 anthropometric measurements in, 879t
 cardiovascular system in, 877t, 881t
 case study of, 876, 878, 883–884
 developmental considerations in,
 850–852, 876, 878t
 gastrointestinal system in, 877t,
 881t–882t
 general survey in, 877t
 genitourinary system in, 877t, 882t
 head-to-toe, 879, 879t–88t
 health history in, 876
 HEENT system in, 877t, 880t–881t
 height in, 879t
 home safety in, 876
 immunizations in, 876
 integumentary system in, 877t, 879t
 musculoskeletal system in, 877t
 neurological system in, 878t, 883t
 physical, 878–883
 relational considerations in, 878t
 respiratory system in, 877t, 881t
 review of systems in, 876, 877t–878t
 screening tests in, 883
 weight in, 879t
 behavioral pain assessment scales for,
 102–107, 105t
 blood pressure in, 494
 breasts in, 530, 532
 cardiovascular system of, 453
 development of, 850–852
 exercise assessment in, 159
 eye of, 328–329, 360
 genitalia of, 615, 653
 hair of, 209, 211
 head of, 274
 health problems in, 875, 884–885
 hearing tests in, 388
 hyperbilirubinemia in, 884–885
 injury prevention for, 161
 lymphatic system in, 494
 musculoskeletal system in, 688–689, 691,
 693
 neck of, 275
 neurologic system of, 749
 nutrition in, 167, 168t
 otitis media in, 370
 pain assessment in, 93–94, 102–107
 paranasal sinuses of, 274
 peripheral-vascular system in, 494
 respiratory system in, 398
 skin of, 209
 sleep assessment in, 159
 spirituality and, 188–189
 stress assessment in, 160
 tonsils of, 274
 visual acuity assessment in, 329
 wellness assessment in, 155
 Inflammatory bowel disease, 606t–607t
 Informing, in patient interview, 10
 Inguinal canal, 653t
 Inguinal hernia, 675t, 681t
 inspection for, 671t
 palpation for, 674t–675t, 676, 677f
 Inhalants, abuse of, 137t
 Inhalation, 396
 Inhibitory amino acids, 89
 Initial Pain Assessment Inventory, 99
 Injury
 in adolescent, 913
 in children, 889
 prevention assessment for, 161–162
 in school-age children, 902
 Inotropes, 444
 Inspection, 20, 67, 67f
 abdominal, 581, 583t–586t
 breast, 541, 543f, 544t–545t
 cardiovascular, 470, 470b, 470t–471t, 472f
 carotid artery, 470b, 471t
 chest, 416, 416f, 416t–418t
 ear, 378t
 eye, 346, 347t–354t
 face, 289, 289t–290t
 female genitourinary system, 630,
 631t–632t, 636t–637t, 637
 femoral hernia, 671t
 gingiva, 299t–300t
 hair, 252, 253t–254t, 254
 head, face, and neck, 289, 289t–290t
 integumentary system, 226, 226t–229t, 230,
 231t–246t, 252, 253t–254t, 254
 jugular vein, 471t
 lingual frenulum, 296
 lip, 297t
 lower extremities, 509t
 male genitourinary system, 667, 668t–671t
 nails, 249, 249t–250t, 252
 neck, 307t, 470, 471t–472t
 nose, 294, 294t–295t
 oral mucosa, 299t–300t
 oropharynx, 303t–304t
 palate, 301t
 paranasal sinus, 291, 292t
 periphera-vascular, 508t, 510b
 precordium, 471–472, 471t–472t, 474, 475
 scalp, 254t
 teeth, 298t
 tongue, 302t–303t
 tonsils, 304t, 305t
 umbilicus, 583t–584t
 upper extremities, 508t
 uvula, 304t
 vaginal wall, 637, 639t
 Wharton's duct, 296
 Inspiration, 81. *See also* Respiration
 Inspiratory center, 396
 Instrumental Activities of Daily Living,
 945b–946b
 Insula, 741t
 Integumentary system, 205–265. *See also* Skin
 abnormalities of, 257t–263t
 in adolescents, 211, 917t, 921t
 anatomy of, 206–208, 206f, 207t, 208f
 assessment of, 36t, 212–263
 in adolescents, 917t, 921t
 age-related disorders in, 213t
 approach to, 222
 biographical data in, 212
 in breast disorders, 536t–537t,
 540t, 542
 in cardiovascular disorders, 459t, 467,
 468t–469t
 cardiovascular system in, 224t, 225
 case study of, 211, 221–222, 255–256
 checklist for, 263–264
 chronic ulceration in, 213
 color in, 226, 226t–227t, 228t–229t
 current health status in, 212–214
 in ear disorders, 373t, 377t, 386
 in eye disorders, 339t, 340
 family history in, 216, 217t
 in female genitourinary disorders, 623t,
 629t–630t, 635
 focused history in, 212b
 in gastrointestinal disorders, 571t,
 579t–580t, 582
 general survey in, 217t, 222–223, 223t
 hair changes in, 214
 in head, face, and neck disorders, 281t,
 287t, 313
 head-to-toe, 223, 223t–224t
 health history in, 212–220
 HEENT system in, 223t–224t, 225
 in infants, 877t, 879t
 inspection in, 226, 226t–246t, 230,
 249t–250t, 252, 253t–254t, 254
 lesions in, 227t–228t, 230, 230f, 231t
 distribution of, 243t–246t
 patterns of, 240t–242t
 primary, 230, 233t–235t
 secondary, 235t–238t
 vascular, 231t–232t
 in lymphatic disorders, 501t, 505t, 507

- Integumentary system (*continued*)
- in male genitourinary disorders, 661t, 666, 667t
 - in menopause assessment, 802t
 - in mental health disorders, 123t, 127t
 - mole in, 212
 - in musculoskeletal disorders, 697t–698t, 703t, 704
 - musculoskeletal system in, 224t, 225
 - nail changes in, 214
 - in neurologic disorders, 757t, 763t–764t, 765
 - neurologic system in, 224t, 225
 - in newborns, 855, 859t–860t
 - nonhealing sore in, 213
 - nursing diagnoses in, 256
 - in nutritional disorders, 173t, 178t
 - odor in, 227t
 - in older adults, 936t, 949t–950t
 - in pain disorders, 104t
 - palpation in, 240, 243t–246t, 247, 247t–248t, 249, 251t, 252
 - past health history in, 214, 214t–215t
 - in patient teaching, 148t, 150t
 - in peripheral-vascular disorders, 501t, 505t
 - physical, 222–256
 - in pregnancy, 827t, 834t
 - pressure ulcers in, 238b, 239t
 - pruritus in, 213
 - psychosocial profile in, 219, 219t–220t
 - rash in, 213
 - reproductive system in, 224t, 225
 - in respiratory disorders, 403t, 413t, 419
 - respiratory system in, 224t, 225
 - review of systems in, 216, 217t–218t
 - in school-age children, 905t, 908t–909t
 - season-related disorders in, 213, 213t
 - symptom analysis in, 212–214
 - tangential lighting in, 230
 - in toddlers and preschoolers, 890t, 894t
 - vascular lesions in, 231t–232t
 - body systems interaction with, 208–209, 210
 - cardiovascular system interaction with, 209, 210, 452, 467
 - drug effects on, 215t–216t
 - ear interaction with, 368
 - endocrine system interaction with, 209, 210
 - eye interaction with, 327
 - gastrointestinal system interaction with, 209, 210, 558, 560
 - genitourinary system interaction with, 209, 210, 616, 655
 - immune system interaction with, 209, 210
 - in infants, 209, 211
 - lymphatic system interaction with, 209, 210
 - in menopausal women, 211
 - musculoskeletal system interaction with, 692
 - neurologic system interaction with, 209, 210, 750
 - in newborns, 850t
 - of newborns, 850t
 - in older adults, 211
 - in pregnancy, 211, 814t–815t
 - reproductive system interaction with, 210
 - respiratory system interaction with, 208, 208f, 210, 397
- Intention tremor, 766b
- Intercostal muscles, 398f
- Interpersonal skills, for assessment, 13
- Interpreter, 9b
- Intertrigo, 245t, 259t
- Interventricular septum, 438f, 439t
- Intervertebral disc, herniation of, 791t
- Interview, 18–19
- communication in, 18–19
 - directive, 18
 - introductory phase of, 19
 - nondirective, 18
 - pitfalls in, 18–19
 - techniques for, 18
 - terminal phase of, 19
 - working phase of, 19
- Intracranial pressure, optic disc effects of, 328
- Introitus, vaginal, 614t, 631t, 633t
- Intuition, 13
- Inversion (body movement), 715f, 716t
- Iranian Americans
- adolescent, 915t
 - childbearing in, 823t
 - mental disorders and, 119t
 - newborn, 853t
 - older adult, 932t
 - pain response in, 96t
 - school-age, 904t
 - spirituality and, 191t
 - views of women and, 618t
- Iris, 322
- anatomy of, 322f, 323f
 - Brushfield spots of, 328–329, 351t
 - function of, 324t
 - inspection of, 351t
- Irish Americans
- adolescent, 915t
 - childbearing in, 822t
 - integumentary system of, 211
 - musculoskeletal health problems in, 694t
 - neurologic disorders in, 751
 - older adult, 932t
 - pain response in, 96t
 - respiratory disease in, 398–399
 - school-age, 904t
 - spirituality and, 192t
- Iritis, 361t
- Iron
- deficiency of, 184t–185t
 - during pregnancy, 169t
- Ischium, 687f
- Ishihara test cards, 341, 341f, 343t
- Islam, 197t
- Isometric maneuvers, for deep tendon reflex testing, 783, 784f
- Isoniazid, breast effects of, 536t
- Isotretinoin, musculoskeletal system effects of, 697t
- Italian Americans, gastrointestinal symptoms in, 562t
- Itching, 209, 213
- J**
- Jaeger cards, 341
- Japanese Americans, cardiovascular disease in, 454
- Jarel Spiritual Well-Being Scale, 199t–200t
- Jaundice, 209, 229t, 579t
- in infant, 855
 - in infants, 884–885
- Jaw, pain in, 276
- Jehovah's Witness, 196t
- Jejunostomy tube, 602t
- Jejunum, 557t
- Jewish Americans
- adolescent, 915t
 - childbearing in, 822t
 - gastrointestinal symptoms in, 562t
 - mental disorders and, 119t
 - newborn, 853t
 - older adult, 932t
 - pain response in, 97t
 - school-age, 904t
 - spirituality and, 192t
 - views of women and, 618t
- Jock itch, 263t, 670t
- Joint(s), 686, 688f, 690t. *See also* Musculoskeletal system
- abnormal findings in, 726t–728t
 - ankle, 725t–726t
 - assessment of, 714–716, 715f, 716t, 718t–728t
 - ball-and-socket, 689f, 691t
 - cervical, 718t
 - clicking of, 716
 - condyloid, 689f, 691t
 - deformity of, 716
 - elbow, 721t, 726t
 - finger, 722t–723t, 727t
 - hinge, 689f, 691t
 - hip, 723t–724t
 - knee, 725t, 727t
 - lumbar, 719t–720t
 - movements of, 715f, 716t
 - pivot, 689f, 691t
 - plane, 689f, 691t
 - range of motion of, 714–716, 715f
 - saddle, 689f, 691t
 - shoulder, 720t
 - sternoclavicular, 718t
 - temporomandibular, 718t
 - thoracic, 719t–720t
 - types of, 689f
 - wrist, 721t–722t
- Judaism, 197t
- Judgment test, in neurologic system assessment, 768t, 771
- Jugular vein, 463, 464f, 493f, 493t
- auscultation of, 478, 478t
 - inspection of, 470, 470b, 470t–471t, 471t, 472f
 - palpation of, 473t
 - pressure in, 449, 450–451, 451f, 470, 470b, 471t, 473f
- K**
- Kaposi's sarcoma, 260t, 300t
- Katz Index, 944b–945b
- Kayser-Fleischer ring, 351t

- Kehr's sign, 597, 598t
Keloid, 211, 237t
Keratinocytes, 207t
Keratohyaline, 207t
Keratosis
 actinic, 211, 212, 258t
 seborrheic, 211, 233t, 243t, 262t
Kernicterus, 884–885
Kernig's sign, 784, 784f
Kidneys, 558f, 559t
 age-related changes in, 930t
 palpation of, 591, 593, 595t
 tenderness of, 588, 591t
Kinesthetics, sensory examination for, 779t–780t
Kinetics, 8
Klinefelter's syndrome, 465
Knee
 deep bending of, 709t
 deviation of, 706t, 708t
 deviations of, 691
 DJD of, 727t
 range of motion of, 725t
 tests for, 729–730, 730f
Knee-jerk reflex, 744–745, 745t, 785t
 in newborns, 867t
Knock knees, 691, 708t
KOH prep, 637, 639t
Koilonychia, 250t
Korean Americans, older adult, 932t
Korotkoff's sounds, 83b, 453b, 516t
Kussmaul respiration, 412, 415b
Kwashiorkor, 185t
Kyphosis, 706t, 708t
 in older adults, 693, 707t
- L**
Labia majora, 612f, 614t
 palpation of, 633t
Labia minora, 612f, 614t
 palpation of, 633t
Labyrinth, 365t
Lachman test, 729–730, 730f
Lacrimal ducts, 322f
Lacrimal glands, 322f, 324t
 inspection of, 349t
 palpation of, 354t
Lactation, 530
Lacto-ovovegetarian diet, 164b
Lactovegetarian diet, 164b
Lambdoid suture, 268f, 271t
Langerhans' cells, 207t
Language, in neurologic system assessment, 768t
Lanugo, 850t, 859t, 860t
Large intestine, 557t
Laryngopharynx, 272t, 394f, 395t
Larynx, 394f, 395t
Lasègue's test, 728, 729f
Lateral spinothalamic tract, 742, 742f, 742t
Latissimus dorsi, 687f
Learning problems, 902, 907
Left ventricular impulse, 474–475
Leg
 circumference of, 514t, 705, 714t
 inspection of, 509t
 length of, 705, 713t–714t
 pain in, 514t–515t
 Legionnaire's disease, 407b
 Lemon, 63t
 Length, in newborns, 857t
 Lens, 322, 323f, 325t. *See also* Eye(s)
 inspection of, 350t–351t
 normal, 351t
 Lentiginosities, 233t, 956t
 Leopold's maneuver, 838t
 Leukemia, gingiva in, 299t
 Leukonychia, 249t
 Leukoplakia, 300t
 Levator palpebrae, 324t
 Level of consciousness, 764, 766t, 769, 769b
 Leydig, interstitial cells of, 650
 Lice, 252, 254
 eyelash, 347t
 pubic, 632t
 Lichen planus, 242t, 300t
 Lichenification, 235t
 Ligament, 686, 690t
 Light touch, 777t–778t
 Limbic system, 89, 113f, 740f, 740t
 Linea alba, 211
 Linea albicantes, 581
 Linea nigra, 211, 583t
 Lip(s), 269f, 271t
 herpes simplex virus infection of, 297t, 317t
 inspection of, 297t
 palpation of, 305t–306t
 Lipids, 164–165
 screening for, 156t
 Lipoma, 234t
 Lipoproteins, 165
 Listening, in patient interview, 10
 Literalness, 19
 Lithium, side effects of, 123t
 Liver, 556f, 557t
 age-related changes in, 930t
 auscultation of, 587t
 of newborns, 850, 851t, 855
 palpation of, 591, 593, 594t
 in pregnancy, 826
 size of, 588, 589t–590t
 Liver span test, 588
 Lochia, 844t
 Loop diuretics, nutrition and, 172t
 Lordosis, 706t, 708t
 in pregnancy, 693, 707t
 in toddlers, 691
 Louis, angle of, 464f, 471t
 LSD, abuse of, 136t
 Lubricant, 66t
 Ludwig's angina, 317t
 Lumbar spine, range of motion of, 719t–720t
 Lumbar vertebrae, 687f
 Lumbosacral disc herniation, 733t
 Lungs. *See also* Respiratory system
 anatomy of, 394f
 auscultation of, 425–430, 425t–426t
 in newborns, 863t
 cancer of, 432t, 433t
 Lunula, 208, 208f
 Lupus erythematosus, 209, 228t
 Lymph, 495b
 Lymph capillaries, 492
 Lymph nodes, 492, 495b
 axillary, 546, 547t–548t
 breast, 529f, 530t, 546, 547t–548t
 cervical. *See* Cervical lymph nodes
 clavicular, 546, 547t–548t
 epitrochlear, 511t, 547t
 inguinal, 515, 515t, 595, 597t, 674t–675t
 upper extremity, 512t, 515
 Lymphadenopathy, 307
 Lymphangitis, 521t
 Lymphatic capillaries, 495b
 Lymphatic system, 491–524
 abnormalities of, 520, 521t, 523t
 anatomy of, 492, 494f, 495b
 assessment of, 504–518
 in adolescents, 918t
 anatomical landmarks in, 504
 approach to, 504
 auscultation in, 515, 516t, 517b
 biographical data in, 497
 in breast disorders, 542
 in cardiovascular disorders, 460t, 467
 cardiovascular system in, 501t, 506t, 507
 case study of, 497, 503–504, 518
 checklist for, 524
 cultural considerations in, 495, 497
 current health status in, 497–498, 499t–500t
 developmental considerations in, 494–495
 drugs in, 499t–500t
 in ear disorders, 373t
 edema in, 498
 endocrine system in, 501t
 in eye disorders, 335t
 family history in, 500, 501t
 fatigue in, 498
 in female genitourinary disorders, 625t, 630t, 635
 focused history in, 497b
 in gastrointestinal disorders, 582
 gastrointestinal system in, 501t, 506t, 507
 general survey in, 501t, 505, 505t, 507
 genitourinary system in, 501t, 507
 in head, face, and neck disorders, 282t, 313
 head-to-toe, 505, 505t–506t
 health history in, 497–503, 497b
 HEENT system in, 501t, 505t, 507
 inspection in, 508t–509t, 510b
 integumentary system in, 501t, 505t, 507
 limb pain in, 498
 in male genitourinary disorders, 666
 in musculoskeletal disorders, 704
 musculoskeletal system in, 501t, 506t, 507
 in neurologic disorders, 758t, 765
 neurological system in, 501t, 506t, 507
 nursing diagnoses in, 518
 in nutritional disorders, 173t
 palpation in, 510, 510b, 511t, 515
 paresthesias in, 498
 past health history in, 498, 498t–499t
 physical, 504–518
 psychosocial profile in, 501, 502t–503t
 reproductive system in, 501t, 507
 in respiratory disorders, 419
 respiratory system in, 501t, 506t, 507
 review of systems in, 501, 501t
 in school-age children, 906t

- Lymphatic system (*continued*)
 symptom analysis in, 498
 in toddlers and preschoolers, 892t
 body systems interaction with, 494, 496
 cardiovascular system interaction with, 452, 494
 drug effects on, 499t–500t
 ear interaction with, 368
 eye interaction with, 327
 female genitourinary system interaction with, 616
 gastrointestinal system interaction with, 560
 head, face, and neck interaction with, 273, 274
 in infants, 494
 male genitourinary system interaction with, 655
 musculoskeletal system interaction with, 692
 neurologic system interaction with, 750
 in older adults, 495
 respiratory system interaction with, 397
 Lymphedema, 509t, 520, 522t
 prevention of, 520
- M**
- Macula, 323f, 324f, 325t
 assessment of, 359t
- Macular degeneration, 361t
 age-related, 360, 361t
- Macule, 233t
- Magnesium hydroxide, nutrition and, 172t
- Magnesium oxide, nutrition and, 172t
- Magnet reflex, 873t
- Malleus, 365t, 366, 366f, 367f
- Malnutrition, 167–170, 183t–185t, 580t
 assessment of. *See* Nutrition, assessment of
 definition of, 164
- Malocclusion, 298t
- Mammary gland. *See* Breast(s)
- Mammography, 549, 959t
- Mandible, 268f, 687f
- Mandibular pressure, 769
- Mania, 141t
- Mannitol, nutrition and, 172t
- Manual compression test, 518b
- Maple syrup urine disease, 874t
- Marasmus, 185t
- Marfan's syndrome, 465
- Marie-Strümpell disease, 732, 732t–733t
- Marijuana, abuse of, 136t
- Marking pen, 62t
- Mask of pregnancy, 211, 228t, 275, 814t, 834t
- Maslow's human needs hierarchy, 21, 21f
- Masseter, 274f
- Mastitis, 552t
- Mastoid process, 268f
- Mathematical skills, in neurologic system
 assessment, 767t, 771
- Maturation crisis, 131
- Maxilla, 268f, 687f
- Maxillary sinus, 268f
 inspection of, 292t
 palpation of, 293t
 percussion of, 293t
 transillumination, 292t
- McBurney's sign, 599t
- McGill Pain Questionnaire, 99
- McMurray's test, 730, 730f
- Mechanical digestion, 558
- Medulla, 741t
- Medulla oblongata, 453
- Meibomian gland, 323f
- Melanoma, 212, 230, 230f, 257, 260t
- Memory, in neurologic system assessment,
 766t–767t, 771
- Ménière's disease, 389t
- Meningeal signs, 784, 784f
- Meninges, 741t
- Meningitis, 791t
- Menopause, 617, 795–809
 assessment of, 797–808
 abdomen in, 805t
 approach to, 800–801
 biographical data in, 796–797
 case study of, 796–797, 798–800, 808
 current health status in, 797
 documentation for, 806t–807t
 family history in, 798
 focused physical assessment in, 801b
 general survey in, 801, 802t
 genitourinary system in, 805t
 head-to-toe, 801, 802t–805t
 health history in, 797–799
 HEENT system in, 802t–803t
 integumentary system in, 802t
 musculoskeletal system in, 804t–805t
 nursing diagnoses in, 808
 past health history in, 797
 physical, 801–807
 psychosocial profile in, 799
 respiratory system in, 804t
 review of systems in, 799f
 definition of, 796–797
 developmental issues in, 796
 integumentary system in, 211
 osteoporosis and, 808
- Menorrhagia, 619
- Menstrual cycle, 613, 613f
 assessment of, 624t
 disorders of, 619
 onset of, 914
 pain and, 619
- Mental health, 111–143
 assessment of, 120–137, 139t–142t
 in adolescent, 913
 in adolescents, 115, 913
 BATHE technique in, 131
 biographical data in, 120t–121t
 case study of, 118, 124, 137
 in children, 114–115, 115t
 developmental level in, 114–115
 drug history in, 122b, 123t
 general survey in, 127t
 head-to-toe, 127t
 health history in, 114, 120, 120t–121t
 maternal history in, 114
 mental status in, 126, 128t–131t
 in older adults, 116–117, 116b
 physical, 126
 in pregnant women, 116, 116b
 psychosocial profile in, 124,
 125t–126t
 review of systems in, 122, 123t–124t
- biological basis of, 112, 112b, 113f
 crisis in, 131, 132b
 cultural perceptions of, 118t–119t
 definition of, 112
 Healthy People 2010 objectives for, 138
 nursing diagnosis for, 137
 prevention of, 112
 theories of, 114
- Mental state, 78
- Mental status, assessment of, 126, 128t–131t,
 751, 752–753, 766t–768t, 770–771,
 771t, 787–788, 788b
 in older adults, 955t
- Mental Status Screening, 770
- Meperidine, abuse of, 136t
- Metabolic syndrome, 182–183
 breast cancer and, 549
- Metaphysis, 688f
- Metatarsals, 687f
- Methadone, abuse of, 136t
- Methylprednisolone, nutrition and, 172t
- Mexican Americans
 childbearing in, 822t
 mental disorders and, 119t
 newborn, 853t
 nutrition and, 171t
 older adult, 932t
 pain response in, 97t
 school-age, 904t
 spirituality and, 192t
 touch and, 9
 views of women and, 618t
- Miadam circumference, 181, 181f
- Micturition, 615, 653
- Midarm muscle circumference, 181
- Midbrain, 741t
- Middle-aged adults. *See also* Older adults
 exercise assessment in, 160
 injury prevention assessment in, 162
 sleep assessment in, 159
 stress assessment in, 161
 wellness assessment in, 158
- Migraine, 791t–792t
- Milia, 211, 859t
- Miliaria, 859t
- Milk lines, 543f
- Minerals, 165
- Mini-Mental State Examination, 770, 787
- Miosis, 352t
- Mitral regurgitation, 480t
- Mitral stenosis, 480t
- Mitral valve, 438f, 439t
- Modified Behavioral Pain Scale, 105
- Mole (nevus), 233t
- Mongolian spots, 211, 859t
- Mononeuropathy, 93
- Mononucleosis, 317t
- Mons pubis, 614t
- Montgomery's glands, 528f, 530t
- Mood disorders, 141t
- Mormonism, 196t
- Moro reflex, 868t
- Morphine, abuse of, 136t
- Motivational level, health and, 155
- Motor cortex, 743–744, 743f
- Motor neurons, 744
- Motor pathways, 743–744, 743f

- Motor vehicle injury, 889
- Mouth, 556t. *See also* Gingivae (gums); Lip(s); Teeth; Tongue
- age-related changes in, 929t
 - anatomy of, 269f
 - assessment of, 37t, 296, 305
 - in adolescents, 917t, 921t
 - in ear disorders, 377t
 - in gastrointestinal disorders, 580t
 - in head, face, and neck disorders, 281t
 - in infants, 880t
 - inspection in, 296, 297t–305t
 - in newborns, 861t–862t
 - in older adults, 937t, 951t
 - palpation in, 296, 305t–306t
 - in school-age children, 909t
 - in toddlers and preschoolers, 894t
 - cancer of, 296b, 300t, 314–315
 - functions of, 271t
 - in older adults, 275
 - in pregnant women, 275
- Movements, 715f. *See also* Joint(s)
- Mucopolysaccharides, 206
- Mucosa
- nasal, 271t
 - inspection of, 294t–295t
 - oral, 271t
 - inspection of, 296, 299t
- Murmurs, 449–450, 449f, 450f
- auscultation of, 480t, 483, 483t–484t–485t, 482b
 - vs. bruit, 478
 - characteristics of, 485b
 - in children, 453, 895t–896t
 - continuous, 449, 451f
 - diastolic, 449, 450f, 483t, 484t–485t, 482b
 - in newborns, 864t
 - in older adult, 952t
 - in pregnant women, 453
 - systolic, 449, 450f, 482t, 483t–484t, 485b
- Murphy's sign, 597–598, 599t
- Muscle(s), 686, 690t. *See also* Musculoskeletal system
- assessment of, 717t
 - strength of, 716t, 717t
- Muscle relaxants, neurologic system effects of, 756t
- Muscle wasting, in cardiac disease, 465
- Musculoskeletal system, 685–736, 690t
- abnormalities of, 732t–734t
 - age-related changes in, 931t
 - anatomy of, 686
 - assessment of, 39t, 693–730
 - in adolescents, 918t, 923t
 - Apley's test in, 730, 730f
 - approach to, 701
 - balance in, 695, 705
 - biographical data in, 694
 - in breast disorders, 541t, 542
 - bulge test in, 729, 729f
 - in cardiovascular disorders, 460t, 467, 470t
 - cardiovascular system in, 698t, 703t, 704
 - case study of, 693, 700–701, 730–731
 - cerebellar function in, 703, 705, 709t–711t
 - checklist for, 734–735
 - coordination in, 695, 705, 710t–711t
 - cultural considerations in, 693, 694t
 - current health status in, 694
 - developmental considerations in, 688–689, 691, 693
 - drugs in, 696t–697t
 - in ear disorders, 373t, 377t, 386
 - endocrine system in, 698t, 704
 - in eye disorders, 334t, 339t, 340
 - family history in, 697, 697t
 - in female genitourinary disorders, 624t, 630t, 635
 - focused history in, 694b
 - gait in, 702–703, 705, 705f, 708t–709t, 711t–713t
 - in gastrointestinal disorders, 572t, 581t, 582
 - gastrointestinal system in, 698t, 703t, 704
 - general survey in, 697t, 702, 704
 - genitourinary system in, 698t, 703t, 704
 - goniometer in, 702f
 - in head, face, and neck disorders, 282t, 288t, 313
 - head-to-toe, 702, 703t
 - health history in, 693–700, 694b
 - HEENT system in, 698t, 703t, 704
 - hematologic system in, 698t, 704
 - immunologic system in, 698t
 - in infants, 877t
 - in integumentary disorders, 218t, 224t, 225
 - integumentary system in, 697t–698t, 703t, 704
 - joints in, 714–716, 715f, 716t, 718t–728t
 - Lachman test in, 729–730, 730f
 - Lasègue's test in, 728, 729f
 - limb measurements in, 705, 713t–714t
 - in lymphatic disorders, 501t, 506t, 507
 - lymphatic system in, 704
 - in male genitourinary disorders, 661t, 666, 668t
 - McMurray's test in, 730, 730f
 - for menopause, 804t–805t
 - in mental health disorders, 124t, 127t
 - movement accuracy in, 705, 711t
 - muscles in, 717t
 - in neurologic disorders, 758t, 764t, 765
 - neurologic system in, 698t
 - in newborns, 866t–867t
 - nursing diagnoses in, 731
 - in nutritional disorders, 173t, 179t
 - in older adults, 939t–940t, 954t–955t
 - in pain disorders, 104t
 - pain in, 695
 - past health history in, 695, 695t–696t
 - patellar ballotement test in, 729, 729f
 - in patient teaching, 148t, 151t
 - in peripheral-vascular disorders, 501t, 506t
 - physical, 701–730
 - in postpartal woman, 845t
 - posture in, 702, 706t–708t
 - in pregnancy, 829t, 841t
 - pronator drift test in, 728, 728f
 - psychosocial profile in, 699, 699t–700t
 - range of motion in, 715f, 718t–726t
 - reproductive system in, 698t, 703t, 704
 - in respiratory disorders, 404t, 414t, 419
 - respiratory system in, 698t, 703t, 704
 - review of systems in, 697, 697t–698t
 - in school-age children, 906t, 911t
 - stiffness in, 695
 - symptom analysis in, 695
 - Thomas test in, 729, 729f
 - in toddlers and preschoolers, 891t, 896t
 - Trendelenburg test in, 729
 - weakness in, 695
 - body systems interaction with, 686, 688, 692
 - cardiovascular system interaction with, 452
 - in children, 688–689, 691, 693
 - drug effects on, 696t–697t
 - ear interaction with, 366–367, 368
 - eye interaction with, 327, 328
 - female genitourinary system interaction with, 614, 616
 - gastrointestinal system interaction with, 558, 560
 - head, face, and neck interaction with, 270, 273
 - hearing and, 366–367, 368
 - in infants, 688–689, 691, 693
 - integumentary system interaction with, 210
 - male genitourinary system interaction with, 653, 655
 - neurologic system interaction with, 686, 688, 692, 748, 750
 - in newborns, 851t
 - in older adults, 693
 - in pregnant women, 693, 817t
 - respiratory system interaction with, 396, 397, 688, 692

Musset's sign, 465

Mydriasis, 352t

Myelin, 739f

Myocardium, 438f, 439t

Myopia, 326, 326f

N

Naegele's rule, 825

Nail pressure, 769

Nails. *See* Fingernails

NANDA-unitary person framework, 21

Narcissistic personality disorder, 140t

Narcotics

 - cardiovascular system effects of, 458t
 - gastrointestinal system effects of, 569t
 - neurologic system effects of, 756t

Nares, 268f, 271t

Narrative documentation, 24

Nasal bone, 268f

Nasal septum, 271t

 - deviated, 295t

Nasal vestibule, 269f, 271t

Nasogastric tube, 601t

Nasolabial fold, 286, 286f

Nasolacrimal duct, 322f

 - inspection of, 349t
 - palpation of, 354t

Nasolacrimal sac, 324t

Nasopharynx, 269f, 272t, 394f, 395t

Nasoscope, 60t

Natal teeth, 861t

- Native Americans. *See also* Navajo Native Americans
- cardiovascular disease in, 454
 - cerumen in, 370
 - gastrointestinal symptoms in, 562t
 - mental disorders and, 119t
 - musculoskeletal health problems in, 694t
 - otitis media in, 370
 - views of women and, 618t
- Nausea, 566
- in abdominal assessment, 566
- Navajo Native Americans
- adolescent, 915t
 - childbearing in, 822t
 - musculoskeletal health problems in, 694t
 - neurologic disorders in, 751
 - newborn, 853t
 - nutrition and, 171t
 - older adult, 932t
 - pain response in, 97t
 - school-age, 904t
 - spirituality and, 192t
- Neck. *See also* HEENT system
- anatomy of, 268, 270f
 - anterior triangle of, 286, 286f
 - assessment of, 275–314, 306–312
 - anatomical landmarks in, 286, 286f
 - auscultation in, 311, 312t, 478, 478t, 516t
 - biographical data in, 275
 - cardiovascular system in, 281t, 288t, 313
 - case study of, 275, 284–285, 312, 314
 - checklist for, 318–319
 - cultural considerations in, 275
 - current health status in, 275–276
 - developmental considerations in, 274–275
 - drugs in, 278t–279t, 279
 - in ear disorders, 377t
 - endocrine system in, 282t, 313
 - in eye disorders, 339t
 - family history in, 279, 279t–280t
 - focused history in, 276b
 - gastrointestinal system in, 281t, 288t, 313
 - general survey in, 281t, 287, 287t, 313
 - genitourinary system in, 282t, 288t, 313
 - head-to-toe, 287, 287t–288t
 - health history in, 275–284, 276b
 - HEENT system in, 281t, 288t, 313
 - hematologic system in, 282t, 313
 - inspection in, 307t, 470, 470t–471t
 - integumentary system in, 281t, 287t, 313
 - lymphatic system in, 282t, 313
 - musculoskeletal system in, 282t, 288t, 313
 - neurologic system in, 288t, 313
 - in newborns, 860t
 - nursing diagnoses in, 314
 - palpation in, 308t–311t, 472, 473t, 511t
 - past health history in, 276, 277t–278t
 - physical, 286–318
 - psychosocial profile in, 283, 283t–284t
 - reproductive system in, 282t, 313
 - respiratory system in, 288t, 313
 - review of systems in, 280, 281t–282t
 - in school-age children, 909t
 - symptom analysis in, 276
 - body systems interaction with, 269–270, 273, 274
 - cardiovascular system interaction with, 270, 273
 - distended veins of, 413t, 414t
 - drug effects on, 278t–279t, 279
 - endocrine system interaction with, 273, 274
 - hematologic system interaction with, 273, 274
 - of infant, 275
 - lymph nodes of. *See* Cervical lymph nodes
 - lymphatic system interaction with, 273, 274
 - mass in, 276
 - musculoskeletal system interaction with, 270, 273
 - neurologic system interaction with, 270, 273
 - pain in, 276
 - posterior triangle of, 286, 286f
 - in pregnant women, 275
 - respiratory system interaction with, 269–270, 273
 - stiffness in, 276
- Neck reflex, 868t–869t
- Necrosis, 209
- Needs, Maslow's hierarchy of, 21, 21f
- Negativism, 888
- Neglect, 762
- Neologisms, 766b
- Neonatal infant Pain Scale, 103
- Neonates. *See* Newborns
- Nerve fibers, in pain transmission, 88f, 89
- Neuritis, optic, 361t
- Neuroglia, 738
- Neuroleptic malignant syndrome, 122
- Neurologic system, 737–794
- abnormalities of, 789t–792t
 - in adolescents, 749, 751
 - afferent pathways of, 738, 739f
 - age-related changes in, 931t
 - anatomy of, 738–746, 740f, 740t–741t, 747f, 748f
 - anterior spinothalamic tract of, 742, 742f
 - ascending tracts of, 739
 - assessment of, 39t, 751–787
 - abstract thinking in, 767t, 771
 - in adolescents, 918t, 923t
 - approach to, 761
 - auditory stimuli in, 764, 769
 - biographical data in, 752
 - in breast disorders, 541t, 542
 - calculative skills in, 767t, 771
 - in cardiovascular disorders, 460t, 467, 470t
 - cardiovascular system in, 757t, 764t, 765
 - case study of, 751, 760–761, 787
 - cerebral function in, 764, 766t–768t
 - checklist for, 793
 - cognitive function in, 766t–768t, 770
 - communication evaluation in, 768t, 771
 - cranial nerve function in, 771, 772t–777t
 - cultural considerations in, 751
 - current health status in, 752
 - developmental considerations in, 749, 751
 - dizziness in, 753
 - documentation for, 769, 783f
 - drugs in, 755t–756t
 - in ear disorders, 373t, 377t, 386
 - endocrine system in, 758t, 765
 - equipment for, 761
 - in eye disorders, 334t, 339t, 340
 - family history in, 756, 756t
 - in female genitourinary disorders, 624t, 630t, 635
 - focused history in, 752b
 - in gastrointestinal disorders, 581t, 582
 - gastrointestinal system in, 757t–758t, 764t, 765
 - general knowledge in, 767t, 771
 - general survey in, 757t, 762, 763t, 765
 - genitourinary system in, 764t, 765
 - Glasgow Coma Scale in, 769, 770t
 - in head, face, and neck disorders, 282t, 288t, 313
 - head-to-toe, 762, 763t–764t
 - headache in, 752
 - health history in, 751–760
 - HEENT system in, 757t, 763t–764t, 765
 - hematologic system in, 758t, 765
 - in infants, 878t, 883t
 - in integumentary disorders, 218t, 224t, 225
 - integumentary system in, 757t, 763t, 765
 - judgment test in, 768t, 771
 - language evaluation in, 768t
 - level of consciousness in, 764, 766t, 769, 769b
 - in lymphatic disorders, 501t, 506t, 507
 - lymphatic system in, 758t, 765
 - in male genitourinary disorders, 662t, 666, 668t
 - mathematical skills in, 767t, 771
 - memory in, 766t–767t, 771
 - meningeal signs in, 784, 784f
 - in mental health disorders, 124t, 127t
 - mental status in, 752–753, 766t–768t, 770–771, 787–788, 788b
 - in musculoskeletal disorders, 698t, 703t, 704
 - musculoskeletal system in, 758t, 764t, 765
 - neurologic system in, 765
 - in newborns, 855, 867t
 - numbness in, 753
 - nursing diagnoses in, 787
 - in nutritional disorders, 173t, 179t
 - in older adults, 940t, 955t–956t
 - orientation in, 766t, 769–770
 - in pain disorders, 104t
 - painful stimuli in, 769
 - paresthesia in, 753
 - past health history in, 753, 753t–754t
 - in patient teaching, 148t, 151t
 - in peripheral-vascular disorders, 501t, 506t
 - physical, 761–787
 - in postpartal woman, 845t
 - in pregnancy, 829t–830t, 841t
 - psychosocial profile in, 758, 758t–759t
 - reflexes in, 782–783, 783f, 784f, 784t–786t
 - reproductive system in, 758t, 764t
 - in respiratory disorders, 404t, 414t, 419
 - respiratory system in, 757t, 764t, 765

- review of systems in, 756, 757t–758t
- in school-age children, 906t, 911t
- sensory system evaluation in, 753, 777, 777t–782t
- stroke scale in, 763b
- symmetry in, 762
- symptom analysis in, 752–753
- syncope in, 753
- thought process in, 767t, 771
- tingling in, 753
- in toddlers and preschoolers, 892t, 897t
- vertigo in, 753
- visual changes in, 753
- vital signs in, 762
- vocabulary in, 767t, 771
- body systems interaction with, 746, 748, 750
- cardiovascular system interaction with, 453, 748, 750
- cells of, 738, 739f
- in children, 749, 751
- descending tracts of, 739
- drug effects on, 755t–756t
- ear interaction with, 367, 368, 369
- efferent pathways of, 738, 739f
- endocrine system interaction with, 748, 750
- eye interaction with, 327, 328, 329f
- female genitourinary system interaction with, 615, 616
- gastrointestinal system interaction with, 559, 560
- head, face, and neck interaction with, 270, 273
- hearing and, 367, 368, 369
- in infants, 749
- integumentary system interaction with, 209, 210
- lateral spinothalamic tract of, 742, 742f
- male genitourinary system interaction with, 653, 655
- motor pathways of, 743–744, 743f
- musculoskeletal system interaction with, 748, 750
- in newborns, 749, 851t
- in older adults, 751
- in pregnant women, 751, 817t
- respiratory system interaction with, 396, 397, 748, 750
- sensory pathways of, 739, 741–742, 742f, 742t
- Neuroma, acoustic, 388t
- Neuromuscular junction, 739f
- Neuron, 738, 739f
- Neuropathic Pain Scale, 99
- Neuropeptides, 89
- Neurotransmitters, 112, 112b, 738
- Nevus (mole), 230, 230f, 240t, 350t
 - change in, 212
 - congenital, 212
 - dysplastic, 212
- Nevus flammeus, 211, 860t
- New Age beliefs, 198t
- Newborns, 849–886. *See also* Infants
 - assessment of, 853–875
 - abdomen in, 864t
 - abdominal circumference in, 857t
 - Apgar score in, 854, 854t
 - approach to, 854–855
 - Barlow-Ortolani maneuver in, 855, 866t
 - blood pressure in, 858t–859t
 - cardiovascular system in, 864t
 - case study of, 853, 874–875
 - chest circumference in, 856t
 - chest in, 863t
 - cultural considerations in, 850, 852, 853t
 - developmental considerations in, 850–852
 - female genitalia in, 865t
 - general survey in, 855
 - head circumference in, 856t
 - head-to-toe, 855, 856t–867t
 - health history in, 854, 854t
 - HEENT system in, 860t–862t
 - integumentary system in, 859t–860t
 - length measurement in, 857t
 - male genitalia in, 865t–866t
 - musculoskeletal system in, 866t–867t
 - neurological system in, 867t
 - physical, 854–875
 - physiological changes in, 850, 850t–851t, 852f
 - posture in, 856t
 - pulse in, 858t
 - rectum in, 865t
 - reflexes in, 867, 868t–873t
 - respiratory system in, 858t, 863t
 - screening tests in, 867, 874t
 - temperature measurement in, 858t
 - weight measurement in, 857t–858t
 - body temperature of, 79
 - cardiovascular system of, 851t
 - circulatory adaptation in, 850, 852f
 - gastrointestinal system of, 851t
 - genitourinary system of, 851t
 - head of, 850t
 - health problems in, 860t, 862t, 875
 - hepatic adaptation of, 850
 - integumentary system of, 850t
 - liver of, 851t
 - musculoskeletal system of, 851t
 - neurologic system of, 749, 851t
 - physiological changes in, 850, 850t–851t
 - respiratory system of, 851t
 - sleep in, 851t
 - thermoregulation in, 851t
 - Niacin, nutrition and, 172t
 - Nicotine
 - abuse of, 137t
 - sleep and, 158
 - Night terrors, 158
 - Nightingale, Florence, 14
 - Nightmares, 158
 - Nipple, 528f, 529f, 530t
 - discharge from, 534
 - inspection of, 545t
 - palpation of, 546, 546t–547t
 - Nissl bodies, 739f
 - Nociception, 88, 88f
 - Nociceptors, 88f, 89
 - Nodule, 234t
 - Noise, hearing loss and, 370
 - Nonfluent aphasia, 766b
 - Nonsteroidal anti-inflammatory drugs
 - ear effects of, 372t
 - gastrointestinal system effects of, 569t
 - integumentary system effects of, 216t
 - lymphatic system effects of, 500t
 - neurologic system effects of, 756t
 - respiratory system effects of, 402t–403t
 - Norepinephrine, 112b
 - Nose. *See also* HEENT system
 - age-related changes in, 929t
 - anatomy of, 268f, 269f, 394f
 - assessment of, 37t, 292, 294–295
 - in adolescents, 921t
 - in infants, 880t
 - inspection in, 294, 294t–295t
 - in newborns, 861t
 - in older adults, 937t, 951t
 - palpation in, 295, 296t
 - in school-age children, 909t
 - in toddlers and preschoolers, 894t
 - congestion of, 276
 - functions of, 271t, 395t
 - of infant, 274
 - polyps of, 295t, 317t
 - Nosebleed, 276, 316t
 - Numbness, in neurologic system assessment, 753
 - Numeric scales, for pain, 99, 100
 - Nurses Assessment of Pain Inventory, 106
 - Nursing diagnosis, 4f, 5, 5b, 6b–7b
 - actual, 5, 5b
 - possible, 5
 - potential, 5
 - Nursing process, 4–5, 4b, 4f
 - Nutrients, 164–167
 - Nutrition, 163–187
 - for adolescents, 168t–169t, 919
 - assessment of, 170–182
 - 24-hour recall in, 175–176
 - anthropometry in, 177, 179–182
 - biographical data in, 170
 - current health status in, 170–171
 - disease-specific, 183t–185t
 - drugs in, 172t
 - family history in, 172
 - focused history in, 175b
 - food intake records in, 176
 - general survey in, 173t, 178t
 - head-to-toe assessment in, 178t–179t
 - health history in, 170–172
 - nutritional history in, 175–176
 - past health history in, 171–172
 - physical, 177–182, 178t–179t
 - psychosocial profile in, 172, 173t–175t
 - review of systems in, 172, 173t
 - case study of, 170, 176–177, 182
 - culture and, 170, 171t
 - definition of, 164
 - disorders of, 183t–185t
 - guidelines for, 165–167, 166f
 - for infants, 168t
 - nursing diagnosis for, 182
 - for older adults, 169–170, 942
 - during pregnancy, 845–846
 - for pregnant women, 169, 169t
 - for preschoolers, 168t, 892
 - for school-age children, 167t, 168t, 907
 - for toddlers, 168t, 892

- Nutritional deficiency, 167–170
 Nutritional history, 175–176, 175b
- O**
- Obesity
 in adolescents, 913, 924
 health effects of, 182–183
 in school-age children, 902
 screening for, 157t
 in toddlers and preschoolers, 897t
 waist-to-hip ratio for, 182
- Object cues, 8
 Objective data, 17
 Objective Pain Score, 105
 Oblique muscles (ocular), 323f, 325t
 Observation, 19–20
 Obsessive-compulsive disorders, 139t, 140t
 Obstetric history, 624t. *See also* Pregnancy
 Obturator muscle test, 600t
 Occipital bone, 268f
 Occipital lobe, 740f, 741t
 Occipitalis, 274f
 Oculocephalic reflex, 772t–773t
 Oculomotor nerve (CN III), 268f, 328, 746t, 747f, 772t–773t
 Oculovestibular reflex, 775t
 Odor, body, 227t
 OLDCART mnemonic, 98b
 Older adults, 927–960
 abuse of, 117, 1036
 advance directives for, 946
 aging-related changes in, 928t–931t, 956t–957t
 assessment of, 73, 933–957
 approach to, 948
 Barthel Index in, 945t
 biographical data in, 934
 cardiovascular system in, 938t, 952t
 case study of, 933, 946–948, 957–958
 clock scoring in, 957b
 cultural considerations in, 932, 932t
 current health status in, 934–935
 disease presentation in, 933
 endocrine system in, 940t
 fall risk in, 941b
 family history in, 935
 female reproductive system in, 953t
 functional, 944, 944b–945b
 gastrointestinal system in, 938t, 953t
 general survey in, 935t, 948
 genitourinary system in, 938t–939t
 head-to-toe assessment of, 948, 949t–957t
 health history in, 933–934
 HEENT system in, 936t–937t, 950t–951t
 hematologic system in, 940t
 immunologic system in, 940t
 incontinence in, 941t
 instrumental activities of daily living in, 945t–946t
 integumentary system in, 936t, 949t–950t
 Katz Index in, 944, 944b–945b
 male reproductive system in, 954t
 medications in, 934t, 935
 multiple disorders in, 933
 musculoskeletal system in, 939t–940t, 954t–955t
 neurological system in, 940t, 955t–956t
 past health history in, 935
 physical, 948–957
 psychosocial profile in, 942–944
 reproductive system in, 939t, 953t–954t
 respiratory system in, 937t, 952t
 review of systems in, 935, 935t–940t
 screening tests in, 958, 959t
 skin lesions in, 956t–957t
 surgery in, 935
 symptoms in, 933, 934–935
 body temperature of, 79
 breasts in, 532
 cardiovascular system in, 453–454
 cognitive impairment in, 958
 digestion in, 561
 drug effects in, 935, 943t
 endocrine system in, 931t
 exercise for, 160, 942
 gastrointestinal system in, 929t–930t
 hearing loss in, 387
 HEENT system in, 928t–929t
 hematologic system in, 931t
 home safety for, 944
 immunological system in, 931t
 injury prevention for, 162
 integumentary system in, 211, 928t
 lymphatic system in, 495
 male genitalia in, 654, 657
 menopause in, 617
 mental health assessment in, 116–117
 mouth of, 275
 musculoskeletal system in, 693, 931t
 neurologic system in, 751
 neurological system in, 931t
 nutrition in, 169–170, 942, 943t
 pain in, 95, 95b, 106t, 107
 peripheral-vascular system in, 495
 reproductive system in, 930t
 respiratory system in, 398, 929t
 self-care in, 943–944
 sleep in, 159, 942–943, 943b
 spirituality in, 190
 stress in, 161
 suicide in, 117b
 support systems for, 944
 taste in, 275
 teeth of, 275
 visual acuity in, 330
 wellness assessment in, 158
- Olfactory bulb, 269f
 Olfactory nerve (CN I), 268f, 271t, 746t, 747f, 772t
- Onycholysis, 250t
 Open snap, 479t
 Opening snap, 482f
 Ophthalmoscope, 59t
 Ophthalmoscopy, 355, 355b, 356t–359t
 apertures for, 356t
 Opiates, abuse of, 136t
 Opium, abuse of, 136t
 Opposition (body movement), 715f, 716t
 Optic chiasm, 326, 326f
 Optic disc, 324f
 anatomy of, 323f, 326, 326f, 355, 355f
 blood vessels of, 358t–359t
 cardiovascular effects on, 328
 color of, 330
 examination of, 355f, 357t–358t
 function of, 325t
 Optic disc cup, 325t, 355, 355f
 Optic fundus, 328
 Optic nerve (CN II), 268f, 326, 326f, 746t, 747f, 772t–773t
 anatomy of, 323f, 329f
 Optic radiation, 326, 326f
 Optic track, 326, 326f
 Oral antidiabetics, integumentary system effects of, 216t
 Oral cancer, 296b, 300t, 303t, 314–315
 Oral contraceptives
 breast effects of, 535t
 eye effects of, 333t
 nutrition and, 172t
 Orbicularis oculi, 274f, 323f, 324t
 Orbicularis oris, 274f
 Orbital fat, 323f
 Orchitis, 673t, 681t
 Organ of Corti, 365t, 367
 Orgasm, male, 650
 Orientation test, 769–770
 Orlistat, nutrition and, 172t
 Oropharynx, 269f, 272t, 394f, 395t
 inspection of, 303t–304t
 Osteoarthritis, 732, 955t
 Osteoblast, 686
 Osteoclast, 686
 Osteogenesis imperfecta, hearing in, 372t
 Osteoporosis, 731, 734t, 808
 screening for, 157t
 Otalgia, 371
 Otitis externa, 380t, 389t
 Otitis media, 370, 381t, 388t
 adhesive, 381t
 serous, 381t, 389t
 Otolaryngology, 60t
 Otoscopy, 379, 379t–381t, 382
 Oucher scale, 99–100
 Ovarian cancer, 644, 646t
 Ovarian cyst, 645t
 Ovary (ovaries), 559t, 612f, 613f, 615t
 bimanual examination of, 641t
 Oxycodone, abuse of, 136t
- P**
- Pain, 87–108
 abdominal, 563–564, 564b, 565b
 acute, 91t, 92
 assessment of, 88, 97–107, 98b
 behavioral pain assessment scales in, 102–107, 105t
 biographical data in, 97
 in children, 94–95, 99–100, 100f, 105–107
 current health status in, 97–98
 family history in, 100
 head-to-toe, 104t
 health history in, 97–98
 in infants, 93–94, 102–107, 105t
 in older adults, 95, 95b
 past health history in, 100
 physical, 102–107, 104t
 postoperative, 102–103, 106–107, 106t
 psychosocial profile in, 100, 101t–102t
 review of systems in, 100

- scales in, 98–100, 100f
- terminology in, 98b
- back, 564, 688
 - in pregnancy, 693
- behavioral response to, 103b
- breast, 534
- cancer-related, 91t, 92
- case study of, 95, 107
- central, 90, 93
- chest, 400, 420t, 455, 487t
- in children, 94–95, 105–107
- chronic, 91t, 92
- classification of, 90–92, 91t
- culture and, 95, 96t–97t
- deafferentation, 93
- deep, 93
- in dementia, 106t, 107
- dental, 276
- ear, 371
- eye, 331
- gate-control theory of, 90, 90f
- genital, 658
- head, 276
- in infants, 93–94, 102–107
- jaw, 276
- limb, 498, 514t
- menstrual, 619
- misconceptions about, 88b
- modulation of, 90, 90f
- mouth, 276
- musculoskeletal, 695
- neck, 276
- neuropathic, 89, 93
- nociceptive, 92–93
- in older adults, 95, 95b
- parietal, 564
- perception of, 89–90
- peripheral sensitization to, 90
- phantom limb, 93
- physiological response to, 103b
- psychological response to, 103b
- quality of, 565b
- radiating, 92, 565b
- rating scales for, 33b, 94, 98–100, 100f
 - behavioral, 102–107, 105t
- reassessment of, 107
- referred, 92, 92b, 92f, 564
- response to, 104t
- scapular, 564
- sensitization to, 90
- sensory examination for, 778t–779t
- shoulder, 564
- somatic, 92, 93
- stimuli of, 88–89, 89t
- superficial, 93
- sympathetically maintained, 93
- threshold of, 89–90
- tolerance for, 90
- transduction of, 88–89, 88f
- transmission of, 88f, 89
- visceral, 92, 563–564
- Pain Assessment in Advanced Dementia Scale,** 107
- Pain Assessment Tool,** 103
- Pain Rating Scale, for infant,** 103–104
- Pain scales, 98–100, 100f**
 - for adults, 99
- behavioral, 102–107, 105t
- for infants, 102–107, 105t, 106t
- multidimensional, 99
- unidimensional, 99
- Palate,** 269f, 272t, 301t
- Pallor,** 229t
- Palmar erythema, in pregnancy,** 495, 814t, 834t
- Palmar grasp reflex,** 783, 869t
- Palpation,** 20, 67–69
 - abdominal, 591–595, 592t–595t
 - abdominal aorta, 591, 591t, 593, 593t
 - ballottement, 69, 70f
 - bladder, 595, 596t
 - breast, 541, 543–544, 543f, 544f, 546, 546t
 - carotid artery, 472, 473t
 - cervical nodes, 306–308, 308t–310t
 - chest, 418, 420t–422t, 474–475
 - deep, 68–69, 69f, 591, 592t
 - ear, 379t
 - epigastric area, 474t
 - eyeball, 354t
 - face, 290, 291t
 - female genitalia, 630, 633t, 634, 637, 640, 640t–641t
 - hair, 254, 255t
 - hand, 722t–723t
 - hand areas for, 68f
 - head, 290, 291t
 - hernia, 674t–675t, 676, 677f
 - inguinal lymph nodes, 595, 597t
 - jugular vein, 473t
 - kidney, 591, 593, 595t
 - labia majora, 633t
 - labia minora, 633t
 - lacrimal gland, 354t
 - light, 67, 69f, 591, 592t
 - lips, 305t–306t
 - liver, 591, 593, 594t
 - lymph node, 306–308, 308t–310t, 511t–512t, 515, 546, 547t–548t, 674t–675t
 - male genitourinary system, 672, 672t–675t, 676, 677f
 - mouth, 296, 305t–306t
 - muscle, 717t
 - nails, 251t, 252
 - nasolacrimal duct, 354t
 - neck, 306–308, 308t–311t, 472, 473t, 511t
 - nipple, 546, 546t–547t
 - nose, 295, 296t
 - paranasal sinus, 292, 293t
 - parotid gland, 306t
 - precordium, 474–475, 474t–475t
 - pulsations, 69f
 - pulses, 510t, 512t–514t. *See also* Pulse(s)
 - salivary gland, 296, 306t
 - scalp, 254, 255t
 - skin, 240, 243t–246t, 247, 247t–248t, 249, 251t, 252
 - spleen, 591, 593, 594t
 - sublingual gland, 306t
 - tactile fremitus, 68f
 - temperature, 68f
 - temporomandibular joint, 291t
 - thyroid gland, 308, 310t–311t
 - tongue, 296, 306t
- vibratory sensations, 68f
- wrist, 721t–722t
- Palpebral fissure,** 286, 286f
- Palpitations,** 456
- Pancreas,** 557t–558t, 558f
 - age-related changes in, 930t
 - in pregnancy, 818t
- Pancreatitis,** 607t
- Panic disorders,** 139t
- Pap test,** 634, 637, 638t
 - equipment for, 65t
 - for older adults, 959t
- Paper clip,** 64t
- Papilledema,** 361t
- Papilloma,** 350t, 551t
- Papules,** 209, 233t
- Paralinguistics,** 7–8
- Paralytic ileus,** 607t
- Paranasal sinuses,** 37t, 268f, 271t, 290–291
 - of infant, 274
 - palpation of, 292, 293t
 - percussion of, 292, 293t
 - transillumination of, 291, 292t
- Paranoid personality disorder,** 140t
- Paraphimosis,** 682t
- Paraphrasia,** 766b
- Parasympathetic nervous system,** 453, 746, 749f
- Parathyroid glands,** 272t
 - in pregnancy, 818t
- Paresthesias,** 456, 498
 - in neurologic system assessment, 753
- Parietal bone,** 268f
- Parietal lobe,** 740f, 741t
- Parietal pain,** 564
- Parkinson's disease,** 792t
- Paronychia,** 250t
- Parotid duct,** 270f
- Parotid gland,** 270f, 272t, 557t
 - palpation of, 306t
- Parotitis,** 302t, 315t
- Past health history,** 28–29, 28t, 33, 33t–34t
 - in abdominal assessment, 566, 567t–568t
 - in adolescent assessment, 916
 - in breast assessment, 534, 534t–535t
 - in cardiovascular system assessment, 456, 456t–457t
 - in ear assessment, 371, 371t
 - in eye assessment, 325, 332
 - in facial assessment, 276, 277t–278t
 - in female genitourinary system assessment, 619, 620t–621t
 - in head assessment, 276, 277t–278t
 - in integumentary system assessment, 214, 214t–215t
 - in lymphatic system assessment, 498, 498t–499t
 - in male genitourinary system assessment, 658–659, 659t
 - in menopause assessment, 797
 - in musculoskeletal system assessment, 695, 695t–696t
 - in neck assessment, 276, 277t–278t
 - in neurologic system assessment, 753, 753t–754t
 - in nutrition assessment, 171–172
 - in older adult assessment, 935

- Past health history (*continued*)
 in pain assessment, 100
 in peripheral-vascular system assessment, 498, 498t–499t
 in pregnancy assessment, 825–826
 in respiratory system assessment, 400, 401t
 in school-age child assessment, 904
 in spirituality assessment, 192–193
 in teaching assessment, 147
 in toddler and preschooler assessment, 890
- Patella, 687f
- Patellar ballottement test, 729, 729f
- Patellar reflex, 744–745, 745t, 785t
 in newborns, 867t
- Pectoralis major, 687f
- Pediatric Anxiety Rating Scale, 115t
- Pediculosis, 252, 254
- Pedigree, 1038–1039
- Peers
 adolescent, 914
 school-age, 903
- Pelvic examination, 65t, 634, 636t–637t
- Penicillins
 integumentary system effects of, 216t
 respiratory system effects of, 403t
- Penis, 650, 653t. *See also* Genitourinary system, male
 detumescence of, 650
 discharge from, 658
 erection of, 650
 inspection of, 668t–671t
 palpation of, 672t, 676
- Penlight, 59t
- Peptic ulcer disease, 579t
- Perception sharing, in patient interview, 10
- Percussion, 20, 69–70, 70f
 abdominal, 588, 588f, 589t–590t
 bladder, 590t–591t
 chest, 422f, 423, 423t–424t
 direct (immediate), 69–70, 70f
 fist (blunt), 70, 71f
 frontal sinus, 293t
 indirect (mediate), 70, 71f
 kidney, 588, 591t
 liver, 588, 589t–590t
 paranasal sinus, 292, 293t
 precordium, 476t, 478
 sounds on, 70t
 splenic, 588, 588f, 590t
- Perfusion-ventilation problems, 414t
- Pericardial friction rub, 449–450, 451f, 479t
- Pericarditis, 488t
 constrictive, 472f
 temperature in, 465
- Pericardium, 438f, 440t
- Perimenopause, 624t, 796–797. *See also* Menopause
- Perineum, 612f, 614t, 631t–632t, 633t
- Periodontitis, 299t, 317t
- Periosteum, 686, 688f
- Peripheral arterial disease, 519, 521t–522t.
See also Peripheral-vascular system, assessment of
- Peripheral nervous system, 745–746, 746t, 747f. *See also* Neurologic system
- Peripheral-vascular system, 491–524
 abnormalities of, 521t–523t
 anatomy of, 492, 492f, 493f, 493t
 assessment of, 504–518
 Allen test in, 517b
 anatomical landmarks in, 504
 ankle-brachial index in, 518b
 approach to, 504
 auscultation in, 515, 516t, 517b
 biographical data in, 497
 cardiovascular system in, 501t, 506t
 case study of, 497, 503–504, 506, 508
 checklist for, 523–524
 color change test in, 518b
 cultural considerations in, 495, 497
 current health status in, 497–498, 499t–500t
 developmental considerations in, 494–495
 edema in, 498
 endocrine system in, 501t
 family history in, 500, 501t
 fatigue in, 498
 focused history in, 497b
 gastrointestinal system in, 501t, 506t
 general survey in, 501t, 505, 505t
 genitourinary system in, 501t
 head-to-toe, 505, 505t–506t
 health history in, 497–503, 497b
 HEENT system in, 501t, 505t
 inspection in, 508t–509t, 510b
 integumentary system in, 501t, 505t
 limb pain in, 498
 manual compression test in, 518b
 musculoskeletal system in, 501t, 506t
 neurological system in, 501t, 506t
 nursing diagnoses in, 519
 palpation in, 510, 510b, 511t, 515
 paresthesias in, 498
 past health history in, 498, 498t–499t
 physical, 504–518
 psychosocial profile in, 501, 502t–503t
 reproductive system in, 501t
 respiratory system in, 501t, 506t
 review of systems in, 501, 501t
 symptom analysis in, 498
 Trendelenburg test in, 518b
 atherosclerosis of, 519, 521t–522t
 body systems interaction with, 494
 in infants, 494
 in older adults, 495
 in pregnancy, 495
- Peristalsis, 581
- Peritoneum, 558
- Peritonitis, 607t
- Person, orientation to, 770
- Personal space, 8
 culture and, 9
- Personality disorders, 140t
- Pes cavus, 728t
- Pes planus, 728t
- Petechiae, 231t
- Peyer's patches, 495b
- Phalanges, 687f
- Phalen's test, 716, 717f
- Phantom limb pain, 93
- Pharyngeal wall, posterior, 269f
- Pharyngitis, 317t
- Pharyngopalatine arch, 269f
- Pharynx, 272t, 395t, 556t
- Phencyclidine, abuse of, 136t–137t
- Phenothiazines
 gastrointestinal system effects of, 569t
 integumentary system effects of, 216t
 musculoskeletal system effects of, 697t
- Phenylketonuria, 874t
- Phenylpropanolamine, sleep and, 159b
- Phimosis, 670t, 682t
- Phobias, 139t
- Physical assessment, 20, 53–86. *See also* *specific systems and patient populations*
 anthropometric measurements for, 83–84, 84f
 approach to, 72–73
 auscultation for, 71–72, 72b. *See also* Auscultation
 of children, 73
 complete, 54–55, 54t
 components of, 54t, 74–85
 body type, 74
 cultural considerations, 78
 developmental considerations, 78
 distress signs, 74
 dress, 77–78
 facial characteristics, 74
 gait, 74
 general survey, 74, 77–78
 hygiene, 77–78
 mental state, 78
 posture, 74, 77
 speech, 77
 of disabled patient, 73–74
 documentation of, 85
 environment for, 72–73
 equipment for, 55, 56t–66t
 focused, 54, 54t, 55
 head-to-toe scan for, 84–85. *See also* Head-to-toe physical assessment
 height for, 83–84, 84f
 inspection for, 67, 67f. *See also* Inspection medical, 55
 nursing, 55
 in nutrition assessment, 177–182, 178t–179t
 of older adults, 73
 in pain assessment, 102–107, 104t
 palpation for, 67–69, 68f, 70f. *See also* Palpation
 percussion for, 69–70, 70f, 70t, 71f. *See also* Percussion
 positions for, 74, 75t–77t
 of pregnant patient, 73
 purpose of, 54, 55b
 in teaching assessment, 149–151, 150t–151t
 vital signs for, 78–83
 blood pressure. *See* Blood pressure
 documentation of, 83
 pulse, 80–81, 80f, 80t, 81f
 respirations, 81
 temperature. *See* Body temperature
 weight for, 83–84, 84f
- Physician, nurse relationship to, 14, 16, 16b
- PIE documentation method, 24
- Pie wedge method, of breast palpation, 543, 543f
- Pinguecula, 349t, 361t

- Pins, 64t
 Pituitary gland, in pregnancy, 818t
 Pityriasis rosea, 244t, 260t
 Place, orientation to, 769–770
 Planning, 4f, 5
 Plantar reflex, 745t, 786t, 869t
 Pleura, 394, 394f, 395t
 Pleural effusion, 433t
 Pneumonia, 433t
 Pneumotaxic center, 396
 Pneumothorax, 433t
 Pocket light, 59t
 Pocket ruler, 61t
 Pocket vision screener, 58t
 Point localization, 782t
 Point of maximum impulse, 474
 Poisoning, in children, 889
 Poker Chip Tool, for pain, 100
 Polish Americans, integumentary system of, 211
 Polycythemia, 414t
 Polyneuropathy, 93
 Polyps, nasal, 295t, 317t
 Pons, 741t
 Popliteal pulse, 492f, 493t, 513t
 Port-wine stain, 211, 232t, 860t
 Portal vein, 493f, 493t
 Position(s), 74, 75t–77t
 for breast assessment, 539, 544t–545t
 for cardiovascular assessment, 466, 466f
 dorsal recumbent, 75t
 for female genitourinary system assessment, 629
 of fetus, 837t–838t
 knee-chest, 76t
 left lateral recumbent, 76t
 prone, 76t
 Sims', 75t
 sitting, 75t
 standing, 77t
 supine, 75t
 Position sense, examination for, 779t–780t
 Post-traumatic syndrome, 139t
 Posterior cruciate ligament, 729
 Posterior tibialis pulse, 492f, 493t, 513t
 Posterior triangle, 286, 286f
 Postoperative Pain Scale, 102–103
 Postpartal examination, 643t–645t, 843, 843t–845t
 Postpartum depression, 116, 116b, 845b
 Posture, 8, 74, 77, 706t–708t
 culture and, 10
 in newborns, 856t
 Potassium hydroxide, 66t
 PQRST, 31, 33
 Precordial impulses, 475t–476t
 Precordium
 auscultation of, 478–485, 478t, 481t–482t, 485b
 inspection of, 471–472, 471t–472t, 474–475
 palpation of, 474–475, 474t–475t
 percussion of, 478, 476t
 Prednisone, nutrition and, 172t
 Preeclampsia, 495
 Pregnancy, 813–847
 abdomen in, 561
 in adolescents, 818–819, 913
 age effects on, 819
 assessment of, 73, 823–845
 abdomen in, 836t–839t
 approach to, 832
 biographical data in, 824
 blood tests in, 833b
 breast in, 828t, 835t–836t
 cardiovascular system in, 828t, 835t
 case study of, 823, 831–832, 842, 843, 845
 cultural considerations in, 819t–823t
 current health status in, 824–825
 deep tendon reflexes in, 841t
 documentation for, 824–825
 endocrine system in, 830t
 estimated date of birth in, 825
 family history in, 826–827
 focused history in, 824b
 gastrointestinal system in, 829t, 836t–839t
 general survey in, 827t, 832–833, 833f
 genitourinary system in, 829t, 840t
 head-to-toe, 833, 834t–841t
 health history in, 823–831
 HEENT system in, 827t–828t, 834t–835t
 hematologic system in, 830t
 immunologic system in, 830t
 integumentary system in, 827t, 834t
 laboratory tests in, 833b
 musculoskeletal assessment in, 839t, 841t
 musculoskeletal system in, 829t
 Naegle's rule in, 825
 neurologic system in, 829t–830t, 841t
 nursing diagnoses in, 842
 past health history in, 825–826
 physical, 832–845
 postpartal, 843, 843t–845t
 psychosocial profile in, 827, 830–831
 reproductive system in, 829t
 respiratory system in, 828t, 835t
 review of systems in, 827, 827t–830t
 symptoms in, 825, 825t
 urine tests in, 824–825, 833b
 back pain in, 693
 blood pressure in, 495
 breasts in, 532, 816t
 cardiac disease in, 826
 cardiovascular system in, 453, 815t–816t
 cervical cancer in, 826
 cervix in, 817t, 840t
 complications of, 846
 diabetes in, 826
 edema in, 839t
 endocrine system in, 818t
 ethnicity and, 819t–823t
 face in, 275
 fetal heart tones in, 837t
 fetal position in, 837t–838t
 fetal presentation in, 838t
 folic acid supplement in, 751
 fundal height in, 836t
 gait in, 693
 gastrointestinal system in, 816t, 826
 genitourinary system in, 817t
 HEENT system in, 815t
 hematologic system in, 818t
 Homans' sign in, 839t
 hypertension in, 826
 immunologic system in, 494–495, 818t
 infectious diseases in, 826
 integumentary system in, 211, 814t–815t
 Leopold's maneuver in, 838t
 liver disease in, 826
 lordosis in, 693, 707t
 mask of, 211, 228t, 275, 814t, 834t
 mental disorders and, 116, 116b
 mouth in, 275
 musculoskeletal system in, 817t
 neck in, 275
 neurologic system in, 751, 817t
 nutrition during, 169, 169t, 845–846
 palmar erythema in, 495, 834t
 peripheral-vascular system in, 495
 physiologic changes of, 814, 814t–818t
 pulmonary disease in, 826
 reproductive organ changes with, 615, 617
 respiratory system in, 398, 815t
 sexually transmitted diseases in, 826
 telangiectasis in, 495
 trimesters of, 814
 vaginitis in, 826
 Prehypertension, 81, 516t
 Preload, 444, 475t–476t
 Premature Infant Pain Profile, 105
 Premenstrual syndrome, 624t
 Prepatellar bursa, 686
 Presbycusis, 369, 389t
 Presbyopia, 330, 361t
 Preschoolers, 887–899
 abuse of, 889, 889b
 activity in, 892
 assessment of, 889–897
 biographical data in, 890
 cardiovascular system in, 891t, 895t–896t
 case study of, 889b, 897t
 current health status in, 890
 family history in, 890
 gastrointestinal system in, 891t, 896t
 general survey in, 890t, 893, 893f, 893t
 genitourinary system in, 891t, 896t
 health history in, 889–890
 HEENT system in, 890t–891t, 894t–895t
 hematologic system in, 892t
 integumentary system in, 890t, 894t
 lymphatic system in, 892t
 musculoskeletal system in, 891t, 896t
 neurologic system in, 892t, 897t
 past health history in, 890
 physical, 893, 893t–897t
 psychosocial profile in, 892–893
 respiratory system in, 891t, 895t
 review of systems in, 890, 890t–892t
 communicable diseases in, 889
 developmental changes in, 888
 emotional changes in, 888
 exercise in, 159–160
 fears in, 889
 health concerns in, 897t
 injury in, 161, 889
 motor vehicle injury in, 889
 negativism in, 888
 neglect of, 889
 nutrition in, 168t, 892
 obesity in, 897t
 physical changes in, 888

- Preschoolers (*continued*)
 physiological anorexia in, 892
 poisoning in, 889
 psychosocial changes in, 888
 sexuality in, 893
 sibling rivalry in, 888, 893
 sleep in, 159, 892
 spirituality in, 189
 stress in, 160, 889, 893
 temper tantrums in, 888
 toilet training for, 888
 visual acuity assessment in, 330
 wellness assessment in, 155
- Pressure equalization tube, 381t
- Pressure ulcers, 230, 238b, 239t
- Preventive healthcare, 17
- Priapism, 670t, 682t
- Primary preventive healthcare, 17
- Primitive reflexes, 783–784
- Problem, collaborative, 5, 5b
- Problem-oriented medical records, 23
- Problem-solving skills, for assessment, 12–13
- Progesterone, breast effects of, 530
- Progestins
 breast effects of, 535t
 female genitourinary system effects of, 622t
- Prolactin, 530
- Pronation (body movement), 715f, 716t
- Pronator drift test, 728, 728f
- Prostate gland, 650, 651f, 652t
 benign hypertrophy of, 653, 680t
 cancer of, 679, 682t
 examination of, 675t, 676–677, 677f
 palpation of, 675t
- Prostate-specific antigen, 679, 959t
- Prostatitis, 682t
- Protein, 164
- Protein-calorie malnutrition, 185t
- Protraction (body movement), 715f, 716t
- Pruritus, 209, 213, 237t
 senile, 211
- Pseudofolliculitis, 211
- Psoriasis, 209, 233t, 236t, 245t, 261t
- Psychodynamic theory, 114
- Psychological state, health and, 154
- Psychomotor skills, for assessment, 13
- Psychosocial profile, 28t, 29, 44, 44t–47t, 916
 in abdominal assessment, 573, 573t–574t
 in adolescent assessment, 916, 919–920
 in breast assessment, 537, 537t–538t
 in cardiovascular system assessment, 460, 460t–462t
 in ear assessment, 373, 373t–374t
 in eye assessment, 335, 335t–336t
 in female genitourinary system assessment, 625, 625t–626t
 in head, face, and neck assessment, 283, 283t–284t
 in integumentary system assessment, 219, 219t–220t
 in lymphatic system assessment, 502, 502t–503t
 in male genitourinary system assessment, 662, 662t–663t
 in menopause assessment, 799
 in mental health disorders, 124, 125t–126t
 in musculoskeletal system assessment, 699, 699t–700t
 in neurologic system assessment, 758, 758t–759t
 in nutrition assessment, 172, 173t–175t
 in older adult assessment, 942–944
 in pain assessment, 100, 101t–102t
 in peripheral-vascular system assessment, 502, 502t–503t
 in pregnancy assessment, 827, 830–831
 in respiratory system assessment, 403, 404t–407t
 in school-age child assessment, 906–908
 in spirituality assessment, 193
 in teaching assessment, 149, 149t–150t
 in toddler and preschooler assessment, 892–893
- Psychotic disorders, 140t–141t
- Pterygium, 349t, 361t, 950t
- Ptosis, 348t
- Puberty. *See also* Adolescents
 female, 617b, 914
 male, 654, 656t, 914
- Pubic hair
 female, 617b
 male, 654, 656t, 668t
- Puddle sign, 596
- Pull-to-sit reflex, 873t
- Pulmonary artery, 438f, 439t
- Pulmonary disease, in pregnancy, 826
- Pulmonary fibrosis, 433t–434t
- Pulmonary semilunar valve, 438f, 439t
- Pulmonary vein, 438f, 439t
- Pulse(s), 80–81, 80f, 80t, 81f, 492f, 493t
 abnormal, 477f
 absent, 511t
 apical, 81f
 brachial, 492f, 493t, 511t
 in cardiovascular system assessment, 465
 carotid, 492f, 493t, 510t
 Corrigan's (water-hammer), 477f
 dorsalis pedis, 492f, 493t, 513t
 femoral, 492f, 493t, 512t
 in newborns, 864t
 in infants, 881t
 large, bounding, 477f
 in newborns, 858t, 864t
 popliteal, 492f, 493t, 513t
 posterior tibialis, 492f, 493t, 513t
 radial, 80, 80f, 492f, 493t, 511t
 sites for, 492f, 493t
 small, weak, 477f
 temporal, 492f, 510t
 ulnar, 492f, 493t, 511t
 unequal, 511t
- Pulse pressure, 81
- Pulsus alternans, 477f
- Pulsus bigeminus, 477f
- Pulsus bisferiens, 477f
- Pulsus paradoxus, 477f
- Punctum, disorders of, 330
- Pupil, 322. *See also* Eye(s)
 accommodation testing of, 353t
 Adie's, 352t
 anatomy of, 322f, 323f
 Argyll Robertson, 353t
 inspection of, 351t–352t
 oval, 353t
 tonic, 352t
- Pupillary light reflex, 329, 346, 352t
- Purkinje fibers, 444f, 445
- Purpura, 231t
 senile, 949t, 956t
- Pursed-lip breathing, 413t
- Pustule, 235t
- Pyrosis, 566
- Q**
- Questions
 closed, 18
 open, 10, 18
- Quick Confusion Scale, 787–788, 788b
- R**
- Radial deviation, 716t
- Radial pulse, 80, 492f, 493t, 511t
- Radius, 687f
- Rales, 425t, 427, 428t, 429
- RAMS test, 710t
- Range of motion
 ankle, 725t–726t
 arm, 721t
 cervical spine, 718t–719t
 elbow, 721t
 finger, 722t–723t
 hand, 722t–723t
 hip, 723t–724t
 knee, 725t
 lumbar spine, 719t–720t
 nonperformance of, 726t
 shoulder, 720t
 thoracic spine, 719t–720t
 wrist, 722t
- Ranvier, nodes of, 739f
- Rash, 213
- Raynaud's phenomenon/disease, 523t
- Reality presentation, in patient interview, 11
- Reasoning, 12b
- Rebound tenderness, abdominal, 601t
- Receptive aphasia, 766b
- Rectal examination
 in female genitourinary system assessment, 642, 642t
 in male genitourinary system assessment, 675t, 676–677, 677f
- Rectocele, 632t
- Rectovaginal examination, 642, 642t
- Rectum, 558f, 612f, 651f
 of infant, 882t
 inspection of, 671t
 of newborn, 865t
 palpation of, 675t
- Rectus abdominis, 687f
- Rectus femoris, 687f
- Rectus muscles (ocular), 323f, 324t–325t
- Red marrow, 686
- Red reflex, 329, 356t–357t
- Redirecting, in patient interview, 10
- Referred pain, 564
- Reflection, 12b
 in patient interview, 10
- Reflex(es), 782–784, 783f, 784f, 784t–786t
 abdominal, 596, 597t–598t, 745t, 786t
 Achilles, 745t, 785t

- anal, 745t, 786t
- Babinski, 784, 786t, 869t–870t
- biceps, 745t, 784t
- blink, 329, 350t
- brachioradialis, 745t, 785t
- bulbocavernosus, 745t, 786t
- cognitive inhibition of, 783
- corneal, 350t, 773t–774t
- corneal light, 329–330, 345t
- crawling, 872t
- cremasteric, 745t, 786t
- crossed extension, 872t
- deep tendon, 744, 745t, 782–783, 783f, 784t–786t
- in pregnancy, 841t
- extrusion, 871t
- flexor (withdrawal), 745
- glabellar, 784, 872t
- grasp, 783, 869t
- isometric maneuvers for, 783, 784f
- magnet, 873t
- Moro, 868t
- newborn/infant, 867, 868t–873t
- oculocephalic, 772t–773t
- oculovestibular, 775t
- in older adults, 751, 956t
- palmar, 783, 869t
- patellar, 744–745, 745t, 785t
- in newborns, 867t
- plantar, 745t, 786t, 869t
- primitive, 783–784
- pull-to-sit, 873t
- pupillary, 329, 346, 352t
- red, 329, 356t–357t
- rooting, 784, 870t
- snout, 784
- spinal, 744–745, 744f, 745t
- startle, 868t
- stepping, 870t
- sucking, 784, 871t
- superficial, 745t, 783, 786t
- swallowing, 871t
- tonic neck, 868t–869t
- triceps, 745t, 785t
- trunk incurvation, 873t
- Reflex arc, 744, 744f
- Reflexive thinking, 12
- Reflux, abdominojugular (hepatojugular), 473t, 476
- Refractoriness, 444
- Religions, 194, 195t–198t. *See also* Spirituality
- Renal artery, auscultation of, 587t
- Reposition (body movement), 715f, 716t
- Reproductive system. *See also* Genitourinary system
- female
 - age-related changes in, 930t
 - anatomy of, 612–613, 612f, 613f
 - assessment of, 38t
 - in adolescents, 918t, 922t–923t
 - in breast disorders, 537t, 541t, 542
 - in cardiovascular disorders, 459t
 - in ear disorders, 373t
 - in eye disorders, 334t, 339t
 - in gastrointestinal disorders, 572t, 581t, 582
 - in head, face, and neck disorders, 282t, 313
 - in integumentary disorders, 218t, 224t, 225
 - in lymphatic disorders, 501t, 507
 - in musculoskeletal disorders, 698t, 704
 - in neurologic disorders, 758t
 - in nutritional disorders, 173t
 - in older adults, 939t, 953t–954t
 - in peripheral-vascular disorders, 501t
 - in pregnancy, 829t. *See also* Pregnancy
 - in respiratory disorders, 404t, 419
 - in school-age children, 911t
 - breast interaction with, 530, 531
 - cardiovascular system interaction with, 452
 - gastrointestinal system interaction with, 560
 - musculoskeletal system interaction with, 692
 - neurologic system interaction with, 750
 - process of, 613, 613f
 - respiratory system interaction with, 397
- male
 - age-related changes in, 930t
 - anatomy of, 650, 651f, 652t–653t
 - assessment of, 38t
 - in adolescents, 918t, 922t–923t
 - in cardiovascular disorders, 460t
 - in eye disorders, 334t, 339t
 - in gastrointestinal disorders, 572t, 581t, 582
 - in head, face, and neck disorders, 282t, 313
 - in integumentary disorders, 218t, 224t, 225
 - in lymphatic disorders, 501t, 507
 - in musculoskeletal disorders, 698t, 704
 - in neurologic disorders, 758t
 - in nutritional disorders, 173t
 - in older adults, 939t, 953t–954t
 - in peripheral-vascular disorders, 501t
 - in respiratory disorders, 404t, 419
 - in school-age children, 911t
 - cardiovascular system interaction with, 452
 - drug effects on, 659t–660t
 - endocrine system interaction with, 653, 655
 - function of, 650–651, 654b
 - musculoskeletal system interaction with, 653, 655, 692
 - neurologic system interaction with, 750
 - neurological system interaction with, 653, 655
 - respiratory system interaction with, 397
 - urinary system interaction with, 651, 653, 655
- Reserpine, breast effects of, 536t
- Resonance, 423
- Respect, 11b
- Respiration, 394, 412
 - apneustic, 415b
 - assessment of, 80t, 81
- Biot's, 412, 415b
- Cheyne-Stokes, 412, 415b
- in infants, 398
- Kussmaul, 412, 415b
- in newborns, 858t
- in older adults, 398
- in pregnant women, 398
- shallow, 412
- Respiratory rate, 80t, 81, 396, 412, 466
- Respiratory system, 393–436
- abnormalities of, 432t–434t
 - acid-base balance and, 395–396
 - age-related changes in, 929t
 - anatomical landmarks of, 409, 409f, 410f, 411f, 412f
 - anatomy of, 394, 394f, 395t, 398f
 - assessment of, 38t, 399–434
 - in adolescents, 917t, 921t
 - anatomical landmarks in, 409, 409f, 410f, 411f, 412f
 - auscultation in, 425–430, 425t–426t
 - biographical data in, 399
 - in breast disorders, 537t, 541t, 542
 - breath sounds in, 425–429, 425t–426t, 427f, 427t, 428t–429t
 - in cardiovascular disorders, 459t, 467, 469t
 - cardiovascular system in, 404t, 414t, 419
 - case study of, 399, 407–408, 430
 - checklist for, 434–435
 - culture and, 398–399
 - current health status in, 399
 - developmental considerations in, 398
 - diaphragmatic excursion in, 423
 - drugs in, 402t–403t
 - in ear disorders, 373t, 377t, 386
 - endocrine system in, 419
 - in eye disorders, 334t, 339t, 340
 - family history in, 400, 402t
 - in female genitourinary disorders, 623t, 630t, 635
 - focused history in, 399b
 - in gastrointestinal disorders, 572t, 580t, 582
 - gastrointestinal system in, 404t, 414t, 419
 - general survey in, 403t, 411–412, 413t, 419
 - genitourinary system in, 404t, 419
 - in head, face, and neck disorders, 281t, 288t, 313
 - head-to-toe, 412, 413t–414t
 - health history in, 399–407
 - HEENT system in, 404t, 413t–414t, 419
 - hematologic system in, 404t, 419
 - immunologic system in, 404t
 - in infants, 877t, 881t
 - inspection in, 416, 416f, 416t–418t
 - in integumentary disorders, 217t, 224t, 225
 - integumentary system in, 403t, 413t, 419
 - in lymphatic disorders, 501t, 506t, 507
 - lymphatic system in, 419
 - in male genitourinary disorders, 661t, 666
 - for menopause, 804t
 - in mental health disorders, 123t, 127t
 - in musculoskeletal disorders, 698t, 703t, 704
 - musculoskeletal system in, 404t, 414t, 419
 - in neurologic disorders, 757t, 764t, 765

- Respiratory system (*continued*)
- neurologic system in, 404t, 414t, 419
 - in newborns, 855, 858t
 - nursing diagnoses in, 430
 - in nutritional disorders, 179t
 - in older adults, 937t, 952t
 - in pain disorders, 104t
 - palpation in, 418, 420t–422t
 - past health history in, 400, 401t
 - in patient teaching, 148t, 150t
 - percussion in, 422f, 423, 423t–424t
 - in peripheral-vascular disorders, 501t, 506t
 - physical, 410–430
 - in pregnancy, 828t, 835t
 - psychosocial profile in, 403, 404t–407t
 - reproductive system in, 404t
 - respiratory patterns in, 412, 415b
 - respiratory rate in, 412
 - review of systems in, 403t–404t
 - in school-age children, 905t, 910t
 - symptom analysis in, 400
 - temperature in, 412
 - in toddlers and preschoolers, 891t, 895t
 - body systems interaction with, 396–397
 - cardiovascular system interaction with, 396, 397, 452, 453
 - drug effects on, 403t–403t
 - dysfunction of, 431
 - ear interaction with, 368
 - eye interaction with, 327
 - female genitourinary system interaction with, 616
 - gastrointestinal system interaction with, 559, 560
 - head, face, and neck interaction with, 269–270, 273
 - in infants, 398
 - integumentary system interaction with, 208, 208f, 210
 - male genitourinary system interaction with, 655
 - musculoskeletal system interaction with, 396, 397, 692
 - neurologic system interaction with, 396, 397, 748, 750
 - in newborns, 851t
 - in older adults, 398
 - in pregnant women, 398, 815t
- Restating, in patient interview, 10
- Reticular activating system, 741t
- Reticulospinal tract, 743–744, 744t
- Retina, 323f, 325t, 326. *See also* Eye(s)
- Retinal detachment, 361t
- Retinitis pigmentosa, 361t
- Retraction (body movement), 715f, 716t
- Review of systems, 28t, 36, 36t–39t
 - in abdominal assessment, 571, 571t–573t
 - in adolescent assessment, 916, 917t–918t
 - in breast assessment, 536, 536t–537t
 - in eye assessment, 334, 334t–335t
 - in female genitourinary system assessment, 622, 622t–625t
 - in head, face, and neck assessment, 280, 281t–282t
 - in infant assessment, 876, 877t–878t
 - in integumentary system assessment, 216, 217t–218t
 - in lymphatic system assessment, 501, 501t
 - in male genitourinary system assessment, 660, 661t–662t
 - in mental health disorders, 122, 123t–124t
 - in musculoskeletal system assessment, 697, 697t–698t
 - in neurologic system assessment, 756, 757t–758t
 - in nutrition assessment, 172, 173t
 - in older adult assessment, 935, 935t–940t
 - in pain assessment, 100
 - in peripheral-vascular system assessment, 501, 501t
 - in pregnancy assessment, 827, 827t–830t
 - in respiratory system assessment, 403t–404t
 - in school-age child assessment, 904, 905t–906t
 - in teaching assessment, 148–149, 148t
 - in toddler and preschooler assessment, 890, 890t–891t
- Rheumatoid arthritis, 732
- Rh(D) incompatibility, screening for, 157t
- Rhinitis
 - allergic, 316t
 - viral, 316t
- Rhonchi, 426t, 428t
- Ribs, 687f
- Right lymphatic duct, 492
- Right ventricular hypertrophy, 472f
- Riley Infant Pain Scale, 105t, 106
- Ringworm, 241t, 263t
- Rinne test, 383, 384t–385t
- Risk-taking behavior, in adolescents, 115, 115b, 919
- Rods, 326
- Roman Catholic Church, 195t
- Romberg sign, 956t
- Romberg test, 385t, 705, 710t
- Rooting reflex, 870t
- Rosacea, 261t
- Rotation (body movement), 715f, 716t
- Round window, 366f
- Rovsing's sign, 600t
- Roy's adaptation theory, 21
- Rubella, 240t
- Rubrospinal tract, 743–744, 744t
- S**
- Sacrum, 687f
- Safety pin, 64t
- Sagittal suture, 268f, 271t
- Saline mount, 637, 638t
- Salivary glands, 270f, 272t, 557t
 - palpation of, 296, 306t
- Sartorius, 687f
- Scale (weight), 62t
- Scalenus, 398f
- Scales (cutaneous), 236t
- Scales (pain), 98–100, 100f, 102–107, 105t, 106t
- Scalp
 - inspection of, 254t
 - palpation of, 254, 255t
- Scapula, 687f
- Scar, surgical, 236t
- Schizoid personality disorder, 140t
- Schizophrenia, 140t–141t
- Schizotypal personality disorder, 140t
- Schlemm's canal, 325t
- School-age children, 901–913
 - assessment of, 904–911
 - biographical data in, 904
 - cardiovascular system in, 906t, 910t
 - case study of, 904, 911
 - cultural considerations in, 903, 903t–904t
 - current health status in, 904
 - family history in, 904
 - gastrointestinal system in, 906t, 911t
 - general survey in, 905t, 908t
 - genitourinary system in, 906t, 911t
 - head-to-toe assessment in, 908, 908t–911t
 - health history in, 904
 - HEENT system in, 905t, 909t–910t
 - hematologic system in, 906t
 - integumentary system in, 905t, 908t–909t
 - lymphatic system in, 906t
 - musculoskeletal system in, 906t, 911t
 - neurologic system in, 906t, 911t
 - past health history in, 904
 - psychosocial profile in, 906–908
 - reproductive system in, 911t
 - respiratory system in, 905t, 910t
 - review of systems in, 904, 905t–906t
 - behavior problems in, 902
 - developmental changes in, 902–903
 - eating disorders in, 902
 - emotional changes in, 903
 - emotional problems in, 902
 - exercise for, 160, 907
 - eye examination in, 360
 - health concerns in, 912
 - homeless, 912
 - immunizations for, 912
 - injury in, 161, 902
 - learning problems in, 902, 907
 - nutrition for, 167t, 168t, 907
 - physical changes in, 903
 - psychosocial changes in, 903
 - relationships of, 907
 - self-concept of, 907
 - sexuality of, 907
 - sleep in, 159, 907
 - spirituality in, 189
 - stress in, 160–161, 907–908
 - substance abuse in, 902
 - visual acuity assessment in, 330
 - wellness assessment in, 157
- Scientific method, 13
- Sclera, 323f, 324t
 - color of, 330
 - inspection of, 350t
 - muddy, 330, 350t
- Scleroderma, 228t, 732
- Scoliosis, 706t, 708t
 - in adolescents, 691, 923t
- Scraper, 65t
- Scratch test, 588, 589t
- Screening tests
 - in infants, 883
 - in newborns, 867, 874t
 - preventive, 156t–157t
- Scrotum, 650, 651f, 652t

- auscultation of, 677, 678t
 - inspection of, 670t
 - palpation of, 673t, 676
 - self-examination of, 676b
 - Season, skin disorders and, 213, 213t
 - Sebaceous cyst, scrotal, 670t
 - Sebaceous glands, 206f, 207t, 208, 211
 - Seborrhea, 261t
 - Seborrheic keratosis, 211, 949t, 956t
 - Sebum, 211
 - Secondary preventive healthcare, 17
 - Sedatives
 - abuse of, 135t
 - male sexual function effects of, 660t
 - neurologic system effects of, 756t
 - respiratory system effects of, 403t
 - Selective serotonin reuptake inhibitors, side effects of, 123t
 - Self, therapeutic use of, 18
 - Self-care, in older adults, 943–944, 944b–94bb, 945t
 - Self-concept
 - in adolescents, 919
 - in school-age children, 907
 - Semicircular canal, 366f
 - Seminal vesicle, 650, 651f, 652t
 - Senile cataract, 330
 - Senile ectasias, 957t
 - Senile purpura, 949t, 956t
 - Sensory cortex, 739, 741–742, 742f, 742t
 - Sensory function. *See also* Neurologic system
 - assessment of, 777, 777t–782t
 - extinction test for, 782t
 - graphesthesia test for, 777, 781t
 - kinesthetics test for, 779t–780t
 - light touch test for, 777t–778t
 - pain tests for, 778t–779t
 - point localization for, 782t
 - stereognosis test for, 777, 780t
 - two-point discrimination for, 781t
 - vibration test for, 779t
 - Sensory pathways, 739, 741–742, 742f, 742t
 - Septum, nasal, 271t
 - deviated, 295t
 - Serotonin, 112b
 - Serous otitis media, 381t, 389t
 - Sexual history
 - in female genitourinary system assessment, 627
 - in male genitourinary system assessment, 663–664
 - Sexuality
 - in adolescents, 919–920
 - in older adults, 939t
 - in school-age children, 907
 - in toddlers and preschoolers, 893
 - Sexually transmitted diseases, 644, 644t, 645t, 658b, 664
 - in adolescents, 913
 - in pregnancy, 826
 - Shoulder
 - range of motion of, 720t
 - referred pain to, 597, 598t
 - strength of, 720t
 - Sibling rivalry, 888
 - Sickle cell anemia, 874t
 - Signs, 17
 - Silence, in patient interview, 10
 - Sinoatrial (SA) node, 444–445, 444f
 - Sinusitis, 316t
 - Situational crisis, 131
 - Sjögren's syndrome, 361t
 - Skene's ducts, 612f
 - Skene's gland, 614t, 633t
 - Skin, 206, 206f, 207t. *See also* Integumentary system
 - abnormalities of, 257t–263t
 - age-related changes in, 928t, 956t–957t
 - age-related disorders of, 213t
 - atrophy of, 238t
 - cancer of, 212–213, 256, 258t, 262t
 - color of, 226, 226t–227t, 228t–229t
 - lesions of, 227t–228t, 230, 230f
 - annular/circular, 241t
 - arciform, 241t
 - clinical description of, 231t
 - color of, 231t
 - confluent, 240t
 - diffuse/generalized, 243t
 - discrete, 240t
 - distribution of, 243t–246t
 - exudative, 231t
 - genital, 658
 - grouped, 240t
 - gyrate, 242t
 - intertriginous, 246t
 - iris-shaped, 241t
 - linear, 241t
 - localized, 244t
 - pain and, 231t
 - pattern of, 240t–242t
 - polycyclic, 242t
 - primary, 230, 233t–235t
 - regional, 244t–246t
 - reticular, 242t
 - round/oval, 240t
 - scattered, 243t
 - season and, 213t
 - secondary, 230, 235t–238t
 - shape of, 231t
 - size of, 231t
 - in sun-exposed areas, 246t
 - surface characteristics of, 231t, 233t–235t
 - texture of, 231t
 - vaginal, 619
 - vascular, 231t–232t
 - moisture of, 247, 247t, 249
 - season-related disorders of, 213t
 - temperature of, 247t, 514t
 - texture of, 248t
 - turgor of, 247, 248t, 249
- Skinfold calipers, 62t
- Skull, 268f, 271t
- Sleep
 - in adolescents, 919
 - assessment of, 158–159, 159
 - in newborns, 855
 - in older adults, 942–943, 943b
 - in school-age children, 907
 - in toddlers and preschoolers, 892
- Sleep latency, 158
- Small intestine, 556f, 557t
- Smegma, 676
- Smooth muscle relaxants, genitourinary
 - eye effects of, 333t
 - head, face, and neck effects of, 279t
- Snellen eye chart, 58t, 330, 341, 342t–343t
- SOAPIE documentation method, 23
- Social anxiety disorder, 139t
- Social theory, 114
- Sodium bicarbonate, nutrition and, 172t
- Soft palate, 269f, 272t, 301t
- Somatosensory cortex, 89
- Somnambulism, 158
- Sore throat, 276
- Sound, 365–366
 - conduction of, 366, 367f
- Spasticity, 709t
- Speculum, 66t
- Speculum examination, of female genitalia, 634, 636t–637t
- Speech, in physical assessment, 77
- Sperm, 650–651, 654b
- Spermatic cord, 559t, 652t
- Spermatocoele, 670t, 682t
- Spermatogenesis, 650–651, 654b
- Sphenoid bone, 268f
- Sphenoid sinus, 268f, 269f
- Sphygmomanometer, 58t
- Spider angioma, 232t
- Spinal cord, 739. *See also* Neurologic system
 - cross-section of, 739
 - roots of, 745–746
 - tumor of, 792t
- Spinal nerve roots, 745–746
- Spinal nerves, 745–746, 748f
- Spinal reflexes, 744–745, 744f, 745t
- Spine, 559t
 - cervical, 718t–719t
 - lumbar, 719t–720t
 - straight leg raising test for, 728, 728f
 - thoracic, 687f, 719t–720t
- Spinothalamic tracts, 742, 742f, 742t
- Spirituality, 187–202
 - assessment of, 190, 192–200, 195t–198t, 199t–200t
 - biographical data in, 192
 - current health status in, 192
 - family history in, 193
 - health history in, 192–193
 - Jarel Spiritual Well-Being Scale in, 199t–200t
 - nursing diagnoses in, 201
 - past health history in, 192–193
 - psychosocial profile in, 193
 - brief review of, 188
 - case study of, 190, 193–194, 200–201
 - culture and, 190, 191t–192t
 - definitions of, 201
 - development of, 188–190
 - perceptions of, 201
- Spirolactone
 - breast effects of, 536t
 - nutrition and, 172t
- Spleen, 495b, 556f, 559t
 - palpation of, 591, 593, 594t
 - rupture of, 607t
 - size of, 588, 588f, 590t
- Splenomegaly, 588, 588f, 590t
- Sprue, 228t

- Squamous cell carcinoma, 262t, 957t
 cutaneous, 212–213
 eyelid, 348t
 Squamous suture, 268f, 271t
 Stapes, 365t, 366, 366f, 367f
 Startle reflex, 868t
 Stasis ulcers, 211
 Stensen's duct, 272t
 assessment of, 301t–302t
 inspection of, 296
 Stepping reflex, 870t
 Stereognosis, 777, 780t
 Sternal angle, 464f, 471t
 Sternal rub, 769
 Sternoclavicular joint, 718t
 Sternocleidomastoid, 274f, 286, 286f, 398f, 687f
 Stethoscope, 56t–57t, 71–72, 72b
 for blood pressure, 82b–83b
 Stiff lung, 433t–434t
 Stiffness
 musculoskeletal, 695
 neck, 276
 Stimulants
 abuse of, 135t
 cardiovascular system effects of, 457t
 musculoskeletal system effects of, 697t
 Stomach, 556f, 556t–557t, 558f
 age-related changes in, 930t
 Stomach cancer, 605t
 Stomatitis
 allergic, 300t
 candidal, 316t
 Stork bite, 211, 859t
 Strabismus, 329, 361t
 Straight leg raising test, 728, 729f
 Stratum basale, 206f, 207t
 Stratum corneum, 206f, 207t
 Stratum germinativum, 206f, 207t
 Stratum granulosum, 206f, 207t
 Stratum lucidum, 206f, 207t
 Stratum spinosum, 206f, 207t
 Strawberry mark, 211
 Streptococcal pharyngitis, 317t
 Stress
 in adolescents, 920
 in children, 889
 management of, 160–161
 in school-age children, 902, 907–908
 sleep and, 159
 in toddlers and preschoolers, 893
 Striae, 581, 583t
 Striae gravidarum, 211, 561, 814t
 Stridor, 426t, 429
 Stroke, 762, 763b, 790t
 risk factors for, 789
 Stroke volume, 80, 444, 475t–476t
 Stycar chart, 58t
 Subacromial bursa, 686
 Subclavian vein, 493f, 493t
 Subconjunctival hemorrhage, 350t
 Subcutaneous tissue, 206, 206f
 Subdural hematoma, 790t
 Subjective findings, 17
 Sublingual gland, 270f, 272t, 557t
 palpation of, 306t
 Submandibular gland, 270f, 272t, 557t
 Substance abuse, 135t–137t
 in adolescents, 913
 assessment of, 133–137, 134b
 in school-age children, 902, 907
 Substance dependence, 133
 Substance intoxication, 134
 Substance withdrawal, 134
 Succussion splash, 586t
 Sucking reflex, 871t
 Sugar, 63t
 Suggesting, in patient interview, 10
 Suicide
 in adolescents, 115b
 assessment of, 132, 133t
 in older adults, 117b
 risk for, 132b, 133t
 Sulfasalazine, nutrition and, 172t
 Sulfonamides, integumentary system effects
 of, 216t
 Summarizing, in patient interview, 11
 Supination, 715f, 716t
 Support, basic, 12b
 Support systems, 154
 Supraorbital pressure, 769
 Surfer's ear, 380t
 Surgery
 in older adults, 935
 scar with, 236t
 Sutures, cranial, 268f, 271t
 Swallowing reflex, in newborns, 871t
 Swan-neck deformity, 734t
 Sweat glands, 206f, 207t, 208, 211
 Swimmer's ear, 380t
 Symmetry, in neurologic system assessment, 762
 Sympathetic nervous system, 453, 746, 749f
 Symphysis, 690t
 Symphysis pubis, 612f
 Symptoms, 17
 Synapse, 113f, 738, 739f
 Synarthrosis, 690t
 Synchondrosis, 690t
 Syncope, 456, 753
 Synovial joints, 689f, 690t, 691t
 Syphilis, 241t, 632t, 669t, 682t
 screening for, 157t
 Systemic lupus erythematosus, 732
 Systole, ventricular, 442–443
- T**
 Tachycardia, 80
 Tachypnea, 81, 412, 415b
 Tactile fremitus, 418, 421t–422t
 Tandem walking, 709t
 Tanning, 228t
 Tape measure, 61t
 Tarsal plate, 323f
 Tarsals, 687f
 Taste, in older adults, 275
 Tay-Sachs disease, hearing in, 372t
 Teaching, 145–151
 assessment for, 147–151
 biographical data in, 147
 current health status in, 147
 family history in, 148
 health history in, 147–149
 past health history in, 147
 physical assessment in, 149–151, 150t–151t
 psychosocial profile in, 149, 149t–150t
 review of systems in, 148–149, 148t
 case study of, 146, 149
 culture and, 147
 developmental level and, 146–147
 effective, 146b
 goals of, 146
 Tearing, 331
 Teeth, 271t
 age-related changes in, 929t
 deciduous, 269f, 274
 inspection of, 298t
 natal, 861t
 in older adults, 275
 permanent, 269f, 274
 in school-age children, 905t
 tetracycline effects on, 298t
 in toddlers and preschoolers, 891t
 Telangiectasia, 232t
 Telangiectasis, in pregnancy, 495
 Telogen, 214
 Temper tantrums, 888
 Temperature
 body. *See* Body temperature
 skin, 247t, 514t
 Temporal bone, 268f
 Temporal lobe, 740f, 740t
 Temporal pulse, 492f, 510t
 Temporalis, 274f
 Temporomandibular joint, 271t, 291t, 718t, 726t
 Tendon, 686, 690t
 Tennis elbow, 726t
 Terry's nails, 249t
 Tertiary preventive healthcare, 17
 Test tubes, 63t
 Testis (testes), 650, 651f, 652t
 cancer of, 679
 palpation of, 673t
 self-examination of, 676b
 torsion of, 673t, 683t
 Testosterone, 650
 Tetracyclines
 dental effects of, 279t, 298t
 integumentary system effects of, 216t
 nutrition and, 172t
 Thalamus, 741t
 Theme identification, in patient interview, 10
 Thenar prominence atrophy, 727t
 Thermometer, 56t
 Thermoregulation, in newborns, 851t
 Thiazides, nutrition and, 172t
 Thomas test, 729, 729f
 Thoracic duct, 492, 494f
 Thoracic spine
 range of motion of, 719t–720t
 vertebrae of, 687f
 Thought processes, in neurologic system
 assessment, 767t, 771
 Thrills, 474t–475t. *See also* Murmurs
 Throat. *See also* HEENT system
 assessment of, 37t, 296
 inspection in, 296, 303t–304t
 sore, 276
 Thrombophlebitis, 523t
 Thrombosis
 arterial, 522t
 deep vein, 514t–515t, 522t

- Thymus, 494, 495b
 Thyroid cartilage, 270f, 395t
 Thyroid gland, 270f, 272t
 auscultation of, 311, 312t
 disorders of, 312b
 palpation of, 308, 310t–311t
 in pregnancy, 818t
 Thyroid hormones, female genitourinary
 system effects of, 622t
 Thyroxine deficiency, 874t
 Tibia, 687f
 Tibialis pulse, 492f, 493t
 Tic douloureux, 775t
 Tidal volume, 81
 Time, orientation to, 769
 Tinea capitis, 244t, 253t, 262t
 Tinea corporis, 263t
 Tinea cruris, 263t, 670t
 Tinea manum, 263t
 Tinea pedis, 263t
 Tinea versicolor, 228t
 Tinel's test, 716, 717f
 Tingling, in neurologic system assessment of,
 753
 Tinnitus, 371
 Toddlers, 887–899
 abuse of, 889, 889b
 activity in, 892
 assessment of, 889–897
 biographical data in, 890
 cardiovascular system in, 891t, 895t–896t
 case study of, 889b, 897t
 current health status in, 890
 family history in, 890
 gastrointestinal system in, 891t, 896t
 general survey in, 890t, 893, 893f, 893t
 genitourinary system in, 891t, 896t
 health history in, 889–890
 HEENT system in, 890t–891t, 894t–895t
 hematologic system in, 892t
 integumentary system in, 890t, 894t
 lymphatic system in, 892t
 musculoskeletal system in, 891t, 896t
 neurologic system in, 892t, 897t
 past health history in, 890
 physical, 893, 893t–897t
 psychosocial profile in, 892–893
 respiratory system in, 891t, 895t
 review of systems in, 890, 890t–892t
 communicable diseases in, 889
 developmental changes in, 888
 emotional changes in, 888
 exercise assessment in, 159–160
 fears in, 889
 health concerns in, 897t
 injury in, 161, 889
 motor vehicle injury in, 889
 negativism in, 888
 neglect of, 889
 nutrition in, 168t, 892
 obesity in, 897t
 physical changes in, 888
 physiological anorexia in, 892
 poisoning in, 889
 psychosocial changes in, 888
 sexuality in, 893
 sibling rivalry in, 888, 893
 sleep in, 159, 892
 spirituality in, 189
 stress in, 160, 889, 893
 temper tantrums in, 888
 toilet training for, 888
 visual acuity in, 329
 wellness assessment in, 155
 Toe tapping, 710t
 Toe walking, 709t
 Toilet training, 888
 Tolbutamide, nutrition and, 172t
 Tongue, 269f, 271t
 black, hairy, 303t
 cancer of, 303t
 geographic, 302t
 inspection of, 302t–303t
 palpation of, 296, 306t
 red, beefy, 303t
 Tongue depressor, 63t
 Tonic neck reflex, 868t–869t
 Tonsillitis, 318t
 Tonsils, 269f, 272t, 495b
 development of, 274–275
 inspection of, 304t, 305t
 Tooth. *See* Teeth
 Torticollis, 307t
 Torus mandible, 301t
 Torus palatinus, 301t, 318t
 Touch
 communicative aspects of, 8–9
 light, 777t–778t
 Trachea, 270f, 394f, 395t
 position of, 420t
 shifts of, 41413t
 Tragus, 366f
 Tranquilizers, male sexual function effects of,
 660t
 Transillumination
 fontanel, 860t
 frontal sinus, 292t
 maxillary sinus, 292t
 paranasal sinuses, 291, 292t
 scrotum, 673t
 Transilluminator, 61t
 Trapezius, 274f
 Trapezius muscle, 286, 286f, 398f
 Trapezius squeeze, 769
 Trapezoids, 687f
 Treacher-Collins syndrome, hearing in, 372t
 Tremors, 766b
 Trendelenburg test, 518b, 729
 Trial-and-error thinking, 13
 Triamterene, nutrition and, 172t
 Triceps brachii, 687f
 Triceps skin fatfold measurement, 180, 180f
 Triceps skinfold calipers, 62t
 Tricuspid regurgitation, 472f
 Tricuspid valve, 438f, 439t
 Tricyclic antidepressants, side effects of, 123t
 Trigeminal nerve (CN V), 746t, 747f,
 773t–774t
 Triple screen, 833b
 Trochlear nerve (CN IV), 329f, 746t, 747f,
 772t–773t
 Trunk incurvation reflex, 873t
 Tuberculosis, 403b, 434t
 Tubes, abdominal, 599, 601t–602t
 Tunica adventitia, 441b, 441f
 Tunica intima, 441b, 441f
 Tunica media, 441b, 441f
 Tuning fork, 60t
 Tuning-fork tests, 382–383, 384t–385t
 Turbinates, 271t
 inspection of, 295t
 Turgor, of skin, 247, 248t, 249
 Twenty-four-hour recall, in nutrition
 assessment, 175–176
 Two-point discrimination, 781t
 Tympanic cavity, 365t
 Tympanic membrane, 365t, 366f, 376f
 assessment of, 380t–381t
 Tympanotomy, 381t
 Tympany, on percussion, 70t
U
 Ulcer(s)
 aphthous, 300t
 chronic, 213
 corneal, 351t
 gastric, 608t
 pressure, 230, 238b, 239t
 stasis, 211, 238t, 509t
 Ulceration, 209
 Ulcerative colitis, 579t, 607t
 Ulna, 687f
 Ulnar deviation, 716t
 Ulnar pulse, 492f, 493t, 511t
 Umbilical cord, 864t
 Umbilicus, inspection of, 583t–584t
 Uremia, 229t
 Uremic frost, 209
 Ureter, 558f, 615t, 652t
 Urethra
 female, 612f, 615t
 male, 650, 651, 651f, 652t
 Urethral meatus
 female, 631t, 633t
 male, 652t, 671t, 672t–673t
 Urethral specimen, in male genitourinary
 system assessment, 676b
 Urinary diversion, 602t
 Urinary incontinence, 938t–939t, 941t
 Urinary system, 614. *See also* Genitourinary
 system
 disorders of, 619
 Urinary tract infection, 614, 646t
 Urination, in newborns, 851t
 Urine tests, in pregnancy assessment,
 824–825, 833b
 Urticaria, 242t, 243t
 Uterus, 559t, 612f, 613f, 615t
 bimanual examination of, 641t
 Uvula, 269f, 272t
 inspection of, 304t
V
 Vagina, 612f, 613f, 614t, 615t, 631t, 633t
 speculum examination of, 636t–637t, 637
 Vaginal bleeding, 619
 Vaginal discharge, 619, 632t, 633t
 Vaginal pool smear, 637, 638t
 Vaginal wall
 bimanual examination of, 640t, 641t
 inspection of, 637, 639t

- Vaginitis
 monilial, 645t
 in pregnancy, 826
- Vaginosis, bacterial, 644t
- Vagus nerve (CN X), 746t, 747f, 775t–776t
- Validation, data, 20
- Valvular heart disease, auscultation for, 480, 480t, 480f, 480t–485t, 485b. *See also* Heart sound(s); Murmurs
- Varicocele, 670t, 683t
- Varicose veins, 509t, 523t
- Vas deferens, 650, 652t
- Vastus medialis, 687f
- Vegetarian diet, 164b
- Veins, 441b, 441f
 pressure in, 450–451, 451f
 varicose, 509t, 523t
- Velcro rales, 426t, 427
- Vellus, 206, 254t
- Vena cava, inferior, 493f, 493t, 559t
- Vena caval syndrome, 816t
- Venous hum, 449, 451f, 478, 587t
- Venous lakes, 956t
- Venous star, 232t
- Ventricle (cardiac)
 left, 438f, 439t
 right, 438f, 439t
- Ventricles (brain), 741t
- Ventricular diastole, 442
- Ventricular systole, 442–443
- Venule, 441b, 441f
- Verbal cues, 10
- Vernix caseosa, 850t, 859t
- Vertebrae, 687f
- Vertical strip method, of breast palpation, 543, 543f
- Vertigo, 371, 753
- Vesicle, 234t
- Vestibular nerve, 366f
- Vestibule, 614t
- Vestibulospinal tract, 743–744, 744t
- Vibration, sensory examination for, 779t
- Vietnamese Americans
 adolescent, 915t
 childbearing in, 823t
 mental disorders and, 119t
 musculoskeletal health problems in, 694t
 newborn, 853t
 nutrition and, 171t
 older adult, 932t
 pain response in, 97t
 school-age, 904t
 spirituality and, 192t
 views of women and, 618t
- Violence, by adolescent, 913–914
- Viral rhinitis, 316t
- Visceral pain, 563–564
- Vision, 322, 326, 326f. *See also* Visual acuity
 accommodation in, 328, 353t
 adaptation in, 328
 blurred, 332
 color, 341, 341f, 343t
 distance, 342t–343t
 double, 331
 loss of, 331
 near, 343t
 in neurologic system assessment, 753
 peripheral, 341–342, 342f
 screening of, 157t
- Visual acuity, assessment of, 341–342, 342t–343t
 in African Americans, 360
 Allen test in, 329
 in children, 360
 in diabetes, 360
 in infants, 329, 360
 in older adults, 330
 in preschool children, 330
 in school-age children, 330, 360
 in toddlers, 329
- Visual acuity charts, 58t
- Visual Analogue Scale, for pain, 99, 100
- Visual fields, 326f, 344t
- Vital signs, 78–83
 blood pressure. *See* Blood pressure
 in cardiovascular system assessment, 465–466
 documentation of, 83
 in neurologic system assessment, 762
 postpartal, 843t
 pulse, 80–81, 80f, 80t, 81f
 respirations, 80t, 81
 temperature. *See* Body temperature
- Vitamin C, during pregnancy, 169t
- Vitamins, 165
- Vitiligo, 229t, 233t, 263t
- Vitreous chamber, 323f
- Vitreous humor, 325t
- Vocabulary test, in neurologic system assessment, 767t, 771
- Vocal cues, 7–8
- Voice sounds, abnormal, 429–430
- Vomiting, 566
- W**
- Waardenburg's syndrome, hearing in, 372t
- Waist-to-hip ratio, 182
- Walking, 709t. *See also* Gait
- Warts
 genital, 632t, 669t, 680t
 plantar, 728t
- Watch tick test, 382, 383t, 775t
- Water
 dietary, 165
 loss of, 165
- Water cup, 64t
- Water-hammer (Corrigan's) pulse, 477f
- Weakness, 695
 in older adults, 693
- Weber test, 384t
- Weight, 83–84, 84f
 in cardiovascular system assessment, 466
 in gastrointestinal system assessment, 565, 565b
 in infants, 879t
 measurement of, 180, 180f
 in newborns, 857t–858t
 during pregnancy, 827t
 sleep and, 158
- Wellness, 153–162
 assessment of, 154–162, 155b
 in adolescents, 157
 in children, 157
 developmental stage and, 155, 157–158
 exercise in, 159–160
 in infants, 155
 injury prevention in, 161–162
 in middle-aged adults, 158
 in older adults, 158
 rest and sleep evaluation in, 158–159, 159b
 stress management in, 160–161
 in young adults, 158
 behavioral factors in, 154–155
 case study of, 154
 plan of action for, 155b
 screening for, 156t–157t
- Wernicke's aphasia, 766b
- Wernicke's area, 740f
- Wet prep, 637, 638t
- Wharton's duct, 272t
 assessment of, 301t–302t
 inspection of, 296
- Wheal, 234t
- Wheezes, 426t, 428t, 429t–430t
- Whisper test, 382, 383t
- Whispered pectoriloquy, 426t, 430
- Williams' syndrome, hearing in, 372t
- Wooden tongue depressor, 63t
- Word-Graphic Rating Scale, for pain, 100
- Wrist
 palpation of, 721t–722t
 Phalen's test for, 716, 717f
 range of motion of, 722t
 Tinel's test, 716, 717f
- X**
- Xanthelasma, 348t
- Xanthoma, 209
- Y**
- Yale-Brown Obsessive Compulsive Scale, 115t
- Yeast infection, 645t
- Yellow marrow, 686
- Young adults
 exercise assessment in, 160
 injury prevention assessment in, 162
 sleep assessment in, 159
 spirituality in, 189–190
 stress assessment in, 161
 wellness assessment in, 158
- Z**
- Zinc, during pregnancy, 169t
- Zonules, function of, 325t
- Zygomatic bone, 268f
- Zygomaticus, 274f
- Zygote, 613